CHAOS AND MICROBIAL SYSTEMS

DE93 011727

Progress Report

July 1989 - July 1990

Mark Kot

*Dept. of Applied Mathematics, FS-20**University of Washington**Seattle, Washington 98195*

July 1990

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-89ER60880

(University of Washington)

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED *ce*

Table of Contents

1.	Brief Project Summary	2
2.	Report of Technical Progress	3
3.	Proposed Research for the Coming Year	5
4.	Appendices	7
	Appendix A: Bistability and Chaos in a Model Microbial System	Reprint removed
	Appendix B: Multidimensional Trees, Range Searching, and a Correlation Dimension Algorithm of Reduced Complexity	Reprint removed
	Appendix C: Box-counting with Base b^n Numerals	Reprint removed
	Appendix D: Experimental Protocols	

§1. Brief Project Summary

The last ten years have seen a host of new results in the field of nonlinear dynamics, most notably regarding the phenomenon called *chaos*. Many recent results have come from mathematicians and from theoretical and experimental physicists; yet much of the early work was motivated by issues in population dynamics. A recurrent theme of this recent research is that *seemingly* random fluctuations often occur as the result of simple deterministic mechanisms. Hence, much of the recent work in nonlinear dynamics has centered on new techniques for identifying order in seemingly chaotic systems. These techniques when applied in ecology have led to new insights for several systems, most notably childhood disease epidemics.

To determine the robustness of these techniques, chaos must, to some extent, be brought into the laboratory. Preliminary investigations of the forced double-Monod equations, a model for a predator and a prey in a chemostat with periodic variation in inflowing substrate concentration, suggest that simple microbial systems may provide the perfect framework for determining the efficacy and relevance of the new nonlinear dynamics in dealing with complex population dynamics. In recognition of this fact, I have proposed research with two main components:

- (1) mathematical analysis and computer simulation of the periodically forced double-Monod equations and of related models; and
- (2) experimental (chemostat) population studies that evaluate the accuracy and generality of the models, and that judge the usefulness of various new techniques of nonlinear dynamics to the study of populations.

My colleagues and I have made a great deal of progress in analyzing the models. We also have experiments underway that will provide us with accurate estimates of the relevant parameters. We hope to continue this theoretical and experimental research and to begin work on the second of the above stated goals.

§2. Report of Technical Progress

A fairly complete picture of the forced double-Monod model is beginning to emerge. The model describes a protozoan predator and a bacterial prey in a chemostat with inflowing substrate that is periodically forced. The behavior of the model depends critically on the degree of forcing. In the absence of forcing, the system exhibits relatively benign behavior: depending on the dilution rate of the chemostat and the inflowing substrate concentration, one observes (a) washout of predator and prey, (b) washout of the predator, (c) coexistence of predator and prey at an overdamped equilibrium, (d) coexistence at an underdamped equilibrium, and (e) coexistence on a limit cycle. In particular, there is no chaos in the absence of forcing. This simplicity disappears as one forces the inflowing substrate.

Complex dynamics arise most readily when forcing is added to a limit cycle. Small-amplitude or high-frequency forcing typically leads to quasiperiodic flow on a torus. As we increase the period or amplitude of forcing, it is not unusual to see periodic solutions reemerging via phase locking (alias frequency locking, mode locking, or resonance). Additional increases in the period or amplitude of forcing may lead to a sequence of period-doubling bifurcations and, in some cases, to chaotic solutions. This particular route to chaos

$$\textit{quasiperiodicity} \rightarrow \textit{phase locking} \rightarrow \textit{period doubling} \rightarrow \textit{chaos}$$

is rather common for the forced double-Monod system.

We have taken stroboscopic sections of the tori that are associated with the quasiperiodic regime, and of the strange attractors that appear in the chaotic regime. These stroboscopic sections are formally equivalent to Poincaré sections and from them we have constructed Poincaré maps: monotone circle maps for the quasiperiodic regime, noninvertible unimodal maps of the interval for the chaotic regime. Each of the two maps betokens a different range of dynamics. In addition, circle maps and unimodal maps of the interval have been studied in their own right; the behavior and bifurcations found in these theoretical studies are an augur of the dynamics to be found in the forced double-Monod system. Detailed bifurcation

diagrams for the forced double-Monod system show that the behavior of this system may be particularly well understood in terms of the circle maps that emerge as Poincaré maps. Phase-locking regions for the model correspond to Arnol'd tongues for the maps. Criticality in the circle maps sets the stage for chaos in the model. This criticality may arise with an increase in the period or amplitude of forcing.

A detailed exposition of these and other results may be found in Appendix A. This appendix is the body of a manuscript that has been submitted, for publication, to the *Journal of Theoretical Biology*.

In addition to a better understanding of the forced double-Monod system, there have been several technical or theoretical spinoffs that have arisen from our examination of this system. The more prominent of these have been fast new methods for computing fractal dimensions (of, for example, the strange attractors of the forced double-Monod system). A student of mine (Stuart Bingham) and I have developed an improved method for computing correlation dimension. This method has been described in a publication in *Physics Letters* (Appendix B). In addition, Stuart has developed and submitted for publication (Appendix C) a new method for computing the box-counting dimension or capacity of a fractal set. This work was completed while he was supported by the first year of this DOE grant.

There has been steady progress on the experimental front. The experimental work is being done in collaboration with Drs. Gary Sayler, a microbial ecologist, and Terry Schultz, a protozoologist, at the University of Tennessee. A chemostat has been set up with pumps to periodically force inflowing substrate; a fraction collector has been set up to collect regular samples. These are all dedicated pieces of equipment. In addition, a flow cytometer has been made available, as needed, to count cell densities in the samples. There are experiments underway to measure the appropriate physiological and ecological parameters of the predator, *Tetrahymena pyriformis*, and of the prey, *Escherichia coli*.

§3. Proposed Research for the Coming Year

Most of our analyses of the forced double-Monod system have been for a set of generalized ciliate/bacterium parameters. The experimental portion of the project is about to provide us with accurate estimates of the maximum specific growth rates of *Tetrahymena pyriformis* and *Escherichia coli*, their yield coefficients, and also their half-saturation constants. With these parameters in hand, we will perform a more detailed and complete analysis of the forced double-Monod system, using an assortment of techniques (e.g. bifurcation diagrams, Poincaré maps, perturbation methods).

At the same time, there have been suggestions that the forced double-Monod model is imperfect. Several scientists have argued that the predator's functional response may depend on predator density as well as prey density: predators' functional responses saturate at sufficiently high prey-to-predator ratios. Others — the detailed references are contained in Appendix A — have pointed out that suspension feeding ciliates often act as "prudent predators" in that they cease to feed at low prey densities. This would intimate some sort of threshold or possibly a type III functional response. Detailed analyses will be performed of the effects that these modifications might have, in the presence of forcing, on population dynamics.

As the experimental phase of the project progresses, we will also have an opportunity to test predictions of the various models and to test the efficacy of various new techniques of nonlinear dynamics. A detailed exposition of the experimental protocols is included as Appendix D. What follows is a sample of some of the questions that we will examine and of some the technique that we will test in answering these questions:

- (1) Preliminary analyses of the forced double-Monod model suggest that small-amplitude and/or high-frequency forcing has little effect on the underlying limit cycle. In general, the limit cycle gives way to quasiperiodic behavior close to the original cycle.

Time series will be collected. Power spectrum will be computed to determine the power associated with the underlying limit cycle and with the forcing. Phase portraits will be reconstructed using Takens' method to see whether it is possible to

reconstruct a torus close the original cycle. A Poincare map will be taken of the reconstructed orbit in and an attempt will be made to construct the associated circle map.

- (2) For high-frequency forcing, there should be a tendency for the various species to do a progressively better job of averaging over exogenous forcing as we move up the food chain.

This will be tested with time series for *E. coli*, *T. pyriformis* and, if possible, glucose.

- (3) As the amplitude and/or period of forcing increases, the system should show greater tendency to phase lock and to undergo period-doubling bifurcations.

This will be tested by comparing stroboscopic samples of the system.

- (4) With further increases in the amplitude and/or period of forcing, the system should exhibit chaotic dynamics.

Time series will be examined for erratic aperiodicity. The power series of the time series will be examined to assess the power at the forcing frequency and to look for signs of broadening or flattening of the spectrum. Phase portraits will again be reconstructed using Takens' method to see if there is any sign of a low dimensional strange attractor. The fractal dimension of the phase portrait will be computed using the correlation dimension algorithm described in Appendix B. Maps will be computed if possible.

The empirical results will be compared and contrasted with the theoretical predictions of the model.

APPENDIX D

Experimental Protocols

The goal of the experimental program is to study the behavior of a holozoic predator and a heterotrophic prey (*Tetrahymena pyriformis* and *Escherichia coli*) in a chemostat in which the inflowing substrate, glucose, is periodically forced, and to determine whether this experimental system can exhibit the chaotic dynamics found in the forced double-Monod system. To attain this goal we need to perform a number of tasks and experiments:

1. Obtain and maintain cultures of *Tetrahymena pyriformis* and *Escherichia coli*.

A strain of *E. coli* (ATCC 25922) was obtained from Denise King (Dept. of Microbiology, UT). Stock cultures of *E. coli* are grown and stored at 9.9° C on slants containing:

Tryptone	5 grams/liter
Glucose	1 gram/liter
Yeast extract	2.5 grams/liter
Agar	20 grams/liter

An asexual strain of *Tetrahymena pyriformis* was provided courtesy of Dr. Terry Schultz (Assoc. Prof., Dept. of Animal Science - Vet Med., UT) . Stock cultures of *Tetrahymena pyriformis* are grown and stored on slants in a long term medium as per the following protocol:

- (1) 150 ml distilled H₂O
1.5 grams liver extract
boil, filter
- (2) to 100 ml of supernate add
0.8 grams dextrin
0.06 grams NaC₂H₃O₂•7H₂O
0.5 grams yeast extract
0.5 grams casein enzymatic hydrolysate
cool to room temperature
pH 7.1 - 7.4

- (3) add 1.6 grams agar
 boil and dissolve
 pour 5 ml/tube with cap
 autoclave, slant tubes before cooling, place tape around cap to avoid
 moisture loss
- (4) Add 3 ml sterile distilled H₂O and 1 drop of *Tetrahymena* from previous
 long term culture.

Store (one tube in incubator, one at room temperature).

Subculture every 2 weeks.

A medium for short term culture (and experimentation with) *Tetrahymena* consists of:

1000 ml of distilled H₂O
20 g protease-peptone
5g dextrose (d-glucose)
1 g yeast extract
1 ml of 3% Fe EDTA solution
pH 7.35

For experiment:

50 ml in 250 ml flask
Top with foam plug
Autoclave and allow to cool before use

For short term culture:

In capped test tubes add 30 ml of medium
Autoclave and cool before use
Short term cultures maintained by subculturing on Monday and Friday.
(Add 1/3 ml from previous culture to each tube)

The short term medium may also be used as an axenic chemostat medium for
T. pyriformis.

2. Estimate growth kinetic constants for *Escherichia coli* from batch culture.

The goal here is to determine the maximum specific growth rate μ_1 , the half saturation constant K_1 , and the yield coefficient Y_1 for our strain of *E. coli*.

Washed cells of *E. coli* will be seeded into a glucose-mineral salts medium (50 ml) and incubated, with shaking (150 rpm) at room temperature. The seeding will be to an initial cell concentration of approximately 10^7 cell/ml. Specific growth rates will be determined by linear regression of corrected counts to the equation $\log H = \mu t + \log H_0$. The curve relating specific growth rate to nutrient concentration is expected to be hyperbolic. From a plot of the reciprocal of growth rate against the reciprocal of substrate concentration one can estimate the maximum growth rate and the half saturation constant. From estimates of the final crop of bacterial cells and the glucose concentration at final crop, we may estimate the yield coefficient..

Medium:

To prevent irreversible complexing, medium constituents are autoclaved in four separate portions (Watson et al., 1981) (all concentrations are final concentrations in the medium). The four portions consist of:

(a)	K_2HPO_4	0.335 g/l
	KH_2PO_4	1.77 g/l
(b)	$(NH_4)_2SO_4$	1.25 g/l
	$MgSO_4 \cdot 7H_2O$	0.1 g/l
	$Na_2EDTA \cdot 2H_2O$	0.0004 g/l
	$NaMoO_4 \cdot 2H_2O$	0.001 g/l
	$FeSO_4 \cdot 7H_2O$	0.001 g/l
(c)	$CaCl_2$	0.02 g/l
	NaCl	0.01 g/l
(d)	glucose	25, 50, 75, 100, 125, 150, 200, 500, etc. mg/liter

Distilled water is distributed among these portions and they are autoclaved separately at 120° C for 30 to 60 minutes, except for the glucose solution which is autoclaved for only 15 minutes. The pH should be about 7.5 after the cooled solutions are mixed. This medium is also used in the chemostat. It provides for the growth of *E. coli* but is insufficient, as such, for *Tetrahymena pyriformis*.

Cell Counts:

Viable cell counts will be determined by dilution and plating onto 0.25 x nutrient agar supplemented with 0.05% glucose and agar (10 g/l). These counts will be corroborated with direct cell counts of samples with a Coulter counter (30 µm orifice) or a Facscan flow cytometer. Cell suspensions will be prepared for counting by diluting samples with saline solution containing 8.5 g/l NaCl and 0.2 g/L Na₂(EDTA) which has been filtered through a 0.22 µm Millipore filter.

Determination of Glucose Concentration:

Glucose concentration will be determined colorimetrically with a glucose oxidase diagnostic kit purchased from Sigma Diagnostics (Procedure #510).

3. Examine the chemostat dynamics of the *E. coli* - glucose subsystem.

The growth constants measured in batch culture for *E. coli* when taken in concert with the Monod model for growth in a chemostat allow us to predict the behavior of *E. coli* in continuous culture. To ascertain the effectiveness of these predictions we will test two hypotheses:

- (a) *E. coli* should wash out at or near $D > \mu_1 S_i / (K_1 + S_i)$ where D is the dilution rate, S_i is the inflowing substrate concentration, and μ_1 and K_1 are as above. For a series of inflowing substrate concentrations, we will increment dilution rate, allowing the system to equilibrate after each increment, until washout occurs. The dilution rate at washout will be compared with the predicted value.
- (b) In the absence of washout, the system should settle down to a steady state with equilibrium concentrations (weight or biomass per unit volume) of glucose and bacteria of

$$S^* = K_1 D / (\mu_1 - D) \quad \text{and} \quad H^* = Y_1 (S_i - S^*).$$

These two relations may be tested as per hypothesis (a). Measuring a series of equilibrium concentrations for different dilution rates also allows us to estimate, independently, the yield coefficient and the functional response [$f(S^*) = D (S_i - S^*)/H^*$]. The latter may be used for an independent estimate of the maximum growth rate and the half saturation constant..

The procedures in sections 2 and 3 will be repeated, if necessary, until we have some confidence in our understanding of and our ability to predict the behavior of the bacterium-substrate subsystem.

Chemostat, Pumps, Microcomputer:

We are currently using a 350 ml bench top chemostat dedicated to this project. We have, already set up and ready to operate, two Gilson 302 precision piston pumps (with 10 ml/min pump heads) to be used with the chemostat, and an IBM XT to control the pumps. Mr. Rodney Bain has kindly provided software that will control the pumps so that the inflowing substrate is constant or periodic.

Wall growth:

The inner surface of the chemostat will be treated with a bacteriostatic silicone agent Protosil-28 (PCR Research Chemical, Inc.) so as to reduce the likelihood of wall growth. The Protosil-28 should be diluted 1:100 with distilled water and the dilute solution will be left in contact with all inner surfaces for at least 20s. Surfaces will be rinsed with distilled water and the chemostat will be air dried for 24 hr. The silicone coating is stable to autoclaving and washing (Ratnam et al., 1982).

Cell Counts:

Samples will be taken from the chemostat on a regular basis using a fraction collector. Bacteria will be fixed in formalin and cell counts will be obtained using a Coulter counter or a Facscan flow cytometer as described in section 2.

Medium:

The chemostat medium is the same as that prescribed in section 2.

4. Estimate growth kinetic constants for *Tetrahymena pyriformis* from batch culture.

The next step is to determine the maximum specific growth rate μ_2 , the half saturation constant K_2 , and the yield coefficient Y_2 for our strain of *T. pyriformis*. All experiments will be started by adding a fixed number of *T. pyriformis* to a suspension of *E. coli* cells in a chemically defined medium which contains no carbon or energy source for *E. coli* growth: Inocula for batch cultures will be collected from *E. coli* and *T. pyriformis* chemostats after they have reached steady state. Measured volumes of cultures from the two chemostats will be added to batch growth vessels and will be diluted with sterile growth medium with no glucose. Batch cultures will be propagated in glass vessels maintained at 26° C and aerated with humidified air. Specific growth rates will be determined by linear regression of corrected counts to the equation $\log P = \mu t + \log P_0$. The curve relating specific growth rate to prey

density is expected to be hyperbolic though we will also consider the possibility, suggested in the literature (Bazin et al., 1983; Curds and Cockburn, 1968) that the functional response is hyperbolic with prey to predator ratio rather than just prey number. From a plot of the reciprocal of growth rate against the reciprocal of prey concentration we will estimate the maximum growth rate and the half saturation constant. From estimates of the final crop of eukaryotic and prokaryotic cells, we will estimate the yield coefficient..

Medium:

The energy free medium is that outlined in section 2, minus the glucose. The medium, with glucose, is used for the *E. coli* chemostat. The *T. pyriformis* chemostat uses the experimental medium described in section 1.

Cell counts:

Bacteria are to be counted as before. *T. pyriformis* will be fixed in formalin and counted with a Coulter counter (100 μ m orifice) or a Facscan flow cytometer

Cell washing:

Watson et al. (1981) have demonstrated that ciliates introduced into a dense suspension of unwashed *E. coli* cells may suffer a temporary decrease in their density due to the presence of harmful metabolites. In batch experiments with high initial *E. coli* densities

($> 5 \times 10^8$ particles/ml), *E. coli* should be washed with 0.85% (w/v) saline solution on a membrane filter (0.45 μm pore size) under sterile conditions. The bacteria should next be resuspended in fresh batch culture medium and counted so that the volume of suspension needed for the inoculation can be calculated.

Washing has been found to have a marked and inconsistent effect on the physiological state of *T. pyriformis* and, in turn, on the behavior of *Tetrahymena* batch cultures; these organisms should not be washed.

Alternative (chemostat-free) protocol:

Flasks (100 ml.) of sterile culture medium and known glucose concentration should be inoculated with *E. coli* and shaken at room temperature for approximately two days. With the bacterial population just entering stationary phase, inoculate with a large number of axenically grown *T. pyriformis*. Shake at room temperature for 5 hr. Measure both bacterial and protozoan at beginning and at end of 5 hr feeding period. See Curds and Cockburn (1968) for details.

5. Examine the *T. pyriformis* - *E. coli* subsystem with a two-stage chemostat.

In section 3, I described how continuous culture of *E. coli* could be used to obtain an independent estimate of *E. coli*'s functional response and growth constants. If *T. pyriformis* is grown in continuous culture on nongrowing bacteria similar estimates may be obtained for *T. pyriformis*. As the growth of *T. pyriformis* shows some dependence on oxygen levels and vessel geometry, parameters obtained in this manner may be more useful than batch estimates in predicting the behavior of the complete *T. pyriformis* - *E. coli* - glucose system.

The experimental system consists of two chemostats linked in series. In the first vessel, the bacterial prey is allowed to come to steady state at a low dilution rate so that essentially all of its rate-limiting (energy-yielding) substrate is consumed. The overflow from this chemostat is then fed into the chemostat that contains the protozoan. The functional response f of *T. pyriformis* may be obtained from the specific rate of predation at equilibrium $f = D (H_1^* - H_2^*) / P^*$, where H_1^* and H_2^* are the equilibrium population densities of the bacterial prey in the first second stages of the continuous culture system, P^* is the equilibrium predator density in the second stage, and D is the dilution rate (cf. Bazin et al., 1983). If the predator's functional response is of the Monod type, the plot of f against H_2^* will generate a rectangular hyperbola. If the functional response is of the Contois type, the hyperbolic relationship appears when f is plotted against the ratio (H_2^* / P^*) . In either case, maximum growth rate, half-saturation constant, and yield coefficient may be estimated from a knowledge of H_2^* , P^* , D , and H_1^* values.

The medium is that described in section 2. The equipment and counting techniques have already been described.

6. Examine the complete *T. pyriformis* - *E. coli* - glucose system in single-stage chemostat with constant inflowing substrate.

The single stage chemostat represents a food chain which converts glucose to *E. coli* and *E. coli* into *T. pyriformis*. With the information garnered from the experiments of sections 1-5 (*E. coli* and *T. pyriformis* maximum growth rates, half-saturation constants, yield coefficients, and form of functional response) we should be able to predict the dynamics of this food chain in terms of the two experimental controls, D , the dilution rate, and S_i the inflowing substrate concentration. In particular, we will construct a two-parameter operating diagram for this system and verify that we can detect the following 5 types of behavior close to their predicted parameter values:

- (a) washout of both predator and prey
- (b) washout of predator only
- (c) coexistence of predator, prey, and substrate at a stable equilibrium with exponential return to equilibrium upon perturbation.
- (d) coexistence of predator, prey, and substrate at a stable equilibrium with damped oscillations upon perturbation.
- (e) limit cycles

The tests will consist of a series of experiments in which inflowing substrate concentration is held fixed (at concentrations between 25 and 250 mg/l) but dilution rate is varied (between 0 and 0.25 per hour). After each change in dilution rate, the system will be allowed to time to pass through its transient; observed dynamics will then be compared with the predicted dynamics. Predator - prey phase portraits will be compared with those constructed using Takens' method. If limit cycles are observed, their period will be compared with that predicted by the model.

7. Examine the complete *T. pyriformis* - *E. coli* - glucose system in single-stage chemostat with periodically forced inflowing substrate.

If a simple system can exhibit limit cycles, what are the effects of external periodic forcing on this limit cycle ? Clearly, the work in this section is very much contingent on the experiments in section 6 being able to turn up periodic dynamics. To the extent that the double-Monod model is accurate, this should be possible. If a limit cycle cannot be found what factors mitigate against its existence ?

Once a small limit cycle is found, we will add periodic forcing to the inflowing substrate. A energy rich and an energy-poor medium will be prepared and the two Gilson 302 precision piston pumps will force the media periodically and out phase with one another. The use of two media guarantees that the inflowing volume remains constant even as the energy content varies periodically. The forcing will be at a frequency above that of underlying limit cycle. The amplitude of forcing is determine readily from the glucose concentrations of the two media.

References

- Bazin, M. J., Curds, C., Dauppe, A., Owen, B.A., Saunders, P.T. 1983. Microbial predation dynamics. *ACS Symp. Series*. **207**: 253-264.
- Curds, C.R. and A. Cockburn. 1968. Studies on the growth and feeding of *Tetrahymena pyriformis* in axenic and monoxenic culture. **54**: 343-358.
- Fredrickson, A.G. 1983. Interactions of microbial populations in mixed culture situations. *ACS Symp. Series*. **207**: 201-227.
- Jost, J.L., Drake, J.F., Fredrickson, A.G. , and Tsuchiya, H.M. 1973. Interactions of *Tetrahymena pyriformis*, *Escherichia coli*, *Azotobacter vinelandii*, and glucose in a minimal medium. *J. Bacteriol.* **113**: 834-840.
- Ratnam, D.A., Pavlou, S., and Fredrickson, A.G. 1982. Effects of attachment of bacteria to chemostat walls in a microbial predator-prey relationship. *Biotech. Bioeng.* **24**: 2675-2694.
- Swift, S.T., Najita, I.Y., Ohtaguchi, K. and Fredrickson, A.G. 1982. Some physiological aspects of the autoecology of the suspension-feeding Protozoan *Tetrahymena pyriformis*. *Microb. Ecol.* **8**: 201-215
- Swift, S.T., Najita, I.Y., Ohtaguchi, K. and Fredrickson, A.G. 1982. Continuous culture of the ciliate *Tetrahymena pyriformis* on *Escherichia coli*. *Biotech. Bioeng.* **24**: 1954-1964.
- Taylor, W.D., Gates, M.A., and Berger, J. 1976. Morphological changes during the growth cycle of axenic and monoxenic *Tetrahymena pyriformis*. *Can. J. Zool.* **54**: 2011-2018.
- Watson, P.J., Ohtaguchi, K., and Fredrickson, A.G. 1981. Kinetics of growth of ciliate *Tetrahymena pyriformis* on *Escherichia coli*. *J. Gen. Microbiol.* **122**, 323-333.

END

DATE
FILMED
5/13/93