PROGRESS REPORT

MARYLAND CONTROLLED FUSION RESEARCH PROGRAM

Submitted to the
Department of Energy
Washington, D.C.

Period Covered
November 1, 1991 - October 31, 1992

Principal Investigators
H. R. Griem and C. S. Liu

Laboratory for Plasma Research
University of Maryland
College Park, MD 20742-3511

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

RECEIVED
JUN 30 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

TASK I: Plasma Theory Principal Investigators: T. M. Antonsen, Jr.,* J. F. Drake,** J. M. Finn, P. Guzdar, A. Hassam,* C. S. Liu,* E. Ott.* Additional Physicists: R. Kleva, F. Waelbroeck)

I. TASK DESCRIPTION

Purpose: The Maryland Theory Program is dedicated to the following tasks:

- (i) To intensify our effort to develop an understanding of the fundamental processes which produce anomalous transport in tokamaks and to propose methods for enhancing confinement.
- (ii) To continue our investigation of tokamak edge plasma turbulence, sheared flow generation and the L/H transition.
- (iii) To continue our investigation of major disruptions with an emphasis on developing techniques for stabilization.
- (iv) To continue collaborations with experimentalists and theorists of major fusion laboratories and university fusion centers for close interplay of theory, experiment, and simulation.
- (v) To provide the theoretical support to the existing and proposed Maryland experimental programs in plasma diagnostics such as laser particle tagging and cyclotron radiation measurements from tokamaks.
- (vi) To continue the education and training of scientists in plasma theory relevant to fusion.

*Professor

**Professor and contact person for this contract.

II. TECHNICAL PROGRESS

In recent years, members of the Maryland Plasma Theory Group have made significant contributions to the national fusion theory program, and, in many cases, these theoretical developments helped to interpret experimental results and to design new experimental programs.

In the following, we summarize the technical progress in four major areas of tokamak research: (a) L/H transition and edge turbulence and transport; (b) active control of microturbulence and transport; (c) major disruptions; and (d) the sawtooth crash.

A. L/H Transition in Tokamaks

The transition of the tokamak to a regime of improved confinement (L/H transition) has led to recognition of the importance of the edge plasma in controlling global confinement. Recent experimental measurements at UCLA and on DIII-D at G. A. have shown that poloidal rotation very likely plays an important role in the onset of the H-mode. Measurements on TEXT indicate that sheared rotation has a stabilizing influence on edge turbulence and locally reduces transport. Measurements at UCLA also point out the importance of poloidal asymmetry in understanding the L/H transition.

During the past year, we have developed a conceptual model of the tokamak edge in both L- and H-mode. In this model we identify four important ingredients: poloidally asymmetric transport driven by the drift-resistive ballooning mode; parallel flows of the order of the sound velocity (c_s) produced by the asymmetric transport; poloidal rotation (Stringer mechanism) produced by these parallel flows; and finally order one ($\tilde{n}/n \sim 1$) turbulence produced by the crossfield gradient of the parallel flows ($V_{||}'$ instability). The nonlinear interaction between these four components determines whether the plasma remains in L-mode or makes a transition to the H-mode. When magnetic pumping is large, the poloidal rotation remains of the order of the diamagnetic level. Poloidally asymmetric transport produces a significant variation of the density along flux surfaces in the edge and results in parallel flows of order c_s . Experimentalists on CCT, TEXT and DIII-D are presently attempting to confirm the presence of these parallel flows in L mode edge plasmas. When the magnetic pumping is weak, the Stringer mechanism (parallel flow) causes the edge plasma to spin, amplifying the existing diamagnetic level shear layer which is present in the L-mode. Poloidal rotation increases to $V_\theta \sim c_s a / qR$ when poloidal rotation competes with the parallel flow in smoothing the edge poloidal asymmetries. In the H phase the poloidal asymmetries and consequently parallel flows are greatly reduced. Transport is suppressed by sheared rotation, allowing steepened edge density gradients.

In the following sections we describe the more detailed calculations we have completed and are presently pursuing to solidify the model presented above. This

work was described as an invited talk at the recent TTF workshop at Oak Ridge and will be presented at the 1992 IAEA meeting.

1. 3D Fluid Simulation of the Nonlinear Drift Resistive Ballooning Modes in Tokamak Edge Plasmas.

The single most important experimental observation for identifying the cause of fluctuations and transport in tokamak edge plasmas is the strong inside-outside asymmetry of the anomalous particle flux (with a larger flux on the outside). This indicates that an instability with an outward ballooning character would be the most likely candidate for driving the observed particle flux. Since most edge plasmas are collisional, the only model which has the necessary requisites is the drift resistive ballooning mode (DRBM). Thus over the last year we have explored the linear stability of these modes¹ and developed a 3D code to study their nonlinear evolution.²

Our calculations have shown that:

1. In the nonlinear phase, these modes have a strong ballooning character. As a consequence, the turbulent transport caused by these modes has a strong poloidal asymmetry.
2. The magnitude of the anomalous transport D^{an} associated with these modes is $2 \times 10^4 \text{ cm}^2/\text{sec}$ which is consistent with experimental observation.
3. These modes can give rise to poloidal flows which are driven both by the Reynold's stress^{3,4,5} as well as the anomalous Stringer mechanism.⁶
4. The radial and poloidal correlation lengths were found to be in reasonable agreement with observations on TEXT.⁷

Thus the 3D simulations of the DRBMs have indicated that these modes have the potential for explaining edge fluctuations and transport in tokamaks.

2. Loss of Static Equilibrium, Flow Generation and Edge Turbulence at the Edge of Tokamaks

We have recently shown from the experimental data on at least four different tokamaks (CCT, TEXT, DIII-D, PBX) that the hydrodynamic time scale for establishment of isobaric equilibria along a flux surface is not short but is in fact comparable to the local cross-field particle diffusion time at the edge.⁸ This observation implies that our standard notions of MHD equilibria need reexamining. One consequence is that if the transport is poloidally asymmetric, large, sonic parallel flows are a necessary and integral part of the equilibrium. Such parallel flows have indeed been observed in the T-10 and DITE tokamaks.

Our 3D simulations of DRBM have succeeded in explaining many aspects of experimental observations of edge turbulence as discussed earlier. However, one of the important features that is not explained is the large levels of density fluctuations observed in the edge. Experiments on TEXT indicate that $\delta n/n_0 \sim 40\%$. However, from our 3D studies $\delta n/n \sim 10\%$. One important secondary instability which can enhance the fluctuation levels to of order unity is the $V_{||}'$ instability. The question that we have studied is how the parallel velocity shear flow is driven. Both experiments and our 3D simulations have shown that the poloidal asymmetric anomalous transport D^{an} is such that $D^{an}/L_n^2 \sim (c_s/\pi q R)$, i.e., the anomalous particle confinement time in the edge is comparable to the sound transit time. As a consequence, the asymmetric particle flux can drive parallel flows which are large and are of the order of c_s (the acoustic velocity). Since this occurs only in the edge region the above-mentioned parallel flow is localized to the edge and hence is capable of driving the $V_{||}'$ instability. We find that for most tokamaks edge parallel flows of order c_s should be observed and preliminary 3D simulations of the $V_{||}'$ instability in a magnetic shearless plasma show that $\delta n/n \sim 0(1)$ can be readily driven.⁸

3. Generation of Sheared Flow and Reduction of Convective Transport in a Two-Dimensional Plasma with Unsheared Magnetic Field

As discussed in Sec. 1, a plausible model for the L-mode, the H-mode, and the L-H transition involves resistive ballooning modes in the edge region. The 3D numerical results described in Sec. 1 indicate that 3D effects are, of course, necessary for the destabilization of the ballooning modes, which lead to the convective vortices. The variation of the field line curvature along the magnetic field lines, which is responsible for the ballooning, determines the unstable range of mode numbers. However, it was found that the nonlinear dynamics, particularly the development of rotation shear and suppression of convective transport by reduction of the ballooning vortices, is essentially two dimensional. That is, the variation in the third direction along the magnetic field, the flow in that direction, and the magnetic shear do not play a decisive role in the dynamics. This result motivates the development of a simple 2D code to study the dynamics of sheared flow generation.

We have written an incompressible 2D slab code with unsheared uniform magnetic field B_z in the third direction (reduced MHD). The boundary conditions were free slip conditions on the velocity at $x = 0, b$ and periodic boundary conditions on y with period "a". In this geometry, x corresponds to the minor radius in the tokamak and y to the poloidal angle. The 2D nature of this code has allowed us to scan the parameter space quite completely. (A single run takes 5-20 minutes on a Cray, as opposed to 100 hours for the 3D code of Sec. 1.) We first considered an initial equilibrium with convective vortices and uniform density similar to those obtained in resistive ballooning. It was found⁴ that there can be a linear instability in which a sheared 1D velocity $v_y = v_y(x)$ can develop. This mode is unstable for viscosity $\mu = 0$ (ideal) if the aspect ratio of the initial vortices satisfies $b/a \gtrsim 1$. A viscous mode, with growth rate scaling as $\mu^{3/4}$, exists for $b/a \lesssim 1$. For long enough

vortices, $b/a \lesssim 0.5$ the mode is stable. Nonlinearly, the mode develops by a vortex reconnection or “peeling” process. The generation of sheared flow is part of this process. In the later stages the shear stretches the y -dependent structures from the initial vortices, leading to a late stage in which the flow is pure x -dependent shear flow $v_y(x)$, which decays slowly due to viscosity.

To prevent the final decay, we have also performed studies with a source driving the initial vortex flow.⁵ We found that the final state consists of stable vortices (of opposite vorticity, i.e., counter-rotating) if the viscosity is large enough that the linear mode is stabilized. For smaller viscosity (higher Reynolds number $R_e \sim 1/\mu$) the time asymptotic state consists of tilted counter-rotating vortices. Past a higher value $R_e > R_c$, there is a bifurcation to a state with a chain of islands (co-rotating vortices) with sheared flow outside the island separatrix. The island width decreases as $R_e \rightarrow \infty$.

These results are in qualitative agreement with those of Sec. 1, although no such detailed scaling with respect to the vortex aspect ratios or the Reynolds number R_e is possible with the 3D code. We have also begun a 2D study of the stability of such convective vortices, but in which the vortices are driven by a temperature difference and the source of vorticity is buoyancy (the Rayleigh-Taylor instability, or Bénard convection).⁹ Preliminary results indicate that the Rayleigh-Taylor instability generates vortices similar to those in the previous study^{4,5} and that a shear flow can develop. Phase mixing in this shear flow completely stabilizes the Rayleigh-Taylor vortices, leading to a pure shear flow. However, when this occurs, there is no source to drive the vortex flow and therefore nothing to drive the shear flow. The latter then decays, until it reaches a sufficiently small value that the Rayleigh-Taylor vortices are destabilized again, leading again to the generation of shear and stabilization of the vortices. For small Rayleigh number $R_a \sim 1/\mu$ this leads to a periodic relaxation oscillation reminiscent of ELMs. During the ELM burst, the transport can be 5-10 times classical (based on the actual transport used in the code, which may still be anomalous) whereas between ELMs the transport is slightly above classical.

The observation of these ELM-like oscillations is made more intriguing by the observation that the edge plasma in the H-mode is at or near the ideal ballooning limit.¹⁰ Therefore, an intriguing scenario is that resistive ballooning modes develop vortices, leading to the enhanced transport observed in the L-mode. If a shear layer develops, a reduction in the transport, as in the H-mode, should occur. Then the pressure gradient can build up until the ideal ballooning threshold is reached. The ideal ballooning modes which occur should periodically generate a stabilizing sheared flow, leading to relaxation oscillations as observed in the 2D simulations.

B. Active Control of Microturbulence and Transport

1. Active Confinement Control — Stabilization of Microturbulence by

Externally Driven Perpendicular Rotation

There is strong experimental evidence that perpendicular velocity shear may suppress microturbulence. In the past year we have been in the forefront of research identifying the concepts and means by which a plasma may be externally torqued to set up sufficient velocity shear to improve confinement.¹¹⁻¹³ Our focus has been broad, examining ways to suppress turbulence in a large fraction of the tokamak, not just the edge. We have achieved results on two key questions.

(1) How much velocity shear is needed? We have completed the linear theory of four microinstabilities in the presence of $v'_\perp$ to obtain the stability criterion. These modes are the resistive-g mode,¹¹ the “ $i - \delta$ ” drift wave¹¹ (or trapped particle mode),¹³ the η_i mode,^{11,13} and the $v'_\parallel$ mode.¹³ In all four cases, surprisingly but desirably, the critical $v'_\parallel$ criterion is the same, namely, $v'_\perp \gtrsim c_s/L_s$ or c_s/R . This result provides a benchmark against which all schemes for externally induced shear can be measured. While the result is linear, it is consistent with the scaling arguments of Biglari, Diamond, and Terry.¹⁴ We have also confirmed the general criterion by numerical simulation for the 3D resistive ballooning mode² as well as the 2D Rayleigh-Taylor instability.¹⁵

(2) Are such shears realistically attainable? We have suggested that NPI could be used to drive velocity shears. We have examined both poloidal injection³ and toroidal injection.^{12,13} In both cases, while the early calculations show that achievement of the above criterion will not be an easy matter, they also show that the proposed schemes are in the realm of feasibility. For poloidal injection, it may be possible to achieve a relatively narrow shear layer ($\lesssim 20\%$ of minor radius) with existing NBI injected poloidally. In toroidal injection, if the toroidal rotation speed can be made to exceed the sound speed, stabilization can be achieved.

The calculations clearly point to more work being needed in the areas of non-linear suppression of microinstability from $v'_\perp$ and the details of poloidal viscous (magnetic pumping) damping. Work is in progress in these areas.

2. Spontaneous Poloidal Spin-Up from Asymmetric Particle Fueling

If a source of plasma is turned on in a tokamak, with a source rate larger on the outboard side of a flux surface than on the inboard side, a spontaneous poloidal spin-up of the plasma can occur. We have recently shown¹⁶ this from a calculation for an arbitrarily shaped, axisymmetric tokamak plasma where the particle (and momentum) sources may be nonaxisymmetric. Our calculation thus applies to particle sources arising from pellets, gas puffing, or centrally neutral beam fueled systems like TFTR supershots. The mechanism for the spin-up is the Stringer mechanism.⁶ The spin-up is destabilized provided the poloidally asymmetric portion of the particle fueling rate exceeds the damping rate of poloidal rotation from magnetic pumping reduced by a factor ϵ , the inverse aspect ratio.

On TFTR, H-modes have been triggered from pellet injection. Preliminary data analysis shows that the spin-up criterion described above may be satisfied. More interesting are the data from TFTR supershots. Here, the transient time of rise in particle density is of order 250 ms while typical magnetic pumping damping times are greater than 100 ms in the interior. If the beam deposition can be made sufficiently poloidally asymmetric, it may be possible to achieve an externally induced spin-up in the plasma interior.

C. Major and Minor Density Limit Disruptions in TFTR

Stable operation of tokamak discharges is limited to line-averaged densities $\bar{n}$ below a critical value $\bar{n}_c \sim B_\theta/q_a R$, where q_a is the safety factor at the plasma edge. When the density limit is exceeded, the temperature profile begins to contract and there follows a series of minor disruptions in which central confinement is temporarily degraded but the discharge recovers, followed by a final major disruption in which confinement is abruptly lost and the discharge terminates. The rapid drop in the plasma current following the energy quench generates a large force on the vacuum vessel which can cause substantial damage. Since ITER will have to operate near the density limit to optimize its performance, it is important to understand the mechanisms of both the major and the minor disruptions.

We have been able to successfully simulate the loss of confinement during disruptions by including a model for line radiation in our resistive MHD code.¹⁷ In our simulations, the initial loss of energy confinement is caused by the injection of a cold plasma bubble from the edge into the center by a $q = 1$ kink mode. Subsequently, a cold plasma bubble has been seen to move from the edge into the center at the onset of disruptions on the TFTR tokamak. As a consequence, we have begun a collaboration with Eric Frederickson and the TFTR experimental group with the goal of understanding disruptions on TFTR.¹⁸ This collaboration has led to a new understanding of the difference between major and minor disruptions, and why a discharge can recover from a minor, but not a major disruption. As the temperature profile contracts, the $q = 1$ instability undergoes transition from an internal to an external kink as the resistivity just outside the $q = 1$ rational surface increases. Early in the contraction phase, the temperature just outside the $q = 1$ surface is relatively high. The time over which the bubble is formed and injected by the $q = 1$ kink is less than the resistive diffusion time across the bubble; the kink is internal. As a consequence, q remains equal to unity in the bubble after injection and although there is a large drop in the central temperature due to the injection of the bubble, there is no drop in the current and the internal inductance is unchanged. In this case the disruption is minor and the discharge recovers. In contrast, when the profile contracts to the extent that the cold radiating edge plasma lies just outside the $q = 1$ rational surface, the resistive diffusion time across the bubble is comparable to or less than the bubble formation time. In this case the instability is essentially an external kink and the bubble carries little current; there is a large drop in the internal inductance suggesting a redistribution of plasma current.

Because of the change in the current profile, the $m/n = 2/1$ mode grows to large amplitude terminating the discharge in a major disruption.

Therefore, the temperature of the plasma just outside the $q = 1$ rational surface determines whether the contracting current and temperature profiles are unstable to an internal kink, which causes a minor disruption from which the discharge can recover, or an external kink, which redistributes the current and causes a major disruption that terminates the discharge. By heating the edge plasma to a temperature of a couple of hundred electron volts, one may be able to prevent the transition from internal to external kink that causes the major disruption. Then, the discharge would just slowly radiate its energy away, rather than losing it abruptly to the vacuum vessel.

D. The Fast Sawtooth Crash Driven by Ion Gyroradius Effects

Experimental measurements of the sawtooth crash time τ_c in the TFTR and JET tokamaks range from 20 to 2000 μsec . The crash time, based on a Sweet-Parker model of resistive reconnection by the $m = 1$ tearing mode is of the order of 2000 μsec , which is short enough to explain the slow crashes, but not the fast crashes.

Previously,¹⁹ we showed that the electric field E induced during magnetic reconnection during the sawtooth crash greatly exceeds the Dreicer field E_D , and, therefore, classical resistivity cannot play a role in this process. Including electron inertia in Ohm's law, we found that during collisionless reconnection the current layer collapses to a thickness which is much smaller than the ion gyroradius ρ_s .

Recently, we have found that the extreme narrowness of this current layer alters the parallel force acting on the electrons.²⁰ The polarization drift of the ions across this very narrow current layer formed during reconnection causes a perturbation in the ion density. The electrons move rapidly along the magnetic field lines to charge neutralize this ion density perturbation, leading to a large parallel gradient in the electron pressure. We have examined the effect of this parallel pressure force on the reconnection process. We have carried out numerical simulations of magnetic field line reconnection using an Ohm's law which includes the effect of the parallel electron pressure gradient. When the width of the resonant current layer is less than ρ_s , we find that the parallel pressure force induces a further contraction in the width of the current layer. Furthermore, not only does the current layer contract in the direction of the plasma inflow to the X-point, but the current layer also shrinks in the direction of the outflowing plasma. At the same time, the plasma flow in the dissipation region detaches from the current layer, the width of the outflowing plasma layer increasing to ρ_s . In the Sweet-Parker model, the length L of the dissipation region around the X-point in the direction of the plasma outflow is given by the equilibrium scale length r_1 . However, our simulations show that actually L is a function of ρ_s and that L is smaller than R_1 . Thus, the reconnection of magnetic field lines does not follow a Sweet-Parker model.

Based upon our numerical simulations, we have derived an expression for the sawtooth crash time given by $\tau_{crash} = (L/\rho_s)\tau_A/\Delta q$, where $\Delta q = 1 - q(0)$ is the amount by which $q(0)$ falls below unity and the Alfvén time $\tau_A = R/c_A$. For a tokamak discharge with $n \sim 5 \times 10^{-3} \text{cm}^{-3}$, $B_\phi \sim 40 \text{kG}$, $r_1/a \sim 0.2$, and $\rho_s/a \sim 5 \times 10^{-3}$, $\tau_{crash} \sim (8/\Delta q)(L/r_1)_{\mu\text{sec}}$. If $\Delta q = 0.2$, then $\tau_{crash} \sim 40(L/r_1)_{\mu\text{sec}}$ with $L < r_1$ for non-Sweet-Parker reconnection. This short crash time generated by the parallel gradient in the electron pressure is consistent with the experimental observations of fast crashes on TFTR and JET.

REFERENCES

1. D. McCarthy, P. Guzdar, J. F. Drake, T. M. Antonsen and A. B. Hassam, UMLPR #92-007 (to appear in Phys. Fluids B).
2. P. N. Guzdar, J. F. Drake, A. B. Hassam, D. McCarthy and C. S. Liu, Proceedings of the First Energy Research Power Super Computer Users Symposium, May 21-22, 1991, Gaithersburg, MD (published by US DOE) article 15 and UMLPF #92-023; P. Guzdar, J. F. Drake, A. B. Hassam and C. S. Liu, submitted to Phys. Fluids B (1992).
3. P. H. Diamond and Y. B. Kim, Phys. Fluids B **3**, 1626 (1991).
4. J. Drake, J. Finn, P. Guzdar, V. Shapiro, V. Shevchenko, F. Waelbroeck, A. Hassam, C. S. Liu, and R. Sagdeev, "Peeling of Convective Cells and the Generation of Sheared Flow," Phys. Fluids B Lett. **4**, 488 (1992).
5. J. Finn, J. Drake, and P. Guzdar, "Instability of Vortex Flow and Generation of Sheared Flow," submitted to Phys. Fluids B (1992).
6. A. B. Hassam, T. M. Antonsen, Jr., J. F. Drake and C. S. Liu, Phys. Rev. Lett. **66**, 309 (1991).
7. A. Wootton et al., Phys. Fluids. B **2**, 2879 (1990).
8. J. F. Drake, A. B. Hassam, P. N. Guzdar, C. S. Liu, D. McCarthy, UMLPF #91-054 (submitted to Nucl. Fusion).
9. J. M. Finn et al., International Sherwood Theory Meeting 1992 (Sante Fe), paper 1D11.
10. P. Gohil, M. A. Mahdavi, L. Lao et al., Phys. Rev. Lett. **61**, 1603 (1988).
11. A. B. Hassam, Comments on Plasma Phys. and Controlled Fusion **14**, 275 (1991).
12. A. B. Hassam and F. L. Waelbroeck, Proceedings of Workshop on New Ideas in Tokamak Confinement, La Jolla, CA (1992).
13. F. L. Waelbroeck, T. M. Antonsen, P. N. Guzdar, and A. B. Hassam (to be published), Phys. Fluids (1992).
14. H. Biglari, P. H. Diamond, and P. W. Terry, Phys. Fluids B **2**, 1 (1990).
15. A. B. Hassam, Phys. Fluids B **4**, 485 (1992).
16. A. B. Hassam and T. M. Antonsen (in preparation).
17. R. G. Kleva and J. F. Drake, Phys. Fluids B **3**, 372 (1991).

18. E. D. Frederickson, K. McGuire, J. F. Drake, R. G. Kleva, and the TFTR MHD Group, Phys. Rev. Lett. (submitted 1992).
19. J. F. Drake and R. G. Kleva, Phys. Rev. Lett. **66**, 1458 (1991).
20. R. G. Kleva, J. F. Drake and F. Waelbroeck, International Sherwood Theory Meeting, 1992 (Santa Fe), paper 2C16.

Reprints + Preprints removed

TASK II: Measurements of Electron Parallel Momentum Distributions Using Cyclotron Wave Transmission Diagnostics
(Principal Investigators: F. Skiff and D. A. Boyd)

A series of proof-of-principal experiments were conducted through a collaboration with the Versator II group at the Massachusetts Institute of Technology. The objective of these experiments was two-fold. First of all, the objective was to explore the sensitivity and resolution of the cyclotron wave transmission technique. In parallel, we also improved the theoretical understanding of the momentum resolution of the diagnostic. The second, closely related, objective was to explore the feasibility of using selective perturbations in connection with the transmission measurements as a form of test-particle diagnostic which we call Vlasov Tagging. The experiments were largely performed in two weeks of machine time over a 20-month period in 1991. Despite the very limited resources, the experiments were a very successfull in achieving both of the objectives. Results have been presented both in conference proceedings and in papers which are currently being submitted for publication.

We rebuilt the Versator cyclotron wave transmission diagnostic with improvements such as low noise amplifiers, matched detector diodes, and RF shielding for the detection circuits. In this way we were able to improve the signal-to-noise of the diagnostic by an order of magnitude. The addition of extensive shielding permitted measurements during the application of lower-hybrid wave power to the plasma to a degree not previously possible. These improvements also led to the recognition and verification that the transmission data could also be used to determine the line-averaged group-velocity of the cyclotron waves. This fact makes the transmission diagnostic less dependent on other diagnostics for calibration. In order to understand the resolution, a theoretical analysis of the diffraction integral which relates the transmitted to the received signal was performed. This analysis demonstrated that the resolution of the diagnostic is actually better on larger machines where the distance between the transmitter and receiver horns is greater. Previous estimates of the resolution were shown to be based on a wrong understanding, though the previous results provided an estimate of the resolution which is close to the actual value for the Versator experiment.

In the experiments, measurements of cyclotron wave transmission were made for two different kinds of plasma discharge. Measurements were made in plasmas where the plasma current was sustained by the continuous application of lower-hybrid waves and in plasmas where there were no lower-hybrid waves or only short pulses to produce selective perturbations. A variety of techniques were tried for the purpose of comparison. Measurements were made with constant cyclotron wave frequency, with frequency dithering, and with broad sweeping of the frequency. As a result of these measurements, conclusions can be drawn not only about the cyclotron wave diagnostic but also about the nature of the lower-hybrid wave interactions observed. These results have been, and are being, distributed through the following

publications:

Conference Proceedings

1. Vlasov Tagging Proof-of-Principle Experiments on Versator II, F. Skiff, D. A. Boyd, and J. A. Colborn, 18th European Physical Society Conference on Controlled Fusion and Plasma Physics, Berlin, June 1991.
2. Feasibility Studies of Vlasov Tagging Using Short Pulses of Lower Hybrid Waves on Versator II, F. Skiff, D. A. Boyd, and J. A. Colborn, Bull. Am. Phys. Soc. **36**, 2241 (1991).
3. A New Kind of Ammeter for the Tokamak de Varennes, D. A. Boyd and F. Skiff, Bull. Am. Phys. Soc. **36**, 2441 (1991).
4. Measurements of Electron Parallel Momentum Distribution Using Cyclotron Wave Transmission, F. Skiff, D. A. Boyd, and J. A. Colborn, 9th Topical Conference on High-Temperature Plasma Diagnostics, Santa Fe, March 1992.

Articles

1. Measurements of Electron Parallel Momentum Distribution Using Cyclotron Wave Transmission, F. Skiff, D. A. Boyd, and J. A. Colborn, in preparation for submission to Nucl. Fusion.
2. Observation and Analysis of a Two-Parallel-Temperature Electron Tail during Lower-Hybrid Current Drive on a Tokamak, J. A. Colborn, F. Skiff, D. A. Boyd, M. Forkolab, J. P. Squire, and J. Villasenor, in preparation for submission to Phys. Rev. Lett.

TASK III: Experimental Study of Electron Transport in Tokamak Plasmas (01-5-27287) (Principal Investigator: R. F. Ellis)

During this time period (Nov 1, 1990 – June 30, 1991) the grant funds were used to support Scott Janz, the University of Maryland graduate student who was working on his dissertation. Mr. Janz made substantial progress during this time, including the following:

- a. Nonthermal emission data from low density ECH discharges was analyzed using a code which he developed to predict X and O mode emission for specific distribution functions. In the process a reflection model was developed. It became clear that a more sophisticated analysis of nonthermal emission was needed.
- b. The horizontal view system was maintained and temperature profiles could be produced regularly. Much work remained in making this process automatic from the point of view of the end user of the temperature profiles.

TASK IV: ECE Diagnostics on DIII-D (FAS 01-5-27295)
(Principal Investigator: R. F. Ellis)

During this time period (May 1, 1991 – October 31, 1991) these supplemental funds were employed to allow a modest University of Maryland contribution to DIII-D, in the area of ECE diagnostics. Contributions were made in two areas:

a. Scott Janz, a University of Maryland graduate student, completed all the substantive work on his Ph.D. thesis, which studied nonthermal electrons produced during ECH heating and current time. A new analysis of combined X and 0 mode ECE led to a reasonable determination of the behavior of these high energy ($E > 100$ keV) electrons including their localization in momentum space and their confinement time.

b. Improvements to the horizontal system were made, including better calibration and lineup and a bit better data analysis. At the end of this period it appeared likely that General Atomic would pick up support of this work in some form.

DATE
FILMED
8/03/92