013502

10F1

[Bifurcations and Patterns in nonlinear dissipative systems] DOE/ER/13738--T2
PROGRESS REPORT FOR DOE

GRANT NUMBER DE-FG03-87ER13738

February 26, 1992

Guenter Ahlers and David S. Cannell

DOE/ER/13738--T2

Department of Physics and Center for Nonlinear Studies

DE92 013502

University of California at Santa Barbara

1. Personnel

At present, three graduate students, four post-doctoral research associates, and the principal investigators are contributing to the research sponsored by this Grant.

The students are Mr. Steven Trainoff who receives salary support from the Grant, Ms. Kristine Lerman who has a University of California Fellowship and requires salary support only during the summer months, and Mr. Mike Dennin who has a DOD Fellowship and requires no support from this grant. Of the post-doctoral researchers, Dr. Marco Dominguez-Lerma receives salary support from this Grant. Dr. Stephen Morris and Dr. Lars Berge have Fellowships from their home countries Canada and Norway respectively. Dr. Eberhard Bodenschatz had a Fellowship until recently from the Deutsche Forschungsgemeinschaft; but that has expired. He is presently paid from University of California sources. Our ability to attract people with fellowship or sabbatical support is largely attributable to the scientific attractiveness of our research group, but is in part also due to our ability to offer a small salary supplement to Fellowships from sources other than this Grant through the Center for Nonlinear Studies on our Campus. Thus, the support provided by the DOE is highly leveraged.

MAILED

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

2. Progress during the last Year and Plans for the next Grant Period.

a. Rayleigh-Bénard convection with an imposed horizontal flow

Mr. Trainoff has continued his study of Rayleigh-Bénard convection in a long, narrow, horizontal cell with the imposed flow parallel to the long side. In his system, rolls with their axes parallel to the short sidewall (transverse rolls) bifurcate from the conduction state for throughflow Reynolds numbers R less than 0.5. For larger R , longitudinal rolls first form. Mr. Trainoff is exploring experimentally the convectively unstable region for this system as a function of R . Sufficiently close to the convective instability this region normally contains no structures because any disturbances are swept away faster than they can grow. He has developed methods for generating pulses of convection rolls which grow as they travel downstream in the system. An interesting aspect of this system is that somewhat further above the convective, but well below the absolute, instability there exist spontaneously occurring noise-sustained structures. This is so because in our system even a very small perturbation has the opportunity to grow to a macroscopic level since the system is very long and thus a large time elapses before the perturbation is swept out of the system. We are now in the process of determining quantitatively the noise level necessary to explain this phenomenon so that a comparison with the influence of thermal noise on this system can be made. In a related but different system studied recently in our laboratory, namely Taylor-vortex flow with axial flow, thermal noise (*i.e.* of intensity $k_B T$) was adequate to account for the experimental results. A preliminary report on our progress will be presented at the March Meeting of the APS in Indianapolis.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

b. Rayleigh-Bénard convection in a gas under Non-Boussinesq conditions

Dr. Bodenschatz and Dr. Morris have continued their study of convection in a gas (CO_2) under pressure (about 20 bar). They each have an apparatus with a pressure stability of 5ppm, and a temperature stability of better than $10^{-3}K$. During the last year they have improved the quality of their convection cells substantially. By making the sample cell extremely thin (0.3 mm), they achieve a very large aspect ratio (radius/height of about 150) without encountering excessive lateral diffusion times which would make it impossible to achieve a stationary state in the system. However, a thin sample requires exceptional uniformity of the spacing between top and bottom plates so as to keep the relative variation of the gap small. Our plates are optically flat (to better than $\lambda/4$). The bottom plate is servoed by three piezo-electric posts until it is parallel to the top plate within about $0.1 \mu m$.

Dr. Morris will continue our study of strong departures from the Boussinesq approximation. Under those conditions, the bifurcation from conduction to convection becomes hysteretic, and the pattern which forms consists of hexagons rather than rolls. Our sample has a large aspect ratio (radius/height), and we find that a perfect lattice consisting of more than 10^4 hexagonal cells forms above the bifurcation. Dr. Morris expects to study various aspects of the hexagons, including for instance the dynamics of isolated defects in an otherwise perfect lattice. We have also discovered that the non-Boussinesq system tends to form stable spirals. Spiral patterns are known to occur in chemical and biological pattern-forming systems, and we will explore the extent to which there may be a communality between these diverse pattern-formation phenomena. Our system has the great advantage that the spirals are perfect, stable indefinitely, and contain many turns.

Dr. Bodenschatz expects to devote his effort to the study of fluctuations below the

bifurcation from conduction to convection in a gas under pressure. These fluctuations are *extremely* weak, and so far have never been observed in Rayleigh-Bénard convection (recently, they were studied in electro-convection in a nematic liquid crystal; see below). The fluctuations are expected to be relatively stronger in a thinner sample. Therefore Dr. Bodenschatz will take advantage of the extremely thin sample ($d = 0.25mm$) which he can utilize in his system, and will work at a higher pressure (45 bar) to make the bifurcation accessible in his sample. Preliminary measurements have revealed the existence of fluctuations below the bifurcation, but their quantitative study remains for the next grant period.

c. Electro-convection in a Nematic liquid crystal

Stimulated by interesting results which we obtained about fluctuations below the first bifurcation in electro-convection and in collaboration with Rehberg and associates in Bayreuth (Germany), we have built an electro-convection apparatus in Santa Barbara. This project is being conducted by Mr. Mike Dennin. We expect to investigate the fluctuation amplitudes and the nature of the bifurcation to Williams rolls as a function of cell thickness and drive-voltage wave-form. We are learning how to prepare and align the director of our samples, and are exploring different nematics which may be more suitable for this work than those used in the past (MBBA).

d. Rayleigh-Bénard convection in a Nematic liquid crystal

Stimulated by theoretical work in the Institute of Professor Lorenz Kramer at the University of Bayreuth in Germany, we started a new program of experimental study of Rayleigh-Bénard convection in a horizontal layer of a nematic liquid crystal heated from below and

in a horizontal magnetic field. The work at Bayreuth has suggested that there should be a richness of bifurcation phenomena in this system, including forward and backward bifurcations, tricritical bifurcations, and Lifshitz points. We were able to acquire at no cost a 12-inch Varian magnet to provide the field. With the expertise of the Quantum Institute at UCSB, Dr. Lars Berge was able to custom design pole pieces for this magnet which will give us a field homogeneity at 1 kG of 0.1 % over a volume of $5 \times 5 \times 5 \text{ cm}^3$. Dr. Berge has built a special non-magnetic convection apparatus which will fit into the magnet gap. He has measured the bifurcation line as a function of the magnetic field. His results agree reasonably, but not perfectly, with the prediction. He has also studied in detail one of the Lifshitz points in this system. Measurements of the angle of the oblique rolls as a function of the magnetic field agree well with general expectations about Lifshitz points. Ours are the first quantitative experimental results for this type of bifurcation point that we know of. Although consistent with the expected general behavior, our experiments differ substantially from the detailed calculations for this particular system. We expect to explore in collaboration with the theorists in Bayreuth to what extent any differences between experiment and theory can be accounted for by uncertainties in the fluid properties, and to what extent they may be indicative of basic flaws in the equations of motion which have been used in the theory.

END

**DATE
FILMED**

6/02/92