AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

Progress Report
THE THEORY OF RELATIVISTIC
HEAVY ION COLLISIONS

DE-FG02-93ER40768

July 1993

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

I. Introduction

**Program and Principal Achievements during 1993
of the Stony Brook- Brookhaven Collaboration on
'Theory of Relativistic Heavy-Ion Collisions'**

The program discussed here was started only few months ago on January 1, 1993. It was initiated by the Nuclear Theory Groups at Brookhaven and Stony Brook, and its primary goals are studies of highly excited matter and its production in nuclear collisions at very high energies. Although both groups are engaged in a wide variety of projects in different areas, it was considered important to pool efforts and resources through the establishment of a permanent existing collaboration in order to help with the analysis of on-going experiments at the AGS and the planning of future experiments at RHIC.

Abstracts describing the contents of completed papers and providing some details of on-going projects are given below. Here, we shall provide a more general orientation on how the project has started and what different participants are doing. Before doing that, it is useful to note how the very existence of this collaboration has already somewhat changed the life of both Groups.

First, some administrative changes have taken place at Stony Brook. The Center for Nuclear Theory has been established as an Organized Research Unit at Stony Brook.¹ Shuryak will serve as Director of the Center. G.E. Brown, A.D. Jackson and J. Verbaarschot are faculty members partially assigned to this Center. C.B. Dover has been appointed Adjunct Professor at Stony Brook. He now supervises one of our graduate students and plans to offer a special graduate course next year.

Second, we have established 'Stony Brook-Brookhaven Discussion Days' once a month. These meetings, held alternately at Stony Brook and BNL, include a review talk (or talks) and long informal discussions. These meetings involve the participation of 10-15 people and take half a working day. This arrangement allows discussion of some major topic in considerable detail. Our initial sense is that these meetings are quite helpful, especially for junior members of the Groups and students who are actively involved in the

¹As an 'Organized Research Unit', the Center is administratively separate from the Department of Physics and reports directly to the Dean of Mathematical and Physical Sciences.

discussions and get a much better understanding of what other people are doing.

Personnel associated with this project include Professor Edward Shuryak as Principle Investigator and Dr. Carl B. Dover and Professor Andrew D. Jackson co-Principle Investigators. The program supports two postdoctoral fellows — T. Schlagel and E. Schnedermann. Both postdocs are physically at Brookhaven, and Schnedermann has partial support from German funds. Schlagel works in collaboration with S. Kahana and Y. Pang on the development of the ARC code. Schnedermann is mainly involved in work with Dover and A. Baltz on the evaluation of production rates of rare species using the coalescence model.

At the moment, the graduate students supported by this project include David Brown (who works with C.Dover), Wen Chen Chang, Chi Ming Hung and Momchil Velkovsky (all working with E.Shuryak).

There are a number of people at Brookhaven and Stony Brook who, although not formally involved in this program, are working on related subjects and in close collaboration with this project. These include Stony Brook faculty G.E. Brown, I. Zahed and J. Verbaarschot, Stony Brook Research Faculty M. Prakash and V. Koch, BNL physicists A. Baltz, S. Kahana and S. Gavin, and postdoctoral students Y.Pang (BNL), Li Xiong (Stony Brook) and T. Schaefer (Stony Brook).

Before turning to a review of the main achievements of the last year, it is useful to indicate the way in which we have selected topics. We have included recent work of people directly involved in (*i.e.*, supported by) this program. Related work performed by others at BNL and Stony Brook has not been included unless performed in direct collaboration with participants. Description of this related work can be found in the Progress Reports of the Stony Brook and Brookhaven Nuclear Theory Groups. There is, nevertheless, a substantial degree of overlap between the topics presented here and in these other Reports. We believe that this 'double counting' is useful in the interests of providing a balanced view of our attack on this problems. We note in this regard that the level of funding of the current proposal is less than 10% of the total funding of Nuclear Theory at BNL and Stony Brook, and it goes mainly to support of young people, postdocs and students.

For clarity of presentation, we have divided the topics of interest into the following main topics:

- The dynamics of nuclear collisions at very high energies (RHIC and LHC).
- The dynamics of nuclear collisions at AGS energies.
- High-temperature QCD and the physics of the quark-gluon plasma.
- The production of strangelets and other rare objects.
- Other activities.

The Dynamics of Nuclear Collisions at Very High Energies

In preparation for large-scale experiments to be running at RHIC and LHC, the most obvious questions are the temperatures and densities which are likely to be reached. We have made significant progress on these issues during the past year. The principal difference from lower energies (AGS and SPS) is that dynamics are no longer dominated by 'soft' hadronic physics. Rather, one enters the realm of 'semi-hard' hard processes which involves partons of 'a few' GeV. Thus, one is entitled to turn to 'partonic cascades' based on perturbative QCD rather than complicated and model-dependent event generators, based on hadronic cross sections. These studies have led to the so called 'hot glue scenario'² of high energy heavy ion collisions. According to this scenario, the gluon equilibration time is quite short (*i.e.*, 0.3–0.5 fm/c) and leads to initial temperatures (of about 400–600 MeV at RHIC) which are twice the temperatures assumed previously. Probably the best test for this scenario is enhanced charm production. Both the STAR collaboration at RHIC (using advanced silicon vertex detectors and the excellent tracking capability of the TPC) and PHENIX (using $e-\mu$ coincidences) are investigating the experimental possibilities to observe it. Stony Brook and Brookhaven theorists are actively involved in discussions with the experimental groups.

We have also demonstrated³ that other possible signatures of a quark-gluon plasma, such as the production of photons and dileptons, are also significantly modified in the 'hot glue scenario'. Briefly, the smaller number

²E. Shuryak, Two-stage equilibration in high energy nuclear collisions, *Phys.Rev.Lett.* **68** (1992) 3270.

³E. Shuryak and Li Xiong, Dilepton and photon production in 'hot glue scenario', *Phys.Rev.Lett.*, **70** (1993).

of charged partons (quarks and anti-quarks) in this scenario is actually more than compensated by the much higher temperatures involved provided that one exploits appropriate 'windows of opportunity' in the various kinematic variables. Continuing projects (with Li Xiong and A.Makhlin) include detailed studies of this scenario and gluon chemical equilibration as well as dilepton, photon and charm production.

We note that this topic has become one of the most active areas of RHIC-related studies. At least two other strong groups (K. Geiger and J. Kapusta in Minnesota and B. Müller with collaborators at Duke) are now working in this direction. The consensus appears to be that this scenario and its main consequences are 'inevitable' if perturbative QCD can be used at all.

The evolution of matter during the first few fm/c considered above is well enough described by simple scale-invariant expansion. It is not so if one think about longer-term matter evolution. After a long and controversial history of applications to high-energy hadron-hadron and hadron-nucleus collisions, relativistic hydrodynamics may finally become a convincing concept when applied to ultrarelativistic nuclear collisions with large projectiles and targets. Schnedermann and Heinz⁴ have proposed a new method, which we call 'global hydrodynamics', for solving the hydrodynamic equations. Our aims are, first, to make calculations easier by using an approximate but efficient scheme and, second, to gain new insight into the dynamics by establishing the global expansion characteristics as a new viewpoint. Integrating over the reaction zone by assuming certain forms of profiles, we arrive at a much simpler problem to be solved. This enables us to do large numbers of computer runs easily and, thus, to perform a systematic scan of the whole range of possible initial conditions following each case individually until freeze-out. We can then take this hydrodynamical simulation as a framework for other computations including hadronic spectra, hadronic abundances, interferometry, and dileptons.

The Dynamics of Nuclear Collisions at AGS Energies

The development of the new event generator, ARC, for the AGS energy range was the primary goal of the group led by S. Kahana. The motivation for this new generator and its principal distinction from other available

⁴E. Schnedermann and U. Heinz, Phys.Rev. **C47** (1993) 1738,

codes (*e.g.*, RQMD One developed by the Frankfurt group) is simple: To construct an ‘honest’ cascade code free from model-dependent concepts (such as strings) and to rely on experimental data for the production of hadrons and resonances in the ‘few’ GeV region of interest. The code is essentially finished, and its major tests have been performed. These include the ‘stopping’ of nucleons, pion production and, especially, the production of positive and negative kaons. The comparisons performed so far show that the overall performance of ARC is very satisfactory. If so, it means that most of the available data can be explained by the ordinary rescattering of hadrons and hadronic resonances.

However, further studies may show more exciting results. One interesting point is related to antiproton production in heavy ion collisions. It seems that antiproton absorption in dense nuclear matter differs from the predictions of simple binary collision processes. As in the case of strangeness, it has been suggested that anomalously high levels of antinucleons are possible for heavy ion collisions in which quark-gluon plasma is formed. Pertinent data exist at the AGS, and previous theoretical analysis has had considerable difficulty in obtaining sufficient numbers of antiprotons. The problem is clear. AGS energies are only slightly above the threshold for $\bar{p}$ production and the annihilation cross section is large for the resultant low energy $\bar{p}$ ’s. Using ARC, a three-body screening mechanism was identified which strongly reduces the absorption of anti-nucleons. This mechanism heralds the appearance of a problem for all rescattering codes at high density. Interactions for a process with cross section σ are triggered by closest approach distances less than $r_0 = (\sigma/\pi)^{1/2}$. If r_0 is greater than the average distance between particles, it will be impossible to ignore third bodies. The large $\bar{p}p$ annihilation cross section at low relative energy precipitates this crisis for $\bar{p}$ production at even modest nuclear densities. A straightforward ‘fix’ is to introduce an annihilation time delay deduced from the $\bar{N}N$ relative velocity and distance. During the delay other particles, generally pions, may collide with the $\bar{N}$ or its targeted nucleon and prevent annihilation. Annihilation is, after all, a causal process taking place only when the nucleon and antinucleon overlap. The end result of our ARC calculations is $\bar{p}$ production rates for pA and AA processes which are in substantial agreement with experiment. These results have recently been published in parallel with the E802 collaboration.⁵

⁵S.H. Kahana, Y. Pang, T. Schlagel and C.B. Dover, Phys.Rev. **C47** (1993) R1356.

High-temperature QCD and the Physics of the Quark-Gluon Plasma

We have made significant efforts to understand recent lattice data⁶ concerning the 'dynamical confinement' of quarks at high temperatures. According to these calculations, $\bar{q}q$ move in a highly correlated way even at very high temperatures, and 'wave functions' describing this correlation are similar to those at $T = 0$. It has been speculated that hadron-like modes exist even at high T , in sharp contrast to a picture of a nearly ideal quark-gluon plasma. These lattice data were understood, and even reproduced, using a two-dimensional Schrödinger equation with an effective quark mass of $M_{\text{eff}} = \pi T$. An analogy with QED suggested that a modified logarithmic Coulomb potential is the cause of these 'bound states'.⁷ A more detailed study⁸ has shown that a linear (quarkonium-like) potential in QCD is needed to explain these data. Even spin-dependent splittings (*e.g.*, the $\pi\rho$ splitting) are accurately reproduced. Contrary to the original interpretation of DeTar and DeGrand, this phenomenon does *not* imply the existence of hadronic modes in the plasma in any ordinary sense. Instead, it represents the existence of a non-trivial *magnetic* interaction between the gluo-magnetic currents in the plasma. Unlike the interaction of charges, the magnetic field is not screened in plasma, and these effects can survive even at high T .

The Production of Strangelets and Other Rare Objects

A phase-space coalescence model has been used in conjunction with ARC to describe composite particles (*e.g.*, deuterons, tritons *et cetera*) observed in relativistic heavy-ion collisions at the AGS. ARC generated proton and neutron distributions were used as input for a coalescence model which tests CM momentum and the distance of closest approach of neutron-proton pairs after freeze-out for the formation of a deuteron (with the appropriate statistical weight). Several sets of preliminary data have been successfully described by the model. The fall-off of deuteron multiplicity with increasing rapidity in the E802 Si+Au data is well described. The E814 data at $p_T = 0$ on deuteron, triton, and ^3He production in the Si+Pb reaction allows the model to be successfully extended to three nucleon coalescence. Finally, the

⁶T. DeGrand *et al.*, at Quark-Matter-91.

⁷T.H. Hansson and I. Zahed, Hadronic Correlators in Hot QCD, SUNY-NTG-91-44.

⁸V. Koch, E. Shuryak, G.E. Brown and A.D. Jackson, Confinement of quarks propagating in the spatial direction in hot QCD, SUNY-NTG-92-4.

same model parameters can be used to describe the 5° E886 data for Au+Pt production of deuterons, tritons and ^3He . The ^4He production seen in this experiment is also well described.

Using this simple coalescence model, we have estimated production rates for stable strange nuclear clusters in Si+Au and Au+Au collisions at Brookhaven AGS energies of 14.6 GeV/A and 11.7 GeV/A, respectively.⁹ As input to the calculations, we use lambda distributions (which are in good agreement with data) as well as proton and neutron distributions from ARC. If a bound $\Lambda\Lambda$ state exists, its production should be as large as about 8% of all central Au+Au events. Predictions have also been made for singly- and doubly-strange hypernuclei with baryon numbers $A = 3 - 6$.

We have estimated the production of strange quark clusters in high energy heavy ion reactions in which a quark-gluon plasma intermediate state is formed. A microscopic fission-chain model (PHASER) has been constructed for the description of QGP rehadronization. Color fluctuations, $\bar{q}q$ pair production and energy fragmentation have been taken into account. The spin-flavor degeneracy factors and the formation factors for multibaryon quark clusters are determined. Baryon, antibaryon and meson yields have been estimated, as well as those of light multi-strange quark clusters.¹⁰

These authors have extended the Bethe-Weizsäcker nuclear mass formula to strange hadronic matter composed of nucleons, lambdas and cascade hyperons.¹¹ The generalized formula contains several volume and symmetry terms constrained by phenomenologically determined Λ -nuclear, Ξ -nuclear and $\Lambda\Lambda$ interaction parameters and by hyperon-hyperon interaction parameters suggested by one boson exchange models. They find that multi- Λ hypernuclei are generally unstable with respect to $\Lambda\Lambda \rightarrow \Xi N$ conversion. For strange hadronic matter we determine, as function of the baryon number A , the line of strong-interaction stability, along which a large strangeness fraction $|S|/A \sim 0.5 - 1.1$ and a low charge fraction $q/A \sim 0.2$ hold, and no fission occurs. The binding energy per baryon increases monotonically to its bulk limit, $B/A \rightarrow 38$ MeV, $|S|/A \rightarrow 1.1$ and $q/A \rightarrow 0$ for a reasonable set of parameters under the assumption that the hyperon species saturate at den-

⁹A.J. Baltz, C.B. Dover, S.H. Kahana, Y. Pang, T. Schlagel, and E. Schnedermann, *Physics Letters B*, submitted for publication.

¹⁰C.B. Dover, Z. Árvay, T. Csörgö, J. Zimányi and U. Heinz, *Z.f.Physik C*, submitted for published.

¹¹C.B. Dover and A. Gal, *Nuclear Physics A*, to be published.

sities similar to those of protons and neutrons in nuclei. Even in the extreme limit of vanishingly small hyperon-hyperon interaction strengths, strange hadronic matter with $B/A \rightarrow 15$ MeV, $|S|/A \rightarrow 0.7$ and $q/A \rightarrow 0$ in the bulk limit should exist. Our mass formula is in semi-quantitative agreement with recent mean-field calculations which implicitly assumed weak hyperon-hyperon interactions.

Stability of multiply strange baryonic systems in the context of a mean field approach was studied, based on an underlying set of phenomenological meson-baryon interactions.¹²

The coupling parameters which determine the mean fields seen by various baryon species in the many-body system are constrained by reproducing the trend of observed binding energies of single particle (i.e., N , Λ , Ξ) states as well as the energy per particle and density of non-strange nuclear matter. Extrapolating this approach to systems of large strangeness, S , we find a broad class of objects composed of p, n , Λ 's and Ξ 's, which are stable against strong decay. In these systems, the presence of filled Λ orbitals blocks the strong decay $\Xi N \rightarrow \Lambda \Lambda$ and leads to a strangeness fraction $f_s = |S|/A \approx 1$, density $\rho \approx 2\rho_0$, and charge Z comparable to that of hypothetical stable strange quark matter ('strangelets'), but with a low binding energy per particle $E_B/A \approx -10$ MeV. We have developed an approximate mass formula which describes the results of the mean field calculations. Such weakly bound multi-strange objects can be stable for arbitrarily large A , unlike ordinary nuclei, since the Coulomb repulsion generated by the protons is cancelled by the presence of a comparable number of Ξ^- 's. We have studied the weak decays of such objects and the possibility of their production in relativistic heavy-ion collisions.

In the course of a general study of multiply strange nuclear systems, the stability of systems formed by adding Ξ hyperons to ${}^6_{\Lambda\Lambda}\text{He}$ was studied.¹³ In this case, much of the 23 MeV energy difference between $\Xi^0 + n$ and $\Lambda + \Lambda$ is cancelled by the 20.6 MeV neutron separation energy from the α particle. In fact, the Ξ^0 needs to be bound by only 1.17 MeV to ensure the stability of the system against decay via strong interactions. This corresponds to a Ξ^0 well depth of around 14 MeV which is less than the estimate of 20–24

¹²C.B. Dover, D.J. Millener, A. Gal, J. Schaffner and H. Stöcker, Phys.Rev.Lett, submitted for publication, and Annals of Physics, submitted for publication.

¹³(C.B. Dover, D.J. Millener and A. Gal, Nuclear Physics A, submitted for publication.

MeV based on the existing candidate Ξ^- -nucleus data. With strangeness -4, baryon number 7 and a density approaching twice that of nuclear matter, this conventional system has quantum numbers approaching those of the hypothetical strangelets with $|S/A| \sim 1$. The expected lifetime should, however, be in the 100 ps range typical of Λ hypernuclei. The system obtained by adding another Ξ^0 should also be stable unless the 1S_0 $\Xi\Xi$ interaction is strongly repulsive. We have estimated the production rates for this object in relativistic heavy ion collisions.

Other Activities

Tunneling phenomena in the QCD vacuum, described semiclassically by 'instantons', have long been known to be very important for hadronic physics. Recently, evidence has accumulated suggesting that instanton-induced effects are of dominant importance in producing the quark condensate and the masses of most hadrons, including pions and nucleons. The 'instanton liquid model'¹⁴ explained these phenomena qualitatively, and a quantitative theory is now being developed. The principal progress is contained in a sequence of papers written by Shuryak and Verbaarschot (and, in one case, with the participation of Thomas Schaefer). These papers studied mesons and baryons in the 'instanton liquid' model of the QCD vacuum. These works describe numerical simulations performed at NRSC CRAYs, and the major results are masses and other parameters of low-energy hadrons. These results significantly increased accuracy of the measurements. For example, for the first time it was demonstrated that this model leads to a nucleon as a bound state of the correct mass *without* confining effects. The nucleon-delta splitting can be reproduced without perturbative gluomagnetic spin-spin interactions. Another important achievement is direct comparison with lattice data on correlation functions recently obtained by the group of John Negele and coworkers at MIT. Rather unexpectedly, good agreement was found between the two calculations — sometimes literally within error bars. The reason for this good agreement remains to be understood, but it suggests that instanton-induced effects provide the major component of quark-quark interactions inside hadrons.

One of the reasons for the importance of instanton-induced dynamics is

¹⁴E. Shuryak, Nucl.Phys. B203 (1982) 93, 116, 140; D.I. Dyakonov and V. Petrov, Nucl.Phys.

that, in addition to predicting chiral restoration at high temperature, it also suggests a non-trivial role for light quark masses and the number of flavors. Future works along this line (*e.g.*, on properties of the chiral restoration phase transitions, hadronic modifications *et cetera*) can only proceed when instanton behaviour at non-zero temperatures is understood in some detail. Analytical and numerical work in this direction has been published¹⁵ and will continue.

As a by-product of work devoted to tunneling in QCD, some work was also done in order to understand possible tunneling in weak interactions. Although the probability for such tunneling is small under normal conditions, it has been proposed that tunneling can become observable in the SSC domain. There, the cross section for of baryon-number violating processes grow rapidly as one proceeds towards the multi-TeV region. We have performed calculations aimed at the evaluation of this probability by determining the 'streamline' set of configurations¹⁶ which describe the interaction between instantons and anti-instantons. These results were used in papers by other authors and also by E. Shuryak and J. Verbaarschot.¹⁷ We have shown that the 'valley' method reproduces well-known results at low energies and predicts rapid growth of the cross section with energy. There are, however, limitations of this method which preclude its use above a critical energy (of about 35 TeV). Nevertheless, the possibility that this phenomenon might become observable remains. Further work is planned to investigate this question.

We are encouraged by the first five months of operation of this Project. Of course, the broad areas of scientific interest have been defined by many years of activity in nuclear theory both at Brookhaven and at Stony Brook. The primary role of the present contract is to provide a vehicle to give greater emphasis and greater coherence to issues of particular interest to RFP-C physics. Thus, quite aside from progress made on specific topics, we are particularly

¹⁵E. Shuryak and J. Verbaarschot, QCD instantons and finite temperature, SUNY-NTG-91-3.

¹⁶J. Verbaarschot, Streamline and Conformal Invariance in Yang-Mills Theories, SUNY-NTG-91-7.

¹⁷E. Shuryak and J. Verbaarschot, On Baryon Number Violation and Non-Perturbative Weak Processes at SSC Energies, SUNY-NTG-91-42.

encouraged by the fact that we have already established a variety of ways to increase the level of interaction between Stony Brook and BNL. This project is literally in its infancy. We expect it will grow in ways which are profitable to both experimental and theoretical nuclear physics.

II. ABSTRACTS OF ONGOING PROJECTS

Strange Cluster Formation in Relativistic Heavy Ion Collisions

A.J. Baltz, C.B. Dover, S.H. Kahana, Y. Pang, T. Schlagel and E. Schnedermann

Using the cascade code ARC to simulate relativistic heavy ion collisions at Brookhaven AGS energies (11.7 – 14.6 GeV/c), we have estimated the production rate of strange clusters ranging from a hypothetical doubly strange ($S = -2$) bound $(\Lambda\Lambda)_b$ dibaryon to the hypernuclei ${}^6_{\Lambda\Lambda}\text{He}$ and ${}^7_{\Xi^0\Lambda\Lambda}\text{He}$. We extrapolate a simplified version of the coalescence model to systems with $S \neq 0$, with parameters adjusted to reproduce data on non-strange composite formation (d , ${}^3\text{He}$, ${}^3\text{H}$, ${}^4\text{He}$) at these energies. We predict that $(\Lambda\Lambda)_b$ and ${}^6_{\Lambda\Lambda}\text{He}$ would be produced with rates dN/dy at mid-rapidity of order 10^{-1} and 5×10^{-6} , respectively, for central Au+Au collisions. Some experimental possibilities for detection of strange clusters are discussed [1].

[1] A.J. Baltz, C.B. Dover, S.H. Kahana, Y. Pang, T. Schlagel and E. Schnedermann, Physics Letters B, submitted for publication.

Deuteron Production

Ekkard Schnedermann, Anthony Baltz, Carl Dover (BNL)

Based on our previous work [1] on cluster formation in relativistic heavy-ion collisions, which employed a thermal framework, we now work on implementing these methods for the relativistic cascade code ARC [2]. The deuteron plays the role of a simple test case, both in experimental and theoretical respect, for the heavier nuclei like 3H , 4He , and even light hypernuclei.

Experimentally the abundance of the deuterons is much larger than in the other cases, so that already a wealth of data from the AGS experiments is available, which can be compared to our computations. Theoretically the deuteron coalescence has already a long history, albeit for much lower energies, from which we can select hints for the appropriate description for the AGS energy range.

The ARC code generally delivers a good description of the hadron spectra from heavy-ion collisions at AGS energies and thus probably implements a good description of the collision dynamics, which can be tested with the deuteron rate. We have improved on the standard coalescence model, which uses sharp cutoffs in space and momentum for the formation of the deuteron, by using the actual wave function of the deuteron. Our first systematic studies already seem to indicate interesting observations.

- [1] C.B. Dover, U. Heinz, E. Schnedermann and J. Zimanyi, Phys.Rev. **C44** (1991) 1636.
- [2] Y. Pang, T.J. Schlagel and S.H. Kahana, Phys.Rev.Lett **68** (1992) 2743.

Work on the relativistic cascade ARC

T.J. Schlagel, S.H. Kahana and Y. Pang)

We are continuing the development of the relativistic cascade code ARC for nucleus–nucleus collisions. The code has been used successfully at AGS energies to help in the understanding of the complex dynamics of nucleus–nucleus collisions. The approach is to include measured hadron–hadron cross sections as the input to study rescattering of particles in the nuclear medium. The experimental hadronic data is parameterized as a function of energy for interactions at and below AGS energies. Recently, these parametrizations have been extended to higher energy and will be used to study heavy ion collisions at CERN and RHIC energies.

Comparisons between the cascade code and experimental proton spectra in Si+Au collisions clearly demonstrate the importance of the formation and propagation through the nuclear medium of baryon resonances in nucleus–nucleus collisions. The low mass baryon resonances Δ and N^* play a very important role in rescattering at AGS energies. Their presence can explain the proton rapidity distribution, the increased proton ‘temperature’, and the seemingly large number of forward protons. A consequence of baryon resonance dominance in heavy ion collisions at AGS energies is the increased production of strange particles over hadron–hadron collisions. Strangeness production is suppressed at these energies due to the high threshold for production compared with pure pion production. In ARC, the resonances retain most of their energy during rescattering and thus have the chance to produce strangeness in subsequent collisions. Further study of strangeness production is being pursued to determine the contributions from the different hadron–hadron channels (*e.g.*, associated and pair production of strange mesons), and the effects of including K^* propagation on the rapidity and transverse mass spectrum.

ARC has also been used to study the evolution of baryon density in Si+Si, Si+Au, and Au+Au collisions at 11.6 GeV/c laboratory momentum. A region of high baryon density is formed in central Au+Au collisions and is, to a certain degree, thermalized. Preliminary experimental data on inclusive particle spectra from E866 agree with our predictions despite the high baryon densities reached. Signatures for interesting physics, such as the formation of a quark-gluon-plasma, may therefore appear in a subtler form. This makes

a detailed phase space calculation, such as ARC, especially important in understanding the phenomenon of high baryon densities. Further calculations on Au+Au collisions are being made to explore observables which may allow experimental triggering on events with high baryon densities. Along these lines, preliminary calculations on including equation of state effects for nucleus-nucleus collisions into the cascade code ARC through a Pauli exclusion at high baryon density suggests exclusion effects will be stronger at laboratory momenta lower than 11.6 GeV/c, where the six-dimensional phase-space density is higher. Other equation of state effects, such as changes in hadronic masses, are also being considered.

Due to correlations in the particle coordinates and momenta, two-particle interferometry does not directly measure the size of the interaction region. In ARC, one has all the information about the coordinate space and momentum space distribution at any given time. This information can be used directly to predict the experimental interferometry results and study the relationship of interferometry measurements to the size and the entropy of these systems. Recently collaborations have begun with some of the experimental groups on trying to understand the interferometry results.

At higher beam energies, the lowest mass resonance production model may no longer be applicable. Particle production in hadron-hadron collisions is mostly through discrete resonances at low energies. This picture may also be applicable as energies increase. We are developing a production model where nearly all inelastic collisions form two resonances from a given mass and width spectrum. At low energies these resonances are known, but at higher energies the resonance mass spectrum becomes continuous. Particle production follows from the cascade of high mass resonances to lower mass resonances through the emission of hadrons. The details of this model, such as the momentum correlations of produced particles, are currently under study. Eventually this model will be applied to the study of nucleus-nucleus collisions.

Antiproton Production from Heavy Ion Collisions at 14.6 GeV/c

S.H. Kahana, Y. Pang, T. Schlagel and C.B. Dover

Antiproton production in heavy ion collisions involves a strong competition between creation and annihilation mechanisms. Due to the high threshold energy, antiprotons mostly arise from the highest energy baryon-baryon collisions. We have identified a three-body screening mechanism which strongly reduces the subsequent absorption of antiprotons in the medium. We have used the relativistic cascade code ARC to study this effect, and find that it is essential to obtaining good agreement with recent $\bar{p}$ production data from Brookhaven experiment E802 [1].

[1] S. Kahana, Y. Pang, T. Schlagel and C.B. Dover, Phys.Rev. **C47** (1993) R1356-59.

Two-stage Equilibration in High Energy Heavy Ion Collisions[†]

E.V. Shuryak

Using (lowest-order) perturbative QCD, we propose a new scenario for high energy heavy ion collisions, the so called 'hot glue' scenario. We have found that such collisions are likely to proceed via two stages. First, a rapid thermal equilibration of the gluon momentum which takes time $\tau_g \sim 1/3$ fm/c. Second, the multiplication of gluons until chemical equilibrium is reached. This takes several times longer. At the same time, the production of quarks is strongly suppressed, so one only has those quarks which were present as partons of the colliding nuclei.

If this is the case, the initial gluon plasma will be much hotter than previously estimated with an initial temperature $T_g \sim 500 \text{ MeV}$. This is to be compared with the value of 250 MeV found in the so-called 'standard scenario'. This temperature change would lead to significant modifications of all proposed 'QGP signals'.

One of the main consequences of the proposed 'hot glue' scenario is the enhanced production of charm due to reaction $gg \rightarrow \bar{c}c$ reaction which was suggested as a signature for the QGP in [1] and was studied in detail in ref.[2]. This is a very sensitive 'thermometer' for the temperature region in question, $T = 400 - 700 \text{ MeV}$ and (in contrast to the production of strangeness) the approximate perturbative evaluation of all relevant rates is possible.

[†]This paper is published in Phys.Rev.Lett. **68** (1992) 3270.

[1] E. Shuryak, Sov.J. of Nucl.Phys. **28** (1978) 408.

[2] A. Shor, Phys.Lett. **215B** (1988) 375, **233B** (1989) 231.

Dilepton and Photon Production in the 'Hot Glue' Scenario

E. Shuryak and L. Xiong

Perturbative analysis of parton kinetics for high energy nuclear collisions shows that thermal equilibration of gluons takes place very quickly. This suggests that their equilibration temperature is quite high [1]. The sources for photon and dilepton production, quarks and antiquarks, however, are produced much more slowly.

In this work we have considered a simple 'minimal' model which includes only incoming quarks and antiquarks. We have determined the dilepton and photon spectra for this minimum charge and find that, in the interesting kinematic region, plasma signals are actually larger than those previously obtained. If this is the case, experimental observations of thermal dileptons and photons from a quark-gluon plasma have a better chance of success.

[1] E. Shuryak, Phys.Rev.Lett. **68** (1992) 3270.

[†]This paper has been published in Phys.Rev.Lett. **70** (1993) 2241.

Parton Multiplication and Chemical Equilibrium

L. Xiong and E. Shuryak

Chemical equilibration of a parton system (relaxation of chemical potential to zero) should take longer time compared to thermal equilibration of parton momenta, because the various particle production processes are of higher order in α , than the elastic scatterings. The rate of production and number of partons produced in an ultrarelativistic heavy ion collision directly affect the dynamics of the quark-gluon plasma and the final particle multiplicity.

We are particularly interested in the gluon number-changing processes, $mg \leftrightarrow ng$. We have evaluated these reaction rates in a thermal equilibrated plasma for arbitrary m and n . In chemically non-equilibrated plasma the net number-changing rate is nonzero and tends towards equilibrium.

Equation for time evolution of the gluon chemical potential is derived, which includes those rates and plasma expansion. We will study how and when the saturation occurs in an expanding plasma with a longitudinal scaling-invariant motion. We evaluate chemical relaxation time and the amount of entropy production from perturbative QCD mechanisms, between thermal and chemical equilibration.

Parton Equilibration in a Cascade Model

L. Xiong and E. Shuryak

The approach to thermal equilibrium and formation of a quark-gluon plasma at RHIC energies is studied by parton cascade model. It starts with structure functions as initial conditions and simulates the parton system within the nucleus and follows the various microscopic interactions among the constituents.

With this model we can obtain a complete picture of the space-time evolution of the collision process and study many interesting features of the parton system before and after the establishment of thermal equilibrium including entropy production, charm production and other plasma signatures.

The difference between our approach and parton cascades developed by K.Geiger is mainly connected with treatment of multi-parton amplitudes: we do not use the notion like 'virtual gluons' and processes like $g \rightarrow gg$, but keep only physical cross sections on mass shell. The work is now in progress and it is too early to conclude whether results agree or not.

Correlation Functions in the QCD Vacuum[†]

E.V. Shuryak

Correlation functions are one of the key tools used in studies of the structure of the QCD vacuum. They can be evaluated in several ways. First, they can be constructed out of the fundamental fields and can be calculated using quantum field theory methods such as lattice gauge theory. One can also obtain many of these functions empirically using the rich phenomenology of hadron physics. They are also very interesting objects to calculate using various quark models of hadronic structure. In short, they are the main tool allowing us to probe interaction between quarks and anti-quarks. Existence of several approaches to evaluate them can make these correlation functions the common ground for the comparison of quite different approaches.

This review paper is focused on available phenomenological information about these correlation functions and emphasizes the most important qualitative observations. We also give quantitative plots of all main known correlation functions which can be directly compared with lattice results.

After that, we review the predictions of various theoretical approaches including lattice numerical simulations, QCD sum rules based on the operator product expansion and calculations based on the 'instanton liquid' model.

[†]This paper is published in Rev.Mod.Phys. 65 (1993) 1.

Quark Propagation in a Random Instanton Vacuum[†]

E.V. Shuryak and J.J.M. Verbaarschot

In this work we report an extensive numerical study of the simplest ensemble of instantons — the random instanton vacuum. This vacuum is characterized by two parameters: (i) The size ρ and (ii) The density $\tilde{n}$ of instantons (plus antiinstantons) in the QCD vacuum which are chosen to reproduce the size of the quark condensate and the gluon condensate.

For simplicity, we first studied the propagation of scalar quarks (or squarks), in which case there is only a nonzero mode contribution to the propagator. It appears that the scalar mesonic correlation function can be described remarkably well by a ‘constituent quark’ model. However, for spinor quarks the propagator has two different components: the chirality non-flip and the chirality parts. The first one can again be described by a relatively constant quark mass of about 300 Mev. The second, spin-flip component is more complicated and does not resemble a free massive propagator.

We have also studied eigenvalues of the Dirac operator. Because the topological charge in this model fluctuates strongly on all length scales, there is an excess of small eigenvalues above the expected Gaussian distribution. This effect leads to a strong breaking of chiral symmetry and leads to relatively large effective quark masses.

Finally, we use these results for the propagator to derive some consequences for heavy-light mesons such as the B -meson. We have found that B -type mesons have masses 615 MeV larger than the quark mass which is in good agreement with phenomenology. Another result is the splitting between the opposite parity states (*e.g.*, pseudoscalar and scalar) which is found to be 575 MeV in this model. Finally, we have found decay constants for heavy-light mesons and have compared the results with other predictions, including lattice results. Again, the agreement is reasonably good.

[†]This paper has been submitted to Nuclear Physics B.

Mesonic Correlation Functions in the Random Instanton Vacuum[†]

E.V. Shuryak and J.J.M. Verbaarschot

First, a general model-independent discussion of correlation functions is given. We derive new inequalities for these functions and mesonic masses including ones stronger than the Weingarten inequality [1]. A rather complete set of mesonic correlation functions is calculated in the random instanton liquid model. The results are compared with phenomenological expectations [2] and lattice results [3]. These include both diagonal and off-diagonal correlators of all strange and light flavored currents as well as the most important unflavored ones. Our results are used to extract the masses and the coupling constants of the corresponding mesons.

The original idea was to use a crude model as a benchmark for further studies of interacting instantons with the parameters (the instanton density and typical radius) as suggested a decade ago [4] without adjustment. However, the results obtained have shown that this model works amazingly well. As was originally the case for the nuclear shell model, it is not currently clear why it reproduces the correlation functions so well. This result is surprising in view of the apparent absence of such ingredients as confining forces and perturbative effects.

In fact, we reproduce not only the qualitative behaviour of all correlators at distances as large as 2 fm, but (in those few cases where quantitative analysis is possible) we find agreement of physical parameters at the 10 percent level. Thus, it is certainly a useful tool. Wider use of this model would seem to be justified. At the same time, in two channels with strong repulsive interaction — η' and δ — the results found are not satisfactory. The model correctly predicts such repulsion but overestimates its magnitude.

[†] This paper is submitted to Nucl.Phys.B.

- [1] D.Weingarten, Phys.Rev.Lett. **51** (1983) 1830.
- [2] E. Shuryak, Rev.Mod.Phys., January, 1993.
- [3] M.C. Chu, J.M. Grandy, S. Huang, and J.W. Negele, Phys.Rev.Lett., to be published.
- [4] E. Shuryak, Nucl.Phys. **B203** (1982) 116, 140, 237.

The Absence of Strong CP Violation in a Correlated Instanton Vacuum

E.V. Shuryak and J.J.M. Verbaarschot

The absence of strong CP violation is one of the unsolved problems of QCD: either the mechanism should exist which can bring the vacuum θ angle to zero (axions?), or fluctuations of the topological charge are completely screened. The latter possibility for *non-zero* quark masses was recently proposed by Samuel [1], where one can find all relevant references.

We study fluctuations of the topological charge by performing numerical simulation of an ensemble of interacting instantons. The main correlations are induced by the zero mode sector of the fermion determinant, but classical correlation are also taken into account. It is shown that, for small volumes, the fluctuations of the topological charge are random while, for larger volumes, they are strongly suppressed (*i.e.*, the topological charge is screened). However, if quark masses grow from zero, this mechanism gradually turns off, which probably contradicts to the the Samuel's proposal.

The inverse screening length determines the η' mass. It is shown that the screening length is inversely proportional to the square root of number of massless flavors which agrees with the Witten-Veneziano formula.

[1] S. Samuel, Mod.Phys.Lett. **A7** (1992) 2007.

Baryonic Correlators
in the Random Instanton Vacuum

T. Schäfer, E.V. Shuryak and J.J.M. Verbaarschot

Recently we have performed a systematic study of QCD correlation functions in the framework of the instanton liquid model. In the case of mesonic correlators we have obtained very good agreement with lattice measurements reported by Chu *et al.* We have extended this analysis to include the nucleon and delta correlation functions.

As a first step, we have considered quark-quark (diquark) systems. We have found that the quark-quark interaction resembles the one between quarks and antiquarks. Similar to the pion and rho channels, the interaction in the scalar isospin $I = 0$ and vector $I = 1$ diquark channels are completely different: The former has a much stronger attractive interaction. As a consequence, the $SU(3)$ octet (nucleon) and decuplet (delta) correlators are also found to be qualitatively different. Using a complete set of all available correlation functions, we also determine masses and coupling constants for the nucleon and delta.

- [1] E.V. Shuryak, J.J.M. Verbaarschot, SUNY-NTG-92-40.
- [2] T. Schäfer, E.V. Shuryak, J.J.M. Verbaarschot, SUNY-NTG-93.
- [3] M.C. Chu, J.M. Grandy, S. Huang and J.W. Negele, Phys.Rev.Lett. **70** (1993) 255.

Random Matrix Theory and Spectral Sum Rules for the Dirac Operator in QCD

E.V. Shuryak and J.J.M. Verbaarschot

Although the theory of strong interactions is formulated in terms of quarks and gluons, at low energies the same physics can be formulated in terms of mesons and baryons. This equivalence results in strong constraints on the spectrum of the Dirac operator. It appears that the existence of a long wave-length limit of the bosonized theory results in a low-lying spectrum of the Dirac operator that is *universal*. In order to study the low-lying part of the spectrum, we have introduced a new limit: the microscopic limit. In this limit the spectrum near zero is enlarged proportional to the space time volume of the system, and the spectrum near zero virtuality converges to a universal function [1]. Its existence is an expression of the spontaneous breaking of chiral symmetry.

The shape and the correlations of the spectrum have been determined [1] with the help of a random matrix model [2] (which we call the ‘chiral random matrix model’) that, in the large N limit, reduces to the low energy limit of the QCD partition function put forward by Leutwyler and Smilga [3]. This equivalence holds for an arbitrary number of flavors and any value of the QCD vacuum angle. In this model, moments of the inverse squares of the eigenvalues of the Dirac operator obey sum rules, which we conjecture to be universal. In other words, the validity of the sum rules depends only on the symmetries of the theory but not on its details. To illustrate this point we have shown that the sum rules hold for an interacting liquid of instantons. The physical interpretation is that the way the thermodynamic limit of the spectral density near zero is approached is universal. However, its value, *i.e.*, the chiral condensate, is not.

[1] J.J.M. Verbaarschot and I. Zahed, Physical Review Letters, to be published.

[2] J.J.M. Verbaarschot and E.V. Shuryak, Nuclear Physics A, to be published.

[3] H. Leutwyler and A. Smilga, Physical Review D, to be published.

The Higgs-Induced Contribution to Baryon Number Violation at Very High Energies[†]

M. Velkovsky and E.V. Shuryak

The probability of baryon number violation at SSC energies due to tunneling in weak interactions is a subject of intensive studies. This process is predicted to be accompanied by multiple production of W and Higgs bosons and can be technically described by the instanton-antiinstanton ($\bar{I}I$) interaction (see references in [1]). We evaluate numerically the Higgs-induced part of this interaction, corresponding to the so called 'streamline' gauge configurations. As this interaction is known [2] to be repulsive at large $\bar{I}I$ distances, the question was raised whether it can produce a 'second barrier' and significantly reduce the tunneling probability.

We have used a variational approach, using the best trial functions we could find. (Still they are *not* the true minimum of the action, in particular, at zero $\bar{I}I$ separation the total action is not exactly zero.) However, we have found that the so-called *complete Yung ansatz* [2] is a much better approximation to such a minimum than the 'simplified' one for the case of parallel relative orientation. The interaction in this case is such, that it definitely rules out the 'second barrier' scenario.

If so, extrapolation of the cross section into the interesting energy region $E \sim 30$ TeV does not suffer from Higgs-induced suppression. Therefore, the possibility that the effect reaches an observable magnitude (before all the approximations do collapse) looks slightly more probable.

[†]This paper will appear in Physics Letters B.

[1] E.V. Shuryak and J. Verbaarschot, Phys.Rev.Lett. **68** (1992) 2576.

[2] A. Yung, Nucl.Phys. **B297** (1988) 47.

RHIC Systems and Self-Organizing Criticality

A.D. Jackson, V. Koch and T. Wettig

The construction of 'event generator' codes has become a popular activity in preparation for RHIC. These codes share the 'anecdotal' feature that many predictions are remarkably insensitive to input parameters. We suspect that something more fundamental may be taking place. There are a number of (schematic) models of interest to condensed matter theorists which display 'self-organizing criticality'. One such model is the 'forest fire' model of ref.[1]. The common feature of these models is the existence of two (or more) disparate time scales. Their common result is that the predictions for many quantities including the average number of 'partons', the distribution of 'parton' clusters and various correlation functions are *strictly* independent of the details of the underlying dynamics. (In the case of ref.[1], the 'partons' are trees.) The only vestige of the dynamics which persists is the existence of several time scales. We are considering whether such models may be applicable to RHIC systems.

Thus, we are looking for (schematic) self-organizing models whose basic assumptions seem *a priori* compatible with the physical conditions of heavy ion collisions. Once such analogies are discovered, various predictions can be obtained immediately. Such models would offer a two-edged sword for RHIC physics. Quantitative, dynamics-independent predictions would be helpful in checking codes and for comparisons with data. However, the existence of predictions independent of the dynamics would complicate the task of unraveling RHIC data to reveal microscopic dynamics.

While such studies are clearly speculative, we view them as important. Condensed matter studies of these models suggest that complex systems manage to filter out virtually all dynamical information. Any convincing demonstration that RHIC systems are self-organizing would provide a 'proof' of the *non-existence* of signatures for any specific microscopic description of RHIC events.

[1] B. Drossel and F. Schwabl, Phys.Rev.Lett. **69** (1993) 1629.

Strange Hadronic Matter

J. Schaffner, C. B. Dover, A. Gal, C. Greiner and H. Stöcker

In an extended mean field theory, we find a large class of bound multi-strange objects, formed from combinations of $\{p, n, \Lambda, \Xi^0, \Xi^-\}$ baryons, which are stable against strong decay. We predict a maximal binding energy per baryon of $E_B/A \approx -21$ MeV, strangeness per baryon $f_s \approx 1 - 1.2$, charge per baryon $f_q \approx -0.1$ to 0.1 , and baryon density 2.5-3 times that of ordinary nuclei. For $A \geq 6$, we obtain stable combinations involving only $\{\Lambda, \Xi^0, \Xi^-\}$ hyperons [1].

[1] J. Schaffner, C. B. Dover, A. Gal, C. Greiner and H. Stöcker, Physical Review Letters, submitted for publication.

A Generalized Bethe-Weizsäcker Mass Formula for Strange Hadronic Matter

C.B. Dover and A. Gal

We extend the Bethe-Weizsäcker nuclear mass formula to strange hadronic matter composed of nucleons, lambdas and cascade hyperons [1]. The generalized formula contains several volume and symmetry terms constrained by phenomenologically determined Λ -nuclear, Ξ -nuclear and $\Lambda\Lambda$ interaction parameters and by hyperon-hyperon ($\Lambda\Lambda$, $\Lambda\Xi$, $\Xi\Xi$) interaction parameters suggested by One-Boson-Exchange models. We confirm that multi- Λ hypernuclei are generally unstable to $\Lambda\Lambda \rightarrow \Xi N$ conversion. For strange hadronic matter we determine, as function of the baryon number A , the line of strong-interaction stability, along which a large strangeness fraction $|S|/A \sim 0.5 - 1.1$ and a low charge fraction $q/A \sim 0.2$ hold, and no fission occurs. The binding energy per baryon increases monotonically to its bulk limit, $B/A \rightarrow 38$ MeV, $|S|/A \rightarrow 1.1$ and $q/A \rightarrow 0$ for the parameters adopted here assuming that the hyperon species saturate at densities similar to those of protons and neutrons in nuclei. Even in the extreme limit of vanishingly small hyperon-hyperon interaction strengths, strange hadronic matter with $B/A \rightarrow 15$ MeV, $|S|/A \rightarrow 0.7$ and $q/A \rightarrow 0$ in the bulk limit should exist and our mass formula reproduces semi-quantitatively recent mean-field calculations which implicitly assumed weak hyperon-hyperon interactions.

[1] C.B. Dover and A. Gal, Nuclear Physics A, to be published.

Production of Strange Clusters by Quark Gluon Plasma Fragmentation

Z. Árvay, J. Zimányi, T. Csörgö, C.B.Dover and U. Heinz

We discuss the production of strange quark clusters in high energy heavy ion reactions in which a Quark-Gluon-Plasma (QGP) intermediate state is excited. A microscopic fission-chain model (PHASER) has been constructed for the description of QGP rehadronization. Color fluctuations, $\bar{Q}Q$ pair production and energy fragmentation have been taken into account. We determined the spin-flavor degeneracy factors and the formation factors for multibaryon quark clusters. Baryon, antibaryon and meson yields have been estimated, as well as those of light multi-strange quark clusters [1].

[1] Z. Árvay, J. Zimányi, T. Csörgö, C.B.Dover and U. Heinz, Zeitschrift für Physik, submitted for publication.

Multiply Strange Nuclear Systems

J. Schaffner, C.B. Dover, A. Gal, C. Greiner, D.J. Millener and H. Stöcker

We investigate the stability of multiply strange baryonic systems, in the context of a mean field approach obtained from an underlying set of phenomenological meson-baryon interactions. The coupling parameters which determine the conventional $\sigma + \omega$ mean fields (Hartree potentials) seen by various baryon species (N , Λ , Ξ) in the many-body system are constrained by reproducing the trend of observed binding energies of single particle (N , Λ , Ξ) states, as well as the energy per particle and density of non-strange nuclear matter. We also consider additional scalar (σ^*) and vector (ϕ) fields which couple strongly to strange baryons. The couplings of these fields are adjusted to produce strong hyperon-hyperon interactions, as suggested by the data on $\Lambda\Lambda$ hypernuclei. Extrapolating this approach to systems of large strangeness S , we find a broad class of objects composed of neutrons, protons, Λ 's and Ξ 's, which are stable against strong decay. In these systems, the presence of filled Λ orbitals blocks the strong decay $\Xi N \rightarrow \Lambda\Lambda$, leading to a strangeness fraction $f_s = |S|/A \approx 1$, density $\rho \approx (2-3)\rho_0$, and charge fraction f_q in the range $-0.1 < q/A < 0.1$, comparable to that of hypothetical stable strange quark matter ("strangelets"), but with a low binding energy per particle $E_B/A \approx -10$ to -20 MeV. We compare with an approximate mass formula which qualitatively describes the results of the mean field calculations. Such weakly bound multi-strange objects can be stable for very large A , unlike ordinary nuclei, since the Coulomb repulsion generated by the protons is largely cancelled by the presence of a comparable number of Ξ^- 's, leading to a small net charge (positive or negative) of order $A^{1/3}$. We comment on the weak decays of such objects and the possibility of their production in relativistic heavy ion collisions [1].

[1] J. Schaffner, C.B. Dover, A. Gal, C. Greiner, D.J. Millener and H. Stöcker, *Annals of Physics*, submitted for publication.

On The Production and Decay of Strangeness $S = -2$ Hypernuclei

C.B. Dover, A. Gal and D.J. Millener

We examine the production and decay mechanisms of doubly strange $S = -2$ hypernuclei, focusing on reactions of the type (K^-, KX) induced by kaons. The production of Ξ and $\Xi^*(1530)$ hyperons in (K^-, K^+) reactions on nuclear targets is considered, including the possibility of observably narrow Ξ single particle bound states. We present some qualitative arguments regarding the baryon (n, p, Λ) decay widths of Ξ states, which indicate that such excitations could be considerably narrower in light nuclei than suggested by the Ξ decay width $\Gamma_{\Xi} \approx 5 - 10$ MeV in nuclear matter. The predicted spectra of low-lying $S = -2$ hypernuclear states are presented. These configurations, dominantly of $\Lambda\Lambda$ character, can be produced in second order processes with K^- beams. We consider the possibility that the very small cross sections for such states (typically of order 1 nb or less) might be enhanced by $\Xi N \leftrightarrow \Lambda\Lambda$ mixing. Some prospects for the study of strong, electromagnetic and weak decays of low-lying $S = -2$ states are pointed out, and favorable cases for experimental study are identified [1].

[1] C.B. Dover, A. Gal and D.J. Millener, Nuclear Physics A, submitted for publication.

**DATE
FILMED**

7 / 30 / 93

END