AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

DOE/ER/40747-10

(updated from DOE/ER/40747-8)

The Partial Snake Experiment at the AGS

for the Funding Period Aug. 15, 1992 - Aug. 14, 1994

S.Y. Lee

Department of Physics

Indiana University, Bloomington, IN 47405

July 15, 1993

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Prepared for

The U.S. Department of Energy

Agreement No. DE-FG02-92ER40747

MASTER

The Annual Progress on the Partial Snake Experiment at the AGS

In the two year funding period from Aug. 15, 1992 - Aug. 14, 1994, our progresses can be summarized as following: (1) The solenoidal partial snake was installed in the AGS synchrotron in May, 1993 and was subsequently successfully tested. (2) The internal polarimeter for the AGS was assembled, tested and installed in the AGS. (3) The 200 MeV polarimeter at the LINAC was tested. (4) The AGS polarized ion source was renovated. (5) The first phase of the polarized beam experiments was successfully accomplished in April 1-8, 1994. of polarized proton acceleration up to $G\gamma \approx 20$. (6) We are currently renovating tune jump quadrupoles in order to accelerate polarized proton up to 25 GeV/c. The full test of these experiments will be in 1995.

During this funding period, the principal investigator has spent about 20% of his research time on this project, which includes the design and manufacturing of the solenoid partial snake, beam dynamics issues of the AGS with the 5% solenoid partial snake and spin dynamics of synchrotrons with snakes in general. The graduate student, Mr. H. Huang has spent 100% of his research time working on spin dynamics. The graduate student is currently staying at BNL for the E880 experiment preparation.

1. Introduction

Depolarizing resonances are known to exist in all circular accelerators with transverse and longitudinal error fields. At certain energies, horizontal fields in accelerator interact coherently with the proton spin thereby depolarizing the beam. Depolarizing resonances occur when the spin tune ν_s , which equals to $G\gamma$ with $G = 1.7928$ and the relativistic Lorentz factor γ , satisfies the resonance condition, i.e. $\nu_s = kP \pm \nu_z \pm n$, where P is the superperiodicity of the accelerator ($P = 12$ for the AGS), ν_z is the vertical betatron tune, and k, n are integers.

Derbenev and Kondratenko¹ proposed to correct these depolarizing resonances by a spin rotator, called a *snake*. A snake is a composition of magnets, which rotates the spin of each particle by 180° without affecting the orbital closed orbit elsewhere. When a snake is introduced into an accelerator, the spin tune becomes $1/2$. Therefore the resonance condition will not occur. For low energy synchrotrons, 100% snake is relatively expensive due to large transverse orbit distortions. Partial snake² was proposed to overcome only imperfection resonances while the intrinsic resonances can be corrected by the traditional tune jump method.

The available intense polarized source, polarimeters, AGS Booster for multi-pulse accumulation, pulsed quadrupole system including power supplies for the betatron tune jump offers us a unique opportunity in the study of spin dynamics at the AGS. The spin dynamics experiment E880 at the AGS in Brookhaven National Laboratory is one of the major research efforts of the Indiana University accelerator physics group. The E880 experiment is recognized to be one of the most important test of the partial snake idea. The experiment, funded by the DoE DE-FG02-92ER40747, involves physicists from the IUCF, Indiana University, the Brookhaven National Laboratory, the Argonne National Laboratory, the Fermilab, the KEK Laboratory in Japan, and TRIUMF in Canada.

2. Progress of the Project and Accomplishments in the Funding Period

1. An excellent engineer design of the room temperature solenoid was completed in June 18, 1992 by Dr. Alan Otter of TRIUMF, and was reviewed thoroughly by engineers (Bill Leonhardt et al.) and physicists (L. Ratner, T. Roser, L. Ahrens et al. at BNL

and S.Y. Lee at Indiana University). The Everson Electric Company won the bid in June 29, 1992 and did a superb job in manufacturing the solenoid. The solenoid was completed in April 1993 [see appendix A], was delivered to BNL in April 1993 and was installed in the AGS ring in May 1993. We are scheduled to make an initial test of the solenoid with unpolarized proton beam in August 1993. The polarized proton experiment is scheduled in April, 1994.

2. The polarimeters and software data acquisition system for the 200 MeV LINAC and the polarimeter at AGS have been completed. A new design for the internal polarimeter mechanism has been finished which conforms with the new high vacuum requirements of the AGS. Fabrication has started and the whole mechanism installed and tested by the end of July 1993. The mechanism will be remotely controlled by LABVIEW software running on a Macintosh PC. A double-arm detector system is being developed at Argonne National Laboratory that will allow the separation of elastic scattering from quasi-elastic scattering in the interaction of the polarized protons with the Nylon target. The detector system was installed in July 1993. High quality cables of about 50 channels used for the detector system are also being installed. The AGS internal polarimeter has been tested in January, 1994.
3. The fast tune jump ($\Delta\nu_x \approx \mp 0.3, \Delta\nu_z \approx \pm 0.15$ in $2.5 \mu s$), which changes the betatron tunes nonadiabatically, is needed to overcome intrinsic resonances and restores to its original setting in about 2.5 ms. Thus betatron tunes cross the linear coupling resonance line during the adiabatic process and cause exchange of emittances. The preferable operational condition corresponds to an equal emittance for the horizontal and vertical planes³. However, the multi-pulse booster injection may violate this optimal condition of equal emittance. Experimental studies of emittance exchange between horizontal and vertical space are important. Recently, linear coupling correction schemes have also been studied³ for the AGS with 5% solenoidal partial snake.
4. The existence of even order snake resonances in the presence of the overlapping intrinsic and imperfection resonances has been observed in the tracking calculations and the linear response theory of spin mapping equation⁴. Because of the spin tune shift of imperfection resonances, each snake resonance was found to split into two. Careful

control of betatron tunes becomes an important task in polarized proton acceleration of high energy accelerators [see ref. 4 and DOE/ER/40747-9].

5. We have derived the tensor polarization transfer matrices for spin-1 particles in synchrotrons. The evolution of the tensor polarization in the presence of an isolated resonance has also been studied⁵. This is the first theoretical effort to study the evolution of the tensor polarization in synchrotrons.
6. A comprehensive review article about the polarized proton acceleration has recently be completed⁶.
7. The first phase experiments were accomplished in April 4-8, 1994. The polarized beam was successfully accelerated in the AGS up to 10.8 GeV kinetic energy without using vertical orbit correctors (see Appendix).

3. The Experimental Goals of E880 Experiment in 1994-1996

We have made important progresses during past two years on the accelerator physics experiments, E880, at the AGS. We list physics goals of the 1994-1996 as follows.

1. Study emittance growth during the tune jump^{3,7}: This is particularly important when the solenoid partial snake is used. The emittance growth or emittance exchange happens when $\epsilon_x \neq \epsilon_z$. In this experiment, we need a fast digitizing system for the betatron motion. The present AGS data acquisition system will be used for turn by turn phase space tracking. The available Ionization profile monitor in the AGS is handy in this task.
2. Study Beam Dynamics on the Effect of the Spin Chromaticity⁶: Making the spin chromaticity zero can avoid multiple crossing of spin resonance due to synchrotron motion. The spin resonance condition is given by $G\gamma - K = 0$, where $K = mP \pm \nu_z$. Expanding the resonance condition in momentum spread, one obtains the achromatic resonance condition as $G\gamma\beta^2 \mp C_z = 0$, where C_z is the vertical chromaticity. An achromatic spin resonance crossing at the AGS corresponds to a vertical chromaticity of +8.8 at the $0 + \nu_z$ resonance, and a vertical chromaticity of -27 at the $36 - \nu_z$ resonance. The vertical chromaticity of the AGS is about -5 at $0 + \nu_z$ and 0 at about

13 GeV/c and +10 at 25 GeV/c. Therefore sextupole excitation is needed to achieve spin chromaticity correction. The initial test for the unpolarized beam is to study the dynamical aperture caused by chromatic sextupoles.

3. The present 10 fast quadrupoles will contribute about 7% emittance growth⁷ in each tune jump provided that these tune jump quadrupoles are properly aligned⁸. A complete spin flip to overcome intrinsic spin resonances may be achieved by coherent vertical betatron oscillations. The problem is that the betatron decoherence of the kicked beam may dilute the emittance. The decoherence of coherent betatron motion depends on the tune shift with amplitude parameters, $\alpha_{xx}, \alpha_{xz}, \alpha_{zz}$, where $\alpha_{xx} = \frac{\partial \nu_x}{\partial J_x}$, $\alpha_{xz} = \frac{\partial \nu_x}{\partial J_z}$, $\alpha_{zz} = \frac{\partial \nu_z}{\partial J_z}$. This experiment can be performed with an unpolarized beam by measuring the emittance dilution due to coherent betatron kicks. Another important issue was the possible horizontal coherent betatron oscillation resulting from linear coupling.
4. At low energy, the 4.7 Tm solenoid corresponds to a powerful snake. It is interesting to check the adiabatic turn on/off the snake. Previous calculations indicate that the AGS lattice can tolerate up to about 60% partial snake without a skew quadrupole correction. Therefore, adiabatic turn on/off of snake is possible at low energy. We will study beam stability with the varying strengths of the solenoid snake with/without skew quadrupole correction. With the possibility of 50° spin rotation angle at $G\gamma \approx 9$, we may be able to study the long standing problem of the second order snake resonance located at $\nu_z = 8.93$. Such test requires polarized beam. This experiment will be scheduled in 1995.

4. Conclusion

In conclusion, we have accomplished most of our goals of our DOE grant DE-FG02-92ER40747 during the funding period of Aug. 15, 1992 to Aug. 14, 1994. We have successfully constructed a solenoid of 4.7 Tm. The solenoidal partial snake was successfully tested. We have also renovated and rebuilt polarized source, polarimeters and data acquisition systems. Our polarized beam experiment is scheduled from April 4 to April

8, 1994. We have also performed theoretical calculations on the beam dynamics associated with polarized beam acceleration. These studies will be tested in our experiments. Our full-scope polarized beam acceleration experiments with tune jump quadrupoles will be ready in 1995. The availability of polarized beam for RHIC collider experiment will enhance greatly the potential of RHIC.

References

1. Ya.S. Derbenev and A.M.Kondratenko, Part. Accel., 8, 115 (1978).
2. T. Roser, " Properties of Partially Excited Siberian Snakes", AIP Conf. Proc. No. 187, p 1442 (1988). S.Y. Lee, E.D. Courant and S. Tepikian, " The spin tune of a partially excited snake", AIP Conf. Proc. No. 145, (1985).
3. H. Huang, et al., Proc. of EPAC, p.726 (1992) [DOE/ER/40747-1]; H. Huang, S.Y. Lee and L. Ratner, "Minimizing Linear coupling in the AGS with skew quads", AGS Technote AD-376 (1993) [DOE/ER/40747-2]. T. Roser et al., Proc. of the 10th High Energy Spin Conference Nagoya, Japan, Nov 14 (1992).
4. S.Y. Lee, "Spin Depolarization Mechanism in synchrotrons with snakes", Phys. Rev. E47, 3631 (1993); [DOE/ER/40747-09] "Even order snake resonances", Proc. of IEEE PAC, May, Washington (1993) [DOE/ER/40747-07].
5. H. Huang, S.Y. Lee, and L. Ratner, "Transfer matrices for spin tensor polarization with and without snakes", IEEE PAC, May, Washington (1993) [DOE/ER/40747-5].
6. S.Y. Lee, Lecture Note at the RCNP Kikuchi school, Osaka Japan Nov. (1992) [DOE/ER/40747-4].

**DATE
FILMED**

8/24/94

END