ESTABLISHMENT OF AN OIL AND GAS DATABASE FOR INCREASED RECOVERY AND CHARACTERIZATION OF OIL AND GAS CARBONATE RESERVOIR HETEROGENEITY

DOE Contract No. DE-FG22-89BC14425

APPENDIX 1 Draft Topical Report on Subtasks 2 and 3 Volume 1 of 4

PREPARED BY
David C. Kopaska-Merkel, Henry E. Moore, Jr., Steven D. Mann,
and Douglas R. Hall

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

FOR

United States Department of Energy
Bartlesville Project Office
Bartlesville, Oklahoma

Project Manager: Chandra M. Nautiyal

June 1992

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

CONTENTS

	Page
Field codes for Smackover oil fields in southwest Alabama	xxvi
Explanation of map symbols and abbreviations	xxvii
Key to core logs	xxviii
Abbreviations	xxix
Volume I	
Appleton field, Smackover oil pool, Escambia County, Alabama	1
Barnett field, Smackover oil pool, Conecuh and Escambia Counties, Alabama	12
Barlow Bend field, Smackover oil pool, Clarke and Monroe Counties, Alabama	25
Barrytown field, Smackover oil pool, Choctaw County, Alabama	33
Big Escambia Creek field, Smackover gas and condensate pool, Escambia County, Alabama	45
Blacksher field, Smackover oil pool, Baldwin County, Alabama	58
Broken Leg Creek field, Smackover oil pool, Escambia County, Alabama	72
Bucatunna Creek field, Smackover oil pool, Choctaw County, Alabama	80
Burnt Corn Creek field, Smackover oil pool, Escambia County, Alabama	95
Chappell Hill field, Smackover oil pool, Choctaw County, Alabama	105
Chatom field, Smackover gas and condensate pool, Washington County, Alabama	120
Choctaw Ridge field, Smackover oil pool, Choctaw County, Alabama	134
Churchula field, Smackover gas and condensate pool, Mobile County, Alabama	145
Cold Creek field, Smackover oil pool, Mobile County, Alabama	158
Copeland field, Smackover gas and condensate pool, Washington County, Alabama ..	166
Crosbys Creek field, Smackover gas and condensate pool, Washington County, Alabama	174
East Barnett field, Smackover oil pool, Conecuh and Escambia Counties, Alabama	185
Volume II	
East Huxford field, Smackover oil pool, Escambia County, Alabama	195
Fanny Church field, Smackover oil pool, Escambia County, Alabama	202
Gin Creek field, Smackover oil pool, Choctaw County, Alabama	210
Gulf Crest field, Smackover oil pool, Mobile County, Alabama	224
Hanberry Church field, Smackover oil pool, Escambia County, Alabama	231
Hatter's Pond field, Smackover gas and condensate pool, Mobile County, Alabama	238
Healing Springs field, Smackover gas and condensate pool, Washington County, Alabama	251
Huxford field, Smackover oil pool, Escambia County, Alabama	264
Little Escambia Creek field, Smackover oil pool, Escambia County, Alabama	277
Little Mill Creek field, Smackover oil pool, Choctaw County, Alabama	294
Little River field, Smackover oil pool, Baldwin and Monroe Counties, Alabama	302
Little Rock field, Smackover gas and condensate pool, Escambia County, Alabama	316
Lovetts Creek field, Smackover oil pool, Clarke and Monroe Counties, Alabama	323
Melvin field, Smackover oil pool, Choctaw County, Alabama	340
Mill Creek field, Smackover oil pool, Choctaw County, Alabama	348
Mineola field, Smackover oil pool, Conecuh County, Alabama	356
Movico field, Smackover oil pool, Mobile and Baldwin Counties, Alabama	363
North Choctaw Ridge field, Smackover oil pool, Choctaw County, Alabama	374
Volume III	
North Smiths Church field, Smackover oil pool, Escambia County, Alabama	388
North Wallers Creek field, Smackover oil pool, Monroe County, Alabama	395
Northeast Barnett field, Smackover oil pool, Conecuh County, Alabama	408
Northwest Range field, Smackover oil pool, Conecuh County, Alabama	415
Pace Creek field, Smackover oil pool, Clarke County, Alabama	422

CONTENTS—CONTINUED

	Page
Palmers Crossroads field, Smackover oil pool, Monroe County, Alabama	429
Perdido field, Smackover oil pool, Escambia and Baldwin Counties, Alabama	437
Puss Cuss Creek field, Smackover oil pool, Choctaw County, Alabama	449
Red Creek field, Smackover gas and condensate pool, Washington County, Alabama	457
Robinson Creek field, Smackover oil pool, Escambia County, Alabama	465
Silas field, Smackover oil pool, Choctaw County, Alabama	472
Sizemore Creek field, Smackover gas and condensate pool, Escambia County, Alabama	486
Smiths Church field, Smackover gas and condensate pool, Escambia County, Alabama	497
South Burnt Corn Creek field, Smackover oil pool, Escambia County, Alabama	504
South Cold Creek field, Smackover oil pool, Mobile County, Alabama	511
South Vocation field, Smackover oil pool, Monroe County, Alabama	519
South Wild Fork Creek field, Smackover gas and condensate pool, Escambia County, Alabama	527
South Womack Hill field, Smackover oil pool, Choctaw and Clarke Counties, Alabama	534
Southeast Chatom field, Smackover gas and condensate pool, Washington County, Alabama	541
Southwest Barrytown field, Smackover oil pool, Choctaw County, Alabama	549
Souwilpa Creek field, Smackover gas and condensate pool, Choctaw County, Alabama	557
Volume IV	
Stave Creek field, Smackover oil pool, Clarke County, Alabama	564
Sugar Ridge field, Smackover oil pool, Choctaw County, Alabama	576
Toxey field, Smackover oil pool, Choctaw County, Alabama	589
Turkey Creek field, Smackover oil pool, Choctaw and Clarke Counties, Alabama	603
Turnerville field, Smackover oil pool, Mobile County, Alabama	611
Uriah field, Smackover oil pool, Monroe County, Alabama	623
Vocation field, Smackover oil pool, Monroe County, Alabama	638
Wallace field, Smackover oil pool, Escambia County, Alabama	650
Waller Creek field, Smackover oil pool, Monroe County, Alabama	657
West Appleton field, Smackover oil pool, Escambia County, Alabama	665
West Barrytown field, Smackover oil pool, Choctaw County, Alabama	672
West Bend field, Smackover oil pool, Choctaw and Clarke Counties, Alabama	686
West Okatuppa Creek field, Smackover oil pool, Choctaw County, Alabama	699
Wild Fork Creek field, Smackover oil pool, Escambia County, Alabama	706
Wimberly Field, Smackover oil pool, Choctaw County, Alabama	714
Womack Hill field, Smackover oil pool, Choctaw and Clarke Counties, Alabama	722
Zion Chapel field, Smackover oil pool, Choctaw County, Alabama	733

ILLUSTRATIONS

	Page
APT-1. Index map of Appleton field, Alabama	2
APT-2. Structure contour map on top of Smackover Formation in Appleton field, Alabama	3
APT-3. Well log, McMillan 2-14 No. 1 well, Permit No. 3854, Appleton field, Alabama	4
APT-4. Structural cross section of the Appleton field, Alabama	5
APT-5. Graph of production history of the Appleton field, Alabama	6
APT-6. Porosity vs. natural log permeability plot of described core from D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama.	7
APT-7. Detailed core log, D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama	9
APT-8. Paragenetic sequence, D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama	11
BAR-1. Index map of Barnett field, Alabama	13
BAR-2. Structure contour map on top of Smackover Formation in Barnett field, Alabama	14
BAR-3. Well log correlated to lithology, porosity, and permeability, Alger Sullivan "C" No. 1 well, Permit No. 1827, Barnett field, Alabama	15
BAR-4. Structural cross section of the Barnett field, Alabama	16
BAR-5. Graph of production history of the Barnett field, Alabama	17
BAR-6. Porosity vs. natural log permeability plots, Barnett field, Alabama	18
BAR-7. Detailed core log, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama	20
BAR-8. Porosity vs. natural log permeability plot, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama	22
BAR-9. Paragenetic sequence, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama	23
BAR-10. Ternary diagram of pore types, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama	24
BB-1. Index map of Barlow Bend field, Alabama	26
BB-2. Structure contour map on top of Smackover Formation in Barlow Bend field, Alabama	27
BB-3. Well log, Henry Gibby Unit 7-15 No. 1 well, Permit No. 4905, Barlow Bend field, Alabama	28
BB-4. Structural cross section of the Barlow Bend field, Alabama	29
BB-5. Graph of production history of the Barlow Bend field, Alabama	30
BB-6. Porosity vs. natural log permeability plot of John R. Mattmuller et al. No. 1 well, Permit No. 2708, Barlow Bend field, Alabama	31
BTN-1. Index map of Barrytown field, Alabama	34
BTN-2. Structure contour map on top of Smackover Formation in Barrytown field, Alabama	35
BTN-3. Well log correlated to lithology and permeability, McIlwain 11-10 well, Permit No. 1818, Barrytown field, Alabama	36
BTN-4. Structural cross section of the Barrytown field, Alabama	37
BTN-5. Graph of production history of the Barrytown field, Alabama	38
BTN-6. Porosity vs. natural log permeability plots, Barrytown field, Alabama	39
BTN-7. Detailed core log, McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama	41

ILLUSTRATIONS—CONTINUED

	Page
BTN-8. Porosity vs. natural log permeability plot for lithofacies 1, McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama	43
BTN-9. Paragenetic sequence, McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama	44
BEC-1. Index map of Big Escambia Creek field, Alabama	46
BEC-2. Structure contour map on top of Smackover Formation in Big Escambia Creek field, Alabama	47
BEC-3. Well log correlated to lithology, porosity, and permeability, Smith 25-7 No. 1 well, Permit 4727, Big Escambia Creek field, Alabama	48
BEC-4. Structural cross section of the Big Escambia Creek field, Alabama	49
BEC-5. Graph of production history of the Big Escambia Creek field, Alabama	50
BEC-6. Porosity vs. natural log permeability plots, Big Escambia Creek field, Alabama	51
BEC-7. Detailed core log, Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama	53
BEC-8. Porosity vs. natural log permeability plots for three lithofacies, Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama	56
BEC-9. Paragenetic sequence, Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama	57
BLS-1. Index map of Blacksher field, Alabama	59
BLS-2. Structure contour map on top of Smackover Formation in Blacksher field, Alabama	60
BLS-3. Well log correlated to lithology, porosity, and permeability, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	61
BLS-4. Structural cross section of the Blacksher field, Alabama	62
BLS-5. Graph of production history of the Blacksher field, Alabama	63
BLS-6. Porosity vs. natural log permeability plots, Blacksher field, Alabama	64
BLS-7. Detailed core log, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	66
BLS-8. Porosity vs. natural log permeability plots for two lithofacies, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	69
BLS-9. Paragenetic sequence, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	70
BLS-10. Ternary diagram of pore types, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	71
BLC-1. Index map of Broken Leg Creek field, Alabama	73
BLC-2. Structure contour map on top of Smackover Formation in Broken Leg Creek field, Alabama	74
BLC-3. Well log, Hawkins 7-16 No. 1 well Permit 6153-B, Broken Leg Creek field, Alabama	75
BLC-4. Structural cross section of the Broken Leg Creek field, Alabama	76
BLC-5. Graph of production history of the Broken Leg Creek field, Alabama	77
BLC-6. Porosity vs. natural log permeability plot of one core, Broken Leg Creek field, Alabama	78
BUC-1. Index map of Bucatunna Creek field, Alabama	81
BUC-2. Structure contour map on top of Smackover Formation in Bucatunna Creek field, Alabama	82
BUC-3. Well log, A. G. Harrell No. 1 well, Permit No. 2470, Bucatunna Creek field, Alabama	83
BUC-4. Structural cross section of the Bucatunna Creek field, Alabama	84
BUC-5. Graph of production history of the Bucatunna Creek field, Alabama	85

ILLUSTRATIONS—CONTINUED

	Page
BUC-6. Porosity vs. natural log permeability plots, Bucatunna Creek field, Alabama	86
BUC-7. Detailed core log, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama	88
BUC-8. Porosity vs. natural log permeability plot for lithofacies 1, Johnson Unit 14-14 No. 1 well, Permit No. 2753-B, Bucatunna Creek field, Alabama	92
BUC-9. Paragenetic sequence, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama	93
BUC-10. Ternary diagram of pore types, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama	94
BCC-1. Index map of Burnt Corn Creek field, Alabama	96
BCC-2. Structure contour map on top of Smackover Formation in Burnt Corn Creek field, Alabama	97
BCC-3. Well log, A. J. Smith 3-14 No. 1 well, Permit No. 4744, Burnt Corn Creek field, Alabama	98
BCC-4. Structural cross section of the Burnt Corn Creek field, Alabama	99
BCC-5. Graph of production history of the Burnt Corn Creek field, Alabama	100
BCC-6. Detailed core log, Hooper Mathews 4-8 No. 1 well, Permit No. 2377, Burnt Corn Creek field, Alabama	102
BCC-7. Paragenetic sequence, Hooper Mathews 4-8 No. 1 well, Permit No. 2377, Burnt Corn Creek field, Alabama	104
CH-1. Index map of Chappell Hill field, Alabama	106
CH-2. Structure contour map on top of Smackover Formation in Chappell Hill field, Alabama	107
CH-3. Well log correlated to lithology, porosity, and permeability, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	108
CH-4. Structural cross section of the Chappell Hill field, Alabama	109
CH-5. Graph of production history of the Chappell Hill field, Alabama	110
CH-6. Porosity vs. natural log permeability plots, Chappell Hill field, Alabama	111
CH-7. Detailed core log, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	113
CH-8. Porosity vs. natural log permeability plots for three lithofacies, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	117
CH-9. Paragenetic sequence, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	118
CH-10. Ternary diagram of pore types, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	119
CTM-1. Index map of Chatom field, Alabama	121
CTM-2. Structure contour map on top of Smackover Formation in Chatom field, Alabama	122
CTM-3. Well log correlated to lithology, porosity, and permeability, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	123
CTM-4. Structural cross section of the Chatom field, Alabama	124
CTM-5. Graph of production history of the Chatom field, Alabama	125
CTM-6. Porosity vs. natural log permeability plots, Chatom field, Alabama	126
CTM-7. Detailed core log, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	128
CTM-8. Porosity vs. natural log permeability plots for two lithofacies, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	131
CTM-9. Paragenetic sequence, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	132

ILLUSTRATIONS—CONTINUED

	Page
CTM-10. Ternary diagram of pore types, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	133
CHR-1. Index map of Choctaw Ridge field, Alabama	135
CHR-2. Structure contour map on top of Smackover Formation in Choctaw Ridge field, Alabama	136
CHR-3. Well log correlated to lithology, Rentz-Trice Unit 35-1 well, Permit No. 1427, Choctaw Ridge field, Alabama	137
CHR-4. Structural cross section of the Choctaw Ridge field, Alabama	138
CHR-5. Graph of production history of the Choctaw Ridge field, Alabama	139
CHR-6. Porosity vs. natural log permeability plot of five cores combined, Choctaw Ridge field, Alabama	140
CHR-7. Detailed core log, Rentz-Trice Unit 35-1 well, Permit No. 1427, Choctaw Ridge field, Alabama	142
CHR-8. Paragenetic sequence, Rentz-Trice Unit 35-1 No. 1 well, Permit No. 1427, Choctaw Ridge field, Alabama	144
CHU-1. Index map of Chunchula field, Alabama	146
CHU-2. Structure contour map on top of Smackover Formation in Chunchula field, Alabama	147
CHU-3. Well log correlated to lithology, porosity, and permeability, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama	148
CHU-4. Structural cross section of the Chunchula field, Alabama	149
CHU-5. Graph of production history of the Chunchula field, Alabama	150
CHU-6. Porosity vs. natural log permeability plots, Chunchula field, Alabama	151
CHU-7. Detailed core log, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama	153
CHU-8. Porosity vs. natural log permeability plot for lithofacies 1, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama	155
CHU-9. Paragenetic sequence, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama	156
CHU-10. Ternary diagram of pore types, samples from several wells, Chunchula field, Alabama	157
CC-1. Index map of Cold Creek field, Alabama	159
CC-2. Structure contour map on top of Smackover Formation in Cold Creek field, Alabama	160
CC-3. Well log correlated to porosity and permeability, C. P. Newman et al. 8-16 No. 1 well, Permit No. 3027, Cold Creek field, Alabama	161
CC-4. Structural cross section of the Cold Creek field, Alabama	162
CC-5. Graph of production history of the Cold Creek field, Alabama	163
CC-6. Porosity vs. natural log permeability plots, Cold Creek field, Alabama	164
CPL-1. Index map of Copeland field, Alabama	167
CPL-2. Structure contour map on top of Smackover Formation in Copeland field, Alabama	168
CPL-3. Well log, Edith Burch 30-13 No. 1 well, Permit No. 3247, Copeland field, Alabama	169
CPL-4. Structural cross section of the Copeland field, Alabama	170
CPL-5. Graph of production history of the Copeland field, Alabama	171
CPL-6. Porosity vs. natural log permeability plot of one core, Copeland field, Alabama	172
CRC-1. Index map of Crosbys Creek field, Alabama	175
CRC-2. Structure contour map on top of Smackover Formation in Crosbys Creek field, Alabama	176

ILLUSTRATIONS—CONTINUED

	Page
CRC-3. Well log, Sarah Middleton 34-11 No. 1 well, Permit No. 5001, Crosbys Creek field, Alabama	177
CRC-4. Structural cross section of the Crosbys Creek field, Alabama	178
CRC-5. Graph of production history of the Crosbys Creek field, Alabama	179
CRC-6. Porosity vs. natural log permeability plot of two cores combined, Crosbys Creek field, Alabama	180
CRC-7. Detailed core log, Darly Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama	182
CRC-8. Paragenetic sequence, Darly Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama	184
EB-1. Index map of East Barnett field, Alabama	186
EB-2. Structure contour map on top of Smackover Formation in East Barnett field, Alabama	187
EB-3. Well log correlated to lithology, Scott Paper Co. et al. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama	188
EB-4. Structural cross section of the East Barnett field, Alabama	189
EB-5. Graph of production history of the East Barnett field, Alabama	190
EB-6. Detailed core log, Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama	192
EB-7. Paragenetic sequence, Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama	194
EH-1. Index map of East Huxford field, Alabama	196
EH-2. Structure contour map on top of Smackover Formation in East Huxford field, Alabama	197
EH-3. Well log, A.T.I.C. 18-13 No. 1 well, Permit No. 6804, East Huxford field, Alabama	198
EH-4. Structural cross section of the East Huxford field, Alabama	199
EH-5. Graph of production history of the East Huxford field, Alabama	200
FC-1. Index map of Fanny Church field, Alabama	203
FC-2. Structure contour map on top of Smackover Formation in Fanny Church field, Alabama	204
FC-3. Well log, D. J. Steely 36-2 well, Permit No. 1869, Fanny Church field, Alabama	205
FC-4. Structural cross section of the Fanny Church field, Alabama	206
FC-5. Graph of production history of the Fanny Church field, Alabama	207
FC-6. Porosity vs. natural log permeability plot of 19 cores combined, Fanny Church field, Alabama	208
GC-1. Index map of Gin Creek field, Alabama	211
GC-2. Structure contour map on top of Smackover Formation in Gin Creek field, Alabama	212
GC-3. Well log, Choctaw Lumber Co. 31-5 No. 1 well, Permit No. 4546, Gin Creek field, Alabama	213
GC-4. Structural cross section of the Gin Creek field, Alabama	214
GC-5. Graph of production history of the Gin Creek field, Alabama	215
GC-6. Porosity vs. natural log permeability plot of described core for Bolinger Heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	216
GC-7. Detailed core log, Bolinger Heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	218
GC-8. Porosity vs. natural log permeability plots for three lithofacies, Bolinger Heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	221

ILLUSTRATIONS—CONTINUED

	Page
GC-9. Paragenetic sequence, Bolinger Heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	222
GC-10. Ternary diagram of pore types, Bolinger heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	223
GUC-1. Index map of Gulf Crest field, Alabama	225
GUC-2. Structure contour map on top of Smackover Formation in Gulf Crest field, Alabama	226
GUC-3. Well log, M. H. Brown Trust 5-2 No. 1 well, Permit No. 5226, Gulf Crest field, Alabama	227
GUC-4. Structural cross section of the Gulf Crest field, Alabama	228
GUC-5. Graph of production history of the Gulf Crest field, Alabama	229
HC-1. Index map of Hanberry Church field, Alabama	232
HC-2. Structure contour map on top of Smackover Formation in Hanberry Church field, Alabama	233
HC-3. Well log, Huxford 34-2 No. 1 well, Permit No. 5178, Hanberry Church field, Alabama	234
HC-4. Structural cross section of the Hanberry Church field, Alabama	235
HC-5. Graph of production history of the Hanberry Church field, Alabama	236
HP-1. Index map of Hatter's Pond field, Alabama	239
HP-2. Structure contour map on top of Smackover Formation in Hatter's Pond field, Alabama	240
HP-3. Well log correlated to lithology, porosity, and permeability, Hatter's Pond Unit 10-3 No. 1 well, Permit No. 6846, Hatter's Pond field, Alabama	241
HP-4. Structural cross section of the Hatter's Pond field, Alabama	242
HP-5. Graph of production history of the Hatter's Pond field, Alabama	243
HP-6. Porosity vs. natural log permeability plot, Hatter's Pond field, Alabama	244
HP-7. Detailed core log, Hatter's Pond Unit 10-3 No. 1 well, Permit No. 6846, Hatter's Pond field, Alabama	246
HP-8. Porosity vs. natural log permeability plots for three lithofacies, Hatter's Pond Unit 10-3 No. 1 well, Permit No. 6846, Hatter's Pond field, Alabama	249
HP-9. Paragenetic sequence, Hatter's Pond Unit 10-3 No. 1 well, Permit No. 6846, Hatter's Pond field, Alabama	250
HS-1. Index map of Healing Springs field, Alabama	252
HS-2. Structure contour map on top of Smackover Formation in Healing Springs field, Alabama	253
HS-3. Well log correlated to porosity and permeability, W. J. Britton No. 1 well, Permit No. 2478, Healing Springs field, Alabama	254
HS-4. Structural cross section of the Healing Springs field, Alabama	255
HS-5. Graph of production history of the Healing Springs field, Alabama	256
HS-6. Porosity vs. natural log permeability plot, Healing Springs field, Alabama	257
HS-7. Detailed core log, W. J. Britton No. 1 well, Permit No. 2478, Healing Springs field, Alabama	259
HS-8. Porosity vs. natural log permeability plots for two lithofacies, W. J. Britton No. 1 well, Permit No. 2478, Healing Springs field, Alabama	262
HS-9. Paragenetic sequence, W. J. Britton No. 1 well, Permit No. 2478, Healing Springs field, Alabama	263
HUX-1. Index map of Huxford field, Alabama	265
HUX-2. Structure contour map on top of Smackover Formation in Huxford field, Alabama	266

ILLUSTRATIONS—CONTINUED

	Page
HUX-3. Well log correlated to lithology, porosity, and permeability, M. H. Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	267
HUX-4. Structural cross section of the Huxford field, Alabama	268
HUX-5. Graph of production history of the Huxford field, Alabama	269
HUX-6. Porosity vs. natural log permeability plots, Huxford field, Alabama	270
HUX-7. Detailed core log, Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	272
HUX-8. Porosity vs. natural log permeability plot for lithofacies 1, M. H. Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	274
HUX-9. Paragenetic sequence, M. H. Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	275
HUX-10. Ternary diagram of pore types, Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	276
LEC-1. Index map of Little Escambia Creek field, Alabama	278
LEC-2. Structure contour map on top of Smackover Formation in Little Escambia Creek field, Alabama	279
LEC-3. Well log correlated to lithology, porosity, and permeability, T. R. Miller Mill Unit 32-3 well, Permit No. 1654, Little Escambia Creek field, Alabama	280
LEC-4. Structural cross section of the Little Escambia Creek field, Alabama	281
LEC-5. Graph of production history of the Little Escambia Creek field, Alabama	282
LEC-6. Porosity vs. natural log permeability plots, Little Escambia Creek field, Alabama	283
LEC-7. Detailed core log, T. R. Miller Mill Unit 32-3 well, Permit No. 1654, Little Escambia Creek field, Alabama	285
LEC-8. Porosity vs. natural log permeability plots for three lithofacies, T. R. Miller Mill Unit 32-3 well, Permit No. 1654, Little Escambia Creek field, Alabama	291
LEC-9. Paragenetic sequence, T. R. Miller Mill Unit 32-3 well, Permit No. 1654, Little Escambia Creek field, Alabama	293
LMC-1. Index map of Little Mill Creek field, Alabama	295
LMC-2. Structure contour map on top of Smackover Formation in Little Mill Creek field, Alabama	296
LMC-3. Well log, Tims 8-2 No. 1 well, Permit No. 2463-B, Little Mill Creek field, Alabama	297
LMC-4. Structural cross section of the Little Mill Creek field, Alabama	298
LMC-5. Graph of production history of the Little Mill Creek field, Alabama	299
LMC-6. Porosity vs. natural log permeability plot of four cores combined, Little Mill Creek field, Alabama	300
LR-1. Index map of Little River field, Alabama	303
LR-2. Structure contour map on top of Smackover Formation in Little River field, Alabama	304
LR-3. Well log correlated to lithology, porosity, and permeability, J. E. Robinson et al. 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	305
LR-4. Structural cross section of the Little River field, Alabama	306
LR-5. Graph of production history of the Little River field, Alabama	307
LR-6. Porosity vs. natural log permeability plots, Little River field, Alabama	308
LR-7. Detailed core log, J. E. Robinson et al. 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	310

ILLUSTRATIONS—CONTINUED

	Page
LR-8. Porosity vs. natural log permeability plot for lithofacies 1, Robinson 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	313
LR-9. Paragenetic sequence, Robinson et al. 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	314
LR-10. Ternary diagram of pore types, Robinson et al. 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	315
LRK-1. Index map of Little Rock field, Alabama	317
LRK-2. Structure contour map on top of Smackover Formation in Little Rock field, Alabama	318
LRK-3. Well log, A.T.I.C. 31-6 No. 1 well, Permit No. 4303-B, Little Rock field, Alabama	319
LRK-4. Structural cross section of the Little Rock field, Alabama	320
LRK-5. Graph of production history of the Little Rock field, Alabama	321
LOV-1. Index map of Lovetts Creek field, Alabama	324
LOV-2. Structure contour map on top of Smackover Formation in Lovetts Creek field, Alabama	325
LOV-3. Well log correlated to porosity and permeability, W. E. Broughton 11-6 No. 1 well, Permit No. 3347, Lovetts Creek field, Alabama	326
LOV-4. Structural cross section of the Lovetts Creek field, Alabama	327
LOV-5. Graph of production history of the Lovetts Creek field, Alabama	328
LOV-6. Porosity vs. natural log permeability plots, Lovetts Creek field, Alabama	329
LOV-7. Detailed core log, W. M. Threat 11-10 No. 1 well, Permit No. 3535, Lovetts Creek field, Alabama	331
LOV-8. Porosity vs. natural log permeability plots for two lithofacies, W. M. Threat 11-10 No. 1 well, Permit No. 3535, Lovetts Creek field, Alabama	336
LOV-9. Paragenetic sequence, W. M. Threat 11-10 No. 1 well, Permit No. 3535, Lovetts Creek field, Alabama	338
LOV-10. Ternary diagram of pore types, W. M. Threat 11-10 No. 1 well, Permit No. 3535, Lovetts Creek field, Alabama	339
MEL-1. Index map of Melvin field, Alabama	341
MEL-2. Structure contour map on top of Smackover Formation in Melvin field, Alabama	342
MEL-3. Well log, Charles M. Land 2-4 No. 1 well, Permit No. 2319, Melvin field, Alabama	343
MEL-4. Structural cross section of the Melvin field, Alabama	344
MEL-5. Graph of production history of the Melvin field, Alabama	345
MEL-6. Porosity vs. natural log permeability plot of one core, Melvin field, Alabama	346
MC-1. Index map of Mill Creek field, Alabama	349
MC-2. Structure contour map on top of Smackover Formation in Mill Creek field, Alabama	350
MC-3. Well log, Trice Unit 6-6 No. 1 well, Permit No. 2145, Mill Creek field, Alabama	351
MC-4. Structural cross section of the Mill Creek field, Alabama	352
MC-5. Graph of production history of the Mill Creek field, Alabama	353
MC-6. Porosity vs. natural log permeability plot of five cores combined, Mill Creek field, Alabama	354
MIN-1. Index map of Mineola field, Alabama	357
MIN-2. Structure contour map on top of Smackover Formation in Mineola field, Alabama	358

ILLUSTRATIONS—CONTINUED

	Page
MIN-3. Well log, Langford 18-14 No. 1 well, Permit No. 8748, Mineola field, Alabama	359
MIN-4. Structural cross section of the Mineola field, Alabama	360
MIN-5. Graph of production history of the Mineola field, Alabama	361
MOV-1. Index map of Movico field, Alabama	364
MOV-2. Structure contour map on top of Smackover Formation in Movico field, Alabama	365
MOV-3. Well log correlated to porosity and permeability, Janice R. Sullivan et al. 9-4 No. 1 well, Permit No. 2126, Movico field, Alabama	366
MOV-4. Structural cross section of the Movico field, Alabama	367
MOV-5. Graph of production history of the Movico field, Alabama	368
MOV-6. Porosity vs. natural log permeability plot of two cores combined, Movico field, Alabama	369
MOV-7. Detailed core log, Sullivan et al. 9-4 No. 1 well, Permit No. 2126, Movico field, Alabama	371
MOV-8. Paragenetic sequence, Sullivan et al. 9-4 No. 1 well, Permit No. 2621, Movico field, Alabama	373
NCR-1. Index map of North Choctaw Ridge field, Alabama	375
NCR-2. Structure contour map on top of Smackover Formation in North Choctaw Ridge field, Alabama	376
NCR-3. Well log correlated to lithology, porosity, and permeability, Oates Unit 29-6 No. 1 well, Permit No. 1878, North Choctaw Ridge field, Alabama	377
NCR-4. Structural cross section of the North Choctaw Ridge field, Alabama	378
NCR-5. Graph of production history of the North Choctaw Ridge field, Alabama	379
NCR-6. Porosity vs. natural log permeability plots, North Choctaw Ridge field, Alabama	380
NCR-7. Detailed core log, Oates Unit 29-6 No. 1 well, Permit No. 1878 North Choctaw Ridge field, Alabama	382
NCR-8. Porosity vs. natural log permeability plots for three lithofacies, Oates Unit 29-6 No. 1 well, Permit No. 1878, North Choctaw Ridge field, Alabama	385
NCR-9. Paragenetic sequence, Oates Unit 29-6 No. 1 well, Permit No. 1878, North Choctaw Ridge field, Alabama	386
NCR-10. Ternary diagram of pore types, Oates Unit 29-6 No. 1 well, Permit No. 1878, North Choctaw Ridge field, Alabama	387
NSC-1. Index map of North Smiths Church field, Alabama	389
NSC-2. Structure contour map on top of Smackover Formation in North Smiths Church field, Alabama	390
NSC-3. Well log, Edge-Wefel Trust 24-5 No. 1 well, Permit No. 6943, North Smiths Church field, Alabama	391
NSC-4. Structural cross section of the North Smiths Church field, Alabama	392
NSC-5. Graph of production history of the North Smiths Church field, Alabama	393
NWC-1. Index map of North Wallers Creek field, Alabama	396
NWC-2. Structure contour map on top of Smackover Formation in North Wallers Creek field, Alabama	397
NWC-3. Well log correlated to lithology, porosity, and permeability, Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	398
NWC-4. Structural cross section of the North Wallers Creek field, Alabama	399
NWC-5. Graph of production history of the North Wallers Creek field, Alabama	400
NWC-6. Porosity vs. natural log permeability plot of described core for Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	401

ILLUSTRATIONS—CONTINUED

	Page
NWC-7. Detailed core log, Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	403
NWC-8. Porosity vs. natural log permeability plot for lithofacies 1, Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	406
NWC-9. Paragenetic sequence, Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	407
NEB-1. Index map of Northeast Barnett field, Alabama	409
NEB-2. Structure contour map on top of Smackover Formation in Northeast Barnett field, Alabama	410
NEB-3. Well log, Scott Paper Co. 25-13 No. 1 well, Permit No. 7214, Northeast Barnett field, Alabama	411
NEB-4. Structural cross section of the Northeast Barnett field, Alabama	412
NEB-5. Graph of production history of the Northeast Barnett field, Alabama	413
NWR-1. Index map of Northwest Range field, Alabama	416
NWR-2. Structure contour map on top of Smackover Formation in Northwest Range field, Alabama	417
NWR-3. Well log, Paramount A.T.I.C. 7-13 No. 1 well, Permit No. 5930, Northwest Range field, Alabama	418
NWR-4. Structural cross section of the Northwest Range field, Alabama	419
NWR-5. Graph of production history of the Northwest Range field, Alabama	420
PC-1. Index map of Pace Creek field, Alabama	423
PC-2. Structure contour map on top of Smackover Formation in Pace Creek field, Alabama	424
PC-3. Well log, Mrs. E. V. Pelham 16-2 No. 1 well, Permit No. 5058, Pace Creek field, Alabama	425
PC-4. Structural cross section of the Pace Creek field, Alabama	426
PC-5. Graph of production history of the Pace Creek field, Alabama	427
PCR-1. Index map of Palmers Crossroads field, Alabama	430
PCR-2. Structure contour map on top of Smackover Formation in Palmers Crossroads field, Alabama	431
PCR-3. Well log, Blacksher 14-11 No. 1 well, Permit No. 5584, Palmers Crossroads field, Alabama	432
PCR-4. Structural cross section of the Palmers Crossroads field, Alabama	433
PCR-5. Graph of production history of the Palmers Crossroads field, Alabama	434
PCR-6. Porosity vs. natural log permeability plot for one core, Palmers Crossroads field, Alabama	435
PER-1. Index map of Perdido field, Alabama	438
PER-2. Structure contour map on top of Smackover Formation in Perdido field, Alabama	439
PER-3. Well log, F. H. Dees 32-11 No. 1 well, Permit No. 3628, Perdido field, Alabama	440
PER-4. Structural cross section of the Perdido field, Alabama	441
PER-5. Graph of production history of the Perdido field, Alabama	442
PER-6. Porosity vs. natural log permeability plots, Perdido field, Alabama	443
PER-7. Detailed core log, Wade Johnson 30-16 well, Permit No. 4696, Perdido field, Alabama	445
PER-8. Porosity vs. natural log permeability plot of lithofacies 1, Wade Johnson 30-16 well, Permit No. 4696, Perdido field, Alabama	447
PER-9. Paragenetic sequence, Wade Johnson 30-16 well, Permit No. 4696, Perdido field, Alabama	448
PCC-1. Index map of Puss Cuss Creek field, Alabama	450

ILLUSTRATIONS—CONTINUED

	Page
PCC-2. Structure contour map on top of Smackover Formation in Puss Cuss Creek field, Alabama	451
PCC-3. Well log, M. W. Smith Lumber Co. No. 1 well, Permit No. 2519, Puss Cuss Creek field, Alabama	452
PCC-4. Structural cross section of the Puss Cuss Creek field, Alabama	453
PCC-5. Graph of production history of the Puss Cuss Creek field, Alabama	454
PCC-6. Porosity vs. natural log permeability plot of one core, Puss Cuss Creek field, Alabama	455
RC-1. Index map of Red Creek field, Alabama	458
RC-2. Structure contour map on top of Smackover Formation in Red Creek field, Alabama	459
RC-3. Well log, L. E. Heinmiller No. 1 well, Permit No. 2721, Red Creek field, Alabama	460
RC-4. Structural cross section of the Red Creek field, Alabama	461
RC-5. Graph of production history of the Red Creek field, Alabama	462
RC-6. Porosity vs. natural log permeability plot of two cores combined, Red Creek field, Alabama	463
RCK-1. Index map of Robinson Creek field, Alabama	466
RCK-2. Structure contour map on top of Smackover Formation in Robinson Creek field, Alabama	467
RCK-3. Well log, A.T.I.C. 17-5 No. 1 well, Permit No. 8835-B, Robinson Creek field, Alabama	468
RCK-4. Structural cross section of the Robinson Creek field, Alabama	469
RCK-5. Graph of production history of the Robinson Creek field, Alabama	470
SIL-1. Index map of Silas field, Alabama	473
SIL-2. Structure contour map on top of Smackover Formation in Silas field, Alabama	474
SIL-3. Well log correlated to porosity and permeability, Chesnut unit 4-15 No. 1 well, Permit No. 2084, Silas field, Alabama	475
SIL-4. Structural cross section of the Silas field, Alabama	476
SIL-5. Graph of production history of the Silas field, Alabama	477
SIL-6. Porosity vs. natural log permeability plots, Silas field, Alabama	478
SIL-7. Detailed core log, Chesnut Unit 9-1 No. 1 well, Permit No. 2205, Silas field, Alabama	480
SIL-8. Porosity vs. natural log permeability plot for lithofacies 1, Chesnut Unit 9-1 No. 1 well, Permit No. 2205, Silas field, Alabama	483
SIL-9. Paragenetic sequence, Chesnut Unit 9-1 No. 1 well, Permit No. 2205, Silas field, Alabama	484
SIL-10. Ternary diagram of pore types, Chesnut Unit 9-1 No. 1 well, Permit No. 2205, Silas field, Alabama	485
SC-1. Index map of Sizemore Creek field, Alabama	487
SC-2. Structure contour map on top of Smackover Formation in Sizemore Creek field, Alabama	488
SC-3. Well log correlated to lithology, A.T.I.C. 33-1 No. 1 well, Permit No. 3988, Sizemore Creek field, Alabama	489
SC-4. Structural cross section of the Sizemore Creek field, Alabama	490
SC-5. Graph of production history of the Sizemore Creek field, Alabama	491
SC-6. Porosity vs. natural log permeability plot of two cores combined, Sizemore Creek field, Alabama	492
SC-7. Detailed core log, A.T.I.C. 33-1 No. 1 well, Permit No. 3988, Sizemore Creek field, Alabama	494

ILLUSTRATIONS—CONTINUED

	Page
SC-8. Paragenetic sequence, A.T.I.C. 33-1 No. 1 well, Permit No. 3988, Sizemore Creek field, Alabama	496
SCH-1. Index map of Smiths Church field, Alabama	498
SCH-2. Structure contour map on top of Smackover Formation in Smiths Church field, Alabama	499
SCH-3. Well log correlated to lithology, Kelly Unit 2-2 No. 1 well, Permit No. 5498, Smiths Church field, Alabama	500
SCH-4. Structural cross section of the Smiths Church field, Alabama	501
SCH-5. Graph of the production history of the Smiths Church field, Alabama	502
SBC-1. Index map of South Burnt Corn Creek field, Alabama	505
SBC-2. Structure contour map on top of Smackover Formation in South Burnt Corn Creek field, Alabama	506
SBC-3. Well log, Earl H. Weaver 22-15 No. 1 well, Permit No. 5272, South Burnt Corn Creek field, Alabama	507
SBC-4. Structural cross section of the South Burnt Corn Creek field, Alabama	508
SBC-5. Graph of production history of the South Burnt Corn Creek field, Alabama	509
SCC-1. Index map of South Cold Creek field, Alabama	512
SCC-2. Structure contour map on top of Smackover Formation in South Cold Creek field, Alabama	513
SCC-3. Well log, Mobile County Board of School Commissioners 16-12 No. 1 well, Permit No. 2071, South Cold Creek field, Alabama	514
SCC-4. Structural cross section of the South Cold Creek field, Alabama	515
SCC-5. Graph of production history of the South Cold Creek field, Alabama	516
SCC-6. Porosity vs. natural log permeability plot of one core, South Cold Creek field, Alabama	517
SV-1. Index map of South Vocation field, Alabama	520
SV-2. Structure contour map on top of Smackover Formation in South Vocation field, Alabama	521
SV-3. Well log, Container Corporation 34-16 No. 1A well, Permit No. 4225, South Vocation field, Alabama	522
SV-4. Structural cross section of the South Vocation field, Alabama	523
SV-5. Graph of production history of the South Vocation field, Alabama	524
SV-6. Porosity vs. natural log permeability plot of one core, South Vocation field, Alabama	525
SWF-1. Index map of South Wild Fork Creek field, Alabama	528
SWF-2. Structure contour map on top of Smackover Formation in South Wild Fork Creek field, Alabama	529
SWF-3. Well log, Huxford Estate 8-4 No. 1 well, Permit No. 5869, South Wild Fork Creek field, Alabama	530
SWF-4. Structural cross section of the South Wild Fork Creek field, Alabama	531
SWF-5. Graph of production history of the South Wild Fork Creek field, Alabama	532
SWH-1. Index map of South Womack Hill field, Alabama	535
SWH-2. Structure contour map on top of Smackover Formation in South Womack Hill field, Alabama	536
SWH-3. Well log, Womack Hill Unit 14-5 No. 1 well, Permit No. 4335-B, South Womack Hill field, Alabama	537
SWH-4. Structural cross section of the South Womack Hill field, Alabama	538
SWH-5. Graph of production history of the South Womack Hill field, Alabama	539
SEC-1. Index map of Southeast Chatom field, Alabama	542

ILLUSTRATIONS—CONTINUED

	Page
SEC-2. Structure contour map on top of Smackover Formation in Southeast Chatom field, Alabama	543
SEC-3. Well log, Scott "A" 33-1 No. 1 well, Permit No. 5100, Southeast Chatom field, Alabama	544
SEC-4. Structural cross section of the Southeast Chatom field, Alabama	545
SEC-5. Graph of production history of the Southeast Chatom field, Alabama	546
SEC-6. Porosity vs. natural log permeability plot of two cores combined, Southeast Chatom field, Alabama	547
SWB-1. Index map of Southwest Barrytown field, Alabama	550
SWB-2. Structure contour map on top of Smackover Formation in Southwest Barrytown field, Alabama	551
SWB-3. Well log, Choctaw Land 16-7 No. 1 well, Permit No. 4750, Southwest Barrytown field, Alabama	552
SWB-4. Structural cross section of the Southwest Barrytown field, Alabama	553
SWB-5. Graph of production history of the Southwest Barrytown field, Alabama	554
SWB-6. Porosity vs. natural log permeability plot of two cores combined, Southwest Barrytown field, Alabama	555
SOU-1. Index map of Souwilpa Creek field, Alabama	558
SOU-2. Structure contour map on top of Smackover Formation in Souwilpa Creek field, Alabama	559
SOU-3. Well log, Pinkerton 8-9 No. 1 well, Permit No. 2731, Souwilpa Creek field, Alabama	560
SOU-4. Structural cross section of the Souwilpa Creek field, Alabama	561
SOU-5. Graph of production history of the Souwilpa Creek field, Alabama	562
STV-1. Index map of Stave Creek field, Alabama	565
STV-2. Structure contour map on top of Smackover Formation in Stave Creek field, Alabama	566
STV-3. Well log correlated to lithology, porosity, and permeability, McCorquodale 25-1 No. 1 well, Permit No. 2718, Stave Creek field, Alabama	567
STV-4. Structural cross section of the Stave Creek field, Alabama	568
STV-5. Graph of production history of the Stave Creek field, Alabama	569
STV-6. Porosity vs. natural log permeability plots, Stave Creek field, Alabama	570
STV-7. Detailed core log, McCorquodale 25-1 No. 1 well, Permit No. 2718, Stave Creek field, Alabama	572
STV-8. Porosity vs. natural log permeability plot for one lithofacies, McCorquodale 25-1 No. 1 well, Permit No. 2718, Stave Creek field, Alabama	574
STV-9. Paragenetic sequence, McCorquodale 25-1 No. 1 well, Permit No. 2718, Stave Creek field, Alabama	575
SUG-1. Index map of Sugar Ridge field, Alabama	577
SUG-2. Structure contour map on top of Smackover Formation in Sugar Ridge field, Alabama	578
SUG-3. Well log correlated to lithology, porosity, and permeability, Lovett Unit 27-6 No. 1 well, Permit No. 1903, Sugar Ridge field, Alabama	579
SUG-4. Structural cross section of the Sugar Ridge field, Alabama	580
SUG-5. Graph of production history of the Sugar Ridge field, Alabama	581
SUG-6. Porosity vs. natural log permeability plots, Sugar Ridge field, Alabama	582
SUG-7. Detailed core log, Lovett 27-6 No. 1 well, Permit No. 1903, Sugar Ridge field, Alabama	584

ILLUSTRATIONS—CONTINUED

	Page
SUG-8. Porosity vs. natural log permeability plot for lithofacies 1, Lovett 27-6 No. 1 well, Permit No. 1903, Sugar Ridge field, Alabama	587
SUG-9. Paragenetic sequence, Lovett 27-6 No. 1 well, Permit No. 1903, Sugar Ridge field, Alabama	588
TOX-1. Index map of Toxey field, Alabama	590
TOX-2. Structure contour map on top of Smackover Formation in Toxey field, Alabama	591
TOX-3. Well log, Jimmerson 4-6 well, Permit No. 1464, Toxey field, Alabama	592
TOX-4. Structural cross section of the Toxey field, Alabama	593
TOX-5. Graph of production history of the Toxey field, Alabama	594
TOX-6. Porosity vs. natural log permeability plot of described core for Bolinger 3-4 well, Permit No. 1412, Toxey field, Alabama	595
TOX-7. Detailed core log, Bolinger No. 13-4 well, Permit No. 1412, Toxey field, Alabama	597
TOX-8. Porosity vs. natural log permeability plot for lithofacies 2, Bolinger 3-4 well, Permit No. 1412, Toxey field, Alabama	600
TOX-9. Paragenetic sequence, Bolinger 3-4 well, Permit No. 1412, Toxey field, Alabama	601
TOX-10. Ternary diagram of pore types, Bolinger 3-4 well, Permit No. 1412, Toxey field, Alabama	602
TC-1. Index map of Turkey Creek field, Alabama	604
TC-2. Structure contour map on top of Smackover Formation in Turkey Creek field, Alabama	605
TC-3. Well log, Chesley Pruet-Alco Land & Timber Co., Inc. No. 1 well, Permit No. 1400, Turkey Creek field, Alabama	606
TC-4. Structural cross section of the Turkey Creek field, Alabama	607
TC-5. Graph of production history of the Turkey Creek field, Alabama	608
TC-6. Porosity vs. natural log permeability plot for two cores combined, Turkey Creek field, Alabama	609
TUR-1. Index map of Turnerville field, Alabama	612
TUR-2. Structure contour map on top of Smackover Formation in Turnerville field, Alabama	613
TUR-3. Well log correlated to lithology, porosity, and permeability, R. J. Newman Jr., et al. 21-11 No. 1 well, Permit No. 4412, Turnerville field, Alabama	614
TUR-4. Structural cross section of the Turnerville field, Alabama	615
TUR-5. Graph of production history of the Turnerville field, Alabama	616
TUR-6. Porosity vs. natural log permeability plots, Turnerville field, Alabama	617
TUR-7. Detailed core log, R. J. Newman Jr. et al. 21-11 No. 1, Permit No. 4412, Turnerville field, Alabama	619
TUR-8. Porosity vs. natural log permeability plot for lithofacies 1, R. J. Newman Jr. et al. 21-11 No. 1 well, Permit No. 4412, Turnerville field, Alabama	621
TUR-9. Paragenetic sequence, R. J. Newman Jr. et al. 21-11 No. 1 well, Permit No. 4412, Turnerville field, Alabama	622
URH-1. Index map of Uriah field, Alabama	624
URH-2. Structure contour map on top of Smackover Formation in Uriah field, Alabama	625
URH-3. Well log, Bibb G. Mims 7-3 No. 1 well, Permit No. 7034, Uriah field, Alabama	626
URH-4. Structural cross section of the Uriah field, Alabama	627

ILLUSTRATIONS—CONTINUED

		Page
URH-5.	Graph of production history of the Uriah field, Alabama	628
URH-6.	Porosity vs. natural log permeability plot of two cores combined, Uriah field, Alabama	629
URH-7.	Detailed core log, Rufus Garrett 12-7 No. 1, Permit No. 1571, Uriah field, Alabama	631
URH-8.	Paragenetic sequence, Rufus Garrett 12-7 No. 1 well, Permit No. 1571, Uriah field, Alabama	637
VOC-1.	Index map of Vocation field, Alabama	639
VOC-2.	Structure contour map on top of Smackover Formation in Vocation field, Alabama	640
VOC-3.	Well log correlated to lithology, B. C. Quimby 27-15 No. 1 well, Permit No. 1599, Vocation field, Alabama	641
VOC-4.	Structural cross section of the Vocation field, Alabama	642
VOC-5.	Graph of production history of the Vocation field, Alabama	643
VOC-6.	Porosity vs. natural log permeability plot of one core, Vocation field, Alabama	644
VOC-7.	Detailed core log, B. C. Quimby 27-15 No. 1 well, Permit No. 1599, Vocation field, Alabama	646
VOC-8.	Paragenetic sequence, B. C. Quimby 27-15 No. 1 well, Permit No. 1599, Vocation field, Alabama	649
WAL-1.	Index map of Wallace field, Alabama	651
WAL-2.	Structure contour map on top of Smackover Formation in Wallace field, Alabama	652
WAL-3.	Well log, Coastal A.T.I.C. 13-6 No. 1 well, Permit No. 5268, Wallace field, Alabama	653
WAL-4.	Structural cross section of the Wallace field, Alabama	654
WAL-5.	Graph of production history of the Wallace field, Alabama	655
WC-1.	Index map of Wallers Creek field, Alabama	658
WC-2.	Structure contour map on top of Smackover Formation in Wallers Creek field, Alabama	659
WC-3.	Well log, Irvin Langford 9-14 No. 3 well, Permit No. 5019, Wallers Creek field, Alabama	660
WC-4.	Structural cross section of the Wallers Creek field, Alabama	661
WC-5.	Graph of production history of the Wallers Creek field, Alabama	662
WC-6.	Porosity vs. natural log permeability plot of three cores combined, Wallers Creek field, Alabama	663
WAP-1.	Index map of West Appleton field, Alabama	666
WAP-2.	Structure contour map on top of Smackover Formation in West Appleton field, Alabama	667
WAP-3.	Well log, P. H. Gallagher 16-3 No. 1 well, Permit No. 5757, West Appleton field, Alabama	668
WAP-4.	Structural cross section of the West Appleton field, Alabama	669
WAP-5.	Graph of production history of the West Appleton field, Alabama	670
WBT-1.	Index map of West Barrytown field, Alabama	673
WBT-2.	Structure contour map on top of Smackover Formation in West Barrytown field, Alabama	674
WBT-3.	Well log correlated to porosity and permeability, Abston Unit 10-6 No. 1 well, Permit No. 2148, West Barrytown field, Alabama	675
WBT-4.	Structural cross section of the West Barrytown field, Alabama	676
WBT-5.	Graph of production history of the West Barrytown field, Alabama	677

ILLUSTRATIONS—CONTINUED

	Page
WBT-6. Porosity vs. natural log permeability plots, West Barrytown field, Alabama	678
WBT-7. Detailed core log, Lewis Unit 4-16 No. 1, Permit No. 1757, West Barrytown field, Alabama	680
WBT-8. Porosity vs. natural log permeability plot for lithofacies 1, Lewis Unit 4-16 No. 1 well, Permit No. 1757, West Barrytown field, Alabama	684
WBT-9. Paragenetic sequence, Lewis Unit 4-16 No. 1 well, Permit No. 1757, West Barrytown field, Alabama	685
WBD-1. Index map of West Bend field, Alabama	687
WBD-2. Structure contour map on top of Smackover Formation in West Bend field, Alabama	688
WBD-3. Well log correlated to lithology, porosity, and permeability, May 27-10 No. 1 well, Permit No. 2647, West Bend field, Alabama	689
WBD-4. Structural cross section of the West Bend field, Alabama	690
WBD-5. Graph of production history of the West Bend field, Alabama	691
WBD-6. Porosity vs. natural log permeability plots, West Bend field, Alabama	692
WBD-7. Detailed core log, May 27-10 No. 1 well, Permit No. 2647, West Bend field, Alabama	694
WBD-8. Porosity vs. natural log permeability plots for two lithofacies, May 27-10 No. 1 well, Permit No. 2647, West Bend field, Alabama	697
WBD-9. Paragenetic sequence, May 27-10 No. 1 well, Permit No. 2647, West Bend field, Alabama	698
WOC-1. Index map of West Okatuppa Creek field, Alabama	700
WOC-2. Structure contour map on top of Smackover Formation in West Okatuppa Creek field, Alabama	701
WOC-3. Well log, Land 6-1 No. 1 well, Permit No. 5776-B, West Okatuppa Creek field, Alabama	702
WOC-4. Structural cross section of the West Okatuppa Creek field, Alabama	703
WOC-5. Graph of production history of the West Okatuppa Creek field, Alabama	704
WFC-1. Index map of Wild Fork Creek field, Alabama	707
WFC-2. Structure contour map on top of Smackover Formation in Wild Fork Creek field, Alabama	708
WFC-3. Well log, I.P.C. 5-10 No. 1 well, Permit No. 5411, Wild Fork Creek field, Alabama	709
WFC-4. Structural cross section of the Wild Fork Creek field, Alabama	710
WFC-5. Graph of production history of the Wild Fork Creek field, Alabama	711
WFC-6. Porosity vs. natural log permeability plot of one core, Wild Fork Creek field, Alabama	712
WIM-1. Index map of Wimberly field, Alabama	715
WIM-2. Structure contour map on top of Smackover Formation in Wimberly field, Alabama	716
WIM-3. Well log, Bolinger Unit 25-9 No. 1 well, Permit No. 2245, Wimberly field, Alabama	717
WIM-4. Structural cross section of the Wimberly field, Alabama	718
WIM-5. Graph of production history of the Wimberly field, Alabama	719
WIM-6. Porosity vs. natural log permeability plot of three cores combined, Wimberly field, Alabama	720
WOM-1. Index map of Womack Hill field, Alabama	723

ILLUSTRATIONS—CONTINUED

	Page
WOM-2. Structure contour map on top of Smackover Formation in Womack Hill field, Alabama	724
WOM-3. Well log, WHFU; Parker-Locke 9-16 No. 1 well, Permit No. 1655, Womack Hill field, Alabama	725
WOM-4. Structural cross section of the Womack Hill field, Alabama	726
WOM-5. Graph of production history of the Womack Hill field, Alabama	727
WOM-6. Porosity vs. natural log permeability plot of 38 wells combined, Womack Hill field, Alabama	728
WOM-7. Detailed core log, Turner 13-25 well, Permit No. 2327, Womack Hill field, Alabama	730
WOM-8. Paragenetic sequence, Turner 13-25 well, Permit No. 2327, Womack Hill field, Alabama	732
ZC-1. Index map of Zion Chapel field, Alabama	734
ZC-2. Structure contour map on top of Smackover Formation in Zion Chapel field, Alabama	735
ZC-3. Well log correlated to lithology, porosity, and permeability, Federal Land Bank Unit 17-11 No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	736
ZC-4. Structural cross section of the Zion Chapel field, Alabama	737
ZC-5. Graph of production history of the Zion Chapel field, Alabama	738
ZC-6. Porosity vs. natural log permeability plots, Zion Chapel field, Alabama	739
ZC-7. Detailed core log, Federal Land Bank No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	741
ZC-8. Porosity vs. log permeability plot for lithofacies 1, Federal Land Bank No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	743
ZC-9. Paragenetic sequence, Federal Land Bank Unit 17-11 No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	744
ZC-10. Ternary diagram of pore types, Federal Land Bank No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	745

TABLES

	Page
APT-1. Reservoir characterization sheet, Smackover pool, Appleton field, Escambia County, Alabama	8
APT-2. Lithofacies description of core from D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama	10
BAR-1. Reservoir characterization sheet, Smackover pool, Barnett field, Conecuh and Escambia Counties, Alabama	19
BAR-2. Lithofacies description of core from Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama	21
BB-1. Reservoir characterization sheet, Smackover pool, Barlow Bend field, Clarke and Monroe Counties, Alabama	32
BTN-1. Reservoir characterization sheet, Smackover pool, Barrytown field, Choctaw County, Alabama	40
BTN-2. Lithofacies description of core from McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama	42
BEC-1. Reservoir characterization sheet, Smackover pool, Big Escambia Creek field, Escambia County, Alabama	52
BEC-2. Lithofacies description of core from Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama	54
BLS-1. Reservoir characterization sheet, Smackover pool, Blacksher field, Baldwin County, Alabama	65
BLS-2. Lithofacies description of core from International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama	67
BLC-1. Reservoir characterization sheet, Smackover pool, Broken Leg Creek field, Escambia County, Alabama	79
BUC-1. Reservoir characterization sheet, Smackover pool, Bucatunna Creek field, Choctaw County, Alabama	87
BUC-2. Lithofacies description of core from Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama	89
BCC-1. Reservoir characterization sheet, Smackover pool, Burnt Corn Creek field, Escambia County, Alabama	101
BCC-2. Lithofacies description of core from Hooper Mathews 4-8 No. 1 well, Permit No. 2377, Burnt Corn Creek field, Alabama	103
CH-1. Reservoir characterization sheet, Smackover pool, Chappell Hill field, Choctaw County, Alabama	112
CH-2. Lithofacies description of core from Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama	114
CTM-1. Reservoir characterization sheet, Smackover pool, Chatom field, Washington County, Alabama	127
CTM-2. Lithofacies description of core from Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama	129
CHR-1. Reservoir characterization sheet, Smackover pool, Choctaw Ridge field, Choctaw County, Alabama	141
CHR-2. Lithofacies description of core from Rentz-Trice Unit 35-1 No. 5 well, Permit No. 1427, Choctaw Ridge field, Alabama	143
CHU-1. Reservoir characterization sheet, Smackover pool, Chunchula field, Mobile County, Alabama	152
CHU-2. Lithofacies description of core from Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama	154
CC-1. Reservoir characterization sheet, Smackover pool, Cold Creek field, Mobile County, Alabama	165

TABLES—CONTINUED

	Page
CPL-1. Reservoir characterization sheet, Smackover pool, Copeland field, Washington County, Alabama	173
CRC-1. Reservoir characterization sheet, Smackover pool, Crosbys Creek field, Washington County, Alabama	181
CRC-2. Lithofacies description of core from Darly Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama	183
EB-1. Reservoir characterization sheet, Smackover pool, East Barnett field, Conecuh County, Alabama	191
EB-2. Lithofacies description of core from Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama	193
EH-1. Reservoir characterization sheet, Smackover pool, East Huxford field, Escambia County, Alabama	201
FC-1. Reservoir characterization sheet, Smackover pool, Fanny Church field, Escambia County, Alabama	209
GC-1. Reservoir characterization sheet, Smackover pool, Gin Creek field, Choctaw County, Alabama	217
GC-2. Lithofacies description of core from Bolinger Heirs 32-5 No. 1 well, Permit No. 3312, Gin Creek field, Alabama	219
GUC-1. Reservoir characterization sheet, Smackover pool, Gulf Crest field, Choctaw County, Alabama	230
HC-1. Reservoir characterization sheet, Smackover pool, Hanberry Church field, Escambia County, Alabama	237
HP-1. Reservoir characterization sheet, Smackover pool, Hatter's Pond field, Mobile County, Alabama	245
HP-2. Lithofacies description of core from Hatter's Pond Unit 10-3 No. 1 well, Permit No. 6846, Hatter's Pond field, Alabama	247
HS-1. Reservoir characterization sheet, Smackover pool, Healing Springs field, Washington County, Alabama	258
HS-2. Lithofacies description of core from W. J. Britton No. 1 well, Permit No. 2478, Healing Springs field, Alabama	260
HUX-1. Reservoir characterization sheet, Smackover pool, Huxford field, Escambia County, Alabama	271
HUX-2. Lithofacies description of core from M. H. Murphy 26-11 No. 1 well, Permit No. 4577, Huxford field, Alabama	273
LEC-1. Reservoir characterization sheet, Smackover pool, Little Escambia Creek field, Escambia County, Alabama	284
LEC-2. Lithofacies description of core from T. R. Miller Mill Unit 32-3 well, Permit No. 1654, Little Escambia Creek field, Alabama	287
LMC-1. Reservoir characterization sheet, Smackover pool, Little Mill Creek field, Choctaw County, Alabama	301
LR-1. Reservoir characterization sheet, Smackover pool, Little River field, Baldwin County, Alabama	309
LR-2. Lithofacies description of core from J. E. Robinson et al. 11-8 No. 1 well, Permit No. 3046, Little River field, Alabama	311
LRK-1. Reservoir characterization sheet, Smackover pool, Little Rock field, Escambia County, Alabama	322
LOV-1. Reservoir characterization sheet, Smackover pool, Lovetts Creek field, Baldwin and Monroe Counties, Alabama	330
LOV-2. Lithofacies description of core from W. M. Threat 11-10 No. 1 well, Permit No. 3535, Lovetts Creek field, Alabama	333

TABLES—CONTINUED

	Page
MEL-1. Reservoir characterization sheet, Smackover pool, Melvin field, Choctaw County, Alabama	347
MC-1. Reservoir characterization sheet, Smackover pool, Mill Creek field, Choctaw County, Alabama	355
MIN-1. Reservoir characterization sheet, Smackover pool, Mineola field, Monroe County, Alabama	362
MOV-1. Reservoir characterization sheet, Smackover pool, Movico field, Baldwin and Mobile Counties, Alabama	370
MOV-2. Lithofacies description of core from Sullivan et al. No. 1 well, Permit No. 2126, Movico field, Alabama	372
NCR-1. Reservoir characterization sheet, Smackover pool, North Choctaw Ridge field, Choctaw County, Alabama	381
NCR-2. Lithofacies description of core from Oates Unit 29-6 No. 1 well, Permit No. 1878, North Choctaw Ridge field, Alabama	383
NSC-1. Reservoir characterization sheet, Smackover pool, North Smiths Church field, Escambia County, Alabama	394
NWC-1. Reservoir characterization sheet, Smackover pool, North Wallers Creek field, Monroe County, Alabama	402
NWC-2. Lithofacies description of core from Irvin Langford 5-10 No. 1 well, Permit No. 7589, North Wallers Creek field, Alabama	404
NEB-1. Reservoir characterization sheet, Smackover pool, Northeast Barnett field, Conecuh County, Alabama	414
NWR-1. Reservoir characterization sheet, Smackover pool, Northwest Range field, Conecuh County, Alabama	421
PC-1. Reservoir characterization sheet, Smackover pool, Pace Creek field, Clarke County, Alabama	428
PCR-1. Reservoir characterization sheet, Smackover pool, Palmers Crossroads field, Monroe and Clarke Counties, Alabama	436
PER-1. Reservoir characterization sheet, Smackover pool, Perdido field, Baldwin and Escambia Counties, Alabama	444
PER-2. Lithofacies description of core from Wade Johnson 30-16 well, Permit No. 4696, Perdido field, Alabama	446
PCC-1. Reservoir characterization sheet, Smackover pool, Puss Cuss Creek field, Choctaw County, Alabama	456
RC-1. Reservoir characterization sheet, Smackover pool, Red Creek field, Washington County, Alabama	464
RCK-1. Reservoir characterization sheet, Smackover pool, Robinson Creek field, Escambia County, Alabama	471
SIL-1. Reservoir characterization sheet, Smackover pool, Silas field, Choctaw County, Alabama	479
SIL-2. Lithofacies description of core from Chesnut Unit 9-1 No. 1 well, Permit No. 2205, Silas field, Alabama	481
SC-1. Reservoir characterization sheet, Smackover pool, Sizemore Creek field, Escambia County, Alabama	493
SC-2. Lithofacies description of core from A.T.I.C. 33-1 No. 1 well, Permit No. 3988, Sizemore Creek field, Alabama	495
SCH-1. Reservoir characterization sheet, Smackover pool, Smiths Church field, Escambia County, Alabama	503
SBC-1. Reservoir characterization sheet, Smackover pool, South Burnt Corn Creek field, Escambia County, Alabama	510

TABLES—CONTINUED

	Page
SCC-1. Reservoir characterization sheet, Smackover pool, South Cold Creek field, Mobile County, Alabama	518
SV-1. Reservoir characterization sheet, Smackover pool, South Vocation field, Monroe County, Alabama	526
SWF-1. Reservoir characterization sheet, Smackover pool, South Wild Fork Creek field, Escambia County, Alabama	533
SWH-1. Reservoir characterization sheet, Smackover pool, South Womack Hill field, Clarke County, Alabama	540
SEC-1. Reservoir characterization sheet, Smackover pool, Southeast Chatom field, Washington County, Alabama	548
SWB-1. Reservoir characterization sheet, Smackover pool, Southwest Barrytown field, Choctaw County, Alabama	556
SOU-1. Reservoir characterization sheet, Smackover pool, Souwilpa Creek field, Choctaw County, Alabama	563
STV-1. Reservoir characterization sheet, Smackover pool, Stave Creek field, Clarke County, Alabama	571
STV-2. Lithofacies description of core from McCorquodale 25-1 No. 1 well, Permit No. 2718, Stave Creek field, Alabama	573
SUG-1. Reservoir characterization sheet, Smackover pool, Sugar Ridge field, Choctaw County, Alabama	583
SUG-2. Lithofacies description of core from Lovett 27-6 No. 1 well, Permit No. 1903, Sugar Ridge field, Alabama	585
TOX-1. Reservoir characterization sheet, Smackover pool, Toxey field, Choctaw County, Alabama	596
TOX-2. Lithofacies description of core from Bolinger 3-4 well, Permit No. 1412, Toxey field, Alabama	598
TC-1. Reservoir characterization sheet, Smackover pool, Turkey Creek field, Choctaw County, Alabama	610
TUR-1. Reservoir characterization sheet, Smackover pool, Turnerville field, Mobile County, Alabama	618
TUR-2. Lithofacies description of core from R. J. Newman et al. 21-11 No. 1 well, Permit No. 4412, Turnerville field, Alabama	620
URH-1. Reservoir characterization sheet, Smackover pool, Uriah field, Monroe County, Alabama	630
URH-2. Lithofacies description of core from Rufus Garrett 12-7 No. 1 well, Permit No. 1571, Uriah field, Alabama	634
VOC-1. Reservoir characterization sheet Smackover pool, Vocation field, Monroe County, Alabama	645
VOC-2. Lithofacies description of core from B. C. Quimby 27-15 No. 1 well, Permit No. 1599, Vocation field, Alabama	647
WAL-1. Reservoir characterization sheet, Smackover pool, Wallace field, Escambia County, Alabama	656
WC-1. Reservoir characterization sheet, Smackover pool, Wallers Creek field, Monroe County, Alabama	664
WAP-1. Reservoir characterization sheet, Smackover pool, West Appleton field, Escambia County, Alabama	671
WBT-1. Reservoir characterization sheet, Smackover pool, West Barrytown field, Choctaw County, Alabama	679
WBT-2. Lithofacies description of core from Lewis Unit 4-16 No. 1 well, Permit No. 1757, West Barrytown field, Alabama	681

TABLES—CONTINUED

		Page
WBD-1.	Reservoir characterization sheet, Smackover pool, West Bend field, Clarke County, Alabama	693
WBD-2.	Lithofacies description of core from May 27-10 No. 1 well, Permit No. 2647, West Bend field, Alabama	695
WOC-1.	Reservoir characterization sheet, Smackover pool, West Okatuppa Creek field, Choctaw County, Alabama	705
WFC-1.	Reservoir characterization sheet, Smackover pool, Wild Fork Creek field, Escambia County, Alabama	713
WIM-1.	Reservoir characterization sheet, Smackover pool, Wimberly field, Choctaw County, Alabama	721
WOM-1.	Reservoir characterization sheet, Smackover pool, Womack Hill field, Choctaw and Clarke Counties, Alabama	729
WOM-2.	Lithofacies description of core from Turner 13-25 well, Permit No. 2327, Womack Hill field, Alabama	731
ZC-1.	Reservoir characterization sheet, Smackover pool, Zion Chapel field, Choctaw County, Alabama	740
ZC-2.	Lithofacies description of core from Federal Land Bank Unit 17-11 No. 1 well, Permit No. 2372, Zion Chapel field, Alabama	742

Field codes for Smackover oil fields in southwest Alabama

FIELD CODE	FIELD NAME	FIELD CODE	FIELD NAME
APP	Appleton field	NEB	Northeast Barnett field
BB	Barlow Bend field	NWR	Northwest Range field
BAR	Barnett field	PC	Pace Creek field
BTN	Barrytown field	PCR	Palmers Crossroads field
BEC	Big Escambia Creek field	PER	Perdido field
BLS	Blacksher field	PCC	Puss Cuss Creek field
BLC	Broken Leg Creek field	RC	Red Creek field
BUC	Bucatunna Creek field	RCK	Robinsons Creek field
BCC	Burnt Corn Creek field	SIL	Silas field
CH	Chappell Hill field	SC	Sizemore Creek field
CTM	Chatom field	SCH	Smiths Church
CHR	Choctaw Ridge field	SBC	South Burnt Corn Creek field
CHU	Churchula field	SCC	South Cold Creek field
CC	Cold Creek field	SV	South Vocation field
CPL	Copeland Gas field	SWF	South Wild Fork Creek field
CRC	Crosby Creek field	SWH	South Womack Hill field
EB	East Barnett field	SEC	Southeast Chatom field
EH	East Huxford field	SWB	Southwest Barrytown field
FC	Fanny Church field	SOU	Souwilpa Creek field
GC	Gin Creek field	STV	Stave Creek field
GUC	Gulf Crest field	SUG	Sugar Ridge field
HC	Hanberry Church field	TOX	Toxey field
HP	Hatter's Pond field	TC	Turkey Creek field
HS	Healing Springs field	TUR	Turnerville field
HUX	Huxford field	URH	Uriah field
LEC	Little Escambia Creek field	VOC	Vocation field
LMC	Little Mill Creek field	WAL	Wallace field
LR	Little River field	WC	Walls Creek field
LRK	Little Rock field	WAP	West Appleton field
LOV	Lovetts Creek field	WBT	West Barrytown field
MEL	Melvin field	WBD	West Bend field
MC	Mill Creek field	WOC	West Okatuppa Creek field
MIN	Mineola field	WFC	Wild Fork Creek field
MOV	Movico field	WIM	Wimberly field
NCR	North Choctaw Ridge field	WOM	Womack Hill field
NSC	North Smiths Church field	ZC	Zion Chapel field
NWC	North Wallers Creek field		

EXPLANATION OF MAP SYMBOLS AND ABBREVIATIONS

WELL SYMBOLS

	DRILLING OR LOCATION		GAS WELL
	SHUT-IN OR TEMPORARILY ABANDONED WELL		SHUT-IN OR TEMPORARILY ABANDONED GAS WELL
	PLUGGED AND ABANDONED WELL		PLUGGED AND ABANDONED GAS WELL
	SALT WATER DISPOSAL WELL		GAS WELL CONVERTED TO SALT WATER DISPOSAL WELL
	ENHANCED RECOVERY WELL		GAS WELL CONVERTED TO ENHANCED RECOVERY WELL
	OIL WELL		GAS-CONDENSATE WELL
	SHUT-IN OR TEMPORARILY ABANDONED OIL WELL		SHUT-IN OR TEMPORARILY ABANDONED GAS-CONDENSATE WELL
	PLUGGED AND ABANDONED OIL WELL		PLUGGED AND ABANDONED GAS-CONDENSATE WELL
	OIL WELL CONVERTED TO SALT WATER DISPOSAL WELL		GAS-CONDENSATE WELL CONVERTED TO SALT WATER DISPOSAL WELL
	OIL WELL CONVERTED TO ENHANCED RECOVERY WELL		GAS-CONDENSATE WELL CONVERTED TO ENHANCED RECOVERY WELL
	CORE DESCRIBED		TYPE LOG

STRUCTURE MAPS

SL	PERMITTED SURFACE LOCATION
BHL	WELL BOTTOM HOLE LOCATION
T4N	TOWNSHIP 4 NORTH
R6E	RANGE 6 EAST
13	SECTION 13 (GENERALLY NUMBERED 1 THROUGH 36; IRREGULAR SECTION NUMBERS MAY EXCEED 36)
1784	STATE PERMIT NUMBER; USED FOR WELL IDENTIFICATION
7354-A	PERMITTED WELL RE-ENTRY AFTER SUSPENSION OF INITIAL OPERATIONS
2763-B	WELL PERMITTED TO BE DIRECTIONALLY DRILLED
---	GWC --- GAS/WATER CONTACT
---	HKW --- HIGHEST KNOWN WATER
---	LKG --- LOWEST KNOWN GAS
---	LKO --- LOWEST KNOWN OIL
---	OWC --- OIL/WATER CONTACT
---	FIELD BOUNDARY
---	COUNTY BOUNDARY
---	STATE BOUNDARY
A — A'	LINE OF CROSS SECTION

TOOTH ON DOWNTOWN SIDE OF FAULT;
WIDTH BETWEEN UPTHROWN LINE AND
DOWNTOWN LINE PROPORTIONAL TO
AMOUNT OF "MISSING" SECTION AND
ASSOCIATED VERTICAL DISPLACEMENT
ALONG FAULT PLANE.

KEY TO CORE LOGS

coral		channel		desiccation crack
pisoid		cross lamination		vadose pisolite
ooid		nodular texture		diagenetic clasts
peloid		thin interval of nodular anhydrite		fenestral fabric
coated grain		laminae		tuberoid
oncooid		microbial laminae		breccia
<i>Favreina</i>		dolostone		boring
mollusk		lime mudstone		anhydrite
gastropod		emersion surface		dehydrated subaq. gypsum
microbial lam.		thick crust		sandstone
domal strom.		soil zone		siltstone
echinoderm		tepee structure		sandy or silty dolostone
encrusting forams				boundstone
intraclast				mudstone
imbrication				wackestone
dolomite		bioturbation unknown		packstone
thin section or multiple samples		bioturbation intensity 1		grainstone
P&P spl.		bioturbation intensity 2
CP spl.		bioturbation intensity 3
vuggy porosity		bioturbation intensity 4
moldic porosity		bioturbation intensity 5
interparticle porosity
intercrystalline porosity
secondary intraparticle porosity
cement-moldic porosity
fossomoldic porosity
fracture porosity				

Bioturbation intensities from
M. L. Droser and D. J. Bottjer,
1986, Jour. Sed. Pet., v. 56, p.
558-559.

particle-size column:

particles

dolomite crystals

ghosts of particles

ABBREVIATIONS

ac	-	acre	mcf	-	thousand cubic feet
API	-	American Petroleum Institute	mcfgd	-	thousand cubic feet of gas per day
bbls	-	barrels	md	-	millidarcies
bcpd	-	barrels of condensate per day	mg l	-	milligrams per liter
bcf	-	billion cubic feet	mi	-	miles
bhp	-	bottom-hole pressure	mm	-	millimeters
bioturb.	-	bioturbation	mmcf	-	million cubic feet
bopd	-	barrels of oil per day	na	-	not available
bopm	-	barrels of oil per month	ooip	-	original oil in place
bpd	-	barrels per day	phi	-	porosity (%)
bwpd	-	barrels of water per day	psi	-	pounds per square inch
c	-	coarse	psia	-	pounds per square absolute
ck	-	choke	psig	-	pounds per square inch gauge
cm	-	centimeters	pvt	-	pressure-volume-temperature
cu ft	-	cubic feet	scf	-	standard cubic feet
FEL	-	from east line	sed.	-	sedimentary
FNL	-	from north line	stb	-	stock tank barrels
FSL	-	from south line	structs.	-	structures
f	-	fine	text.	-	texture
ft	-	feet	tds	-	total dissolved solids
ftp	-	flowing tubing pressure	tr	-	trace
FWL	-	from west line	µm	-	micrometers(microns)
h	-	interval thickness	vc	-	very coarse
intens.	-	intensity	vf	-	very fine
k	-	kilovolts	xc	-	extra coarse
m	-	medium	xf	-	extra fine

APPLETON FIELD
SMACKOVER OIL POOL
ESCAMBIA COUNTY, ALABAMA

Figure APT-1.--Index map of Appleton field, Alabama.

Figure APT-2.--Structure contour map on top of Smackover Formation in Appleton field, Alabama.

Figure APT-3.--Well log, McMillan 2-14 No. 1 well, Permit No. 3854, Appleton field, Alabama.

Figure APT-4.--Structural cross section of the Appleton field, Alabama. (See figures APT-1 and APT-2 for line of section.)

Figure APT-5.--Graph of production history of the Appleton field, Alabama.

Figure APT-6.--Porosity vs. natural log permeability plot of described core from D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama. Equation applies to regression line.

Table APT-1.--Reservoir characterization sheet, Smackover pool,
Appleton field, Escambia County, Alabama

Discovery well

Operator: Texaco, U.S.A.
Permit number: 3854
Well Name: D.W. McMillan Trust 2-14 #1
Location: Sec. 2, T3N-R9E
Completion date: 10/19/83
Perforations: 12,902-12,938 ft
Initial flow rate: 661 bopd, 1,060 Mcfd,
16/64

Field data (as of 12/31/90)

Number of wells within field limits: 7
Producing wells in pool: 4
Production wells in field: 4
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 2
Salt water disposal wells: 1
Enhanced recovery wells: 0
Dry holes: 2
Drilling wells: 0
Unit spacing (acres): Unitized
Total field area (acres): ~840
Cumulative production
Condensate (bbl): 994,987
Gas (Mcf): 1,885,911
Water (bbl): 1,015,865

Reservoir data

Trap type: Structural
Reservoir structure: Anticline
Proven productive area (acres): 473
Average net pay thickness (feet): 39
Hydrocarbon type: Oil
Oil gravity (API): 52
Initial reservoir pressure (psia): 6,285
Current reservoir pressure (psia): 5,700
Reservoir drive: Solution gas/water
Type of enhanced recovery: none
Mean porosity (core) (%): 5.94
Porosity standard deviation (%): 3.75
Maximum porosity (%): 1.00
Minimum porosity (%): 14.30
Number of porosity analyses: 30
Geometric mean perm (core) (md): 0.69
Maximum permeability (md): 18.80
Minimum permeability (md): 0.01
Number of permeability analyses: 10
Dykstra-Parsons coefficient: na

Figure APT-7.--Detailed core log, D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama.

Table APT-2.--Lithofacies description of core from D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama (see figure APT-7)

Lithofacies 1: Microbial doloboundstone, dolomitized karst, peloid dolograinstone, other rock types

Pore types:

Intercrystalline: 46%

Moldic: 37% [much of this is shelter porosity identifiable only in thin section]

Vuggy: 17% [much of this is shelter porosity identifiable only in thin section]

Depositional Environment: Backbarrier lagoon, rapid upward shoaling to restricted peritidal setting followed by exposure.

Diagenesis: Mature karstification of microbial boundstone with development of vadose pisolite and of solution pipes. Early karst-associated fractures filled with calcite cement. Nonfabric selective dolomitization (second episode; early very finely crystalline dolomite is restricted to nonporous intervals higher in the core) and associated dissolution modified the primary pore system, which was dominated by shelter porosity in boundstone and by interparticle porosity in pellet grainstone (primary fabric is partially preserved where dolomitization was incomplete). Most outsize pores (e.g., shelter pores) are lined with euhedral dolomite cement crystals which apparently represent a third episode of dolomitization that took place after pore-system formation.

Percent of reservoir: 100

Depth interval(s) (ft): 12,971-12,988

Figure APT-8.--Paragenetic sequence, D. W. McMillan 2-15 No. 5 well, Permit No. 6247, Appleton field, Alabama.

BARNETT FIELD
SMACKOVER OIL POOL
CONECUH AND ESCAMBIA COUNTIES, ALABAMA

Figure BAR-1.--Index map of Barnett field, Alabama.

Figure BAR-2.--Structure contour map on top of Smackover Formation in Barnett field, Alabama.

Figure BAR-3.--Well log correlated to lithology, porosity, and permeability, Alger Sullivan "C" No. 1 well, Permit No. 1827, Barnett field, Alabama.

Figure BAR-4.--Structural cross section of the Barnett field, Alabama. (See figures BAR-1 and BAR-2 for line of section.)

Figure BAR-5.--Graph of production history of the Barnett field, Alabama.

Figure BAR-6.--Porosity vs. natural log permeability plots, Barnett field, Alabama. (a) Plot of six cores combined, and (b) plot of described core for Alger Sullivan "C" No. 1 well, Permit No. 1827-A.

Table BAR-1.--Reservoir characterization sheet, Smackover pool, Barnett field, Conecuh and Escambia Counties, Alabama

Discovery well

Operator: Wheless Ind. - Jack E. Stack
 Permit number: 1827-A
 Well Name: Alger Sullivan #1
 Location: Sec. 35, T4N-R7E
 Completion date: 6/08/75
 Perforations: 13,430-13,476 ft
 Initial flow rate: 523 bopd, 740 Mcfd,
 12/64 ck, 1,985 psi

Field data (as of 12/31/90)

Number of wells within field limits: 9
 Producing wells in pool: 2
 Production wells in field: 2
 Shut-in/Temp. abandoned wells: 0
 Plugged and abandoned wells: 5
 Salt water disposal wells: 1
 Enhanced recovery wells: 0
 Dry holes: 0
 Drilling wells: 1
 Unit spacing (acres): unitized
 Total field area (acres): 5,760
 Cumulative production
 Oil (bbl): 324,333
 Gas (Mcf): 370,660
 Water (bbl): 97,363

Reservoir data

Trap type: Structural / stratigraphic
 Reservoir structure: Anticline
 Proven productive area (acres): 1,237
 Average net pay thickness (feet): 12
 Hydrocarbon type: Oil
 Oil gravity (API): 49
 Initial reservoir pressure (psia): 6,618
 Current reservoir pressure (psia): 2,800
 Reservoir drive: Solution gas
 Type of enhanced recovery: none
 Mean porosity (core) (%): 6.00
 Porosity standard deviation (%): 3.30
 Maximum porosity (%): 20.20
 Minimum porosity (%): 1.00
 Number of porosity analyses: 273
 Geometric mean perm (core) (md): 0.14
 Maximum permeability (md): 1670.00
 Minimum permeability (md): 0.01
 Number of permeability analyses: 273
 Dykstra-Parsons coefficient: na

Figure BAR-7.--Detailed core log, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama.

Table BAR-2.--Lithofacies description of core from Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama (see figure BAR-7)

Lithofacies 1: Peloidal dolostone with intercalated dolomitic siltstone and dolomitic sandstone [only the carbonates are permeable]

Pore types:

Intercrystalline: 56%
Secondary intraparticle: 22%
Moldic: 19%
Vuggy: 3%

Mean porosity (%): 8.1

Maximum porosity (%): 14.5

Minimum porosity (%): 2.9

Porosity standard deviation (%): 3.4

Number of analyses: 24

Geometric mean permeability (md): 0.53

Maximum permeability (md): 21.00

Minimum permeability (md): 0.03

Number of analyses: 24

Depositional environment: Tidal flat and shallow lagoon near subaerially exposed source of quartzose sediment.

Diagenesis: Penecontemporaneous growth of displacive anhydrite nodules, pervasive dolomitization, leaching of remnant calcium carbonate, fracturing before precipitation of anhydrite cement, formation of authigenic anhydrite and pore-filling anhydrite cement, pressure solution, minor pyrite authigenesis.

Percent of reservoir: 100

Depth interval(s) (ft): 13,419-13,468

Figure BAR-8.--Porosity vs. natural log permeability plot, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama. Equation applies to regression line. (See table BAR-2 for description of lithofacies.) Data set excludes all samples with porosity values and permeability values below reservoir cutoffs at 6% and 0.1 md, respectively.

Figure BAR-9.--Paragenetic sequence, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama.

Figure BAR-10.--Ternary diagram of pore types, Alger Sullivan "C" No. 1 well, Permit No. 1827-A, Barnett field, Alabama. Percentages of moldic pores indicated. Sample depths are indicated to nearest 0.1 foot.

BARLOW BEND FIELD
SMACKOVER OIL POOL
CLARKE AND MONROE COUNTIES, ALABAMA

Figure BB-1.--Index map of Barlow Bend field, Alabama.

Figure BB-2.--Structure contour map on top of Smackover Formation in Barlow Bend field, Alabama.

Figure BB-3.--Well log, Henry Gibby Unit 7-15 No. 1 well, Permit No. 4905, Barlow Bend field, Alabama.

Figure BB-4.--Structural cross section of the Barlow Bend field, Alabama. (See figures BB-1 and BB-2 for line of section.)

Figure BB-5.--Graph of production history of the Barlow Bend field, Alabama.

Figure BB-6.--Porosity vs. natural log permeability plot of John R. Mattmuller et al. No. 1 well, Permit No. 2708, Barlow Bend field, Alabama. Equation applies to regression line.

Table BB-1.--Reservoir characterization sheet, Smackover pool,
Barlow Bend field, Clarke and Monroe Counties, Alabama

Discovery well

Operator: Kerr-McGee Corp.
Permit number: 4905
Well Name: Henry Gibby Unit 7-15 #1
Location: Sec. 7, T6N-R5E
Completion date: 06/09/86
Perforations: 12,740-12,758 ft
Initial flow rate: 838 bopd, 652 Mcfd,
21/64 ck, 770 psi

Field data (as of 12/31/90)

Number of wells within field limits: 2
Producing wells in pool: 1
Production wells in field: 1
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 1
Salt water disposal wells: 0
Enhanced recovery wells: 0
Dry holes: 0
Drilling wells: 0
Unit spacing (acres): 160
Total field area (acres): 640
Cumulative production
Oil (bbl): 28,089

Reservoir data

Trap type: Structural
Reservoir structure: Anticline
Proven productive area (acres): 71
Average net pay thickness (feet): 44
Hydrocarbon type: Oil
Oil gravity (API): 35
Initial reservoir pressure (psia): 6,097
Current reservoir pressure (psia): na
Reservoir drive: Solution gas / water
Type of enhanced recovery: none
Mean porosity (core) (%): 9.80
Porosity standard deviation (%): 5.23
Maximum porosity (%): 24.60
Minimum porosity (%): 3.90
Number of porosity analyses: 63
Geometric mean perm (core) (md): 0.99
Maximum permeability (md): 476.00
Minimum permeability (md): 0.02
Number of permeability analyses: 49
Dykstra-Parsons coefficient: na

BARRYTOWN FIELD
SMACKOVER OIL POOL
CHOCTAW COUNTY, ALABAMA

Figure BTN-1.--Index map of Barrytown field, Alabama.

Figure BTN-2.--Structure contour map on top of Snackover Formation in Barrytown field, Alabama.

Figure BTN-3.--Well log correlated to lithology and permeability, McIlwain 11-10 well, Permit No. 1818, Barrytown field, Alabama.

Figure BTN-4.--Structural cross section of the Barrytown field, Alabama. (See figures BTN-1 and BTN-2 for line of section.)

Figure BTN-5.--Graph of production history of the Barrytown field, Alabama.

Figure BTN-6.--Porosity vs. natural log permeability plots, Barrytown field, Alabama. (a) Plot of two cores combined, and (b) plot of described core for McIlwain 11-10 well, Permit No. 1818. Equations apply to regression lines.

Table BTN-1.--Reservoir characterization sheet, Smackover pool,
Barrytown field, Choctaw County, Alabama

Discovery well

Operator: Pruet & Hughes Co.

Permit number: 1696

Well Name: A.J. Phillips 12-2 #1

Location: Sec. 12, T10N-R3W

Completion date: 03/25/72

Perforations: 11,830-11,840, 11,848-11,860 ft

Initial flow rate: 264 bopd, 132 Mcfd,
1/4 ck, 600 psi

Field data (as of 12/31/90)

Number of wells within field limits: 9

Producing wells in pool: 3

Production wells in field: 3

Shut-in/Temp. abandoned wells: 0

Plugged and abandoned wells: 4

Salt water disposal wells: 1

Enhanced recovery wells: 0

Dry holes: 0

Drilling wells: 1

Unit spacing (acres): 120

Total field area (acres): 840

Cumulative production

Oil (bbl): 3,171,350

Gas (Mcf): 1,335,890

Water (bbl): 2,571,118

Reservoir data

Trap type: Structural / stratigraphic

Reservoir structure: Faulted anticline

Proven productive area (acres): 554

Average net pay thickness (feet): 43

Hydrocarbon type: Oil

Oil gravity (API): 45

Initial reservoir pressure (psia): na

Current reservoir pressure (psia): na

Reservoir drive: Solution gas / water

Type of enhanced recovery: none

Mean porosity (core) (%): 15.44

Porosity standard deviation (%): 4.14

Maximum porosity (%): 24.20

Minimum porosity (%): 5.70

Number of porosity analyses: 48

Geometric mean perm (core) (md): 6.61

Maximum permeability (md): 109.00

Minimum permeability (md): 1.50

Number of permeability analyses: 38

Dykstra-Parsons coefficient: na

Figure BTN-7.--Detailed core log, Mcllwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama.

Table BTN-2.--Lithofacies description of core from McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama (see figure BTN-7)

Lithofacies 1: Dolomitic pellet grainstone with minor crystalline dolostone

Pore types:

Moldic: 30%
Fracture: 28%
Intercrystalline: 25%
Secondary Intraparticle: 17%

Mean porosity (%): 23.3

Maximum porosity (%): 27.3

Minimum porosity (%): 20.7

Porosity standard deviation (%): 1.8

Number of analyses: 13

Geometric mean permeability (md): 16.65

Maximum permeability (md): 59.40

Minimum permeability (md): 5.4

Number of analyses: 13

Depositional Environment: Tidal flat, mostly subtidal.

Diagenesis: Early formation of lithified nodules and crusts, and the fracturing of these nodules or crusts, partial dolomitization, minor leaching of particles.

Percent of reservoir: 100

Depth interval(s) (ft): 11,824 -11,838.5

Figure BTN-8.--Porosity vs. natural log permeability plot for lithofacies 1, McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama. Equation applies to regression line. (See table BTN-2 for descriptions of lithofacies.

Figure BTN-9.--Paragenetic sequence, McIlwain 11-10 No. 1 well, Permit No. 1818, Barrytown field, Alabama.

BIG ESCAMBIA CREEK FIELD
SMACKOVER GAS AND CONDENSATE POOL
ESCAMBIA COUNTY, ALABAMA

Figure BEC-1.--Index map of Big Escambia Creek field, Alabama.

Figure BEC-2.--Structure contour map on top of Smackover Formation in Big Escambia Creek field, Alabama.

Figure BEC-3.--Well log correlated to lithology, porosity, and permeability, Smith 25-7 No. 1 well, Permit 4727, Big Escambia Creek field, Alabama.

Figure BEC-4.--Structural cross section of the Big Escambia Creek field, Alabama. (See figures BEC-1 and BEC-2 for line of section.)

Figure BEC-5.--Graph of production history of the Big Escambia Creek field, Alabama.

Figure BEC-6.--Porosity vs. natural log permeability plots, Big Escambia Creek field, Alabama. (a) Plot of 27 cores combined, and (b) plot of described core for Smith 25-7 No. 1 well, Permit No. 4727. Equations apply to regression lines.

Table BEC-1.--Reservoir characterization sheet, Smackover pool,
Big Escambia Creek field, Escambia County, Alabama

Discovery well

Operator: Mallard Exploration, Inc.

Permit number: 1649

Well Name: Int. Paper Co. et al Unit 2-6 #1

Location: Sec. 2, T1N-R7E

Completion date: 12/17/71

Perforations: 15,090-15,106 , 15,142-15,178 ft

Initial flow rate: 332 bcpd, 5,891 mcf,
17/64 ck, 3700 psi

Field data (as of 12/31/90)

Number of wells within field limits: 52

Producing wells in pool: 26

Production wells in field: 26

Shut-in/Temp. abandoned wells: 1

Plugged and abandoned wells: 19

Salt water disposal wells: 2

Enhanced recovery wells: 0

Dry holes: 4

Drilling wells: 0

Unit spacing (acres): 640

Total field area (acres): 20,480

Cumulative production

Condensate (bbl): 38,623,880

Gas (mcf): 552,626,786

Water (bbl): 1,252,215

Reservoir data

Trap type: Structural / Stratigraphic

Reservoir structure: Faulted Monocline

Proven productive area (acres): 15,858

Average net pay thickness (feet): 63

Hydrocarbon type: Gas Condensate

Oil gravity (API): 46

Initial reservoir pressure (psia): 7,662

Current reservoir pressure (psia): 3,258

Reservoir drive: Depletion

Type of enhanced recovery: none

Mean porosity (core) (%): 12.00

Porosity standard deviation: 5.20

Maximum porosity (%): 34.10

Minimum porosity (%): 0.20

Number of porosity analyses: 1,648

Geometric mean perm (core) (md): 0.64

Maximum permeability (md): 344

Minimum permeability (md): 0.01

Number of permeability analyses: 1,648

Dykstra-Parsons coefficient: na

Figure BEC-7--Detailed core log, Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama.

Table BEC-2.--Lithofacies description of core from Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama (see figure BEC-7)

Lithofacies 1: Peloid dolowackestone

Pore types:

Intercrystalline: 65%
 Vuggy: 16%
 Secondary intraparticle: 10%
 Moldic: 8%
 Fracture: 1%

Mean porosity (%): 17.8

Maximum porosity (%): 25.0

Minimum porosity (%): 6.0

Porosity standard deviation (%): 5.8

Number of analyses: 34

Geometric mean permeability (md): 21.82

Maximum permeability (md): 344.00

Minimum permeability (md): 0.03

Number of analyses: 34

Depositional Environment: Shallow subtidal restricted low-energy shelf, probably a sheltered lagoon behind a high-energy shoal.

Diagenesis: Dolomitization, leaching, late calcite cementation and lesser dolomite cementation in vugs (and molds). Pressure solution seems to predate at least some dolomitization, and follows fracture formation.

Percent of Reservoir: 61

Depth interval(s) (ft): 15,687-15,721.5

Lithofacies 2: Oncoidal peloid dolopackstone/grainstone, dominated by thin shoaling-upward cycles

Pore types:

Intercrystalline: 35%
 Secondary intraparticle: 35%
 Vuggy: 19%
 Moldic: 11%

Mean porosity (%): 11.5

Maximum porosity (%): 13.0

Minimum porosity (%): 8.8

Porosity standard deviation (%): 1.3

Number of analyses: 12

Table BEC-2.--Lithofacies description of core from Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama (see figure BEC-7) -- Continued

Geometric mean permeability (md): 2.65

Maximum permeability (md): 9.20

Minimum permeability (md): 0.45

Number of analyses: 12

Depositional Environment: Moderate energy shallow-marine restricted shelf, probably lower tidal flat in partially sheltered area, local beach development.

Diagenesis: Dolomitization, leaching, late calcite cementation and anhydrite cementation in molds and vugs. Pressure solution seems to predate at least some dolomitization.

Percent of reservoir: 22

Depth interval(s) (ft): 15,663-15,676

Lithofacies 3: Partially dolomitized coated-grain peloid wackestone and packstone

Pore types:

Secondary intraparticle: 43%

Vuggy: 27%

Intercrystalline: 23%

Moldic: 7%

Mean porosity (%): 7.7

Maximum porosity (%): 11.0

Minimum porosity (%): 5.4

Porosity standard deviation (%): 1.5

Number of analyses: 11

Geometric mean permeability (md): 0.14

Maximum permeability (md): 0.43

Minimum permeability (md): 0.03

Number of analyses: 11

Depositional Environment: Low energy to moderate energy shallow marine restricted shelf, probable lagoon behind high-energy barrier (some *Favreina* pellets observed).

Diagenesis: Dolomitization, leaching, late calcite cementation in vugs.

Percent of reservoir: 17

Depth interval(s) (ft): 15,676-15,687

Figure BEC-8.--Porosity vs. natural log permeability plots for three lithofacies, Smith 25-7 No. 1 well, Permit No. 4727 Big Escambia Creek field, Alabama. (a) Lithofacies 1, (b) lithofacies 2, (c) lithofacies 3. Equations for regression lines are printed for each lithofacies. (See table BEC-2 for descriptions of lithofacies.)

Figure BEC-9.--Paragenetic sequence, Smith 25-7 No. 1 well, Permit No. 4727, Big Escambia Creek field, Alabama.

BLACKSHER FIELD
SMACKOVER OIL POOL
BALDWIN COUNTY, ALABAMA

Figure BLS-1.--Index map of Blacksher field, Alabama.

Figure BLS-2.--Structure contour map on top of Smackover Formation in Blacksher field, Alabama.

Figure BLS-3.--Well log correlated to lithology, porosity, and permeability, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama.

Figure BLS-4.--Structural cross section of the Blacksher field, Alabama. (See figures BLS-1 and BLS-2 for line of section.)

Figure BLS-5.--Graph of production history of the Blacksher field, Alabama.

Figure BLS-6.--Porosity vs. natural log permeability plots, Blacksher field, Alabama. (a) Plot of eight cores combined, and (b) plot of described core for International Paper Co. 20-5 No. 1 well, Permit No. 5242. Equations apply to regression lines.

Table BLS-1.--Reservoir characterization sheet, Smackover pool,
Blacksher field, Baldwin County, Alabama

Discovery well

Operator: Shell Oil Co.

Permit number: 2925

Well Name: Wefel #2

Location: Sec. 29, T3N-R4E

Completion date: 07/27/80

Perforations: 15,585-15,600, 15,617-15,624 ft

Initial flow rate: 120 bopd, 173 Mcfd,
24/64 ck, 329 psi

Field data (as of 12/31/90)

Number of wells within field limits: 8

Producing wells in pool: 3

Production wells in field: 3

Shut-in/Temp. abandoned wells: 0

Plugged and abandoned wells: 4

Salt water disposal wells: 1

Enhanced recovery wells: 0

Dry holes: 0

Drilling wells: 0

Unit spacing (acres): 160

Total field area (acres): 2,400

Cumulative production

Oil (bbl): 1,897,279

Gas (Mcf): 1,879,390

Water (bbl): 955,084

Reservoir data

Trap type: Structural / stratigraphic

Reservoir structure: Anticline

Proven productive area (acres): 1,123

Average net pay thickness (feet): 12

Hydrocarbon type: Oil

Oil gravity (API): 43

Initial reservoir pressure (psia): 8,015

Current reservoir pressure (psia): na

Reservoir drive: Solution gas / water

Type of enhanced recovery: none

Mean porosity (core) (%): 11.60

Porosity standard deviation (%): 4.30

Maximum porosity (%): 25.30

Minimum porosity (%): 6.10

Number of porosity analyses: 294

Geometric mean perm (core) (md): 3.29

Maximum permeability (md): 627.00

Minimum permeability (md): 0.10

Number of permeability analyses: 294

Dykstra-Parsons coefficient: na

Figure BLS-7.--Detailed core log, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama.

Table BLS-2.--Lithofacies description of core from International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama (see figure BLS-7)

Lithofacies 1: Nonskeletal (ooid-dominated) dolograinstone and dolomitic grainstone

Pore types:

Secondary intraparticle: 49 %
Moldic: 29 %
Interparticle: 18 %
Vuggy: 4 %

Mean porosity (%): 15.1

Maximum porosity (%): 19.6

Minimum porosity (%): 8.9

Porosity standard deviation (%): 3.7

Number of analyses: 13

Geometric mean permeability (md): 5.33

Maximum permeability (md): 94.80

Minimum permeability (md): 0.14

Number of analyses: 13

Depositional Environment: High-energy beach with dune-filled runnel in upper part and soil horizon in lower part. Beach deposits overlie a shallow brine pond, and underlie thick brine-pond deposits.

Diagenesis: Early carbonate pore-lining cementation, particle dissolution, pervasive dolomitization, replacement of carbonate by anhydrite, formation of anhydrite cement.

Percent of reservoir: 44

Depth interval(s) (ft): 15,557-15,572

Lithofacies 2: Peloid dolograinstone and peloid grainstone; crystalline dolostone; oncoidal fenestral peloid grainstone.

Pore types:

Intercrystalline: 51%
Moldic: 25%
Secondary intraparticle: 15%
Fracture: 8%

Mean porosity (%): 15.0

Maximum porosity (%): 25.3

Minimum porosity (%): 6.2

Porosity standard deviation (%): 7.0

Number of analyses: 10

Table BLS-2.--Lithofacies description of core from International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama (see figure BLS-7) -- Continued

Geometric mean permeability (md): 25.14

Maximum permeability (md): 353.00

Minimum permeability (md): 0.77

Number of analyses: 10

Depositional Environment: Barrier complex with both subtidal (cross-laminated) and supratidal (fenestral) elements. On top of this barrier developed the brine pond overlain by the beach deposits of lithofacies 1.

Diagenesis: Pervasive dolomitization of upper part. Anhydrite is abundant as replacive laths and nodules, displacive nodules, and as pore-filling cement. Anhydrite formation was followed by minor dolomitization of anhydrite.

Percent of reservoir: 56

Depth interval(s) (ft): 15,578-15,597

Figure BLS-8.--Porosity vs. natural log permeability plots for lithofacies, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama. (a) Lithofacies 1 and (b) lithofacies 2. Equations for regression lines are printed for each lithofacies. (See table BLS-2 for descriptions of lithofacies.) Data set excludes all samples with porosity values and permeability values below reservoir cutoffs at 6% and 0.1 md, respectively.

Figure BLS-9.--Paragenetic sequence, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama.

Figure BLS-10.--Ternary diagram of pore types, International Paper Co. 20-5 No. 1 well, Permit No. 5242, Blacksher field, Alabama. Percentages of moldic pores indicated. Sample depths are indicated to nearest 0.1 foot.

BROKEN LEG CREEK FIELD
SMACKOVER OIL POOL
ESCAMBIA COUNTY, ALABAMA

Figure BLC-1.--Index map of Broken Leg Creek field, Alabama.

Figure BLC-2.--Structure contour map on top of Smackover Formation in Broken Leg Creek field, Alabama.

Figure BLC-3.--Well log, Hawkins 7-16 No. 1 well, Permit No. 6153-B, Broken Leg Creek field, Alabama.

Figure BLC-4.--Structural cross section of the Broken Leg Creek field, Alabama. (See figures BLC-1 and BLC-2 for line of section.)

Figure BLC-5.--Graph of production history of the Broken Leg Creek field, Alabama.

Figure BLC-6.--Porosity vs. natural log permeability plot of one core, Broken Leg Creek field, Alabama. Equation applies to regression line.

Table BLC-1.--Reservoir characterization sheet, Smackover pool,
Broken Leg Creek field, Escambia County, Alabama

Discovery well

Operator: Coastal Oil & Gas Corp.
Permit number: 5838-B
Well Name: Container Corp. 7-15 #1
Location: Sec. 7, T3N-R8E
Completion date: 10/07/88
Perforations: 14,067-14,108 ft
Initial flow rate: 792 bopd, 779 Mcfd,
14/64 ck, 2,440 psi

Field data (as of 12/31/90)

Number of wells within field limits: 3
Producing wells in pool: 2
Production wells in field: 2
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 1
Salt water disposal wells: 0
Enhanced recovery wells: 0
Dry holes: 0
Drilling wells: 0
Unit spacing (acres): 160
Total field area (acres): 480
Cumulative production
Oil (bbl): 249,838
Gas (Mcf): 180,916
Water (bbl): 155,512

Reservoir data

Trap type: Structural
Reservoir structure: Anticline
Proven productive area (acres): 263
Average net pay thickness (feet): 26
Hydrocarbon type: Oil
Oil gravity (API): 41
Initial reservoir pressure (psia): 6,725
Current reservoir pressure (psia): ~5,000
Reservoir drive: Solution gas / water
Type of enhanced recovery: none
Mean porosity (core) (%): 9.53
Porosity standard deviation (%): 3.91
Maximum porosity (%): 16.50
Minimum porosity (%): 2.60
Number of porosity analyses: 47
Geometric mean perm (core) (md): 1.32
Maximum permeability (md): 0.02
Minimum permeability (md): 0.02
Number of permeability analyses: 39
Dykstra-Parsons coefficient: na

BUCATUNNA CREEK FIELD
SMACKOVER OIL POOL
CHOCTAW COUNTY, ALABAMA

Figure BUC-1.--Index map of Bucatunna Creek field, Alabama.

Figure BUC-2.--Structure contour map on top of Smackover Formation in Bucatunna Creek field, Alabama.

Figure BUC-3.--Well log, A. G. Harrell No. 1 well, Permit No. 2470, Bucatunna Creek field, Alabama.

Figure BUC-4.--Structural cross section of the Bucatunna field, Alabama. (See figures BUC-1 and BUC-2 for line of section.)

Figure BUC-5.--Graph of production history of the Bucatunna Creek field, Alabama.

Figure BUC-6.--Porosity vs. natural log permeability plots, Bucatunna Creek field, Alabama. (a) Plot of two cores combined, and (b) plot of described core for Johnson Unit 14-14 No. 1 well, Permit No. 2753-B. Equations apply to regression lines.

Table BUC-1.--Reservoir characterization sheet, Smackover pool,
Bucatumna Creek field, Choctaw County, Alabama

Discovery well

Operator: Sandefer & Andress, Inc.
Permit number: 2470
Well Name: A. G. Harrel #1
Location: Sec. 15, T11N-R5W
Completion date: 3/31/78
Perforations: 12,071-12,100 ft
Initial flow rate: 338 bopd, 99 Mcfd,
12/64 ck, 775 psi

Field data (as of 12/31/90)

Number of wells within field limits: 6
Producing wells in pool: 1
Production wells in field: 1
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 2
Salt water disposal wells: 1
Enhanced recovery wells: 0
Dry holes: 2
Drilling wells: 0
Unit spacing (acres): 80
Total field area (acres): ~240
Cumulative production
Condensate (bbl): 1,134,435
Gas (Mcf): 425,410
Water (bbl): 461,027

Reservoir data

Trap type: Structural
Reservoir structure: Faulted anticline
Proven productive area (acres): 126
Average net pay thickness (feet): 45
Hydrocarbon type: Oil
Oil gravity (API): 36.6
Initial reservoir pressure (psia): 5,596
Current reservoir pressure (psia): 3,285
Reservoir drive: Solution gas / water
Type of enhanced recovery: none
Mean porosity (core) (%): 17.00
Porosity standard deviation (%): 6.80
Maximum porosity (%): 31.20
Minimum porosity (%): 2.90
Number of porosity analyses: 59
Geometric mean perm (core) (md): 1.23
Maximum permeability (md): 30.60
Minimum permeability (md): 0.07
Number of permeability analyses: 59
Dykstra-Parsons coefficient: na

Figure BUC-7.--Detailed core log, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama.

Table BUC-2.--Lithofacies description of core from Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama (see figure BUC-7)

Lithofacies 1: Oncoidal, pelletal, and oolitic grainstone

Pore types:

Secondary intraparticle: 52% (including all 3 particle types: oncoids, pellets, ooids)
 Particle moldic: 48% (all 3 particle types)
 Vuggy: trace

Mean porosity (%): 19.2

Maximum porosity (%): 24.3

Minimum porosity (%): 10.6

Porosity standard deviation (%): 3.8

Number of analyses: 25

Geometric mean permeability (md): 2.39

Maximum permeability (md): 30.60

Minimum permeability (md): 0.16

Number of analyses: 25

Depositional Environment: Flood-tidal delta, periodically inactive, with resultant growth of abundant oncoids.

Diagenesis: The core is brown in color and was interpreted as dolostone in the core-analysis report. Actually, dolomite composes <10 percent of this lithofacies. Early marine rim cement forms a rigid framework in much of this lithofacies, but interparticle interpenetration and porosity occlusion by overcompaction are important locally, and in the uppermost 4 feet of this lithofacies. Late pressure solution formed stylolites and associated replacive anhydrite, including a 2-cm stylocumulate, without substantial porosity reduction. Leaching of unstable particles generated the pore system during or after the major episode of compaction.

Percent of reservoir: 52

Depth interval(s) (ft): 12,256-12,272

Lithofacies 2: Ooid grainstone

Pore types:

Secondary intraparticle: 45%
 Interparticle: 40%
 Moldic: 5%
 Solution-modified: 5%
 Vuggy: 5%
 Intercrystalline: trace
 Fracture: trace

Mean porosity: 10.1

Table BUC-2.--Lithofacies description of core from Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama (see figure BUC-7) -- Continued

Maximum porosity: 11.9

Minimum porosity: 7.7

Porosity standard deviation: 1.8

Number of analyses: 5

Geometric mean permeability (md): 0.42

Maximum permeability (md): 1.61

Minimum permeability (md): 0.07

Number of analyses: 5

Depositional Environment: Flood-tidal delta.

Diagenesis: Reservoir originally resembled that of lithofacies 1, but substantial amounts of primary interparticle porosity have been preserved in lithofacies 2. In addition, this lithofacies has been strongly overprinted by stylolitization, dissolution (mostly fabric selective) and growth of replacive anhydrite laths and stylolites. Early overcompaction was more extensive than in lithofacies 1. Early formation of micritic meniscus cement indicates a brief period of subaerial exposure, probably before deposition of the overlying strata of lithofacies 4, which were deposited in a coastal sabkha. Formation of this early cement was followed by precipitation of marine-phreatic aragonitic rim cement which left many pores still open but which destroyed interparticle pore throats.

Percent of Reservoir: 23

Depth interval(s) (ft): 12,249-12,256

Lithofacies 3: Oolitic pellet grainstone and pelletal ooid grainstone

Pore types:

Pelmoldic: 50%

Oomoldic: 45%

Interparticle: 5%

Mean porosity (%): 25.2

Maximum porosity (%): 31.2

Minimum porosity (%): 18.1

Porosity standard deviation (%): 4.1

Number of analyses: 8

Geometric mean permeability (md): 2.94

Maximum permeability (md): 12.80

Minimum permeability (md): 0.91

Number of analyses: 8

Table BUC-2.--Lithofacies description of core from Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama (see figure BUC-7) -- Continued

Depositional Environment: Backshoal subtidal lagoon grading up into washover or flood-tidal delta.

Diagenesis: Early marine carbonate rim cementation was followed by slight local overcompaction, pore-filling carbonate cementation, and preservation of minor interparticle porosity at the base of the reservoir. Subsequent leaching of particles generated the present reservoir pore system; minor dolomitization and growth of replacive anhydrite laths did not materially affect the pore system.

Percent of Reservoir: 13

Depth interval(s) (ft): 12,272-12,276

Lithofacies 4: Silty anhydritic ooid dolograinstone

Pore types:

Oomoldic: dominant*
 Pelmoldic: minor
 Secondary intraparticle: minor

*Accurate percentages indeterminable.

Mean porosity (%): 13.0

Maximum porosity (%): 24.3

Minimum porosity (%): 7.9

Porosity standard deviation (%): 7.6

Number of analyses: 4

Geometric mean permeability (md): 0.95

Maximum permeability (md): 6.60

Minimum permeability (md): 0.41

Number of analyses: 4

Depositional Environment: Tidal flat/coastal sabkha.

Diagenesis: Early diagenesis included local overcompaction with particle deformation (flattening); generation of the pore system consisted of pervasive dolomitization of cement and of some particles, followed by leaching of nondolomitized particles. Subsequent growth of replacive nodular anhydrite, and anhydrite cementation in some oomolds, partially occluded the reservoir; subhorizontal nodules composed of euhedral dolomite crystals and very small anhydrite laths in subequal proportions may be caliche crusts.

Percent of Reservoir: 12, but includes intercalated nodular anhydrite

Depth interval(s) (ft): 12,242-12,246

Figure BUC-8.--Porosity vs. natural log permeability plot for lithofacies 1, Johnson Unit 14-14 No. 1 well, Permit No. 2753-B, Bucatunna Creek field, Alabama. The other three lithofacies were represented by too few samples to yield interpretable graphs. Equation applies to regression line. (See table BUC-2 for descriptions of lithofacies.) Data set excludes all samples with porosity values and permeability values below reservoir cutoffs at 6% and 0.1 md, respectively.

Figure BUC-9.--Paragenetic sequence, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama.

Figure BUC-10.--Ternary diagram of pore types, Johnson Unit 14-14 No. 1 well, Permit No. 2753, Bucatunna Creek field, Alabama. Percentages of moldic pores indicated. Sample depths are indicated to nearest 0.1 foot.

BURNT CORN CREEK FIELD
SMACKOVER OIL POOL
ESCAMBIA COUNTY, ALABAMA

Figure BCC-1.--Index map of Burnt Corn Creek field, Alabama.

Figure BCC-2.--Structure contour map on top of Smackover Formation in Burnt Corn Creek field, Alabama.

Figure BCC-3.--Well log, A.J. Smith 3-14 No. 1 well, Permit No. 4744, Burnt Corn Creek field, Alabama.

Figure BCC-4.--Structural cross section of the Burnt Corn Creek field, Alabama. (See figures BCC-1 and BCC-2 for line of section.)

Figure BCC-5.--Graph of production history of the Burnt Corn Creek field, Alabama.

Table BCC-1.--Reservoir characterization sheet, Smackover pool,
Burnt Corn Creek field, Escambia County, Alabama

Discovery well

Operator: Smackco, Ltd.
Permit number: 4744
Well Name: A.J. Smith 3-14 #1
Location: Sec. 3, T3N-R9E
Completion date: 11/06/85
Perforations: 12,920-12,980 ft
Initial flow rate: 120 bopd, 135 Mcfd,
18/64 ck, 255 psi

Field data (as of 12/31/90)

Number of wells within field limits: 1
Producing wells in pool: 0
Production wells in field: 0
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 1
Salt water disposal wells: 0
Enhanced recovery wells: 0
Dry holes: 0
Drilling wells: 0
Unit spacing (acres): 160
Total field area (acres): 160
Cumulative production
Oil (bbl): 10,911
Gas (Mcf): 9,038
Water (bbl): 27,710

Reservoir data

Trap type: Structural / stratigraphic
Reservoir structure: Anticline
Proven productive area (acres): 162
Average net pay thickness (feet): na
Hydrocarbon type: Oil
Oil gravity (API): 49
Initial reservoir pressure (psia): 6,123
Current reservoir pressure (psia): 5,909
Reservoir drive: Solution gas / water
Type of enhanced recovery: none
Mean porosity (core) (%): na
Porosity standard deviation (%): na
Maximum porosity (%): na
Minimum porosity (%): na
Number of porosity analyses: na
Geometric mean perm (core) (md): na
Permeability standard deviation (%): na
Maximum permeability (md): na
Minimum permeability (md): na
Number of permeability analyses: na
Dykstra-Parsons coefficient: na

Figure BCC-6.--Detailed core log, Hooper Mathews 4-8 No. 1 well, Permit No. 2377, Burnt Corn Creek field, Alabama.

Table BCC-2.--Lithofacies description of core from Hooper Mathews 4-8 No. 1 well, Permit No. 2377,
Burnt Corn Creek field, Alabama (see figure BCC-7)

Lithofacies 1: Very fine to fine planar-s dolostone (hypidiotopic dolostone)

Pore types:

Intercrystalline: 55%

Outsize: 45%

Depositional Environment: Subtidal flat.

Diagenesis: Pervasive and fabric destructive dolomitization is the most obvious diagenetic process. Uncommon and small fractures are filled with anhydrite cement. Late calcite cement fills a few intercrystalline pores locally.

Percent of reservoir: 100

Depth interval(s) (ft): 13,013.5 -13,020

Figure BCC-7.--Paragenetic sequence, Hooper Mathews 4-8 No. 1 well, Permit No. 2377, Burnt Corn Creek field, Alabama.

CHAPPELL HILL FIELD
SMACKOVER OIL POOL
CHOCTAW COUNTY, ALABAMA

Figure CH-1.--Index map of Chappell Hill field, Alabama.

Figure CH-2.--Structure contour map on top of Smackover Formation in Chappell Hill field, Alabama.

Figure CH-3.--Well log correlated to lithology, porosity, and permeability, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama.

Figure CH-4.--Structural cross section of the Chappell Hill field, Alabama. (See figures CH-1 and CH-2 for line of section.)

Figure CH-5.--Graph of production history of the Chappell Hill field, Alabama.

Figure CH-6.--Porosity vs. natural log permeability plots, Chappell Hill field, Alabama. (a) Plot of three cores combined, and (b) plot of described core for Bolinger 26-7 No. 1 well, Permit No. 2426. Equations apply to regression lines.

Table CH-1.--Reservoir characterization sheet, Smackover pool,
Chappell Hill field, Choctaw County, Alabama

Discovery well

Operator: Midroc Oil Co. - H.V. Watkins, Jr.

Permit number: 2371

Well Name: Jenkins 26-5 #1

Location: Sec. 26, T11N-R3W

Completion date: 09/18/77

Perforations: 11,412-11,422 ft

Initial flow rate: 125 bopd, 16/64 ck,
160 psi

Field data (as of 12/31/90)

Number of wells within field limits: 4

Producing wells in pool: 3

Production wells in field: 3

Shut-in/Temp. abandoned wells: 0

Plugged and abandoned wells: 0

Salt water disposal wells: 1

Enhanced recovery wells: 0

Dry holes: 0

Drilling wells: 0

Unit spacing (acres): 120

Total field area (acres): 960

Cumulative production

Oil (bbl): 1,808,901

Gas (Mcf): 764,644

Water (bbl): 721,390

Reservoir data

Trap type: Structural

Reservoir structure: Faulted anticline

Proven productive area (acres): 256

Average net pay thickness (feet): 33

Hydrocarbon type: Oil

Oil gravity (API): 38

Initial reservoir pressure (psia): 5,178

Current reservoir pressure (psia): 3,451

Reservoir drive: Solution gas / water

Type of enhanced recovery: none

Mean porosity (core) (%): 16.20

Porosity standard deviation (%): 4.93

Maximum porosity (%): 30.10

Minimum porosity (%): 6.40

Number of porosity analyses: 109

Geometric mean perm (core) (md): 3.64

Maximum permeability (md): 3,605.00

Minimum permeability (md): 0.3

Number of permeability analyses: 109

Dykstra-Parsons coefficient: na

Figure CH-7.--Detailed core log, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama.

Table CH-2.-- Lithofacies description of core from Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama (see figure CH-7)

Lithofacies 1: Pellet grainstone

Pore types:

Moldic: major*
 Intercrystalline: major
 Interparticle: minor-major
 Secondary intraparticle: minor

*Accurate percentages indeterminable.

Mean porosity (%): 16.7

Maximum porosity (%): 22.4

Minimum porosity (%): 12.3

Porosity standard deviation (%): 2.8

Number of analyses: 19

Geometric mean permeability (md): 2.11

Maximum permeability (md): 11.80

Minimum permeability (md): 0.40

Number of analyses: 19

Depositional Environment: Restricted lagoon.

Diagenesis: Minor early dolomitization, growth of lenticular gypsum in upper part, fair preservation of primary fabric.

Percent of reservoir: 41

Depth interval(s) (ft): 11,395-11,419

Lithofacies 2: *Favreina*-bearing small-pellet grainstone and *Favreina*-bearing small-pellet dolograinstone

Pore types:

Moldic: 43%
 Secondary intraparticle: 36%
 Intercrystalline: 8%
 Vuggy: 7%
 Fracture: 4%
 Cement moldic: 2%

Mean porosity (%): 12.7

Maximum porosity (%): 18.3

Minimum porosity (%): 7.5

Porosity standard deviation (%): 2.7

Table CH-2.-- Lithofacies description of core from Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama (see figure CH-7) -- Continued

Number of analyses: 17

Geometric mean permeability (md): 1.38

Maximum permeability (md): 7.50

Minimum permeability (md): 0.40

Number of analyses: 17

Depositional environment: Tidal flat.

Diagenesis: Pore system created by simultaneous dolomitization and leaching before deep burial. Fluids may have moved downward or laterally and may have been of mixed origin (mixing-zone dolomitization).

Percent of Reservoir: 33

Depth interval(s) (ft): 11,419-11,439.5

Lithofacies 3: Pellet packstone-grainstone

Pore Types:

Moldic: 100%

Mean porosity (%): 10.2

Maximum porosity (%): 13.9

Minimum porosity (%): 6.4

Porosity standard deviation (%): 2.1

Number of analyses: 11

Geometric mean permeability (md): 1.10

Maximum permeability (md): 3.00

Minimum permeability (md): 0.50

Number of analyses: 11

Depositional environment: Shallow lagoon, increasing restriction upward.

Diagenesis: Leaching and associated minor dolomitization created the pore system.

Percent of reservoir: 18

Depth interval(s) (ft): 11,439.5-11,450

Table CH-2.-- Lithofacies description of core from Bolinger 26-7 No. 1 well, Permit No. 2426,
Chappell Hill field, Alabama (see figure CH-7) -- Continued

Lithofacies 4: Very finely crystalline dolostone

Pore types:

Intercrystalline: 70%

Vuggy: 30%

Mean porosity (%): 20.5

Maximum porosity (%): 30.1

Minimum porosity (%): 14.8

Porosity standard deviation (%): 6.4

Number of analyses: 5

Geometric mean permeability (md): 24.38

Maximum permeability (md): 265.00

Minimum permeability (md): 2.20

Number of analyses: 5

Depositional environment: Tidal flat.

Diagenesis: Early, pervasive dolomitization.

Percent of reservoir: 8

Depth interval(s): 11,390-11,395

Figure CH-8.--Porosity vs. natural log permeability plots for lithofacies. Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama. (a) Lithofacies 1, (b) lithofacies 2, and (c) lithofacies 3. Equations for regression lines are printed for each lithofacies. (See table CH-2 for descriptions of lithofacies.)

Figure CH-9.--Paragenetic sequence, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama.

Figure CH-10.--Ternary diagram of pore types, Bolinger 26-7 No. 1 well, Permit No. 2426, Chappell Hill field, Alabama. Percentages of moldic pores indicated. Sample depths are indicated to nearest 0.1 foot.

CHATOM FIELD
SMACKOVER GAS AND CONDENSATE POOL
WASHINGTON COUNTY, ALABAMA

Figure CTM-1.--Index map of Chatom field, Alabama.

Figure CTM-2.--Structure contour map on top of Smackover Formation in Chatom field, Alabama. (Lowest known gas and highest known water levels taken from Phillips Petroleum Co., Exhibit 1A, Docket Nos. 5-17-881 through 5-17-883, State Oil and Gas Board of Alabama.)

Figure CTM-3.--Well log correlated to lithology, porosity, and permeability, Chatom Unit 20-14
No. 1-04 well, Permit No. 7044, Chatom field, Alabama.

Figure 41.--Structural cross section of Chatom field, Alabama. Highest known water and lowest known gas taken from Phillips Petroleum Co., Exhibit 1A, Docket No. 5-17-881 through 5-17-883, State Oil and Gas Board of Alabama. (See figure 40 for line of section.)

Figure CTM-5.--Graph of production history of the Chatom field, Alabama.

Figure CTM-6.--Porosity vs. natural log permeability plots, Chatom field, Alabama. (a) Plot of five cores combined, and (b) plot of described core for Chatom Unit 20-14 No. 1-04 well, Permit No. 7044. Equations apply to regression lines.

Table CTM-1--Reservoir characterization sheet, Smackover pool,
Chatom field, Washington County, Alabama

Discovery well

Operator: Phillips Petroleum Co.
Permit number: 1543
Well Name: Williams "AA" #1
Location: Sec. 20, T6N-R4W
Completion date: 08/12/70
Perforations: 15,999-16,114 ft
Initial flow rate: 988 bcpd, 4,029 Mcfd,
24/64 ck, 1,933 psi

- **Field data (as of 12/31/90)**

Number of wells within field limits: 14
Producing wells in pool: 6
Production wells in field: 6
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 3
Salt water disposal wells: 1
Enhanced recovery wells: 2
Dry holes: 2
Drilling wells: 0
Unit spacing (acres): Unitized
Total field area (acres): 7,680
Cumulative production
Condensate (bbl): 14,212,601
Gas (Mcf): 109,780,337
Water (bbl): 963,358

Reservoir data

Trap type: Structural

Reservoir structure: Anticline

Proven productive area (acres): 2,080

Average net pay thickness (feet): 56

Hydrocarbon type: Gas condensate

Oil gravity (API): 55

Initial reservoir pressure (psia): 7,903

Current reservoir pressure (psia): na

Reservoir drive: Depletion

Type of enhanced recovery: Gas injection

Mean porosity (core) (%): 14.70

Porosity standard deviation (%): 10.20

Maximum porosity (%): 37.20

Minimum porosity (%): 0.13

Number of porosity analyses: 331

Geometric mean perm (core) (md): 1.21

Maximum permeability (md): 196.00

Minimum permeability (md): 0.02

Number of permeability analyses: 331

Dykstra-Parsons coefficient: na

Figure CTM-7.--Detailed core log, Chatom Unit 20-14 No. 1 1-04, Permit No. 7044, Chatom field, Alabama.

Table CTM-2.--Lithofacies description of core from Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama (see figure CTM-7)

Lithofacies 1: Pelmoldic and crystalline dolostone

Pore types:

Intercrystalline: 62%
Moldic: 22%
Secondary intraparticle: 16%

Mean porosity (%): 18.0

Maximum porosity (%): 22.9

Minimum porosity (%): 8.7

Porosity standard deviation (%): 4.4

Number of analyses: 11

Geometric mean permeability (md): 2.83

Maximum permeability (md): 6.58

Minimum permeability (md): 0.55

Number of analyses: 11

Depositional environment: Shallowing upward shallow subtidal lagoon to ooid shoal.

Diagenesis: Dolomitization and leaching, porosity occlusion by anhydrite and calcite cements.

Percent of reservoir: 62

Depth interval(s) (ft): 16,073-16,086

Lithofacies 2: Pellet dolograinstone

Pore types:

Moldic: 90%
Intercrystalline: 10%

Mean porosity (%): 28.4

Maximum porosity (%): 37.2

Minimum porosity (%): 8.7

Porosity standard deviation (%): 10.6

Number of analyses: 8

Geometric mean permeability (md): 11.13

Maximum permeability (md): 88.00

Minimum permeability (md): 0.02

Number of analyses: 8

Table CTM-2.--Lithofacies description of core from Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama (see figure CTM-7) -- Continued

Depositional environment: Microtidal flat.

Diagenesis: Dolomitization and leaching.

Percent of reservoir: 38

Depth interval(s) (ft): 16,018-16,025.5

Figure CTM-8.--Porosity vs. natural log permeability plots for two lithofacies, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama. (a) Lithofacies 1, and (b) lithofacies 2. Equations for regression lines are printed for each lithofacies. (See table CTM-2 for descriptions of lithofacies.) Data set excludes all samples with porosity values and permeability values below reservoir cutoffs at 6% and 0.1 md, respectively.

Figure CTM-9.--Paragenetic sequence, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama.

Figure CTM-10.--Ternary diagram of pore types, Chatom Unit 20-14 No. 1-04 well, Permit No. 7044, Chatom field, Alabama. Percentages of moldic pores indicated. Sample depths are indicated to nearest 0.1 foot.

CHOCTAW RIDGE FIELD
SMACKOVER OIL POOL
CHOCTAW COUNTY, ALABAMA

Figure CHR-1.--Index map of Choctaw Ridge field, Alabama.

Figure CHR-2.--Structure contour map on top of Smackover Formation in Choctaw Ridge field, Alabama.

Figure CHR-3.--Well log correlated to lithology, Rentz-Trice Unit 35-1 well, Permit No. 1427, Choctaw Ridge field, Alabama.

Figure CHR-4.--Structural cross section of the Choctaw Ridge field, Alabama. (See figures CHR-1 and CHR-2 for line of section.)

Figure CHR-5.--Graph of production history of the Choctaw Ridge field, Alabama.

Figure CHR-6.--Porosity vs. natural log permeability plot of five cores combined, Choctaw Ridge field, Alabama. Equation applies to regression line.

Table CHR-1.--Reservoir characterization sheet, Smackover pool,
Choctaw Ridge field, Choctaw County, Alabama

Discovery well

Operator: Pruet & Hughes Co.
Permit number: 1413
Well Name: Trice #1
Location: Sec. 26, T11N-R4W
Completion date: 06/19/67
Perforations: 11,940-11,952 , 11,963-11,964 ft
Initial flow rate: 840 bopd, 558 Mcfd,
1/4 ck, 1,500 psi

Field data (as of 12/31/90)

Number of wells within field limits: 12
Producing wells in pool: 3
Production wells in field: 3
Shut-in/Temp. abandoned wells: 0
Plugged and abandoned wells: 5
Salt water disposal wells: 1
Enhanced recovery wells: 0
Dry holes: 3
Drilling wells: 0
Unit spacing (acres): Unitized
Total field area (acres): 1,040
Cumulative production
Oil (bbl): 3,608,435
Gas (Mcf): 1,853,154
Water (bbl): 74,593

Reservoir data

Trap type: Structural
Reservoir structure: Faulted anticline
Proven productive area (acres): 549
Average net pay thickness (feet): 22
Hydrocarbon type: Oil
Oil gravity (API): 41
Initial reservoir pressure (psia): 5,650
Current reservoir pressure (psia): na
Reservoir drive: Solution gas / water
Type of enhanced recovery: none
Mean porosity (core) (%): 19.39
Porosity standard deviation (%): 4.95
Maximum porosity (%): 31.10
Minimum porosity (%): 7.50
Number of porosity analyses: 87
Geometric mean perm (core) (md): 21.44
Maximum permeability (md): 1410.00
Minimum permeability (md): 1.90
Number of permeability analyses: 87
Dykstra-Parsons coefficient: na

Figure CHR-7.--Detailed core log, Rentz-Trice Unit 35-1 well, Permit No. 1427, Choctaw Ridge field, Alabama.

Table CHR-2.--Lithofacies description of core from Rentz-Trice Unit 35-1 No. 5 well, Permit No. 1427, Choctaw Ridge field, Alabama (see figure CHR-7)

Lithofacies 1: Oolitic and pelletal dolomitic grainstone and dolograinstone

Pore types:

Moldic: 61%

Secondary Intraparticle: 36%

Interparticle: 3%

Depositional Environment: High-energy subtidal shoal.

Diagenesis: Minor penecontemporaneous gypsum formation was followed by interparticle calcium-carbonate cementation. Next, partial fabric-selective dolomitization and particle dissolution created the secondary pore system that dominates the reservoir. Several episodes of fracturing enhanced reservoir permeability. Stylolitization, anhydrite replacement and cementation, and dedolomitization also affected the reservoir.

Percent of reservoir: 100

Depth interval(s) (ft): 11,969-11,996

Figure CHR-8.--Paragenetic sequence, Rentz-Trice Unit 35-1 No. 1 well, Permit No. 1427, Choctaw Ridge field, Alabama.

CHUNCHULA FIELD
SMACKOVER GAS AND CONDENSATE POOL
MOBILE COUNTY, ALABAMA

Figure CHU-1.--Index map of Churchill field, Alabama.

Figure CHU-2.--Structure contour map on top of Smackover Formation in Churchula field, Alabama.

Figure CHU-3.--Well log correlated to lithology, porosity, and permeability, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Churchula field, Alabama.

Figure CHU-4.--Structural cross section of the Chunchula field, Alabama. (See figures CHU-1 and CHU-2 for line of section.)

Figure CHU-5.--Graph of production history of the Chunichula field, Alabama.

Figure CHU-6.--Porosity vs. natural log permeability plot, Churchula field, Alabama. (a) Plot of 65 cores combined, and (b) plot of described core for Annie M. Hill et al. No. 1 well, Permit No. 2218. Equations apply to regression lines.

Trap type: Structural / stratigraphic

Reservoir structure: Anticline

Proven productive area (acres): 22,113

Average net pay thickness (feet): 34

Hydrocarbon type: Oil

Oil gravity (API): 60

Initial reservoir pressure (psia): 9,255

Current reservoir pressure (psia): 4,317

Reservoir drive: Depletion

Type of enhanced recovery: Gas injection

Mean porosity (core) (%): 10.90

Porosity standard deviation (%): 4.60

Maximum porosity (%): 26.20

Minimum porosity (%): 1.10

Number of porosity analyses: 2,396

Geometric mean perm (core) (md): 0.65

Maximum permeability (md): 392.00

Minimum permeability (md): 0.02

Number of permeability analyses: 2,396

Dykstra-Parsons coefficient: na

Figure CHU-7.--Detailed core log, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Churchula field, Alabama.

Table CHU-2.--Lithofacies description of core from Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Chunchula field, Alabama (see figure CHU-7)

Lithofacies 1: Crystalline dolostone, locally anhydritic

Pore types:

Intercrystalline: 99%

Moldic: 1%

Mean porosity (%): 9.9

Maximum porosity (%): 14.6

Minimum porosity (%): 3.1

Porosity standard deviation (%): 3.9

Number of analyses: 15

Geometric mean permeability (md): 1.04

Maximum permeability (md): 5.90

Minimum permeability (md): 0.09

Number of analyses: 15

Depositional Environment: Shallow subtidal marine and/or peritidal hypersaline.

Diagenesis: Early growth of anhydrite nodules and replacement laths, pervasive dolomitization that was not controlled by pre-existing rock fabric.

Percent of reservoir: 100

Depth interval(s) (ft): 18,444-18,508

Figure CHU-8.--Porosity vs. natural log permeability plot for lithofacies 1, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Churchula field, Alabama. Equation applies to regression line. (See table CHU-2 for description of lithofacies.) Data set excludes all samples with porosity values and permeability values below reservoir cutoffs at 6% and 0.1 md, respectively.

Figure CHU-9.--Paragenetic sequence, Annie M. Hill et al. 10-10 No. 1 well, Permit No. 2218, Churchula field, Alabama.

Figure CHU-10.--Ternary diagram of pore types, samples from several wells, Churchula field, Alabama. Percentages of moldic pores indicated.

COLD CREEK FIELD
SMACKOVER OIL POOL
MOBILE COUNTY, ALABAMA

Figure CC-1.--Index map of Cold Creek field, Alabama.

Figure CC-2.--Structure contour map on top of Smackover Formation in Cold Creek field, Alabama.

Figure CC-3.--Well log correlated to porosity and permeability, C. P. Newman et al. 8-16 No. 1 well, Permit No. 3027, Cold Creek field, Alabama.

Figure CC-4.--Structural cross section of the Cold Creek field, Alabama. (See figures CC-1 and CC-2 for line of section.)

Figure CC-5.--Graph of production history of the Cold Creek field, Alabama.

Figure CC-6.--Porosity vs. natural log permeability plots, Cold Creek field, Alabama. (a) Plot of three cores combined, and (b) plot of described core for C. P. Newman et al. 8-16 No. 1 well, Permit No. 3027. Equations apply to regression lines.

Table CC-1.--Reservoir characterization sheet, Smackover pool,
Cold Creek field, Mobile County, Alabama

Discovery well

Operator: Union Oil Co. of California

Permit number: 2461

Well Name: R.J. Newman et al 9-11 #1

Location: Sec. 9, T1S-R1W

Completion date: 04/18/78

Perforations: 18,425-18,443 ft

Initial flow rate: 500 bopd, 320 Mcfd,
14/64 ck, 810 psi

Field data (as of 12/31/90)

Number of wells within field limits: 3

Producing wells in pool: 2

Production wells in field: 2

Shut-in/Temp. abandoned wells: 0

Plugged and abandoned wells: 1

Salt water disposal wells: 0

Enhanced recovery wells: 0

Dry holes: 0

Drilling wells: 0

Unit spacing (acres): 160

Total field area (acres): 480

Cumulative production

Oil (bbl): 999,846

Gas (Mcf): 1,111,743

Water (bbl): 3,499

Reservoir data

Trap type: Structural / stratigraphic

Reservoir structure: Anticline

Proven productive area (acres): 625

Average net pay thickness (feet): 20

Hydrocarbon type: Oil

Oil gravity (API): 59

Initial reservoir pressure (psia): 8,911

Current reservoir pressure (psia): 4,827

Reservoir drive: Solution gas

Type of enhanced recovery: na

Mean porosity (core) (%): 4.25

Porosity standard deviation (%): 3.51

Maximum porosity (%): 25.50

Minimum porosity (%): 0.50

Number of porosity analyses: 512

Geometric mean perm (core) (md): 0.26

Maximum permeability (md): 327.00

Minimum permeability (md): 0.01

Number of permeability analyses: 201

Dykstra-Parsons coefficient: na

COPELAND FIELD
SMACKOVER GAS AND CONDENSATE POOL
WASHINGTON COUNTY, ALABAMA

Figure CPL-1.--Index map of Copeland field, Alabama.

Figure CPL-2.--Structure contour map on top of Smackover Formation in Copeland field, Alabama

Figure CPL-3.--Well log, Edith Burch 30-13 No. 1 well, Permit No. 3247, Copeland field, Alabama.

Figure CPL-4.--Structural cross section of the Copeland field, Alabama. (See figures CPL-1 and CPL-2 for line of section.)

Figure CPL-5.--Graph of production history of the Copeland field, Alabama.

Figure CPL-6.--Porosity vs. natural log permeability plot of one core, Copeland field, Alabama. Equation applies to regression lines.

Table CPL-1.--Reservoir characterization sheet, Smackover pool,
Copeland field, Washington County, Alabama

Discovery well

Operator: Amoco Production Co.

Permit number: 1670

Well Name: J.J. Laurie Unit #1

Location: Sec. 6, T6N-R4W

Completion date: 02/10/72

Perforations: 16,614-16,630, 16,800-16,835 ft

Initial flow rate: 31 bcfd, 1,519 Mcfd,
30/64 ck, 800 psi

Field data (as of 12/31/90)

Number of wells within field limits: 4

Producing wells in pool: 2

Production wells in field: 2

Shut-in/Temp. abandoned wells: 1

Plugged and abandoned wells: 0

Salt water disposal wells: 0

Enhanced recovery wells: 0

Dry holes: 1

Drilling wells: 0

Unit spacing (acres): 640

Total field area (acres): 6,240

Cumulative production

Condensate (bbl): 813,586

Gas (Mcf): 13,993,943

Water (bbl): 1,131

Reservoir data

Trap type: Structural

Reservoir structure: Anticline

Proven productive area (acres): 1,703

Average net pay thickness (feet): 14

Hydrocarbon type: Gas condensate

Oil gravity (API): 41

Initial reservoir pressure (psia): 7,899

Current reservoir pressure (psia): 3,015

Reservoir drive: Depletion

Type of enhanced recovery: none

Mean porosity (core) (%): 15.40

Porosity standard deviation: 5.74

Maximum porosity (%): 26.40

Minimum porosity (%): 6.70

Number of porosity analyses: 34

Geometric mean perm (core) (md): 1.80

Maximum permeability (md): 52.00

Minimum permeability (md): 0.06

Number of permeability analyses: 34

Dykstra-Parsons coefficient: na

CROSBYS CREEK FIELD
SMACKOVER GAS AND CONDENSATE POOL
WASHINGTON COUNTY, ALABAMA

Figure CRC-1.--Index map of Crosbys Creek field, Alabama.

Figure CRC-2.--Structure contour map on top of Smackover Formation in Crosbys Creek field, Alabama.

Figure CRC-3.--Well log, Sarah Middleton 34-11 No. 1 well, Permit No. 5001, Crosbys Creek field, Alabama.

Figure CRC-4--Structural cross section of the Crosbys Creek field, Alabama.
(See figures CRC-1 and CRC-2 for line of section.)

Figure CRC-5.--Graph of production history of the Crosbys Creek field, Alabama.

Figure CRC-6.--Porosity vs. natural log permeability plot of two cores combined, Crosbys Creek field, Alabama. Equation applies to regression line.

Table CRC-1.--Reservoir characterization sheet, Smackover pool,
Crosbys Creek field, Washington County, Alabama

Discovery well

Operator: Hughes Eastern Petroleum Ltd.

Permit number: 5001

Well Name: Sarah Middleton 34-11 #1

Location: Sec. 34, T8N-R4W

Completion date: 10/26/88

Perforations: 16,394-16,420 ft

Initial flow rate: 823 bc/d, 2,621 Mc/d,
20/64 c/k, 1,975 psi

Field data (as of 12/31/90)

Number of wells within field limits: 3

Producing wells in pool: 2

Production wells in field: 2

Shut-in/Temp. abandoned wells: 0

Plugged and abandoned wells: 0

Salt water disposal wells: 0

Enhanced recovery wells: 0

Dry holes: 1

Drilling wells: 0

Unit spacing (acres): 640

Total field area (acres): 1,280

Cumulative production

Condensate (bbl): 688,145

Gas (Mcf): 1,328,588

Water (bbl): 2,150

Reservoir data

Trap type: Structural

Reservoir structure: Anticline

Proven productive area (acres): 486

Average net pay thickness (feet): 22

Hydrocarbon type: Gas condensate

Oil gravity (API): na

Initial reservoir pressure (psia): 7,710

Current reservoir pressure (psia): 5,387

Reservoir drive: Depletion

Type of enhanced recovery: none

Mean porosity (core) (%): 21.40

Porosity standard deviation (%): 9.20

Maximum porosity (%): 33.70

Minimum porosity (%): 6.30

Number of porosity analyses: 58

Geometric mean perm (core) (md): 0.59

Maximum permeability (md): 17.00

Minimum permeability (md): 0.10

Number of permeability analyses: 58

Dykstra-Parsons coefficient: na

Figure CRC-7.--Detailed core log, Daryl Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama.

Table CRC-2.--Lithofacies description of core from Darly Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama (see figure CRC-7)

Lithofacies 1: Pellet dolograinstone

Pore types:

Moldic: 93%

Interparticle: 7%

Depositional Environment: High-energy shoreface and/or beach. The abundance of low-angle cross stratification suggests either beach laminae or hummocky cross stratification; the presence of abundant trough cross strata and of local fenestrae suggests beach deposition.

Diagenesis: Early (marine?) pore-rimming cementation, pervasive fabric-selective dolomitization, leaching of unstable particles. Subsequent minor fracturing, calcite cementation, anhydrite nodule growth (replacive), stylolitization.

Percent of reservoir: 100

Depth interval(s) (ft): 16,381-16,407.5

Figure CRC-8.--Paragenetic sequence, Darly Dees 3-3 No. 1 well, Permit No. 5115, Crosbys Creek field, Alabama.

EAST BARNETT FIELD
SMACKOVER OIL POOL
CONECUH AND ESCAMBIA COUNTIES, ALABAMA

Figure EB-1.--Index map of East Barnett field, Alabama.

Figure EB-2.--Structure contour map on top of Smackover Formation in East Barnett field, Alabama.

Figure EB-3.--Well log correlated to lithology, Scott Paper Co. et al.
1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama.

Figure EB-4.--Structural cross section of the East Barnett field, Alabama. (See figures EB-1 and EB-2 for line of section.)

Figure EB-5.--Graph of production history of the East Barnett field, Alabama.

Table EB-1.--Reservoir characterization sheet, Smackover pool,
East Barnett field, Conecuh County, Alabama

Discovery well

Operator: Coastal Oil & Gas Corp.
 Permit number: 5568
 Well Name: Grissett 36-16 #1
 Location: Sec. 36, T4N-R7E
 Completion date: 04/27/88
 Perforations: 13,580-13,659 ft
 Initial flow rate: 606 bopd, 1,022 Mcfd,
 12/64 ck

Field data (as of 12/31/90)

Number of wells within field limits: 3
 Producing wells in pool: 3
 Production wells in field: 3
 Shut-in/Temp. abandoned wells: 0
 Plugged and abandoned wells: 0
 Salt water disposal wells: 0
 Enhanced recovery wells: 0
 Dry holes: 0
 Drilling wells: 0
 Unit spacing (acres): 160
 Total field area (acres): 480
 Cumulative production
 Oil (bbl): 565,798
 Gas (Mcf): 626,177
 Water (bbl): 107,003

Reservoir data

Trap type: Structural
 Reservoir structure: Anticline
 Proven productive area (acres): 337
 Average net pay thickness (feet): 58
 Hydrocarbon type: Oil
 Oil gravity (API): 45.1
 Initial reservoir pressure (psia): 6,589
 Current reservoir pressure (psia): 6,222
 Reservoir drive: Solution gas
 Type of enhanced recovery: none
 Mean porosity (core) (%): na
 Porosity standard deviation (%): na
 Maximum porosity (%): na
 Minimum porosity (%): na
 Number of porosity analyses: na
 Geometric mean perm (core) (md): na
 Maximum permeability (md): na
 Minimum permeability (md): na
 Number of permeability analyses: na
 Dykstra-Parsons coefficient: na

Figure EB-6.--Detailed core log, Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama.

Table EB-2.--Lithofacies description of core from Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama (see figure EB-7)

Lithofacies 1: Idiomatic dolostone (planar-e dolostone)

Pore types:

Intercrystalline: 83%

Moldic: 15%

Vuggy: 1%

Secondary Intraparticle: 1%

Depositional Environment: Shallow marine and (?) peritidal.

Diagenesis: Pervasive, nonselective dolomitization; minor anhydrite replacement.

Percent of reservoir: 100

Depth interval(s) (ft): 13,656-13,707.5

Figure EB-7.--Paragenetic sequence, Scott Paper Co. 1-1 No. 1 well, Permit No. 5740, East Barnett field, Alabama.

END

**DATE
FILMED**

7 / 12 / 93