DISCLAIMER

DOE/ER/61035--T1

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DOE/ER/61035--T1

DE92 019048

**FINAL TECHNICAL REPORT
TO THE HEALTH EFFECTS RESEARCH DIVISION
OFFICE OF HEALTH AND ENVIRONMENTAL RESEARCH
U.S. DEPARTMENT OF ENERGY
(UCD 472-504)**

DOE Grant DE-FG03-890-ER61035

**Coordination Chemistry of the $^{212}\text{Pb}/^{212}\text{Bi}$ Nuclear
Transformation:
Alpha-Emitting Radiopharmaceuticals**

Norris J. Parks, Principal Investigator
Wesley R. Harris, Co-investigator
Carl L. Keen, Co-investigator
Stephan R. Cooper, Co-investigator

Received by OSTT

AUG 17 1992

Laboratory for Energy-related Health Research Division
Institute of Toxicology and Environmental Health
University of California
Davis, California 95616

July, 1992

MAILED

OK

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

FINAL TECHNICAL REPORT
TO THE
HEALTH EFFECTS RESEARCH DIVISION
OFFICE OF HEALTH AND ENVIRONMENTAL RESEARCH
U.S. DEPARTMENT OF ENERGY
DOE Grant DE-FG03-890-ER61035
July 1992

**Coordination Chemistry of the $^{212}\text{Pb}/^{212}\text{Bi}$ Nuclear
Transformation:
Alpha-Emitting Radiopharmaceuticals**

Introduction

This project has attracted some extremely capable men and women as undergraduate and postgraduate scholars whose enthusiastic efforts moved various subdivisions of a complex problem forward in parallel. These individuals included collaborating scientists in the Chemistry department at U. C. Davis (W. K. Musker), two post doctoral fellow (P. K. Bharadwaj and Cheri Zidenberg-Cherr) and several undergraduate researchers who have gone on to graduate or professional school and who co-authored at least one publication.

The subdivisions were (a) the synthesis of prototypical thiolate and dithiocarbamate based hexacoordinate complexes, (b) radiochemical engineering for generation of no-carrier-added lead and bismuth radioelements, (c) the first isolation of bismuth-binding proteins from in vivo studies with cyclotron produced $^{205}/^{206}\text{Bi}$ tracer, and (d) initial development of transport mechanisms for the intracellular radiobiological study of alpha emitting bismuth, and (e) the initiation of

chemical equilibrium studies and biochemical pathways with cyclotron-produced, no-carrier-added, ^{203}Pb ($T_{1/2} = 51\text{hr}$).

A series of nuclear reaction cross-section measurements were made to evaluate the production of $^{203}/^{205}/^{206}\text{Bi}$ from natural lead via p,xn reactions (Lagunas-Solar et al. 87). Production of $^{205}/^{206}\text{Bi}$ ($T_{1/2} = 15.3$ and 6.2 days, respectively) over the proton energy range of 20 to 40 MeV was greater than theoretically expected. Consequently, chemically and biologically useful quantities of $^{205}/^{206}\text{Bi}$ were produced in pilot runs. Automation of this targetry system and subsequent radiochemical separations is being improved because it reduces both personnel exposure and the number of radiochemists required to safely perform the separation. By appropriate timing of the anion exchange separations now being employed, ^{203}Pb can also be extracted as essentially no-carrier-added (NCA) material because it is obtained after the initial Pb/Bi separation as a decay product of ^{203}Bi . ^{203}Pb and the various bismuth radionuclides are readily detectable gamma-emitters. They have been the mainstay of the developmental chemical and biological kinetic experiments in addition to revealing the radioanalytical problems associated with NCA Bi under physiological conditions. The originally planned ^{232}U based generator for ^{212}Pb and ^{212}Bi has been superseded by a ^{228}Th system. The system provides for a carrier gas to remove ^{220}Rn ($T_{1/2} = 51\text{sec}$) from the source solution and a flow through a bubbler train at sufficiently low flow rates that ^{212}Pb ($T_{1/2} = 10\text{hr}$) ions are formed and captured in the solvent or reaction media of our choice. With this generator system two phase distribution experiments can be performed in situ. One of the critical tasks of this project was to invent specific experimental designs by which the heretofore unstudied nucleogenic reactions of NCA ^{212}Bi could be unequivocally characterized.

After extensive evaluation by the PI's and collaborators, we decided that the most information content per experiment (i.e., cost-effective) would emanate from a series of competitive binding experiments in organic-aqueous two-phase systems. The candidate complexes of interest are being synthetically tailored by attachment of polar substituents to be either hydrophilic or hydrophobic. This provided us with

- (1) the flexibility to control the chemical properties and electronic character of the "cage" wall surrounding the excited molecule undergoing the nuclear transformation;
- (2) to employ both hydrophilic and hydrophobic scavengers of thermalized Bi ions that have escaped the parent complex by virtue of "hot" or epithermal ion-molecule reaction (the objective here is to measure the extent to which Bi or bismuthyl ions might displace parent ^{212}Pb and thus confound the measurement of ^{212}Bi intramolecular retention);
- (3) to begin early on to examine the results of the nucleogenic reactions in solutions held at physiological pH and containing the naturally occurring ligands for Bi that have been identified in the cellular biochemistry work.

We initiated studies on the chelating abilities of 1,1-dithio ligands. These ligands were found to be very attractive on two accounts: 1) they can strongly bind both Pb and Bi in aqueous media in a wide pH range, and 2) they can bind Pb and Bi discriminately against essential transition metals (Bharadwaj et al. 87,88). We have extended this line of synthesis and were able to prepare ligands containing two or more

dithiogroups per ligand. These ligands also bind strongly both Pb and Bi and serve as an alternative to tri-succinyl substituted compounds originally proposed. Synthesis of succinyl substituted compounds were in progress.

In the area of radiochemical development, we improved our ability to generate no-carrier-added $^{212}\text{Pb}/^{212}\text{Bi}$ reagents that are free from possible contamination by ion exchange columns and have the highest potential specific activity. We can now use precipitated ^{228}Th colloids supported in filter assemblies through which carrier gases of choice transport the ^{220}Rn to liquid nitrogen traps. We developed new emulsion scintillation spectroscopy methods (Avila 1992) for mixed alpha and beta particle emitters and in particular, radiochemical assays of ^{212}Pb (10.6h) and ^{212}Bi (1.01h).

Later, we initiated an additional phase which added three types of experiments to what we are already doing. One was the direct testing of the experimental methods which had been developed to study the intramolecular dynamics of in situ ^{212}Bi formation by nucleogenesis. Because these reactions involved quantities of material that were 10 orders of magnitude lower in concentration than typically encountered in macroscopic studies, we started with the general methods developed for chemical effects of nuclear transformations (CENT) and "hot-atom chemistry" research. We found that the commonly available radio-high performance liquid chromatograph (RPLC) system built of stainless-steel, glass or titanium components were reactive with our no-carrier-added (NCA) materials. This made evaluation of the nucleogenic chemistry complicated at best, impossible at worst. We sought a workable, cost-effective analytical method that still had the capability of providing information about the presence of different elemental and chemical species. Our interest in

such separations stemmed from a variety of limitations of the two-phase, competitive binding experiments. This effort had two components. One was work with inorganic ion and coordination complex electrophoresis which has useful separating properties but is too slow for some objectives to be readily achieved. Radioassay scanners were being assembled and improved this situation. The other was the ongoing design and installation of a radio HPLC system constructed entirely of fluorocarbon polymer material of the general type that solved several separation problems with NCA quantities of labelled material.

The second experiment added was the cellular radiobiology of alpha particles. This was performed with cell culture systems and the use of fluorescent labelled liposomes (Pikul 87) which carry the $^{212}\text{Pb}/^{212}\text{Bi}$ into the cell; this, in turn, can lead to precise quantitation, via fluorescent microscopy, of alpha energy deposition vs. response distribution for a given cell population. We have a melanoma model in vitro and in vivo for this work. This radiobiology work provides the benchmark against which the efficacy of radiotherapeutic alpha emitter molecules will be tested. We can now also commence in vivo studies of the chemical competition between bismuth binding proteins (and other naturally occurring ligands) and synthetic ligands designed to retain both parent lead and daughter ^{212}Bi after the nuclear transformation.

The third experiment added was in vivo studies of radiobiological aspects of ^{212}Bi radiopharmaceuticals. The whole-body clearance, organ distribution, and subcellular distribution of no-carrier-added and carrier-added intraperitoneally administered bismuth radio tracer ($^{205}/^{206}\text{Bi}$) has been determined in Sprague-Dawley rats. Differences in clearance rate kinetics were observed for this study

with administration of neutral solutions of tracer in carbonate buffer as compared to other studies with other chemical forms. The final organ distribution was not strongly dependent on administered chemical form. We provided definitive evidence that bismuth does indeed enter subcellular organelles such as the nucleus and mitochondria which had 30-50% and 10-25%, respectively, of activity in kidney tissue (Zidenberg- Cherr 87,89). The kidneys were the main sink for radiotracer with uptake ranging from 20-50% of total body activity. The calculated energy deposition by recoil nuclei after alpha emission of potentially therapeutically useful ^{212}Bi was found to equal or exceed the alpha energy deposition per organelle if the source is inside the cell nucleus or mitochondria.

Nucleogenic Reactions

We obtained evidence consistent with the hypothesis that nucleogenic intramolecular reaction caused by the nuclear transformation of ^{212}Pb to ^{212}Bi within a coordination complex can occur without decomposition. Using short, 1-2cm ion exchange columns and atmospheric pressure eluant systems, we found that metal complexes of proposed substituents such as dimercaptosuccinic acid (DMSA) of a flexible backbone (typically polyamines), dithiocarboxylates (Bharadwaj and Musker 87), or the tripodal thiaethers released only 40 to 60% of the ^{212}Bi formed by ^{212}Pb decay. This was inferred from retention of a positively charged bismuth, species being on the column and passage of a neutral species; the latter being consistent with the successful formation and survival of a neutral radiobismuth complex. Electrophoresis separations of charged ^{212}Pb macrobicyclic kryptofix complexes and uncomplexed ^{212}Pb and ^{212}Bi lead to the conclusion that this type of complex (with only low electron-density

single sigma bonds) was decomposing at least 80% of the time. No macroscopic bismuth complex of this molecule has ever been formed. If we can verify that at least a small fraction of the fragile ^{212}Bi macrobicyclic cage complex forms nucleogenically, it will help to calibrate the low end of the intramolecular energy range with which we are dealing. With rapid radio HPLC using multiphase ion chromatography and the ability to monitor macroscopic free ligand and complex concentration either in separation processes or equilibrium measurements, we have moved to the next plateau of experimental inquiry. In particular, the extent of thermal reactions that might confound interpretation of the chemical effects of ^{212}Pb to ^{212}Bi nuclear transformation may be quantitated.

Coordination Chemistry of the $^{212}\text{Pb}/^{212}\text{Bi}$ Nuclear Transformation: Alpha Emitting Radiopharmaceuticals

Requirements for a Pb Chelator. The absence of systematic study of Pb(II) coordination chemistry forces effective design of chelators to rely upon broad-based principles for guidance. The hard-soft acid-base theory of Basolo and Pearson suggests that a soft acid cation such as Pb(II) should prefer coordination through soft base donor groups. The ideal chelating agent would combine chemical robustness with low affinity (as measured by the stability constant, K) for biologically important metal ions (e.g., Ca(II) , Mg(II) , Zn(II)).

Thiolates and thioethers are two obvious possibilities. Thiolates have two serious disadvantages: 1) they tend to form coordination polymers and 2) they undergo oxidation to disulfides; consequently we chose initially to evaluate thioethers as donor groups.

First Approach. Motivation for the first generation of ligands came from earlier work on crown thioethers. To accomodate the large Pb(II) ion (ionic radius = 1.19Å), we used the route indicated in Figure 3 to synthesize the previously unreported crown thioether, 24S6. This ligand showed low affinity for Pb(II); crystallographic work (Rawle et al. 86) showed that it handily envelopes the much smaller Ni(II) ion at nearly optimal Ni-S distances. This result suggested that perhaps the cavity size was inappropriate for Pb(II).

Second Approach. Rather than pursue the cavity size question on an empirical basis, we decided to evaluate the intrinsic affinity of Pb(II) for hexakis(thioether) coordination through use of a less stereochemically demanding ligand. We synthesized the previously unreported supertripodal ligand .

The advantages of this class of ligands transcend questions regarding thioether coordination. The facile synthesis of this three-dimensionally articulated ligand provided a general route into pockets that incorporate a wide variety of donor groups. In addition, replacement of the apical methyl group by e.g., oligomethylene sulfonate, would confer water solubility on the ligand- a potentially crucial capability,

In any event, the supertripodal hexakis(thioether) ligand displayed a rich coordination chemistry with a variety of transition metal ions (Thorne et al. 86,87), but little affinity for Pb(II). Consequently, thioethers appeared to be intrinsically ill-suited as ligands for Pb(II).

This low affinity might arise because thioethers fail to neutralize the charge on the lead cation. Most of the Pb(II) complexes in the literature are either neutral or monoanionic.

Third Approach. This realization suggested that the ligands should incorporate some (at most three) anionic groups. To do this within the context of the supertripodal approach (to retain the advantages summarized above), we designed some new ligands

These ligands comprise sterically encumbered tertiary thiolate groups in an attempt to shut down polymer formation by preventing approach of a second Pb(II) ion. This steric protection could also reduce the ease of oxidation of the thiolates by adventitious metal ions (e.g., Cu(II)).

Additional work - Synthesis of Third Generation Ligands. Continuing work focuses on synthesis of the triamino and trithioether trithiolates. We have recently prepared the latter target molecule, but we are now trying to overcome the indifferent regioselectivity of the opening of the thiirane by the thiolate nucleophile. Reduction in temperature mitigates, but does not remove, this problem which is currently under investigation.

Radio HPLC

We assembled a radio HPLC system designed entirely of fluorocarbon polymer material. To this system we have attached three different detectors: (1) We installed a Waters 990 Photodiode Array (PDA) UV-VIS detector system. This system was used to directly analyze the effluent from an HPLC and has advanced data processing capability. The PDA can simultaneously measure absorbance values from 190 to 600 nm and produce a 3D display representing time, absorbance, and wavelength. This detector is capable of making quantitative measurements of the ligands that are being studied.

(2) We also use an IBM-PC based pulse-height analyzer for gamma-ray spectrometry with a NaI(Tl) detector array. The Nucleus Personal

Computer Analyzer (PCA) is a multichannel scalar. This system allows us to directly and simultaneously measure gamma-ray emissions from lead, bismuth or other radionuclides, thereby identifying them. The PCA system measures effluent directly from the HPLC or the PDA. Plans are to display gamma-ray spectra in the same format as the UV data from the PDA, where this is useful for following the various nucleogenic reaction of ^{212}Pb and short-lived daughters.

(3) The third detector system was a Packard Trace II Model 7150 Liquid Scintillation Radioactivity Flow Monitor (RFM). This is a radioactivity flow monitor primarily intended for the detection of low energy beta emitting radionuclides in the effluent from a high performance liquid chromatograph. However, the detector geometry is a small bore (approx. 0.03in.) coil of tubing between two phototubes. The coil contains HPLC effluent mixed with scintillation cocktail. Our previous work with liquid scintillation beta spectroscopy showed that this arrangement should permit detection efficiency for alphas approaching 100% and facile correction for beta background if the signal output from the RFM is routed to the pulse height analyzer. This capability derives from fact that most of the energy for energetic betas will be deposited outside the scintillator containing coil and the short 40-80 micrometer range alphas will deposit most of their energy inside the coil. In the standard configuration for this system, two counting channels can be monitored, thereby allowing us to detect both single and dual label systems. This ability permits us to monitor both lead and bismuth radioisotopes simultaneously in a quantitative fashion.

All three of the above-mentioned detectors were integrated with the HPLC system. The system as a whole (HPLC and detectors) allows us to quantitatively measure the interaction between radionuclides and

ligands of choice and meets the time requirement for separations that are fast with respect to the short half-lives of the ^{212}Bi and daughters.

Protein Chemistry

It was necessary to determine what protein(s) may compete for bismuth in radiopharmaceuticals and to determine whether radiobismuth derived from metabolized pharmaceuticals will associate with low molecular weight cytosolic proteins such as those which have been previously described. The present investigation was designed to determine the influence of cellular Bi concentration on the localization of Bi in the kidney cytosol taken up from a tracer dose of the element.

In this study, localization of Bi tracer was determined in kidneys of Bi-exposed and non-exposed rats. The results show that following its repeated injection, Bi accumulates mainly in the kidney followed by the liver, spleen, pancreas, bone, and brain. Earlier work by our laboratory using intraperitoneal administration and Vienet et al., using IV injection of inorganic ^{205}Bi have also shown a similar pattern of tissue distribution.

The results indicate that the cellular concentration of Bi influences its molecular localization. Using both gel exclusion and ion exchange chromatography we found evidence that in exposed rats, where kidney Bi concentrations are approximately $5\mu\text{g/g}$, Bi is associated with a metallothionein-like protein. However, rats with no prior exposure to Bi, and thus non-detectable levels of this element in the kidney, did not contain detectable levels of metallothionein; in these

animals, Bi radioactivity was associated with the specific higher molecular weight and hemoglobin fraction of the cell cytosol.

Metallothionein(s), contained in the soluble fraction of the cell, are low molecular weight (6,000-10,000 daltons), cysteine-rich proteins which exist in several isoforms. These proteins can bind both essential metals such as Zn and Cu and nonessential metal ions. The biological significance of the binding of metals to metallothionein has not been conclusively determined; however, it has been suggested that it may function as a detoxification protein binding such metals as cadmium and mercury as well as zinc under conditions of zinc toxicosis.

In conclusion, results from this investigation indicate that Bi exposure may alter the metabolism of Cu and Zn and additionally, that Bi in the kidney, at a concentration of 15 μ g/g tissue, induces the synthesis of metallothionein. Consequently, when Bi is administered in tracer quantities, such as that amount proposed for use as a radiopharmaceutical, the fate of any dissociated Bi will be influenced by prior exposure of the subject to metallothionein inducing substances. Competitive studies to evaluate the ability of various complexing agents to impede formation of Pb and Bi metallothionein products are in progress.

We initiated studies investigating the metabolism of trace amounts of ^{212}Pb in normal rats. Tissue and cellular localization of no carrier added ^{212}Pb has been determined in rats injected with either free ^{212}Pb (group 1) or ^{212}Pb complexed with dithiocarboxycyclic acid (DTC, group 2). Rats were killed 4h post-injection. In both groups, 20% of retained activity (RA) was in blood (95% in the RBC fraction). The majority of the remaining RA was in 4 tissues; however, the

order of magnitude differed between the groups. In group 1 the pattern of RA per gram tissue found was: liver>kidney>spleen>pancreas; in group 2 the pattern was: pancreas>spleen>liver>kidney. In both groups the distribution of ^{212}Pb in liver cellular fractions was: 600g fraction, 53%; nuclear, 2%; mitochondrial, 5%; cytosolic, 40%. Kidney ^{212}Pb distribution was: 600g fraction, 38%; nuclear, 2%; mitochondrial, 10%; cytosolic, 50%. These results show that the relative tissue uptake of trace amounts of ^{212}Pb is dependent on the form in which the Pb is presented to the animal. In contrast, complexing ^{212}Pb with DTC did not influence its subcellular localization relative to "free" ^{212}Pb .

Cellular Radiation Biology

The following text summarizes work supported in part by this project with allotransplantation of canine myelomonocytic leukemic cell SK9ML-2 that was established at our laboratory by T.K. Kawakami and G. R. Cain from a dog with spontaneous-occurring nonlymphocytic leukemia was transplanted in nonlethally irradiated pups. Allotransplantation was confirmed by chromosomal analysis. The clinical course of the disease was acute, and a diagnosis of acute myelomonocytic leukemia (equivalent to M4 of the French-American-British system of classification) was made based on the cytomorphological and cytochemical characteristics. Splenomegaly, hepatomegaly, and diffuse lymphadenopathy were observed at necropsy of all leukemic pups.

The dog is a large random-bred animal and is an ideal biomedical model for lymphohematopoietic disease (Wilson and Shifrine, 1980). However, the dog is not widely used for clinical and therapeutic studies of acute myelogenous leukemia, primarily because the

spontaneous disease in the dog is rare. Investigations of myeloid leukemia in the dog are primarily limited to individual or small group case reports emphasizing clinical manifestations or diagnostic methods (Christopher et al. 1986; Couto 1985; Jain et al. 1981).

We have established a canine myeloid leukemic cell line in order to have available a uniform and continuous source of myeloid leukemic cells for use in cellular and molecular studies. This cell line was developed from leukemic cells obtained from a male adult dog with spontaneous acute nonlymphocytic leukemia. In order to determine the leukemogenicity and course of clinical disease in vivo, we transplanted the cell line SK9ML-2 into nonlethally irradiated newborn dogs. This report describes the clinicopathologic response to allotransplantation of the cell line SK9ML-2 and 3 subclones into newborn nonlethally irradiated dogs.

Materials and methods

Animals.

Newborn beagle pups were obtained from a closed colony at the Laboratory for Energy-Related Health Research. All pups were nursed by their dam during the experiment. All dogs were housed indoors or in outdoor facilities approved and conducted under the guidelines of the Animal Use and Care Administration Committee, University of California, Davis, CA.

Radiation Exposure

Newborn pups received 100 cGy whole body gamma irradiation at a dose rate of 20 cGy/minute in a ^{137}Cs irradiator (Mark I, model 30, J.L. Shepard and Assoc., Glendale, CA). Dosimetry was quantitated with LiF thermoluminescent dosimetry chips (TLD-100, Harshaw Chemical Co., Solon, OH).

Inoculum

The cell inoculum was SK9ML-2 cells obtained from an actively growing liquid culture. The cells were grown in vitro in Liebovitz medium (L-15, GIBCO, Santa Clara, CA), supplemented with 15% defined fetal bovine serum (HyClone Lab, Logan UT), 2.0 mM glutamine, 1.0 mM sodium pyruvate, 5×10^{-7} M hydrocortisone acetate (Sigma Chemical Co, St Louis, MO), and 1×10^{-5} M monothioglycerol (Sigma). The cells were harvested and washed in phosphate buffered saline (PBS) by centrifugation ($1000 \times g$ for 10 minutes), resuspended in 1 ml of PBS, and injected intraperitoneally ($1-2 \times 10^8$ cells/pup). Single cells for cloning were isolated in soft agarose medium containing 0.32% agarose (Sea Plaque, FMC, Rockland, ME) in complete L-15 medium supplemented with 50% autologous conditioned medium. Seven newborn pups (3 females and 4 males) were inoculated with SK9ML-2 cells or subclones from actively growing cultures. The pups were observed daily and blood samples were collected 7 days after injection and when clinical signs of weakness and lassitude developed.

Cell Count

All cell counts were obtained with an electronic cell counter (Coulter ZB1, Hialeha, FL). Wrights-Giemsa stained peripheral blood smears were used for determining differential blood cell counts.

Chromosome analysis

Chromosome analyses of leukemic cells from affected pups were obtained to determine the number of metacentric sex chromosomes of the leukemic cells in female recipients. The chromosome preparations were performed as previously described (Kawakami, et al. 1989).

Cytochemical stains

Cytochemical stains for alkaline phosphatase, myeloperoxidase, non-specific esterase, and Sudan black B were used to differentiate immature and blastic cells in the blood films. The staining methods were as previously described (Jain, 1981); (Yam, et al. 1971).

Gross and histopathology

A complete gross necropsy examination was performed on all dogs with transplanted myeloid leukemia. Tissues for histologic examination were fixed in buffered formalin, embedded in paraffin, sectioned at 6 μ m, and stained with hematoxylin and eosin (HE).

Results

SK9ML-2 was the uncloned cell culture that was utilized after the leukemic bone marrow cells from the original canine donor had been maintained in continuous liquid culture for more than 20 weeks.

SK9ML2.0, SK9ML-2.6 and SK9ML2.8 were subcloned cells developed from single clonal cells after the cell line SK9ML-2 had been established. The leukemic cells, uncloned or cloned, were capable of producing transplantable myeloid leukemia except in dog 87L23U (Table 1).

The clinical course of the disease was acute. Blood samples obtained seven days after injection were normal. During the second week post injection, pups were initially clinically normal but leukocyte counts were markedly elevated. Dog 87L23X was found dead on the thirteenth day post inoculation. All remaining leukemic pups were subsequently euthanized on days 13 to 17 due to weakening clinical condition. A single dog (87L23U) remained normal clinically and

hematologically and was released from the study six months post inoculation.

Chromosomal analyses of leukemic cells were of male (donor) genotype. Wright's-Giemsa-stained blood smears showed predominantly monomorphic immature cells (approximately 95%) with round nuclei and high nuclear-cytoplasmic ratio (fig 1). Strongly positive (4+) reactions for alkaline phosphatase were observed for 89-100% of leukemic cells. Nonspecific esterase reaction was positive for 90-100% of the cells, 6-10% of the cells were weakly positive (1+) for Sudan black B staining, and 1-8% of the cells were weakly positive for the peroxidase reaction. A diagnosis of acute myelomonocytic leukemia (equivalent to M4 of the French-American-British system of classification) was made based on the cytomorphological and cytochemical characteristics.

Splenomegaly, hepatomegaly, and diffuse lymphadenopathy were observed at necropsy of all leukemic pups. Hepatic tissues were pale and had prominent reticular lobulation, spleens were reddened and granular on cross sectioning, and lymph nodes were enlarged, pale, and smooth surfaced. Histologic sections of bone marrow from multiple sites had complete effacement of normal hematopoietic structures by a monotonous blastic leukemic cellular infiltrate. Similar leukemic infiltrates were seen diffusely throughout the red pulp of the spleen, in liver sinusoids, and throughout cortical and medullary zones of all lymph nodes. Leukemic cells were also present in the meninges and choroid plexus of the brain, in intralveolar septa of the lung and in scattered perivascular foci in other organs.

Implications

The results from this study indicate that canine myeloid leukemic cell line SK9ML-2 and its subclones are allotransplantable to nonlethally irradiated dogs. This proved the leukemogenicity of the cloned cells and provides a means of reproducing acute myelomonocytic leukemia in the dog. Examination of chromosomal spreads revealed the single metacentric sex chromosome of the male donor indicating that the leukemia was produced by cell transplantation.

The reason for the failure of the one clonal culture (SKML-2.6) to produce leukemia is not known. It is possible that this clonal cell culture had lost the properties for leukemogenicity. This may be similar to immortalized murine cell cultures that fail to produce leukemia in isogenic hosts (Metcalf, 1985). Another possibility is that this pup retained some degree of immune responsiveness. Further study on additional subclones of non-transplantable and transplantable myelomonocytic leukemic cell lines may provide insight to the mechanism of leukemogenicity of canine myeloid leukemia cells.

Fetal allotransplantation of noncloned radiation-induced canine myeloid leukemia cells has been previously reported (Shifrine, et al. 1971). The advantage of the present cell line model to the fetal model is that the various cellular parameters, e.g. sensitivity to drug, radiation, immunologic, and gene therapy, can be evaluated *in vitro* and subsequently applied *in vivo*. In addition, the cell line model does not involve a surgical procedure or require special facilities or expertise. The availability of an allotransplantable canine myelomonocytic leukemic cell line should permit studies that are not possible with spontaneously occurring disease.

Further studies modifying the clinical expression of the transplantable leukemia will allow additional investigations of the usefulness of this myeloid leukemia cell line in understanding the biology and in the development of therapeutic modalities for acute myeloproliferative disorders.

Table 1 Allotransplantation of canine myelomonocytic leukemia in newborn dogs injected with uncloned and cloned myelomonocytic cell cultures.

Animal Number Days	Inoculum * Age at Death	Diagnosis	White Blood Count x 10 ³ Cells/mm ³	
85K98C 13	SK9ML-2	AMML **	> 500	
85K99H	SK9ML-2	AMML	213	17
87L23W 14	SK9ML-2.0	AMML	51.6	
87L23X 13	SK9ML-2.0	AMML	--	
87L23U NA***	SK9ML-2.6	Normal	8.6	
87L23V	SK9ML-2.8	AMML	88.2	13
87L23L 14	SK9ML-2.8	AMML	308	

* Inoculum = Uncloned cells SK9ML-2 ; Cloned cells SK9ML-2.0, SK9ML-2.6, and SK9ML-2.8.

** AMML = Acute myelomonocytic leukemia.

*** NA = not applicable. This animal was clinically diagnosed normal at six month of age and was released from the project.

Canine leukocyte integrins and CA1.4E9 murine monoclonal antibody.

The major characteristics of the canine CD18 leukocyte antigen family, defined by a murine monoclonal antibody CA1.4E9 have been described (Moore et al.1990).The overall molecular organization of this family of leukocyte integrins is similar to the human counterpart and consists of a common beta subunit of 95 kD (CD18) and non-covalently linked alpha subunits of 150 kD (CD11c), 165 kD (CD11b) and 180 kD (CD11a). Conservation of CD18 epitopes between dog, human, feline, equine and bovine is observed.

CD18 expression in lymphoid tissue has a characteristic pattern which is based on differences in the intensity of expression observed in different cell types. The expression of LeuCAMs in canine epidermotropic lymphosarcoma (mycosis fungoides) has been explored since LeuCAMs might be expected to play a major role in the tissue tropism of this tumor. Intense CD18 expression is observed with malignant T cells which infiltrate the epidermis: these cells differ phenotypically from dermal T cells in that they often lack Thy-1 expression.

Immunoperoxidase intensely stains (based on CA1.E9 antibody) epitheliotropic malignant T-cells. With flow cytometry, we found similar intense staining of the transplantable myeloid leukemia, SK9ML-2, described in the previous section on leukemia transplantation. We found that DTPA conjugates (Brechtel et al. 1986) for Bi212 complexation are formed without degrading CA1.4E9..

Immunoperoxidase stain (based on the CA1.4E9 monoclonal antibody) of canine gingiva gave intense CD18 antigen staining by epitheliotropic malignant T-cells in a case of mycosis fungoides . Flow cytometry showed

similar intense staining against our transplantable canine myeloid leukemia, SK9ML-2 (Kawakami and Cain; described above.).

High performance liquid chromatography of CA.4E9 after linking the ^{212}Bi and ^{111}In chelator, para isothionato-benzyl DTPA showed only one major peak. The chromatogram, made with a TSK-3000SW column and saline eluant, showed negligible degradation or dimerization.

Preliminary Studies: CA1.4E9 CONJUGATION AND RADIOLABELLING

Conjugation:

Numerous conjugation reactions were performed to conjugate p-SCN-benzyl-DTPA chelator (Brechtel et al. 1986) with Bovine IgG (as a model Ab) and CA1.4E9. The average yield of conjugation was about 50%. Usually using 6 chelators/antibody in the conjugation reactions, and this usually yielded about 3 chelators/antibody.

The most successful conjugations used Ab concentrations of about $4 \times 10^{-5}\text{M}$ and chelator concentrations of $2.4 \times 10^{-4}\text{M}$. pH is 9-9.5 held by 0.05M bicarbonate in the reaction. Reaction times were about 12-18 hours at room temperature. These extended times are not necessary because the p-SCN moiety on the unreacted DTPA is usually 100% hydrolyzed after 3-4 hours. We have not investigated the shorter reaction times.

The extinction coefficient of p-SCN-benzyl-DTPA (designated MX-DTPA when methylsubstituted) is 15A.U./mg/mL. The actual concentrations of the p-SCN-benzyl chelators (MX-DTPA, DTPA) cannot be determined by weight due to unknown quantities of impurities (NaHCO_3 and Na in unlabelled compound provided by

Coulter Immunology and, in the case of the ^{14}C labelled MX-DTPA, cesium -- a gift from Otto Gansow, NCI/NIH). The ^{14}C labelled DTPA concentration can be determined by counting it (Sp.Act.= 1.43×10^6 dpm/ μmole).

We have used two batches of MX-DTPA conjugated CA1.4E9 available (MXI and MXII). These are in 0.15M NaCl. There is approximately enough Ab for about 100 radiolabelling experiments under the conditions that we use. These batch of Ab conjugate was found to have >95% immunoreactivity of the unconjugated CA1.4E9 by fluorescence activated cell sorting (FACS).

Radiolabelling:

Our current generator is capable of producing about 0.1 mCi to 0.2 mCi of pure bismuth in 500 μL HI. The current generator design emanates Rn-220 (Thoron) gas from ca. 4 mCi Uranium-232 parent barium stearate cake in a filter housing. This output is collected at 77K in a teflon bottle and is a very no-carrier-added product. The efficiency increases several fold when the cake is humidified but the ice build up in collection has led us to use the less efficient mode for the time-being. We have another 100 mCi U-232 on hand for a larger generator. Also, the short-lived Ra-224 based Bi212 generators from Argonne National Laboratory are available for these experiments

Biological system compatible procedures for purification and neutralization of the Bi after washing the Pb-Bi equilibrium solution from the collection bottle have been determined by trial runs. The pure Bi-212 solutions are adjusted to pH=3 and then ~2M acetate buffer (~pH=4.5) can be used buffer the solution up to pH=4-4.5 and reacted with the Ab-MXDTPA conjugate. Separation post-rxn is accomplished by HPLC using a Pharmacia FPLC size exclusion column

loaded with Sephadex G-25 super fine gel. The antibody has been checked for immunoreactivity post-radiolabelling and found to have >95% activity relative to virgin CA1.4E9.

Reaction with the SK9ML-2 Canine Leukemia.

Radiolabelled CA1.4E9 was reacted with SK9ML-2 cells. There were about 200,000 Ab/cell in the reaction and the cells, after washing, had about 10% of the activity on them after a 1/2 hour reaction. This implies that each ML-2 cell has bound approximately 20,000 CA1.4E9 to antigenic sites. This means that a Ab/Bi radiolabelling ratio of 2000 would deliver about 10 Bi-212/cell.

Cellular Microdosimetry.

What is the probability distribution relating the number of alpha particles that are necessary to achieve adequate reproductive death of SK9ML-2 populations from marrow harvest and cell line cultures?

There is no firm theoretical or experimental guidance. We have by personal communication, surveyed opinions of the cognoscenti regarding alpha particle cellular radiation biology and found estimates of 2 to 20 nuclear hits. We do not know whether the Bi212 tagged CA1.4E9 will be endocytosed by SK9ML-2 and lead to the effects of the mass 212 recoil atom being an important factor in radiation cytotoxicity. We have addressed the recoil atom issue (Zidenberg-Cherr 87) in the past and found that an energy density of ca.200-300 eV/atom imparted to surrounding molecular structures would obtain. This is in comparison to about 32 eV/atom from Bi212 alpha particles.

The SK9ML-2 cells have a radius of 7-10 μm , have a nucleus occupying well over half the cell volume, and are dividing with a generation time of 20-30 hours. Ten alpha sources per cell will give ≈ 4 nuclear hits. This will be a target starting point within our experimental capability now.

Summary Conclusions

We believe that the in vitro use of $^{212}\text{Pb}/^{212}\text{Bi}$ chelates attached to antibodies against adhesion antigens as a purging agent for leukaemic marrow has substantial promise. The recently discovered new classes of compounds such as the completely caged and pi electron rich C-60 and larger buckminster fullerenes offer promise of fulfilling the original concept of this proposal. We anticipate the continued exploration of our original concept advanced in 1985, namely the chemical creation of an approximately 11 hour half period alpha emitting moiety for biological applications, with various newly discovered compounds, including the fullerenne class of compounds.

BIBLIOGRAPHY

- Avila, M. J., G. Barrall, S. Raaka, and N. J. Parks.(1992) *Emulston Scintillation Spectroscopy of Mixed Alpha and Beta Particle emitters: Radiochemical Assays of ^{212}Pb (10.6h) and ^{212}Bi (1.01h).* Applied Radiat. Isot.43:753-760.
- Bharadwaj, B.K. and W.K. Musker (1987) *A 1,1-Dithiocarboxylate Ligand with an Easily Derivattivable Group. Synthesis and Structure of Tris[2-(ethylamino)cyclopent-1-ene-1-dithiocarboxylate]bismuth(III).* J. Inorg. Chem. 26(9).
- Bharadwaj, P.K., B.W. Arbuckle, and W.K. Musker (1988) *Molecular and Crystal Structure of Lead(II) and Bismuth(III) Complex of Dipropyldithiocarbamate Ligand.* Inorg. Chim. Acta. 142:243-246.
- Brechbiel, M.W., O.A. Gansow, R.W. Atcher, J. Schlom, J. Esteban, D.E. Simpson and D. Colcher (1986) *Synthesis of 1-(p-Isothionatobenzyl) derivatives of DTPA and EDTA. Antibody Labeling and Tumor Imaging Studies.* Inorg. Chem. 25:2772-2781.
- Broadbeck, R., G.J. Newton, A.C. Mishra, N.J. Parks (1989) *Radiochemistry and Design of ^{220}Rn Generators Based on ^{232}U and ^{228}Th .* Appl. Radiat. Isot. submitted.
- Carlson, T.A. (1979) Chemical Effects of Nuclear Transformat on in Inorganic Systems. (G. Harbottle and A. Maddock, Eds.), North Holland, Amsterdam, pp.14-37.
- Christopher, M.M., Metz, A.L., Polzin, D., and Hayden, D.W. (1986). *Acute myelomonocytic leukemia with neurologic manifestations in the dog.* *Vet Pathol* 23, 140-147.
- Cooper, S.R. (1988) *Crown Thioether Chemistry.* Acc. Chem. Res. 21:141.
- Cooper, S.R., S.C. Rawle, J.R. Hartman, E.J. Hintsa and G.A. Admans (1988) *Crown Thioether Chemistry: Homoleptic Hexakis(Thioether) Complexes of Nickel(II).* Inorg. Chem. 27:1209.
- Cooper, S.R., Rawle, S.C. (1989) *Crown Thioether Chemistry, Structure and Bonding,*(Springer-Verlag, New York)in press.

Couto, C.B. (1985). Clinicopathologic aspects of acute leukemias in the dog. *J Am Vet Med Assoc* 186, 681-685.

Harris, W.R. and K.N. Raymond (1979) *Ferric Ion Sequestering Agents. 3. The Spectrophotometric and Potentiometric Evaluation of Two New Enterobactin Analogues: 1,5,9-N,N',N''-Tris(2,3-dihydroxybenzoyl)-cyclotriazatridecane and 1,3,5-N,N',N''-Tris (2,3-dihydroxybenzoyl)triaminomethylbenzene.* *J. Am. Chem. Soc.* 101(22):6534-6541.

Harris, W.R., K.N. Raymond, and F.L. Weigl (1981) *Ferric Ion Sequestering Agents. 6. The Spectrophotometric and Potentiometric Evaluation of Sulfonated Triccatecholate Ligands.* *J. Am. Chem. Soc.* 103(10):2667-2675.

Jain, N.C., Madewell, B.R., Weller, R.E. and Geissler, M.C. (1981). Clinical-pathological findings and cytochemical characterization of myelomonocytic leukemia in 5 dogs. *J Comp Path* 91, 17-31

Kawakami, T.G. and Cain, R.G. (1992). Allotransplantation of canine myelomonocytic leukemic cell SK9ML-2. Submitted to *Hematology Oncology*

Kawakami, T.G., Cain, G.R. and Taylor, N. (1989). Establishment and partial characterization of radiation-induced canine monocytic leukemic cell line (RK9ML-1). *Leukemia Res* 13, 709-714.

Metcalf, D. (1985). Multi-CSF-dependent colony formation by cells of a murine hemopoietic cell line: specificity and action of multi-CSF. *Blood* 65, 357-362.

Keen, C.L., B. Lönnerdal, and L.S. Hurley (1981) *Developmental Changes in Zinc and Copper in Mouse and Rat Tissues. IN Trace Element Metabolism in Man and Animals (TEMA-4)* (J.M. Gawthorne, J.McC. Howell, and C.L. White, Eds.), Griffin Press Limited, Netly, South Australia, pp.287-290.

Lagunas-Solar, M.C., O.F. Corvacho, L. Nagahara, A.C. Mishra, and N.J. Parks (1987) *Cyclotron Production of No-Carrier-Added ^{206}Bi (6.24d) and ^{205}Bi (15.3d) as Tracers for Biological Studies and for the Development of Alpha-Emitting Radiotherapeutic Agents.* *Int. J. Radiat. and Appl. Isotopes* (in press).

Lloyd, E.L., M.A. Gemmell, C.B. Henning, D.S. Gemmell and B.J. Zabransky (1979) *Cell Survival Following Multiple-Track Alpha Particle Irradiation.* *Int. J. Radiat. Biol.* 35(1):23-31.

Mishra, A.C. and N.J. Parks (1987) *Beta Spectroscopy with Liquid Scintillation systems: Applications in dosimetry of ^{90}Sr and ^{90}Y* . Appl. Radiat. Isot.(Int. J. Appl. Instr., Part A) 38(6):455-461. (1987).

Moore, P.F., Rossitto, P.V. and Danilenko D.M. (1990). Canine leukocyte integrins: Characterization of a CD18 homologue. *Tissue Antigens* 36, 211-220.

Pikul, S.S., N.J. Parks and P.D. Schneider (1987) *In Vitro Killing of Melanoma By Liposome Delivered Intracellular Alpha Irradiation*. Arch. Surg. 122:1417-1420.

Rawle, S.C., J.R. Hartman, D.J. Watkin and S.R. Cooper (1986) *Crown Thioether Chemistry. The Nickel(II) Complex of Hexathia-24-crown-6*. J. Chem. Soc., Chem. Commun. 1986(14):1083-1084.

Scheinberg, D.A., M. Strand, O A. Gansow (1982) *Tumor Imaging With Radioactive Metal Chelates Conjugated to Monoclonal Antibodies*. Science 215:1511-1513.

Shifrine, M., Bulgin, M.S., Dollarhide, N.E., Wolf, H.G., Taylor, N.J. and Wilson, F.D. (1971). Transplantation of radiation-induced canine myelomonocytic leukaemia. *Nature* (London) 232, 405-406.

Thorne, C.M., S.C. Rawle, G.A. Admans, and S.R. Cooper (1986) *Synthesis and Structure of the Nickel II Complex of 1,1,1-Tris[(2-methylthioethylthio)-methyl]ethane, an all Sulfur Ligand that Promotes Hexakis-thioether Coordination*. J. Inorg. Chem. 25:3848-3851.

Thorne, C.M., S.C. Rawle, G.A. Admans and S.R. Cooper (1987) *A New Approach to Hexakis(thioether) Co-ordination: the Cobalt(II) Complex of 1,1,1-tris[(2-methylthioethylthio)-methyl]ethane*. J. Chem. Soc., Chem. Commun. 1987(D 4):306-307.

Thorne, C.M., S.C. Rawle, G.A. Admans and S.R. Cooper (1989) *Supertripodal Chelating Agents. 1,1,1-Tris(((2-(methylthio)ethyl)thio)methyl)ethane, and its Complexes with Fe(II), Co(II), and Ni(II)*. in preparation.

Wahl, A.C. and W.R. Daniels (1958) *Emanating Power of Barium Stearate for 3.9 Second Actinon (^{219}Rn)*. J. Inorg. Nucl. Chem. 6:278-287.

Yam, L.T., Li, C.Y., and Croby, W.H. (1971). Cytochemical identification of monocytes and granulocytes. *Amer J Clin Pathol* 55, 283-290

Yeh, S.M., C.F. Meares and D.A. Goodwin (1979) *Decomposition Rates of Radiopharmaceutical Indium Chelates in Serum*. J. Radioanal. Chem. 53:327-336.

Zidenberg-Cherr, S., N.J. Parks and C.L. Keen (1987) *Tissue and Subcellular Distribution of Bismuth Radiotracer in the Rat: Consideration of Cytotoxicity and Microdosimetry for Bismuth Radiopharmaceuticals*. Radiat. Res. 111:119-129.

Zidenberg-Cherr, S., M.S. Clegg, N.J. Parks and C.L. Keen (1989) *Localization of Bismuth Radiotracer in Rat Kidney Following Exposure to Bismuth*. Biol. Trace Element Res. 19: 185-194.

Zucchini, G.L. and A.M. Friedman (1982) *Isotopic Generation for ^{212}Pb and ^{212}Bi* . Int. J. Nucl. Med. Biol. 9(1):83-84.

END

**DATE
FILMED
9/16/92**