DOE/PC/79938--T7

DE92 017806

**STUDIES OF COUPLED CHEMICAL AND CATALYTIC COAL CONVERSION
METHODS**

LEON M. STOCK, PRINCIPAL INVESTIGATOR

**KUNTAL CHATTERJEE, CARLOS CHENG, MICHAEL ETTINGER, FRANCIS
FLORES, SAO JIRALERSPONG, MIKIO MIYAKE AND JOHN MUNTEAN**

RESEARCH STAFF

DECEMBER 1991

**THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
5735 SOUTH ELLIS AVENUE
CHICAGO, ILLINOIS 60637**

**PREPARED FOR THE
UNITED STATES DEPARTMENT OF ENERGY
UNDER CONTRACT NO. AG22-87PC79938**

ELEMENT	Page
TABLE OF CONTENTS	ii
LIST OF TABLES	iii
LIST OF FIGURES	v
EXECUTIVE SUMMARY	1
SECTION I. CHEMICAL CONVERSION STUDIES	
Introduction	4
Experimental	6
Results	10
Discussion	28
Conclusion	49
SECTION II. CATALYTIC CONVERSION STUDIES	
Introduction	53
Experimental	53
Results and	
Discussion	55
Conclusion	78
REFERENCES	79
APPENDIX	
I. LIST OF PUBLICATIONS.....	82

LIST OF TABLES

Table	Page
1. Analytical data for the coal samples	11
2. Effect of O-alkylation and O-acylation on the solubility of Illinois No. 6 coal	12
3. Effect of O-alkylation on the solubility of Wyodak coal	13
4. Analytical data for the high rank coal sample	14
5. Influence of the nature of base on base promoted butylation of Lower Kittanning coal upon its solubility	15
6. Effect of concentration of base used for C-butylation on the solubility of Lower Kittanning coal	16
7. Influence of the alkyl group on pyridine solubility of Lower Kittanning coal	17
8. Influence of the alkyl group on pyridine solubility of the O-methylated Wyodak coal	18
9. Results for the C-alkylation reactions of PSOC 1197 with Lochmann's base	21
10. Results for the C-alkylation reactions of O-methyl Wyodak coal with Lochmann's base	22
11. Results for the C-alkylation reactions of O-methyl Illinois No. 6 coal with Lochmann's base	23
12. Effect of the reaction conditions on pyridine solubility of O-methyl Illinois No. 6 coal	24
13. Results for C-butylation of O-methylated Illinois No. 6 coal in tetrahydrofuran	25
14. Influence of the solvent of C-butylation on solubility of the O-methylated Illinois No. 6 coal	26

15. Results of solubilization of Illinois No. 6 coal due to metal alkoxide treatments	27
16. Hydrocarbons and their pKa values	35

LIST OF FIGURES

Figure	Page
1. Various possible structures for the true nature of the "magic metalating agent" - the Lochmann's base	20
2. Heteronuclear correlation spectrum of the pyridine soluble part of O-methyl ^{13}C -Illinois No.6 coal. The total spectrum is flanked below by carbon projection and vertically along the contour plot, by proton projection	30
3. Herteronuclear correlation spectrum of the pyridine soluble part of O-methyl ^{13}C -Illinois No.6 coal. A more selective spectrum which has only the region of interest	31
4. Solid state ^{13}C spectra of Wyodak coal: (a) O-methyl ^{13}C -Wyodak coal; (b) pristine Wyodak coal; and (c) the difference spectrum	32
5. ^{13}C NMR spectra of the Lower Kittanning seam coal PSOC 1197 (bottom) and the product obtained in the sodium amide promoted methylation with methyl- ^{13}C iodide (top)	37
6. ^2H NMR spectrum of the product (pyridine soluble part) from the reaction of O-methyl Illinois No. 6 coal with Lochmann's base and deuterium oxide	41
7. ^{13}C -NMR spectrum of the product (pyridine soluble part) from the reaction of O-methyl Illinois No. 6 coal with Lochmann's base and methyl- ^{13}C -iodide. The region of interest is between 10 to 60 ppm	42
8. ^{13}C NMR Spectra of Wilsonville Coal Liquid and Coal Liquid- $\text{Cr}(\text{CO})_3$...	56
9. Synthetic Methods For Arenechromium Tricarbonyl Complexes	57
10. Infrared Spectra of Wilsonville Coal Liquid and Coal Liquid- $\text{Cr}(\text{CO})_3$	59
11. Infrared Spectra of Benzene- $\text{Cr}(\text{CO})_3$ and $\text{Cr}(\text{CO})_6$	60
12. Arenechromium Tricarbonyl Catalyzed Hydrogenation Reactions	61
13. Cr-C Bond Lengths in Mono- and Polycyclic Arenechromium Tricarbonyl Complexes	61

14. 1H NMR Spectra of Napthalene and Napthalene- $Cr(CO)_3$	63
15. 1H and 2H NMR Spectra of Napthalene- $Cr(CO)_3$ Catalyzed Deuteration of Methyl 2,4-Hexadienoate Reaction Product	64
16. 2H NMR Spectra of Unreacted Coal Liquid and Coal Liquid- $Cr(CO)_3$ Deuteration Reaction Product	66
17. 2H NMR Spectra of Unreacted Coal Liquid and Napthalene- $Cr(CO)_3$ Catalyzed Deuteration of Coal Liquid Reaction Product	67
18. Deuteration Reactions and Results Under Mild Conditions	69
19. 1H and 2H NMR Spectra of Coal Liquid- $Cr(CO)_3$ Catalyzed Deuteration of Methyl 2,4-Hexadienoate Reaction Product	71
20. Deuteration Reactions and Results Under Moderate Conditions	72
21. 2H NMR Spectra of $Cr(CO)_6$ Catalyzed Deuteration of Methyl 2,4-Hexadienoate Reaction Product	73
22. Attempted Complexation Reactions of Pocahontas Coal With $Cr(CO)_6$	73
23. Infrared Spectra of Pocahontas Coal Before And After Reaction With $Cr(CO)_6$	74
24. Infrared Spectra of O-Methyl Illinois No. 6 Coal and [Dioxane Extract of O-Methyl Illinois No. 6 Coal]- $Cr(CO)_3$	76
25. Infrared Spectrum of Illinois No. 6 Coal- $Cr(CO)_3$	77

EXECUTIVE SUMMARY

Objectives.

The objective of this research was to convert coal into a soluble substance under mild conditions. The strategy involved two steps, first to breakdown the macromolecular network of coal, and second to add hydrogen catalytically.

We investigated different basic reagents that could, in principle, break down coal's structure and alkylation strategies that might enhance its solubility. We examined O- and C-alkylation, the importance of the strength of the base, the character of the added alkyl groups and other reaction parameters. This work provided new information concerning the way in which hydrogen bonding, polarization interactions between aromatic structures and covalent bonding could be disrupted and solubility enhanced.

Coal liquefaction and upgrading reactions are usually catalyzed by supported and dispersed heterogeneous catalysts. The limiting factor in such systems is the number of coal molecules that are able to come into contact with the catalyst. Further, the severe systems under which coal products are refined demand that catalysts be robust. The objective of our research was to explore new organochromium chemistry that might be feasible for the hydrogenation of coal under mild conditions.

Accomplishments.

Three different coal samples, Lower Kittanning coal, Illinois No. 6 coal and Wyodak coal, were studied in the course of this program.

O-Alkylation of Illinois No. 6 coal increased the pyridine solubility from 27% to 34%, and that of Wyodak coal from 9 to 39%. Presumably, the O-alkylation reaction is successful because alkylation disrupts all intermolecular hydrogen bonding, but it is clear that other binding forces in the oxygen-rich subbituminous and bituminous coals need to be overcome to achieve high solubilization.

Amide-promoted C-alkylation successfully converted more than 80% of the high rank coal to pyridine soluble material. The success of the reaction depended upon the base strength and the

size of the alkyl group. Indeed, the dimensions of the alkyl groups have a profound effect, and we infer that these groups, even in small number, can disrupt the stacking pattern of the coal molecules within the solid, and thereby diminish the polarization forces that bind the coal. C-Alkylation of the O-methylated low rank coals does not have as large an influence on solubility. Even though the O- and C-alkylation of a low rank coal inhibits all hydrogen bonding and may disrupt polarization forces, solubility is not increased appreciably.

Inasmuch as base strength is significant, we studied C-alkylation with a super base, Lochmann's base. This procedure almost quantitatively solubilized the high rank coal. Studies with normally unreactive pure hydrocarbons including 1,1,3-triphenylpropane, 9-benzyl-9,10-dihydroanthracene and 9,10-diphenylanthracene revealed that Lochmann's base can accomplish selective carbon-carbon bond cleavage prior to alkylation. This base was also responsible for significantly reducing the sulfur content of Illinois No. 6 coal.

Lochmann's base was less successful for solubilizing the two lower rank coals. We suspected that condensation reactions of the carbanions that were formed from the coal by this base might be responsible; accordingly, we worked with other bases that could be used in proton rich media. The work with potassium alkoxides in refluxing butanol and octanol was successful. The best results were realized with Illinois No. 6 coal which gave more than 70% soluble material when treated with sodium or potassium benzoxide in refluxing benzyl alcohol.

Generally speaking, the goals of the research in this area were all realized.

Arenechromium tricarbonyl complexes have exhibited diverse chemical properties including the ability to homogeneously catalyze the hydrogenation of dienes under very mild conditions. Research was directed at trying to use such complexes to hydrogenate molecules in coal products. It was demonstrated that arenechromium tricarbonyl complexes can be prepared from the aromatic molecules in coal liquids, coal extracts, and solid coal. Results showed that, despite the difficulties of carrying out this type of sensitive chemistry on coal materials, high yields of coal-arenechromium tricarbonyl complexes were achieved. The coal-arenechromium tricarbonyl complexes made from the coal liquid displayed catalytic activity in the hydrogenation of dienes

under moderate conditions. The coal liquid arenechromium tricarbonyl complex made from the Wilsonville coal liquid catalyzed the deuteration of methyl 2,4-hexadienoate to methyl hex-3-enoate- d_2 at 200°C under 500 psi of dideuterium. Naphthalenechromium tricarbonyl was synthesized and catalyzed the same reaction under very mild conditions of 45°C under 150 psi dideuterium in the presence of the coal liquid. It was hoped that the complexed coal liquid would be able to self catalyze the hydrogenation of other coal molecules. This objective was not realized presumably because the coal-arenechromium tricarbonyl complexes are only able to hydrogenate dienes under the mild conditions used in this work. Also, it is postulated that only monocyclic arenechromium tricarbonyl complexes were formed with the coal liquid as catalysis occurred only under more moderate conditions than those required for the naphthalenechromium tricarbonyl complex. However, it is encouraging that the coal-arenechromium tricarbonyl complexes in fact performed the same hydrogenation chemistry as the pure complexes, and that the arenechromium tricarbonyl complexes exhibited a robustness capable of withstanding severe conditions of coal systems.

While supported and dispersed heterogeneous catalysts are predominately used in the hydrogenation and upgrading of coal products, many problems associated with the use of solid catalysts in such systems certainly exist such as catalyst poisoning as well as deactivation due to coke deposits. The results of this study support the idea that homogenous catalytic systems may be feasible for use in the refining of coal products. This study has shown that such chemistry is possible with coal molecules. The use of these monomolecular reagents can increase the effective concentration of the catalyst by many orders of magnitude when compared to supported and dispersed solid catalysts. The many advantages of homogenous systems warrant further investigation of their use in coal systems, and may provide an economical and chemically efficient method of refining fossil fuel materials.

SECTION I. CHEMICAL CONVERSION STUDIES
PREPARED BY KUNTAL CHATTERJEE

INTRODUCTION

For nearly 20 years, attempts have been made to solubilize coal in common organic solvents. However, most chemical reactions that have been investigated for the transformation of coal to soluble products disrupt strong covalent linkages.^{1,2} The reductive alkylation procedure developed by Sternberg and his associates deserves mention in this context.³ In his approach, the coal sample was treated with alkali metals in the presence of naphthalene in tetrahydrofuran to form anionic coal fragments, which were subsequently alkylated. For example, alkylation of the anionic material from Illinois No. 6 coal with various different alkyl halides gave products that had 7-9 alkyl groups per 100 mol carbon. The alkylated coal was much more soluble than the original coal. However, apart from alkylation reactions, other significant chemistry occurred to fragment the macromolecule. Specifically labile carbon-oxygen and carbon-carbon bonds are cleaved under the experimental conditions.¹ These cleavage reactions, together with the reduction reactions of the aromatic compounds in the coal, disrupted the coal structure sufficiently to achieve a high level of solubility.¹ More recently, interest has been directed to simple alkylation reactions that can alter the nonbonded intermolecular interactions that contribute to the binding of large coal macromolecules in the solid state.¹ These strategies are based on the idea that the introduction of an O-alkyl group can eliminate hydrogen bonding interactions and that the introduction of either O- or C-alkyl groups can disrupt intermolecular polarization forces. Whereas O-alkylation does not generally enhance the solubility of either high or low rank coals to a significant degree,^{1,4} C-Alkylation is especially effective for some high rank coals. Ignasiak and his coworkers demonstrated that after three repeated ethylation reactions, up to 14 ethyl groups per 100 mol carbon could be introduced into an 83.2% carbon coal, and that up to 64% of the ethylated product of an 87.3% C coal could be solubilized in chloroform.^{5,6} But the results obtained in several different laboratories have not been entirely consistent.⁴⁻¹⁰

In the present work, C-alkylation reactions of one high rank coals, one high volatile bituminous coal and one subbituminous coal were studied. The work with the high rank bituminous coal with its low oxygen content and very low phenolic content was successful. With the low rank coals which are rich in phenolic groups, effects of O-alkylation were studied first. Subsequently, these phenolic groups were methylated to obviate the complications of the high negative charge density in the coal macromolecule, and the coals were C-alkylated with strongly basic reagents. This strategy enabled the exclusive assessment of C-alkylation on the solubilization of the coals. Also, new results

were obtained to prove that strong bases can cleave carbon-carbon bonds thereby forming small fragments in hydrocarbons.

EXPERIMENTAL

Materials

The low-volatile, bituminous, Lower Kittanning coal, PSOC 1197, was obtained from the Pennsylvania State University Sample Bank. The two low rank coals, an Illinois No. 6 bituminous coal, APCSP 3, and a Wyodak subbituminous coal, APCSP 2, were supplied by the Premium Sample Program of Argonne National Laboratory. The analytical data for these coals are summarized in Table 1 and 4. All the coal samples were dried at 110 °C under vacuum for 48 hours prior to use.

Commercial anhydrous liquid ammonia (Matheson) was dried by passage through potassium hydroxide and barium oxide. Tetrahydrofuran was freshly distilled from potassium benzophenone ketyl before use. *n*-Heptane (Aldrich) was shaken with concentrated sulfuric acid, and subsequently distilled. The purified solvents were stored in the presence of molecular sieve 5A. Pyridine was purified by distillation over barium oxide. The alkyl halides, methyl, *n*-butyl and *n*-octyl iodides (Aldrich) were dried by molecular sieve 5A. The other chemicals such as *n*-butyllithium (1.6 M solution in hexane) potassium *tert*-butoxide, sodium amide, tetrabutylammonium hydroxide, benzyl alcohol, *n*-octanol, *n*- and *tert*-butanol, sodium metal, potassium metal, ammonium chloride and methanol were used as received from the Aldrich Chemical Company.

Reactions Procedures

Alkylations with potassium and tetrabutylammonium hydroxide. Alkylations were done by the procedure of Liotta¹¹ and Ettinger and coworkers.¹² Typically, the coal (1g) was stirred in freshly distilled tetrahydrofuran (25 ml) for 20 minutes. A solution of aqueous potassium hydroxide (1.5 M, 3.0 ml) was added dropwise to the suspension and the mixture was allowed to stir for two hours. The alkylating agent (4.5-5.0 mmoles) in tetrahydrofuran (5 ml) was added, and the mixture was stirred under dinitrogen for 48 hours. At the end of this time, the reaction mixture was mildly basic. Aqueous 2N hydrochloric acid was used to acidify the suspension, and the tetrahydrofuran was then removed in a rotary evaporator. The residual was collected on a Nucleopore polycarbonate membrane filter (pore size 0.8 μ m) and washed with 50% aqueous methanol until the washings were free of halide ions.

The halide ions were detected with aqueous silver nitrate. Finally, the samples were dried under vacuum at 110 °C for 24 hours.

The number of alkyl groups that were introduced per 100 mol C was estimated from the increase in weight of the product.

Alkylations with potassium trimethylsilanolate. Alkylation reactions with potassium trimethylsilanolate were carried out in a manner similar to that of potassium hydroxide. The major advantage with potassium trimethylsilanolate is that the reactions could be carried out under anhydrous conditions. The coal sample (1.0 g) was stirred in freshly distilled tetrahydrofuran (25 mL) for 20 minutes in dinitrogen. To this mixture, potassium trimethylsilanolate (0.64 g, 5 mmols) was added and allowed to stir for two hours. The alkylating or acylating agent (5-6 mmole) in tetrahydrofuran (5 ml) was added and the mixture was stirred in dinitrogen for 48 hours. All the subsequent proceedings (washing and drying the product) were performed as had been done previously for potassium hydroxide.

Alkylations with sodium amide in liquid ammonia. Typically, liquid ammonia (200 mL) was condensed in a flame dried flask and sodium amide (45 mmol) was added. After 15 minutes, the coal sample (1 g) was added, and the mixture was agitated vigorously for 6 h at -75 °C. The alkyl halide (60 mmol) in tetrahydrofuran (100 mL) was added dropwise to the mixture over a period of 30 min. The supply of the coolant was stopped and the temperature was allowed to rise gradually to room temperature as the ammonia evaporated. Then additional tetrahydrofuran (100 mL) was added to the reaction mixture. After 48 h, any remaining base was destroyed by the addition of ammonium chloride and methanol. The solvent and the excess alkyl halide were evaporated by using a rotary evaporator. The product was subsequently washed with acidified aqueous methanol (methanol / water = 1/3 v/v) and repeatedly with aqueous methanol (more than 20 L). All the reactions and the washing procedures were carried out in dinitrogen atmosphere. After the thorough washings the product was collected and dried to a constant weight at 110 °C under vacuum for 48 h. The number of alkyl groups that were introduced per 100 mol C was estimated from the increase in weight of the product.

The other reactions were conducted by suitable modifications of this procedure. Different amide bases, instead of sodium amide, were used in the same way. It should be noted that the same base concentration (45 mmol/g of coal) was used in most experiments.

Alkylations with Lochmann's base. The alkylation reactions of the coal samples were carried out essentially as reported by Bates and his coworkers.¹³ *n*-Heptane (170 mL) potassium *tert*-butoxide (5.04 g, 45 mmoles) and *n*-butyllithium (30 mL 1.6 M solution in hexane, 48 mmoles) were added to a flame dried flask. The mixture was stirred at room temperature for 15 minutes, the coal sample (1 g) was added, and the solution was refluxed for 6 hours. The initial black solution turned brown when refluxing began. *n*-Butyl iodide (11.0 g, 60 mmoles) in *n*-heptane (70 mL) was added dropwise to this coal anion suspension, after cooling the flask at 0 °C. The mixture was stirred for 48 hours at room temperature to ensure complete alkylation of the coal anions. The residual base was quenched by added ammonium chloride and methanol. The solvents and the excess *n*-butyl iodide were removed by a rotary evaporator. The product was carefully collected and washed with dilute aqueous hydrochloric acid (2 L), aqueous methanol (3:1, v/v, 20 L), and *n*-hexane (4 L). The product was dried to constant weight at 110 °C under vacuum for 48 hours. The number of alkyl groups that were introduced per 100 mol carbon was estimated from the increase in weight of the product.

Various other reactions were tried by changing the solvent and, except for the change in solvent, everything else the same. In one particular experiment, instead of the addition of the alkyl halide to the coal anion, the coal anion was added to a cold solution in the alkyl halide (inverse addition). Some experiments were carried out using tetrahydrofuran solvent, and the temperature was changed from 67 °C to -46 °C.

Reaction of coal with Lochmann's base. The coal samples were treated with Lochmann's base in the same manner as described in the previous paragraph. Only, instead of adding the alkylating agent after the base treatment, the anionic mixture was quenched with methanol and dilute acid. Washing and drying the products were carried out exactly as mentioned earlier. It should be noted that the recovery was always about 95±2% which eliminates the possibility of reagent adduction.

Reaction of coal with potassium alkoxides in refluxing alcohols. Role of bond cleavage reactions in coal samples to give a more soluble material was also studied with metal alkoxides. In particular, the low rank Illinois No. 6 coal (APCSP 3) was treated with potassium alkoxides and quenched with dilute acid. The alcohol (50 mL) and desired amount of potassium metal (40 mmol) were added to a flame dried flask. The mixture was stirred at room temperature for 30 minutes. Note that all the metal does not dissolve in the alcohol inspite of the huge excess of alcohol. The coal sample (1 g) was added and

the solution was refluxed for 24 hours. Finally, the reaction mixture was cooled to 0-5 °C and was acidified with methanol and dilute hydrochloric acid. The acidic mixture was stirred for 48 hours at room temperature. Washing and drying the products were carried out exactly as mentioned earlier. The product recovered was about 95%, when all the alcohol was removed before washing. A part of the product was sent for elemental analysis.

Reaction of dibenzofuran with Lochmann's base. The reaction of dibenzofuran was conducted in a manner similar to coal. Dibenzofuran (0.84 g, 5 mmol) was allowed to react with *n*-butyllithium (15 mL 1.6 M solution in hexane, 24 mmol) and potassium *tert*-butoxide (2.52 g, 22.5 mmol) in refluxing heptane (100 mL) for 6 hours. Methyl iodide (7.1 g, 50 mmol) was added at 0 °C and the mixture was stirred for 48 hours at room temperature. The excess base was neutralized with ammonium chloride and methanol, the solvent and the remaining methyl iodide were evaporated, and the product was extracted into ether which was washed with water and dried over anhydrous magnesium sulfate. Ammonium chloride and methanol, rather than methyl iodide, were used to neutralize the reaction mixture in another experiment. The product distributions were established by conventional GC-MS procedures.

Reaction of hydrocarbons with Lochmann's base. The reactions were conducted in a manner similar to coal. *n*-Heptane (45 - 55 mL), potassium *tert*-butoxide (2.52 g, 22.5 mmol) and *n*-butyllithium (15 mL 1.6 M solution in hexane, 24 mmol) were added to a flame dried flask. The mixture was stirred at room temperature for 15 min, the pure compound (8-10 mmol) was added and the solution was refluxed for 6 hours. The color of the solution changed from yellow to dark red or black on addition of the hydrocarbon. After 6 hours of reflux, the reaction mixture was cooled to 0 °C and dilute hydrochloric acid (1N) was added. The reaction was usually vigorous. The product was extracted into ether which was washed with water and dried over anhydrous magnesium sulfate. The product distributions were established by conventional GC and GC-MS procedures.

Pyridine extraction of the coal samples. In order to test the solubilization of the products in pyridine, a portion of each of the coal samples was subjected to Soxhlet extraction. Pyridine extractions were carried out until the extracts were colorless. The pyridine-soluble portion of the product was isolated by removal of the pyridine with a rotary evaporator followed by drying over warm oil bath (100 °C) at high vacuum. The solid thus obtained was the soluble part of the product and was subjected to other studies. The insoluble residue was washed with methanol : water (4:1, v/v) by Soxhlet extraction to remove any

residual pyridine. After thorough washings, the residue was dried over a warm oil bath at high vacuum. From the weights of the soluble and insoluble residues, the solubility of the coal was determined.

The solubility of the coal was also verified in some experiments by filtering the solution through a nucleopore polycarbonate membrane filter (pore size 0.8 μm); the residue collected on the filter was about 2-6 mg per gram coal. In other representative cases, the isolated soluble solid coal was redissolved in pyridine and a readily filterable solution was obtained. These facts establish that the products were truly soluble in pyridine.

Analysis of products.

Solid-state ^{13}C -NMR spectra of the coal samples were measured at Argonne National Laboratory by using a Bruker Model CXP-100 spectrometer. Solution phase ^{13}C and ^2H -NMR spectra of the extracts of the alkylated coals were recorded with Nicolet, Varian or GE spectrometers. Potassium bromide pellets of the coal products (2-4 wt%) were dried at 90 °C for 24 hours before recording the infrared spectra with a Nicolet Model 200A spectrometer. The molecular weight measurements were obtained by the VPO method using a Hitachi Perkin-Elmer 115 instrument. The observations for three different experiments using chloroform were extrapolated to infinite dilution in the usual way.

The analysis of the products that were obtained from pure compounds was conducted by GC-MS using Hewlett Packard 5970 series mass detector (70 eV) connected to 5890 gas chromatograph (OV-101 capillary column, 15 m x 0.25 mm, 100-180 °C, 5 °C/min). The ^1H -NMR spectrum of the product in chloroform-*d* was recorded on the University of Chicago 500 MHz spectrometer.

RESULTS

Alkylations with mild bases

In order to study the influence of O-alkylation on solubilization, the two low rank coals, Illinois No. 6 and Wyodak coal were selected. The elemental analysis data for both the coal samples were obtained from the Premium Sample Program of Argonne National Laboratory and are shown in Table 1.

TABLE 1
ANALYTICAL DATA FOR THE COAL SAMPLES^a

	Illinois No. 6 APCSP 3	Wyodak APCSP 2
ultimate analysis, wt % daf		
C	77.8	75.0
H	5.7	5.4
N	1.4	1.1
S	2.6	0.5
O (by difference)	12.5	18.0
ash	16.2	9.2
maceral analysis, wt%		
Vitrinite	88.0	89.0
Inertinite	9.0	10.0
Liptinite	3.0	1.0

^aThe data were provided by the Argonne National Laboratory Premium Sample Program.

Illinois No. 6 and Wyodak coal contains five and eight phenolic hydroxyl groups per 100 mol carbon, respectively.¹¹ These phenolic groups were exhaustively alkylated, either by the method of Liotta¹¹ or by Ettinger's method.¹² With Liotta's method, tetrabutylammonium hydroxide was used as the base and with Ettinger's method, potassium hydroxide was the base. The results with the two different bases were comparable, except for the minor retention of tetrabutylammonium salts in Liotta's method.¹² The alkylated products were characterized by spectroscopy. For example, the infrared spectra of the products of both the coals showed substantial decrease in the hydroxyl stretching frequencies near 3400 cm⁻¹, together with increases in the carbon-hydrogen stretching frequencies near 2850-2900 cm⁻¹. The total number of alkyl groups introduced per 100 mol carbon was determined from the increase in weight and from the elemental analysis of the product.

In order to verify the influence of O-acylation reactions instead of O-alkylation, some acylation reactions were performed on the Illinois No. 6 coal. Instead of using aqueous potassium hydroxide or aqueous tetrabutylammonium hydroxide as base, which would hydrolyze the acylating agent, these reactions were carried out in anhydrous tetrahydrofuran using potassium trimethylsilanolate as base. Potassium trimethylsilanolate is known to be an active reagent that can substitute for aqueous potassium hydroxide. To establish the effect of O-alkylation on the solubility of the product, the coal samples were alkylated with various different alkyl halides. The results of the effect of O-alkylation and O-acylation on the solubility of the two coals, Illinois No. 6 and Wyodak coal, are shown in Tables 2 and 3.

TABLE 2
EFFECT OF O-ALKYLATION AND O-ACYLATION
ON THE SOLUBILITY OF ILLINOIS NO. 6 COAL

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db ^a	solubility, wt% in pyridine
1. pristine coal	-	-	27
2. coal, tetrabutylammonium hydroxide, THF, 25 °C, 48 hrs.	CH ₃ I	113	34
3. coal, aqueous potassium hydroxide, THF, 25 °C, 48 hrs.	CH ₃ I	109	35
4. coal, aqueous potassium hydroxide, THF, 25 °C, 48 hrs.	C ₄ H ₉ I	112	38
5. coal, aqueous potassium hydroxide, THF, 25 °C, 48 hrs.	C ₈ H ₁₇ I	117	33
6. coal, aqueous potassium hydroxide, THF, 25 °C, 48 hrs.	C ₁₆ H ₃₃ I	121	36
7. coal, potassium trimethylsilanolate, THF, 25 °C, 48 hrs.	C ₃ H ₇ COCl	114	38
8. coal, potassium trimethylsilanolate, THF, 25 °C, 48 hrs.	C ₇ H ₁₅ COCl	119	36

^aThe increase in weight is directly related to the number of alkyl/acyl groups that were introduced in the coal.

TABLE 3
EFFECT OF O-ALKYLATION ON THE SOLUBILITY OF WYODAK COAL

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db ^a	solubility, wt% in pyridine
1. pristine coal	-	-	9
2. coal, tetrabutylammonium hydroxide, THF, 25 °C, 48 hrs.	CH ₃ I	119	35
3. coal, tetrabutylammonium hydroxide, THF, 25 °C, 48 hrs.	C ₄ H ₉ I	123	39
4. coal, tetrabutylammonium hydroxide, THF, 25 °C, 48 hrs.	C ₈ H ₁₇ I	125	37
5. coal, tetrabutylammonium hydroxide, THF, 25 °C, 48 hrs.	C ₁₆ H ₃₃ I	131	39

^aThe increase in weight is directly related to the number of alkyl groups that were introduced in the coal.

The O-alkylation completely disrupts the hydrogen bonding interactions that are present in coal. Thus the coal fragments that were bonded by only hydrogen bonding can dissociate and are extracted into pyridine. Although there is only a 7-10% increase in solubilization by the O-alkylation of Illinois No. 6 coal, the solubility of Wyodak coal increases from 9% (raw coal) to about 35%. Evidently hydrogen bonding interactions between the phenolic constituents in Wyodak coal are more important than those in Illinois No. 6 coal.

The results of O-acylation are not very different from O-alkylation. The products obtained have the same solubility as the O-alkylated ones. Disruption of hydrogen bonds in the coals clearly enhance their solubility, and the gain with the Wyodak coal is especially large. However, we sought an approach that would lead to even greater solubility and studied stronger bases, which could abstract protons from carbon acids as

well as oxygen acids. The coal anions were quenched with several different electrophiles including labeled reagents to determine the sites of alkylation.

Alkylations with amide bases

Amides were the most reasonable first choice for C-alkylation under mild conditions. The high rank coal taken for our study was the Lower Kittanning coal, PSOC 1197, from the Pennsylvania State University Sample Bank. The elemental analysis data for the coal sample is shown in Table 4. Only negligible amounts of phenolic groups are present in this coal, and all the alkylation reactions take place on carbon atoms.

TABLE 4
ANALYTICAL DATA FOR THE HIGH RANK COAL SAMPLES.

Lower Kittanning Coal PSOC 1197 ^a	
ultimate analysis, wt% daf	
C	89.6
H	4.8
N	1.8
S	1.2
O (by difference)	2.6
ash	10.3
maceral analysis, wt%	
Vitrinite	94.4
Inertinite	5.6
Liptinite	0.0

^aThe data were provided by the Pennsylvania State University Sample Program.

The first reactions were carried out to evaluate the utility of amide bases for alkylation reactions and the results are presented in Table 5.

TABLE 5

INFLUENCE OF THE NATURE OF BASE ON BASE PROMOTED BUTYLATION OF LOWER KITTANNING COAL UPON ITS SOLUBILITY

base used	pKa	solvent	reaction temperature, °C	alkylated coal recovered, wt%, db ^a	solubility in pyridine, wt% ^b
NaN[(CH ₃) ₃ Si] ₂	26	NH ₃	-75	105	16
NaNH ₂	35	NH ₃	-75	109	50
LiN[(CH ₃) ₂ CH] ₂	36	NH ₃	-75	117	53
<i>n</i> -BuLi/TMEDA	42	THF	25	102	10

^aThe increase in weight is directly related to the number of butyl groups that were introduced in the coal.

^bApproximately 5 wt% of the original coal could be extracted in pyridine.

From the preliminary experiments performed, the results of which are presented in Table 5, we decided to use sodium amide in liquid ammonia as the base for subsequent alkylations. The next few reactions were carried out to find the optimum concentration of the base to be used for alkylation. Once again, the high rank coal was used for the reaction and all the coal anions were treated with butyl iodide to maintain uniformity. The concentration of the base used for alkylation, sodium amide, was varied from 0 to 200 mmole per gram of coal, and the solubilization of the alkylated products were observed. Table 6 represents the results for PSOC 1197 coal.

TABLE 6

EFFECT OF CONCENTRATION OF BASE USED FOR C-BUTYLATION
ON THE SOLUBILITY OF LOWER KITTANNING COAL.

concentration of NaNH ₂ , mmoles per gram coal	alkylated coal recovered, wt%, db ^a	solubility, wt% (daf)	
		Pyridine	CHCl ₃
0	-	5	-
15	102	20	-
45	105	50	26 ^b
100	121	60	-
200	124	61	28 ^c

^aThe increase in weight is directly related to the number of alkyl groups that were introduced in the coal.

^bAverage molecular weight was 6600.

^cAverage molecular weight was 4500.

The results in Table 6 indicate that 45 millimole of base per gram of coal yields the best results. Beyond 45 millimole of base per gram of coal, the solubilization is essentially independent of the base concentration. Thus, we used only 45 mmols of base per gram of coal, in all our subsequent experiments.

Once the optimum conditions for the alkylation reactions were established, the coal sample was alkylated with different alkyl groups to study the relationship between the nature of the alkyl group that was introduced and the solubility. As the results given in the next few tables will show, the length of the alkyl group had a major influence on the solubility, particularly for the high rank coals. The observations for the Lower Kittanning coal, PSOC 1197, are shown in Table 7.

TABLE 7
INFLUENCE OF THE ALKYL GROUP ON PYRIDINE SOLUBILITY
OF LOWER KITTANNING COAL

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db	pyridine solubility, wt%
pristine coal	-	-	5
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs.	NH ₄ Cl	-	9
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	CH ₃ I	119	20
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	C ₄ H ₉ I	107	50
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	C ₈ H ₁₇ I	113	90

As we had mentioned earlier, in order to study the exclusive effect of C-alkylation on the solubilization of some low rank coals, the O-methylated coals were used as starting materials. The more acidic oxygen sites were blocked by forming ethers so that only carbanions would be formed during base treatment. Only the subbituminous Wyodak coal was studied for the sodium amide promoted alkylation reactions. Since the results of solubilization were not very encouraging, the other low rank coal, Illinois No. 6, was not investigated.

The results of C-alkylation reactions on the O-methyl Wyodak coal are shown in Table 8.

TABLE 8
INFLUENCE OF THE ALKYL GROUP ON PYRIDINE SOLUBILITY OF THE O-METHYLATED WYODAK COAL.

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db ^a	pyridine solubility, wt%
pristine coal	-	-	9
O-methyl coal	-	-	35
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs.	NH ₄ Cl	-	25
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	CH ₃ I	119	31
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	C ₄ H ₉ I	124	38
NaNH ₂ , liquid NH ₃ , -75 °C, 6 hrs	C ₈ H ₁₇ I	123	43

^aThe increase in weight is directly related to the number of alkyl groups that were introduced in the coal.

The C-octylated Wyodak coal was 43% soluble in pyridine, but C-alkylation does not increase the solubility for this coal greatly beyond what can be achieved by O-alkylation (35% solubilization by O-methylation).

After the completion of the experiments with sodium amide systems, we shifted our attention towards other stronger bases. Alkylolithium bases were examined but the initial experiments were unsuccessful.¹⁴ However, after additional preliminary work, we found that a 1:1 mixture of *n*-butyllithium and potassium *tert*-butoxide (called Lochmann's base for convenience) was more effective.¹⁵ The next paragraphs will describe our reactions with Lochmann's base on the four different coal samples.

Alkylations with Lochmann's base

The use of alkylolithium bases for alkylation of coal is not new. Lazarov and his coworkers used *n*-butyllithium in tetrahydrofuran for the alkylation of some carbon-rich

Bulgarian and German coals.⁸ They reported that the maximum solubility in tetrahydrofuran, 38% was obtained for a bituminous coal containing about 88% C. Chambers and his coworkers performed C-alkylation reactions on the Lower Kittanning coal, PSOC 1197, with three alkyllithium bases with nominal pKa values between 19 and 31.^{9,16} They carried out the reactions in a selective way to investigate the distribution of carbon acids in this coal, and only low solubilities were realized for the thrice-alkylated reaction products.^{9,16}

Our initial attempts to use *n*-butyllithium in tetrahydrofuran or *n*-butyllithium with tetramethylethylene diamine in tetrahydrofuran failed miserably, and the reactions produced much less soluble material than desired.¹⁴ In retrospect, we realized that these attempts were doomed because of the rather high reactivity of the base with tetrahydrofuran. After additional preliminary work, we found that a 1:1 mixture of *n*-butyllithium and potassium *tert*-butoxide (Lochmann's base) in heptane was more effective.¹⁵ *n*-Butyllithium is itself a strong base; however, it has been shown that potassium *tert*-butoxide enhances its basicity in an unprecedented manner.^{17,18} When treated with this base mixture, hydrocarbons in the low acidity range, pKa 35-50, undergo clean hydrogen-metal exchange.¹⁷ The reasons for the success of this reagent have not been established, and several viable interpretations have been advanced by other workers in the field. Specifically, it has been suggested that the alkylpotassium compound, which can be formed in an exchange reaction and can be isolated under suitable conditions, is the active organometallic reagent.¹⁸ Schlosser, however, has shown that such compounds, for example *n*-butylpotassium, are inferior to Lochmann's base.¹⁷ He alternatively has suggested that the alkoxide anion enhances the reactivity of the alkyllithium by the formation of mixed clusters of the organometallic compound and the metallic alkoxide as shown in Figure 1.¹⁷ It has also been proposed that the alkoxide anion kinetically enhances the proton transfer reactions of the thermodynamically strong alkyllithium base.¹⁹

FIGURE 1.
POSSIBLE STRUCTURES FOR THE TRUE REAGENT IN THE LOCHMANN'S BASE

We were led to investigate Lochmann's base by the work of Bates and his associates, who exploited the high basicity of this reagent to convert dimethylphenols to diethylanisoles.¹³ In this present work, Lochmann's base has been applied for the C-alkylation of the one high rank bituminous coal, one low rank bituminous and a subbituminous coal. The work with the higher rank coal with low oxygen content and very low phenolic content, was successful. Both the unmodified lower rank coals were studied with this base. Subsequently, for the reasons mentioned earlier, the O-methylated derivatives of these low rank coals were also subjected to C-alkylation with Lochmann's base. The results for each of the different coal samples will be given separately.

The first set of reactions were performed on the Lower Kittanning coal, PSOC 1197, and the results are shown in Table 9. From our previous experience with sodium amide, only 45 mmols of the base per gram of coal was used. The base was treated with the coal sample for about 6 hours in refluxing heptane, prior to the addition of the alkylating agent. The coal samples were dried at 110 °C under vacuum for 48 hours prior to use. All the reactions were carried out in dry dinitrogen atmosphere.

TABLE 9
RESULTS FOR THE C-ALKYLATION REACTIONS OF PSOC 1197
WITH LOCHMANN'S BASE

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db	pyridine solubility, wt%
pristine coal	-	-	5
<i>n</i> -C ₄ H ₉ Li, THF, 25 °C, 6 hrs.	C ₄ H ₉ I	105	10
<i>n</i> -C ₄ H ₉ Li + K ^t OC ₄ H ₉ (1:1),heptane, 98 °C, 6 hrs	NH ₄ Cl	-	39
<i>n</i> -C ₄ H ₉ Li + K ^t OC ₄ H ₉ (1:1),heptane, 98 °C, 6 hrs	C ₄ H ₉ I	112	90
<i>n</i> -C ₄ H ₉ Li + K ^t OC ₄ H ₉ (1:1),heptane, 98 °C, 6 hrs	C ₈ H ₁₇ I	117	92

After carrying out the alkylation reactions with Lochmann's base on the high rank coal, we turned our attention towards the lower rank coals. In order to avoid the complications of high negative charge density on the phenolic groups, the low rank coals were methylated prior to C-alkylation. The O-methylated coal samples were then subsequently treated with Lochmann's base followed by the alkyl halide.

First, we shall discuss the results on the Wyodak coal sample. The coal was treated with Lochmann's base in refluxing heptane for six hours, then treated with the alkylating agent at 0-5 °C. The results are given in Table 10.

TABLE 10
RESULTS FOR THE C-ALKYLATION REACTIONS OF O-METHYL WYODAK
COAL WITH LOCHMANN'S BASE.

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db ^a	pyridine solubility, wt%
pristine coal	-	-	9
O-methylated coal	-	-	35
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs.	NH ₄ Cl	-	23
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs	C ₄ H ₉ I	108	16
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1),heptane, 98 °C, 6 hrs	C ₈ H ₁₇ I	111	20
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1),heptane, 98 °C, 6 hrs	C ₈ H ₁₇ I	113	44 ^b

^aThe increase in weight is directly related to the number of alkyl groups that were introduced in the coal.

^bAbout 100 mmoles of base, instead of the usual 45 mmoles, was used in this experiment.

Contrary to the results obtained for the high rank coals, the results for the Wyodak coal are not uniform, and in some cases, quite surprisingly, the products are less soluble than the starting materials. Clearly, such observations imply base promoted coupling and

condensation reactions that cross link the coal structure. However, we did find that C-octylation produced a coal that was 44% soluble in pyridine.

The next reactions with Lochmann's base were carried out on the Illinois No. 6 coal sample. Once again the O-methylated coal was used as the starting material. All the reactions were performed with 45 mmols of the base per gram of coal; the coal was refluxed with the base in heptane for six hours and then quenched with the electrophile at 0-5 °C. The results for various electrophiles added to the coal anion are shown in Table 11. In this case, methylation, butylation or octylation provided products that were 39 to 46% soluble in pyridine.

TABLE 11

RESULTS FOR THE C-ALKYLATION REACTIONS OF O-METHYL ILLINOIS NO. 6 COAL WITH LOCHMANN'S BASE.

reaction conditions: base, solvent, temperature, time	electrophile	alkylated coal recovered, wt%, db ^a	pyridine solubility, wt%
pristine coal	-	-	27
O-methylated coal	-	-	34
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs.	NH ₄ Cl	-	40
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs	CH ₃ I	108	46
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs	C ₄ H ₉ I	115	39
<i>n</i> -C ₄ H ₉ Li+K'OC ₄ H ₉ (1:1), heptane, 98 °C, 6 hrs	C ₈ H ₁₇ I	119	43

^aThe increase in weight is directly related to the number of alkyl groups that were introduced in the coal.

Thus after the completion of the series of reactions with Lochmann's base on all the three different coal samples, we were only partially successful in the solubilization of coal in pyridine. The high ranking coal was comparatively easy to solubilize; almost quantitative solubilization was achieved with Lower Kittanning coal. However, the low rank coals approached only 45% solubility. In order to obtain a greater degree of alkylated product together with high solubilization, other reactions were studied with

Lochmann's base at other reaction conditions. The starting material was the O-methylated Illinois No. 6 coal and in all the experiments, the coal anion was quenched with butyl iodide to maintain uniformity. The results are given in Table 12.

TABLE 12
EFFECT OF THE REACTION CONDITIONS ON PYRIDINE SOLUBILITY OF
O-METHYL ILLINOIS NO. 6 COAL

solvent of reaction	reaction temperature (°C)	nature of change in reaction condition	butyl groups per 10 ³ mol C	pyridine solubility wt%
heptane	98	alkyl halide added to coal anion.	4.5	39
pentane	25	alkyl halide added to coal anion.	4.0	41
heptane	98	coal anion added to the solution of alkyl halide (inverse addition).	4.5	32
heptane	98	alkyl halide added to the coal anion, as usual, but the reaction mixture was quenched after 1 hr. instead of the usual 48 hrs.	3.5	33

Use of pentane as the solvent of reaction instead of heptane, does not have a very significant effect on the solubility of the product. However, if the alkylation reaction is stopped after one hour, the product obtained is only 33% soluble in pyridine compared to about 40% solubility obtained if it were carried out for the usual 48 hours. Even when the coal anion is added to the alkyl halide (inverse addition), the degree of alkylation does not increase considerably. Next, a few reactions were conducted with THF solvent at different reaction temperatures. Even though the reactions were carried out at -46 to 67 °C, the products obtained did not show a temperature dependence. The pyridine solubility of the products remained at 33±3% as shown in Table 13.

TABLE 13
RESULTS FOR C-BUTYLATION OF O-METHYLATED ILLINOIS NO. 6 COAL IN
TETRAHYDROFURAN

reaction conditions: base, solvent, time	reaction temperature, °C	butylated coal recovered, wt%, db	pyridine solubility, wt%
$n\text{-C}_4\text{H}_9\text{Li} + \text{K}'\text{OC}_4\text{H}_9(1:1)$, THF, 6 hrs	67	114	33
$n\text{-C}_4\text{H}_9\text{Li} + \text{K}'\text{OC}_4\text{H}_9(1:1)$, THF, 6 hrs	25	117	36
$n\text{-C}_4\text{H}_9\text{Li} + \text{K}'\text{OC}_4\text{H}_9(1:1)$, THF, 6 hrs	-46	113	30

Thus, significant changes in the reaction conditions for alkylation have little effects on the degree of alkylation and solubilization of this coal.

The next reactions were performed by changing the solvent of reaction. All the reactions that were performed earlier with Lochmann's base were carried out in hydrocarbon solvents. Solvents such as tetrahydrofuran are well known for coal swelling. Hence, if the reactions could be carried out in tetrahydrofuran, probably more reactive sites in coal would be exposed for alkylation. However, with the ether solvents such as tetrahydrofuran, undesirable reactions such as α -lithiation followed by ring cleavage, consume the base.²⁰

Nevertheless, three reactions at different temperatures with tetrahydrofuran were performed on the O-methyl Illinois No. 6 coal. The results, which were shown in Table 13, can also mean that tetrahydrofuran is an inappropriate solvent. However, the decomposition of tetrahydrofuran can be avoided by the use of a 2,2,5,5-tetrasubstituted

tetrahydrofuran. Inasmuch as the four hydrogen atoms α to the oxygen atom are absent, the base cannot decompose this ether. 2,2,5,5-Tetramethyl tetrahydrofuran should have all the properties of the tetrahydrofuran. Similar arguments can also be made for tetrahydropyran as solvent. Though it has α -hydrogen atoms, which can be removed by base, the anion does not decompose as rapidly because it does not produce a stable product. Hence, both these solvents were studied with the Lochmann's base to alkylate the O-methyl Illinois No. 6 coal. The results are given in Table 14. The results with heptane and tetrahydrofuran are also included for comparison.

TABLE 14
INFLUENCE OF THE SOLVENT OF C-BUTYLATION ON SOLUBILITY
OF THE O-METHYL ILLINOIS NO. 6 COAL

solvent of alkylation	reaction temperature	butylated coal recovered, wt%, db	pyridine solubility, wt%
heptane	98 °C	118	39
tetrahydrofuran	67 °C	113	33
tetramethyl- tetrahydrofuran	112 °C	117	45
tetrahydropyran	88 °C	119	51

The solubility of the products that were obtained with the novel ether solvents were higher than those obtained with heptane as solvent, and 51% solubilization was achieved in tetrahydropyran. We infer that the reason that both 2,2,5,5-tetramethyltetrahydrofuran and tetrahydropyran gave such encouraging results is because of their stability in strongly basic media. Due to the similarity of these solvents to tetrahydrofuran, they can permeate the coal structure and help transport the reagents into the insoluble solid matrix. Similar results were also obtained in different experiments when the coal sample was C-octylated in these solvents. For example, it was shown that C-octylation in tetrahydropyran gave 52% solubilization compared to only 43% in heptane. Thus, it can be concluded that the C-alkylation reactions with Lochmann's base are more successful in solvents like tetrahydropyran rather than the hydrocarbon solvents.

Several lines of evidence suggested that the bases that were being used might be too powerful and cause undesirable condensation side reactions. In particular, for the low rank coals, treatment with Lochmann's base reduced the solubility in some experiments. Further, the results

also implied that deep seated hydrogen transfer reactions accompanied the preparations of the coal anions. For example, Lochmann's base might abstract an aromatic hydrogen from a coal molecule, the reactive anion, in turn, might engage in a variety of condensation reactions. It therefore appeared desirable to conduct coal anion preparations in the presence of proton donor solvents. This feature was best realized by the use of sodium and potassium alkoxides in refluxing alcohols. The results for a series of experiments are presented in Table 15.

TABLE 15
RESULTS FOR ALKOXIDE TREATMENT OF ILLINOIS NO. 6 COAL IN REFLUXING ALCOHOLS

reaction conditions: base, solvent, time	reaction temperature, °C	coal recovered, wt%, db	pyridine solubility, wt%
raw coal	-	-	27
NaOMe (40 mmol/g), MeOH, 24 h, protonation	65	86	33
KOn-Bu (40 mmol/g), <i>n</i> -BuOH, 24 h, protonation	118	76	42
KOr-Bu (40 mmol/g), <i>i</i> -BuOH, 24 h, protonation	83	83	42
KOn-Oct (40 mmol/g), <i>n</i> -OcOH, 24 h, protonation	196	79	53
KOBz (40 mmol/g), BzOH, 24 h, protonation	205	43 ^a	82
NaOBz (40 mmol/g), BzOH, 24 h, protonation	205	51 ^a	74
KOBz (40 mmol/g), BzOH, 24 h, protonation	205	95	66

^aThe recovery of the products after the reaction was low since a significant portion of the coal was soluble in the alcohol, which was removed during washing and filtration.

The results presented in this Table indicate that the concept was successful.

DISCUSSION

Alkylations with hydroxide bases

All the mild bases used for alkylation such as potassium hydroxide and tetrabutylammonium hydroxide are known to alkylate selectively on the acidic oxygen atoms present in coal. However, evidence for some C-alkylation reactions with tetrabutylammonium hydroxide at long reaction times has been reported.²¹ But the extent of C-alkylation of coal is very modest, and it is only a minor side reaction. The O-alkylation reactions were performed only on the low rank coals in this study.

As shown in Table 2, the solubility of the O-methyl derivative of Illinois No. 6 coal increases from 27% (for the raw coal) to 34%. Almost all the O-alkylated or O-acylated derivatives show a definite increase in solubility. The solubility does not depend appreciably on any other factors such as the length of the alkyl group. The O-alkylation reaction merely disrupts the hydrogen bonding interactions present in the coal. Thus, the molecular fragments in the coal macromolecules that are linked only by hydrogen bonding are dissociated and some fragments can be extracted in pyridine. Obviously the disruption of hydrogen bonds did not depend significantly upon the length of the alkyl group. Although there is only a 7-10% increase in solubilization by O-alkylation for Illinois No. 6 coal, this number is significantly larger for Wyodak coal as noted in Table 3. The solubility of the coal increases from 9% (raw coal) to about 35% - 39% for the O-alkylated coals. Evidently hydrogen bonding interactions between the macromolecular components are more important in Wyodak coal, than in the Illinois No. 6 coal. As evident from Table 1, Wyodak coal has about 18 oxygen atoms per 100 mol carbon, compared to only 12 for Illinois No. 6 coal. Hence, as can be predicted, the influence of hydrogen bonds in Wyodak coal would be of more significance than the Illinois No. 6 coal.

The site of alkylation was studied by using ¹³C-labeled compounds. At least three different oxygen centers, hindered phenols, unhindered phenols and carboxylic acids, were alkylated during the O-alkylation. Our results regarding the nature of the hydroxyl functional groups in Illinois No. 6 coal that were alkylated, are compatible with the results obtained by Stock and Willis.²² The intention was to locate the sites of alkylation in a more direct way than had been done in the previous study.²² Heteronuclear two-

dimensional (2D) spectroscopy would provide a method for the assignment of introduced methyl groups in the carbon and proton spectra of the methylated coal. The HETCOR spectra were not obtained as a routine run. The number of acquisitions had to be optimized because every time the run was carried out overnight, the computer could not store the data. The lower number of acquisition did not yield a good carbon projection. However, a compromise between the number of acquisitions and good carbon resolution was obtained after a few trials, and the plots are shown in Figures 2 and 3.²³

The coal was alkylated with tetrabutylammonium hydroxide and methyl- ^{13}C -iodide. The use of such two-dimensional NMR techniques is not common in coal chemistry because of the complex molecular structures. Since the interest was on the sites of O-alkylation, the resonances corresponding to aryl and alkyl methyl ethers were considered. These resonances, although very close in the carbon spectrum, are in different regions in the proton spectrum. The 2D contour, cross-correlation spectrum of the protons coupled to the labeled carbon nuclei provided evidence of the formation of aryl and alkyl ethers. Cross-correlation of the proton-carbon resonances for non-hindered aryl methyl ethers was apparent from the cross peaks at $x = 55$ to 56 ppm and $y = 3.5$ to 4.1 ppm. Cross-correlation of the proton-carbon resonances for hindered aryl methyl ethers was apparent by the cross peaks at $x = 59$ ppm and $y = 3.5$ ppm. Further, there was considerable intensity near 49 to 50 ppm in the 1D carbon spectrum, which could be assigned to the methyl esters. Thus, we see that three different oxygen sites are being alkylated by tetrabutylammonium hydroxide in the Illinois No. 6 coal. The poor resolution made it impossible to study cross-correlation associated with C-methylation.

The other low rank, high oxygen coal, Wyodak subbituminous coal, was similarly alkylated with tetrabutylammonium hydroxide and methyl- ^{13}C -iodide to determine the oxygen sites being alkylated. The solid state ^{13}C NMR spectrum of the alkylated product is shown in Figure 4. For the sake of convenience, the ^{13}C spectrum of the raw coal and the difference spectrum were also plotted. As evident from the figure, once again similar to the Illinois No. 6 coal, three different oxygen centers are methylated; the signals range in the frequencies between 50 to 60 ppm. Though quantitative values for the number of alkyl groups introduced in the coal could not be determined by this method, the position of the introduced methyl group could be assigned unambiguously.

FIGURE 2.
HETERONUCLEAR CORRELATION SPECTRUM OF THE PYRIDINE SOLUBLE PART
OF O-METHYL ^{13}C -ILLINOIS NO. 6 COAL. THE TOTAL SPECTRUM IS FLANKED
BELOW BY CARBON PROJECTION AND VERTICALLY ALONG THE CONTOUR PLOT
BY PROTON PROJECTION.

FIGURE 3.
HETERONUCLEAR CORRELATION SPECTRUM OF THE PYRIDINE SOLUBLE PART
OF O-METHYL ^{13}C -ILLINOIS NO. 6 COAL. THIS MORE SELECTIVE SPECTRUM
SHOWS ONLY THE REGION OF INTEREST.

FIGURE 4.
SOLID STATE ^{13}C SPECTRA OF WYODAK COAL: (A) O-METHYL ^{13}C WYODAK
COAL, (B) PRISTINE WYODAK COAL, AND (C) THE DIFFERENCE SPECTRUM.

Alkylations with amide bases

After the completion of the alkylations with mild bases, we turned our attention towards the stronger bases. These reactions were performed on all the coal samples. Different solubilization results were obtained for different products, as had been shown in the results section, and a thorough discussion of all the results is given below. Once again, in some reactions the coal anions were quenched with labeled reagents to determine the site of alkylation. We shall start our discussion with the results obtained for amide bases.

First, we shall discuss the results obtained by the alkylation reactions on the Lower Kittanning coal, PSOC 1197. As evidenced from Table 5, sodium amide in liquid ammonia at -75°C seems to be the best choice of base. The weakest base, sodium hexamethyldisilazide, $\text{pK}_a = 26$,²⁴ provides only 16% soluble material, but the stronger bases, sodium amide, $\text{pK}_a = 35$ ²⁵ and lithium diisopropyl amide, $\text{pK}_a = 36$,²⁴ provides much greater yields of soluble products. These observations imply that the strength of the base is very significant in the solubilization reaction. The use of a strongly basic reagent, which would have also formed the coal carbanions irreversibly, the *n*-butyllithium / tetramethylethylenediamine, $\text{pK}_a = 42$, in tetrahydrofuran, turned out to be a failure. The reason for the failure appear to be the competitive reactions of the reactive coal anions that are formed in kinetically controlled reactions, with the solvent. The solvent, tetrahydrofuran, is consumed via the formation of tetrahydrofuran anion, which decomposes irreversibly as shown earlier.

Once it was established that sodium amide in liquid ammonia could be used for alkylation reactions, the concentration of the base needed for the reaction became important. Table 6 shows the effect of concentration of base used for alkylation on the solubility. The results for 45, 100 and 200 mmol of base per gram of the coal gave essentially the same solubility in pyridine and in chloroform. In addition, we found that the number average molecular weight of the material in the chloroform extracts decreased from 6600 with 45 mmol of the base to 4500 with 200 mmol of the base. This observation suggests that high concentrations of base favor bimolecular anionization reactions that lead to fragmentation. Since our main intention was to assess exclusively the role of C-alkylation on solubilization, a low concentration of the base, 45 mmol, was used in all our subsequent reactions. Other additional reactions involving carbanion formations and their subsequent bimolecular fragmentation reactions were studied and will be discussed later.

A few reactions were carried out with the sodium amide system by changing the reaction temperature; it was observed that a large change in the temperature had only a very small influence on the course of reaction. In sharp contrast, the length of the alkyl group introduced in the coal has a major effect on the outcome of the reaction. The results for protonation and alkylation of the Lower Kittanning coal, PSOC 1197, are presented in Table 7.

When the reaction was terminated with ammonium chloride, the product was only 9% soluble in pyridine. This result implies that the reaction of PSOC 1197 with a modest concentration of sodium amide does not lead to hydrocarbon elimination reactions or other base catalyzed fragmentation reactions.

This feature of chemistry, though not observed with moderate concentrations of sodium amide, has been found to take place with much stronger base or with very high concentrations of sodium amide. More evidence for base promoted fragmentation reactions have been obtained and this information will be discussed later.

The chain length of the alkyl group can have an enormous influence on the conversion of coal to pyridine-soluble products. This influence is especially dramatic for the PSOC 1197 coal. On octylation, this coal provided a product that was 90% soluble in pyridine. It implies that the structural bulk of the added alkyl group is a key factor in determining the extractability of the alkylation product. However, similar enhancement in solubility is not observed for the other coals. Wyodak coal shows no influence of the alkyl groups. The O-methyl Wyodak coal was used as the starting material, which had about 35% soluble material, and the octylated product had only 43% soluble material. Thus, C-alkylation does not seem to influence the solubilization of the Wyodak coal.

Inasmuch as sodium amide is not sufficiently basic to abstract protons from aromatic compounds, we presume that it abstracts protons from aliphatic structural elements with $pK_a < 35$. Examination of the pK_a data for hydrocarbons that may have structural elements representative of coal molecules, Table 16, suggests that C-alkylation occurs on fluorenylic structures, hydroaromatic compounds and on benzylic structural elements. Similar suggestions have also been proposed by Gawlak and his associates⁶ and in one of our publications on this topic.¹⁰ This feature of chemistry was probed by the

TABLE 16

HYDROCARBONS AND THEIR pKa VALUES

Compound	pKa	Compound	pKa
19.9		19.6
23.0	(Ph) ₂ CH ₂	33.4
		(Ph) ₃ CH	31.5
19.8		30.6
21.4		31.8

examination of the ^{13}C -NMR spectra of the methylation products obtained for PSOC 1197 by using methyl- ^{13}C -iodide as shown in Figure 5.

Alkylations with Lochmann's base

We began our work on Lochmann's base with the Lower Kittanning coal, PSOC 1197, and the results are presented in Table 11. It is evident from the Tables 8 and 11 that the butylated products obtained through the reaction of *n*-butyllithium in tetrahydrofuran and through the reaction of sodium amide in ammonia were 10 and 50% soluble in pyridine, respectively. In contrast, butylation with Lochmann's base in heptane provided a 90% soluble material. The reason for the effectiveness of this base was explored by the determination of the degree of solubilization that could be realized by the treatment of coal with base and protonation. Unreacted coal is only 5% soluble in pyridine, and the reaction with sodium amide in liquid ammonia followed by protonation yielded a product that was 9% soluble in pyridine (Table 8). On the basis of this observation, it was concluded that modest concentrations of sodium amide did not significantly alter the molecular framework through hydrocarbon elimination reactions or other base catalyzed fragmentation reactions. However, when the reaction was performed with Lochmann's base in heptane and the intermediate was treated with ammonium chloride in methanol, the recovered coal was 39% soluble in pyridine. Clearly, the base alone must have altered the molecular structure either through base-catalyzed rearrangement reactions or more probably, through hydrocarbon elimination reactions that could reduce the dimensions of the macromolecule. More information regarding this aspect of bond cleavage will be discussed later.

However, the much higher degree of solubility (92%) that is achieved after treatment with base and C-octylation presumably results from the disruption of intermolecular polarization forces by the long alkyl groups, as discussed previously for the sodium amide system. The alkyl groups are probably placed in strategic positions to disrupt the stacking patterns of the coal. When the alkyl fragment is sufficiently large, the polarization bonding forces between the structural fragments of the coal can be disrupted and the mixture solubilized.

We subsequently explored the scope of the reactions of Lochmann's base by the study of the two lower ranking coals. Before we studied the other coal samples, a

FIGURE 5.
 ^{13}C NMR SPECTRA OF THE LOWER KITTANNING COAL PSOC 1197 (BOTTOM)
AND THE PRODUCT OBTAINED IN THE SODIUM AMIDE PROMOTED
METHYLATION WITH METHYL- ^{13}C IODIDE (TOP).

representative compound was chosen, which was treated with Lochmann's base under coal conditions. The low rank coals contain large amounts of oxygen atoms, and a significant portion of them occur in heterocycles.²⁶ Hence, we studied dibenzofuran to investigate the chemical reactions that might occur with these substances in the coal under the influence of Lochmann's base. In the first experiment, dibenzofuran was treated with Lochmann's base in refluxing heptane for six hours and the reaction mixture was quenched with ammonium chloride and methanol. Analysis of the product showed that 95% of the starting material was recovered, and about 5% of the cleavage product, 2-phenylphenol, was obtained. The cleavage product, 2-phenylphenol, was extracted with sodium bicarbonate and was identified.

In the next experiment, the reaction mixture was treated with methyl iodide. Since the reactions were performed in a similar manner to that of coal, excess base as well as excess methyl iodide was present. Many products, including mono-, di-, tri-, and even heptamethyl derivatives were observed in the GC-MS analysis. A reaction pathway for the permethylation reaction is shown in Scheme 1. Initially, a hydrogen atom is abstracted from the aromatic ring to provide the ortho-metallated compound. The initial methylation reaction yields 2-methyldibenzofuran. The methyl hydrogen atoms are benzylic and more acidic than the aromatic hydrogen atoms. Thus, subsequent anionization occurs at the benzylic position and the second methyl group is introduced selectively at the benzylic carbon atom.

Scheme 1. Proposed mechanism for the introduction of more than one methyl group in dibenzofuran.

Finally, the alkylation reactions were carried out for the two low rank coals, Illinois No. 6 and Wyodak coal. Since the main emphasis was to study the influence of C-alkylation on solubilization, the O-methylated coals were used as the starting materials for most of the reactions. The results for the subbituminous Wyodak coal are shown in Table 13.

Surprisingly, sodium amide provides a more soluble product than Lochmann's base for both pristine and O-methylated Wyodak coal. The pristine coal is only 9% soluble in pyridine. Lochmann's base promoted octylation of Wyodak coal produces 33% soluble material, but mere O-octylation with tetrabutylammonium hydroxide produces even more soluble material, 37%. With sodium amide, the product is about 43% soluble in pyridine.

However, as Table 13 shows, greater concentration of the base (100 mmols per gram of coal instead of the usual 45 mmols) leads to a slight increase in solubility of the product. The solubility of the product increases to 44% by a single octylation with 100 mmols of the base per gram of coal. As much as the number of introduced alkyl group does not change significantly, this enhancement in solubility has to be due to bond cleavage reactions. Smaller fragments are thus formed due to carbon-carbon and carbon-heteroatom bond cleavage, which are further extracted by pyridine.

The other low rank coal bituminous Illinois No. 6 coal, APCSP 3, was used in many subsequent reactions. Once again, the O-methylated coals were used as starting materials for most of the reactions. The results for the C-alkylation reactions are shown in Table 14. The pyridine solubility of the products that were obtained after the reaction mixtures were quenched with electrophiles ranging from ammonium chloride to octyl iodide were not very different. There is no correlation between the nature of the alkyl group and solubility. Illinois No. 6 coal behaves entirely differently than Lower Kittanning coal in the sense that the length of the alkyl group does not alter solubility. The results that have been obtained for the C-alkylation of these two coals parallel the observations of Wachowska and her collaborators for the reductive alkylation of high and low rank coals.²⁷

We were surprised by the observation that no more than 2 to 4 C-alkyl groups were introduced into the coal, especially because dibenzofuran so readily experienced multiple alkylation. This compound and many other activated aromatic compounds are readily metallated by strong bases and alkylated.²⁸ Several experiments were conducted to investigate this feature of the reaction with coal. First, there was no evidence for the metallation of aromatic compounds in Illinois No. 6 coal. When the coal anion was quenched with deuterium oxide, no deuterium signal was observed in the aromatic region of the ²H-NMR spectrum of the products that were soluble in pyridine (Figure 6). Second, if the alkylation reaction proceeds with coal as it does with pure dibenzofuran, Scheme 1, further anionization is inevitable and ethyl and isopropyl derivatives should be produced. For one thing, the degree of alkylation would be much higher than what was obtained. Hence, we infer that although dibenzofuran undergoes metallation with Lochmann's base, such reactions do not take place in coal. This is quite reasonable because coal has other more acidic protons which are readily abstracted by the strong base, in preference to the aromatic protons.

The nature of the carbon acids in Illinois No. 6 coal was studied by methylation with methyl-¹³C iodide. The ¹³C NMR spectrum of the pyridine extract of the product, which contains about 46% of the organic matter, is shown in Figure 7.

Unfortunately, the resonance of the methyl-¹³C group span a broad range, and the spectroscopic results do not provide precise information on the sites of C-methylation. Nevertheless, certain aspects of the spectrum deserve mention. The very sharp signals near 47, 29, 20 and 12 ppm arise from the presence of residual tetrabutylammonium salts.

FIGURE 6.
 ^2H NMR SPECTRUM OF THE PRODUCT (PYRIDINE SOLUBLE PART) FROM THE
 REACTION OF O-METHYL ILLINOIS NO. 6 COAL WITH LOCHMANN'S BASE AND
 DEUTERIUM OXIDE.

FIGURE 7.
 ^{13}C NMR SPECTRUM OF THE PRODUCT (PYRIDINE SOLUBLE PART) FROM THE
REACTION OF O-METHYL ILLINOIS NO. 6 COAL WITH LOCHMANN'S BASE AND
METHYL- ^{13}C -IODIDE. THE REGION OF INTEREST IS BETWEEN 10 TO 60 PPM.

Most of the ^{13}C signal intensity appears from 11 to 28 ppm. Elements of this broad signal are consistent with Chamber's view^{9,16} that benzylic and fluorenylic positions are selectively methylated. Particularly strong signals occur near 18 and 25 ppm. Although it is unwise to make structural assignments on the basis of chemical shift data alone, these resonances are compatible with the products of methylation of fluorenyl derivatives, the resonance of the methyl group in 9-methylfluorene is at 18 ppm, and with the products of methylation of α -alkylbenzyl derivatives, the resonance of the methyl group in the simplest α -alkylbenzyl derivative, isopropylbenzene, is at 24 ppm. The broad band of resonance between 15 and 30 ppm are consistent with the methylation of a variety of compounds ranging from, for example, diarylmethanes, the resonance of the methyl group in 1,1-diphenylethane is at 22 ppm, to dihydrophenanthrenes, the resonance of the methyl group in 9-methyl-9,10-dihydrophenanthrene is at 20 ppm. The absence of prominent signals near 27 and 31 ppm, which could arise from 9,9-dimethylfluorene and 2,2-diphenylpropane, respectively, implies that dialkylation is not a significant reaction. In spite of the presence of excess base in the reaction medium, dimethylation does not appear to occur.

Previous studies have also shown that benzylic alkylations are prominent in reductive alkylation reactions.²⁹ One very significant difference that distinguishes Lochmann's base promoted C-alkylation from reductive C-alkylation deserves mention. During reductive alkylation, aromatic compounds are converted to dihydro derivatives which undergo methylation to yield substances that contain methyl groups bonded to quaternary carbon atoms of carbocycles as illustrated below.

Compounds of this kind exhibit resonances near 30 ppm. Such absorptions are minor in the spectrum of the C-alkylation products in Figure 7.

The resonance near 54 and 59 ppm reveal that O-methylation of the O-methylated Illinois No. 6 coal also takes place. About 4 methyl groups per 100 mol C are introduced during the C-alkylation, and the ratio of C- to O-methylation is approximately

7. These results suggest that about 0.5 new O-methyl groups per 100 mol C are introduced into the O-methyl Illinois No. 6 coal. Previous studies on anisole showed that it was stable to Lochmann's base and that orthometallation rather than carbon-oxygen bond cleavage takes place.¹³ Hence, O-methyl bond cleavage followed by remethylation with the labeled alkylating agent can be ruled out. The treatment of dibenzofuran with Lochmann's base under the conditions of coal alkylation reactions led to the formation of 5% 2-phenylphenol. Thus, we infer that new labeled O-methyl groups can arise through base promoted carbon-oxygen bond cleavage reactions in diaryl ethers and heterocycles. It is pertinent that reductive alkylation reactions cleave about two carbon-oxygen bonds per 100 mol C in Illinois No. 6 coal,²⁹ and that the products of reductive alkylation are considerably more soluble than those obtained from Lochmann's base alkylation. These results imply that the solubilization of Illinois No. 6 coal is significantly enhanced by carbon-oxygen bond cleavage and that such covalent bond cleavage reactions are more important for solubilization than the disruption of polarization forces or hydrogen bonding interactions by the introduction of O- and C-alkyl groups.

One of the most notable features of the work with Lochmann's base and other C-alkylation reactions centers on the observation that greater success is realized with higher rank coals. Lazarov and his associates pointed out this fact in their work with the European coals,⁷ and our work with Lower Kittanning coal supports this contention. There is a basic difference in the reactivity of the high and the low rank coals.

A few points about the nature of the Lochmann's base promoted C-alkylation need to be mentioned before we discuss the reason for this difference. The coal reaction system is quite complex, and an adequate interpretation of the results requires an appreciation of several aspects of chemistry. First, irreversible reactions occur between the carbon acids in the solid coal and the bases to provide butane and the coal anions which remain in the solid phase. Second, grafting of the base to coal through nucleophilic addition to saturated and unsaturated carbonyl compounds can occur. Third, concurrent proton transfer reactions take place between the anions that are formed first and the other carbon acids in the coal and promote the formation of an equilibrated mixture of the most thermodynamically stable coal anions. Further, the addition of the alkylating agent initiates a new set of reactions between the anions and the electrophile. These reactions are rapid, but proton transfer reactions of the carbon acids in coal may continue during this step. Under these circumstances, both kinetic and thermodynamic factors could influence the outcome of the solubilization experiment. Proton transfers occur reversibly,

but alkylation is an irreversible reaction. The situation is illustrated in Scheme 2 for two different carbon acids in coal.

Scheme 2. Curtin-Hammett analysis of coal alkylation.

The physical organic analysis of this problem is embodied in the Curtin-Hammett principle.³⁰ If the rates of the alkylation reactions with rate constants k_2 and k_3 are slow relative to the rates of proton transfer with rate constants k_1 and k_{-1} , then the ratio of the individual alkylated products, $\text{CoalR}_1\text{CH}_3/\text{CoalR}_2\text{CH}_3$, would reflect the relative activation energies for the alkylation reactions. However, if the rates of the alkylation reactions are fast relative to the rates of proton transfer, then the product ratio would depend on the relative thermodynamic stability of the carbanions. The more stable the anion, the greater the yield of its alkylation product. The simple situation for a homogeneous solution is greatly complicated by the heterogeneity of coal. Nevertheless, our results and Chamber's results appear to be best accommodated by an explanation that is based upon the reaction pathway in Scheme 2 and that presumes carbanion equilibration. For example, we were unable to detect products of alkylation at aromatic carbon atoms even though the bases that were used in the reaction were capable of abstracting protons from such positions. Thus, it is reasonable to presume that aryllithiums may be formed in initial reactions, but are converted to more stable substances in subsequent equilibrium controlled processes.

Reactions with alkoxide bases.

Because the carbanions that are produced by the very strong bases can also cause condensation reactions, we sought to reduce the carbanion concentration by using an entirely different system, alkali metal alkoxides in alcohols, and treated the low rank Illinois No. 6 coal with this base. Potassium or sodium alkoxides were used in different alcohols at

reflux temperatures for a definite period of time and then the products were treated with acid. The results are shown in Table 15.

Significant work has been done with alcohols as solvents and reagents for the liquefaction of coal. Use of low boiling alcohols under super critical conditions or in hydrogen donor solvents in the presence and absence of bases has been investigated by a number of workers. For the sake of brevity, we should only mention the recent summary of this work by Kuznetsov *et al.*³² They have discussed the work of Bartle, Ross, Makabe, Bimer, Winans, Mayo, Sternberg, and others.³²

It is well known that some investigators have neglected the possibility that alcohols adduct with coal molecules. In these new experiments, we measured the conversion exclusively on the basis of the weight of the insoluble coal residue.

The reaction of Illinois No. 6 coal with sodium methoxide in refluxing methanol yielded a product which was only 33% soluble in pyridine. Treatment with potassium *n*-butoxide and *t*-butoxides in *n*- and *t*-butanol with boiling points 118 and 83 °C, respectively, increased the solubility to 42%. There was no significant difference between the effectiveness of the two alcohols. Potassium *n*-octoxide in *n*-octanol (b.p.196 °C) gave even more soluble product, 53%. None of these treatments were as effective as potassium or sodium benzoate in refluxing benzyl alcohol (b.p.205 °C). More than 70% soluble material was obtained in this way.

It is quite evident that the strategy was successful. We believe that the success of this reaction originates in the addition of hydrogen from the alcohol and the subtle base catalyzed carbon-carbon elimination reactions as well as base-catalyzed retrocondensation reactions that depolymerize the low rank coals.

Role of Base-catalyzed bond cleavage reactions in pure hydrocarbons

We inferred that the high ranking coal samples undergo hydrocarbon elimination reactions when treated with Lochmann's base, thereby yielding a more soluble product. To verify that base promoted elimination reactions can take place with hydrocarbons, we performed reactions with selected pure compounds. Five different hydrocarbons, 1,1'-binaphthyl, bibenzyl, 1,1,3-triphenylpropane, 9-benzyl-9,10-dihydroanthracene and 9,10-diphenylanthracene were studied.

All the compounds were treated with Lochmann's base and the anionic system was quenched with dilute aqueous acid. The first two compounds, 1,1'-binaphthyl and bibenzyl were ideal examples of compounds linked through two sp^2 carbon atoms and two sp^3 carbon atoms, respectively. It was observed that both the compounds were recovered unchanged at the end of the reaction. Evidently, no bond cleavage reaction took place. However, two other compounds, 1,1,3-triphenylpropane and 9-benzyl-9,10-dihydroanthracene, did undergo decomposition. When 1,1,3-triphenylpropane was treated with Lochmann's base in heptane at 98°C for 6 hours followed by protonation, only 20% of the starting material was recovered. The products that were identified by GC were toluene (23%) and diphenylmethane (24%). It is evident that two different acidic benzylic protons are abstracted and that the two carbanions formed in reactions A and B undergo decomposition via β -elimination to give styrene together with another stable benzylic carbanion.

When 9-benzyl-9,10-dihydroanthracene was treated with the base, about 76% of it was recovered at the end of the reaction. The decomposition products were toluene (18%) and anthracene (22%). No other compounds were observed in the GC trace of the product mixture. Once again, we infer that the base abstracts a benzylic proton from the dihydroanthracene ring and that the carbanion undergoes vinylogous β -elimination.

Finally, when 9,10-diphenylanthracene was treated with the base, equally interesting results were obtained. Only 10% of this compound was recovered at the end of the reaction and the major products that could be characterized were 9-phenylanthracene, 9-butyl-10-phenylanthracene, 9,10-dihydro-9,10-diphenylanthracene and 9-butyl-9,10-diphenyl-9,10-dihydroanthracene. Many other minor products were formed and were characterized by the interpretation of their GC retention times and their mass spectrum. These substances are shown in Scheme 3.

Scheme 3. Products from the reaction of 9,10-diphenylanthracene with Lochmann's base. Certain identifications are tentative.

These results indicate that strongly basic reagents can cleave Calkyl-Calkyl bonds if stable benzylic carbanions can be formed. We presume that such β -elimination reactions can take place in coal. Thus, anionic depolymerization is possible, provided the structure permits. Also, Caryl-Caryl bonds can undergo cleavage due to nucleophilic attack of butyl group (from the base). Many of the reaction products of 9,10-diphenylanthracene are butylated. However, the reasonable mass balances indicate that such nucleophilic addition is minor with coal.

CONCLUSIONS

O-Alkylation reactions of the low rank coals increase their solubility, but the increase in the pyridine-solubility of Illinois No. 6 coal from 27% for the raw coal to 34% for the O-alkylated coals is not large. The change for the more phenolic, subbituminous Wyodak coal is significantly greater from 9% to 39%. Presumably, the O-alkylation reactions prevent inter- and intramolecular hydrogen bonding between structural elements of the coal. This disruption of hydrogen bonding is apparently responsible for the greater solubility of the Wyodak coal, but, even so, only 39% of the material becomes soluble and it is clear that other binding forces need to be overcome to achieve high solubilization.

Amide-promoted C-alkylation reactions successfully converted the high rank Lower Kittanning coal into soluble products. The stronger bases with pKa values near 35 are more effective than the weaker bases with pKa values near 25. The extractability of the products, in general, parallels the number of added alkyl groups, which, in turn, parallels the base strength. At least for the high ranking coals, the dimensions of the alkyl group have a very profound influence on the degree of solubilization. We infer that the alkyl groups are placed in strategic positions to disrupt the stacking patterns of the coal. When the alkyl fragment is sufficiently large, the polarization bonding forces between the structural fragments of the coal can be disrupted and the mixture solubilized. In view of the fact that the methyl group has essentially the same steric requirements as a sheet of aromatic molecules, it has little influence on the solubility of the coal. On the other hand, molecular modeling confirms that the larger butyl and octyl groups have much greater steric requirements that cannot be relieved by simple rotations. In this situation, favorable polarization interactions with the solvent exceed those between the molecules in the solid, and solubilization is realized.

Insofar as the lower rank coal samples are concerned, C-alkylation does not enhance the solubility of the products greatly. The introduction of a large alkyl group changes the solubility of the product, but the size of the alkyl group is not the deciding factor. The octyl groups have essentially the same effect as the methyl groups. Even though the added alkyl groups inhibit all hydrogen bonding interactions and may disrupt polarization forces, high solubility is not achieved.

C-Alkylation reactions with Lochmann's base converted the high rank coal almost quantitatively to soluble products. A very important fact to be noted for the Lochmann's base is that significant carbon-carbon bond cleavage reactions can take place. Such elimination reactions, together with the disruption of intermolecular forces, appear to be responsible for the high solubilization of the products. It should be pointed out that such hydrocarbon elimination reactions appear to be almost negligible with the sodium amide system. Our spectroscopic results suggest that the relatively more acidic protons, from benzylic and fluorenylic carbons in coal, are abstracted by the base, and are subsequently alkylated.

Analysis of the results suggested that the base-catalyzed reactions of the low rank coals (and possibly high rank coals) appeared to be less successful than they really were because the very strong bases promoted condensation reactions of the coal carbanions with other coal molecules. Accordingly, we examined the reactions of the low rank coals in media where the base concentration was high, but protons were available to trap the carbanions that would be found in elimination reactions. The results at temperatures near 200 °C with potassium benzoxide were very good with more than 80% of the Illinois No. 6 coal entering solution.

A major point regarding the differences in solubility of the low- and high-rank coals deserves mention. The differences in solubility of the alkylated high- and low-rank coals presumably depend upon an array of factors, including major differences in molecular structures of the original coals and the array of inter- and intramolecular forces that operate in the liquid and solid state. We believe that the differences in the effectiveness of C-alkylation for the solubilization of low- and high-rank coals originate in the differences in molecular structure. It is very well known that many properties of coal are a discontinuous function of rank³³ and we advanced the view that had previously been suggested by Wender³⁴ that the success of C-alkylation depended upon the degree of coalification, with higher rank coals more

susceptible to solubilization as a consequence of their lessened structural complexity.¹⁰ In point of fact, Ouchi and coworkers pointed out that the average molecular weights observed for the high-rank coals seem to be small than the corresponding values for the low rank coals, and proposed that the high ranking coals had more highly condensed aromatic rings but a lower degree of polymerization.³⁵ More recently, Larsen and Wei have also concluded from their work that the amount of pyridine extract and average molecular weight increases with the rank up to a carbon content of 86%.³⁶ On the basis of the increase in the quantity of the extractable material with rank and change in molecular weight of the extracts, they also conclude that the coalification process is a net depolymerization.³⁶

Our results are compatible with this concept. The lower degree of solubilization of the low rank coals is compatible with the notion that the macromolecular network is larger and more complex. In this situation, a low degree of C-alkylation is inadequate for the achievement of solubilization because covalent bonding and molecular dimensions dictate the outcome rather than intermolecular interactions such as hydrogen bonding or polarization forces. Thus, for the low rank coals, conventional bond cleavage is required for depolymerization and solubilization. In contrast, the lower degree of polymerization of high rank coals reduces the challenge considerably. In some situations, it may only be necessary to disrupt the intermolecular polarization forces between the aromatic and hydroaromatic molecules to achieve solubilization. As has been mentioned earlier, destruction of such interactions leads to almost quantitative solubilization for the high rank coals.

SECTION II. CATALYTIC COAL CONVERSION STUDIES
PREPARED BY MICHAEL ETTINGER

INTRODUCTION

When an arene is coordinated to $\text{Cr}(\text{CO})_3$, profound changes in its reactivity occur^{37,38}. Ring and benzylic hydrogen atoms exhibit enhanced acidity, and typical arene reactions such as electrophilic aromatic substitutions are quenched. In contrast, the ring carbon atoms are activated toward attack by nucleophiles³⁷. These changes in reactivity are manifestations of the forceful electron withdrawing nature of the $\text{Cr}(\text{CO})_3$ component, and of the ability of this group to stabilize charged intermediates. Most importantly, arenechromium tricarbonyl complexes have been proven to be good hydrogenation catalysts for the hydrogenation of dienes³⁹. Most simple complexes such as toluenechromium tricarbonyl require the use of high temperatures and high dihydrogen pressures for effective hydrogenation reactivity, but arenechromium tricarbonyl complexes with naphthalene, anthracene, and phenanthrene as the arene show remarkable activity under extremely mild conditions. The synthesis and chemistry of coal arene groups to which $\text{Cr}(\text{CO})_3$ groups have been attached will provide interesting information as to whether arenechromium tricarbonyl complexes made with coal molecules are active hydrogenation catalysts for the hydrogenation of dienes analogously to those made with pure aromatic molecules, or for the hydrogenation of coal molecules. Many of the aspects and difficulties of synthesis of organometallic complexes using coal molecules are addressed including the complexation of solid Illinois No. 6 coal.

EXPERIMENTAL

Materials

Chromium hexacarbonyl was obtained from Aldrich, Strem, and Pressure Chemical and was used as received. Potassium hydroxide, naphthalene, and 2,4-hexadienoic acid, ethanol, 1-butanol were used as received. Tetrahydrofuran, dioxane, dibutyl ether, hexane, and acetone were freshly distilled from sodium benzophenone ketyl immediately before use. All solvents were deoxygenated via standard freeze-pump-thaw cycles. Illinois No. 6 coal (APCS #3) and Pocahontas coal (APCS #5) were obtained through the Argonne Premium Coal Sample Program

and were dried for 12 hours at 110°C under vacuum. The coal liquids were obtained through Amoco Research Center from the Wilsonville Pilot Plant, run 250D. Dideuterium (99.5%) was obtained from Linde and was used as received.

Procedures

Triaminechromium tricarbonyl. Chromium hexacarbonyl (17.6g, 0.08mol), KOH (32.0g), ethanol (80ml), n-butanol (80ml) and water (25ml) were placed in a flame dried 500mL three neck RB flask fitted with a stirrer and a reflux condenser. The contents of the flask were warmed gradually to 150°C. Chromium hexacarbonyl, which sublimed initially, was washed out later by the solvent and no more sublimation took place. The reaction solution gradually turned red. After 4h of refluxing, the mixture was cooled, 150mL of concentrated aqueous ammonia solution was added and the mixture was slowly stirred for two hours. A yellow precipitate was filtered through a glass filter, washed with aqueous ammonia, methanol, and ether.

Napthalenechromium tricarbonyl. $C_{10}H_8Cr(CO)_3$ was prepared by a slightly modified procedure⁴³ to that described by Pauson⁴⁰. A mixture of naphthalene (5.3g, 0.042mol) and $Cr(CO)_6$ (4.4g, 0.020mol) were refluxed in dibutyl ether (100 mL), hexane (10mL), and THF (1mL) for 70 hrs under nitrogen. The heating bath was maintained at 160°C for 15 hrs and at 145°C for the remaining time. Solvent and excess naphthalene were removed under vacuum and the product purified over a column of silica gel with a 1:1:4 $CHCl_3$: CH_2Cl_2 : Hexane.

Methyl-2,4-hexadienoate. 2,4-Hexadienoic acid was refluxed in acidic methanol for six hours. The methyl ester product was washed sequentially with water, aqueous sodium bicarbonate, water, and was then distilled under vacuum.

Coal liquid arenechromium tricarbonyl. $(NH_3)_3Cr(CO)_3$ (4.68g, 0.025mol), coal liquid (5g), and dioxane (25mL) were loaded into a 100 mL flask provided with a magnetic stirrer, and a reflux condenser. The solution was refluxed in an oil bath for 48 hrs. The reaction mixture was filtered

through Celite in a glass filter and the flask and Celite were washed with ether. The product was purified over a column of silica gel with a 1:1:4 CHCl_3 : CH_2Cl_2 : Hexane.

Micro-hydrogenation reactions. Hydrogenation reactions were carried out in glass vials using a 10mL micro hydrogenation apparatus. Typically, the reactants and dried, deoxygenated acetone were loaded into the glass vial and it was evacuated and flushed with nitrogen three times, dideuterium three times and then brought up to the final pressure at the desired temperature. At the end of the reaction time, the mixture was filtered through Celite in a glass filter and concentrated under vacuum.

Hydrogenation reactions. Hydrogenation reactions employing more severe conditions using heptane as a solvent were carried out in a Parr 71 mL high pressure and temperature reactor fitted with a thick walled glass liner which reduced the volume to 30 mL. Typically, the reactor was flushed with nitrogen and then pressurized with dideuterium. The reactor was then placed in a fluidized sandbath and oscillated at 60 rpm.

RESULTS AND DISCUSSION

In the first stage of the investigation, experiments were carried out using a coal liquid distilled at approximately 850°C obtained from run 250D at the Wilsonville pilot plant using Illinois No. 6 coal. In contrast to solid coals and heavy distillates, this material is soluble in all common organic solvents. Because the ^{13}C chemical shifts of aromatic carbon atoms complexed to a $\text{Cr}(\text{CO})_3$ group move upfield from 100-129ppm into the range of 80-100ppm, any complexation of the coal arene groups will be evident using this coal liquid because the baseline is very clean in the 80-100ppm region. The ^{13}C NMR spectrum of the unreacted coal liquid is shown in Figure 8.

The first stage of research centered on the conversion of the aromatic hydrocarbons in a coal liquid into arenechromium tricarbonyl compounds. Synthesis of arenechromium tricarbonyl complexes are generally achieved through one of the methods shown in Figure 9. Reactions to

FIGURE 8.
 ^{13}C NMR SPECTRA OF WILSONVILLE COAL LIQUID (TOP)
AND COAL LIQUID- $\text{Cr}(\text{CO})_3$ (BOTTOM).

FIGURE 9.
SYNTHETIC METHODS FOR ARENECHROMIUM TRICARBONYL COMPLEXES.

form complexes with the coal liquid arenes using the method of Mahaffy and Pauson⁴⁰ in which the arene is refluxed in dioxane with chromium hexacarbonyl required long reaction times and large excesses of chromium hexacarbonyl to achieve acceptable levels of complexation. To alleviate these problems for use with the coal liquid, the use of an alternative reagent, triaminechromium tricarbonyl, was investigated. This reagent is easy to synthesize⁴¹ and has been used to obtain a number of organochromium complexes⁴². Reactions of the coal liquid with triaminechromium tricarbonyl proved to be very successful. High yields of complexation with the coal liquid arene molecules were obtained to the point where the complexed coal liquid product was orange colored and crystalline at room temperature. The ¹³C NMR spectrum of one such product is shown in Figure 8. The carbonyl resonance can be seen at 234 ppm along with the shift in the aromatic carbons to 80-100 ppm. The infrared spectra of the unreacted and complexed coal liquid are shown in Figure 10. The characteristic Cr(CO)₃ carbonyl stretch is displayed and will be very useful in diagnosing complexation in solid coals. Note that the carbonyl stretch of the complexed Cr(CO)₃ unit is distinctly different than the carbonyl stretch associated with carbonyl groups of chromium hexacarbonyl, Figure 11. In addition, as shown in Figure 9, chromium hexacarbonyl in a dibutyl ether: hexane: tetrahydrofuran solvent mixture can be used for the synthesis of polycyclic arenechromium tricarbonyl complexes⁴³.

The second stage of the investigation of the chemistry of the coal liquid arenechromium tricarbonyl complexes will study the hydrogenation capacity of the complexes. Studies have been carried out using simple arenechromium tricarbonyl complexes^{39,43} as catalysts in the hydrogenation of methyl 2,4-hexadienoate in which the severity of reaction conditions varied from 150°C at 700 psi dihydrogen pressure for toluenechromium tricarbonyl to 40°C at 60 psi dihydrogen pressure for naphthalene and phenanthrene chromium tricarbonyl, Figure 12. The proposed explanation for the dramatic increase in activity using polycyclic arenechromium tricarbonyl complexes is that there exist two Cr-C bonds with the bridgehead carbon atoms out of the six Cr-C bonds that are longer than the others with bond orders of less than one⁴⁵, Figure 13. The longer bonds should therefore be more easily displaced by the incoming diene ligand in the

FIGURE 10.
INFRARED SPECTRA OF WILSONVILLE COAL LIQUID (TOP)
AND COAL LIQUID-Cr(CO)₃ (BOTTOM).

FIGURE 11.
INFRARED SPECTRA OF BENZENE- $\text{Cr}(\text{CO})_3$ (TOP) AND $\text{Cr}(\text{CO})_6$ (BOTTOM).

FIGURE 12.
ARENECHROMIUM TRICARBONYL CATALYZED HYDROGENATION REACTIONS.

FIGURE 13.
Cr-C BOND LENGTHS IN MONO- AND POLYCYCLIC ARENECHROMIUM
TRICARBONYL COMPLEXES.

Mean Cr-C Bond Distances for Selected Pairs

	<u>1.4</u>	<u>2.3</u>	<u>5.6</u>	<u>(5.6)-(1.4)</u>
Benzene	2.223	2.217	2.224	
Napthalene	2.200	2.202	2.321	+0.120
Phenanthrene	2.210	2.208	2.289	+0.080

rate determining step. This advantage is not realized in the case of the monocyclic arenechromium tricarbonyl complexes such as benzenechromium tricarbonyl where all of the Cr-C bonds are equal length with bond orders close to unity. The effect of solvent was also examined. Coordinating solvents such as tetrahydrofuran or acetone increased the rate of the hydrogenation reaction presumably by facilitating the initial ligand substitution step³⁹.

Experiments were carried using the coal liquid arenechromium tricarbonyl complexes under hydrogenation conditions hoping to demonstrate that the arene molecules in coal or coal liquids can be complexed with chromium tricarbonyl groups to yield catalytically active molecules. In addition, the complexation of the polycyclic aromatics known to exist in coal might yield the more active, although more sensitive, polycyclic arenechromium tricarbonyl complexes. Low severity hydrogenation reactions were carried out in a glass micro hydrogenation apparatus at pressures from 80 psi to 200 psi and temperatures less than 50°C using dideuterium gas. The use of dideuterium gas allows the addition of any deuterium to the reactants to be detected by deuterium NMR. The first reactions were carried out with methyl 2,4-hexadienoate with dideuterium catalyzed by naphthalenechromium tricarbonyl in acetone as described in the literature in order to insure that the procedure and apparatus were satisfactory. The ¹H NMR spectrum of the naphthalenechromium tricarbonyl complex is shown in Figure 14. The reaction yielded the desired product of methyl hex-3-enoate-2,5-d₂. A similar reaction was carried out in the presence of small amount of coal liquid to test whether or not the coal liquid would poison the catalyst. The heteroatoms and metal atoms found in fossil fuels have been found to be notorious in deactivating catalysts. This reaction also proved to be successful yielding a 63% conversion of methyl 2,4-hexadienoate to methyl hex-3-enoate-2,5-d₂. The ¹H and ²H NMR spectra of the products are shown in Figure 15. It is evident and promising that the presence of the coal liquid does nothing significant to inhibit the catalyst.

The next set of experiments were carried out using the coal liquid arenechromium tricarbonyl complexed material as the prospective catalyst. Attempts were made to directly hydrogenate the coal liquid arenechromium tricarbonyl complexed material in hope that the arenechromium

FIGURE 14.
 ^1H NMR SPECTRA OF NAPHTHALENE (TOP) AND NAPHTHALENE- $\text{Cr}(\text{CO})_3$ (BOTTOM).

FIGURE 15.
 ^1H (TOP) AND ^2H (BOTTOM) NMR SPECTRA OF NAPHTHALENE- $\text{Cr}(\text{CO})_3$ CATALYZED
 DEUTERATION OF METHYL 2,4-HEXADIENOATE REACTION PRODUCT.

tricarbonyl complexes would catalyze the hydrogenation of other components in the coal liquid. However, this proved to be unsuccessful. In part, this could be due to the fact that all of the possible active sites for hydrogenation available in the coal liquid may of been rendered unreactive by complexation with $\text{Cr}(\text{CO})_3$. Therefore, experiments were carried out with unreacted coal liquid as the substrate along with the coal liquid arenechromium tricarbonyl complex as the catalyst. A comparison can be made of the ^2H NMR spectra of the unreacted coal liquid and the coal liquid arenechromium tricarbonyl reaction product which, unfortunately, demonstrates that no significant deuteration had taken place, Figure 16. Two possible reasons for this are that the coal liquid arenechromium tricarbonyl complex is not an active catalyst at the employed conditions, or that there are no suitable sites for hydrogenation in the unreacted coal liquid.

In order to test the latter hypothesis of whether hydrogenation substrates exist in the coal liquid, an experiment was carried out using naphthalenechromium tricarbonyl, a very active arenechromium tricarbonyl catalyst under mild conditions, and unreacted coal liquid as the substrate. A comparison of the ^2H NMR spectra of the unreacted coal liquid before and after the reaction, Figure 17, shows that a small level of dideuterium addition has taken place. This is concluded from the observation that the relative aromatic deuterium content has decreased indicating that aliphatic deuterium content has increased, and, that the two major resonances of the unreacted coal liquid at approximately 1.54 ppm and 1.16 ppm are no longer the dominant resonances. The major resonances in the reacted coal liquid have shifted to approximately 2.27 ppm and 1.45 ppm. In addition, new peaks have appeared at 3.57 ppm and 2.71 ppm. Thus, at a preliminary level, it can be concluded that there are some hydrogenation sites available in the coal liquid.

In order to test the catalytic activity under mild conditions of the coal liquid arenechromium tricarbonyl complex, experiments were carried out using methyl 2,4-hexadienoate, a known reactive substrate, and the coal liquid arenechromium tricarbonyl complex as the catalyst. The desired product was not obtained in any significant yield.

Although the activity of the coal liquid arenechromium tricarbonyl complex as a hydrogenation catalyst appears to be small for methyl 2,4-hexadienoate and unreacted coal liquids under the mild

FIGURE 10.

^2H NMR SPECTRA OF UNREACTED COAL LIQUID (TOP) AND
COAL LIQUID- $\text{Cr}(\text{CO})_3$ DEUTERATION REACTION PRODUCT (BOTTOM).

FIGURE 17.
 ^2H NMR SPECTRA OF UNREACTED COAL LIQUID (TOP) AND NAPHTHALENE- $\text{Cr}(\text{CO})_3$
CATALYZED DEUTERATION OF COAL LIQUID REACTION PRODUCT (BOTTOM).

conditions employed, it may well prove to be viable under moderately more severe conditions. A summary of the results of the above reactions is shown in Figure 18. One possible explanation postulated for the lack of activity of the coal liquid arenechromium tricarbonyl complexes under the mild conditions is that only monocyclic aromatic complexes were being formed with the coal liquid. Monocyclic arenechromium tricarbonyl complexes require higher temperatures for the activation step to occur which is the thermal liberation of the arene from a hexahapto to a tetrahapto bonded ligand. In addition, the solvent used in the reactions may have inhibited the deuteration of certain substrates in the coal liquids. The use of a coordinating solvent such as acetone assists in the partial liberation of the arene ligand by weakly occupying the empty coordination site, therefore, increasing catalytic activity for the hydrogenation of molecules, such as 1,3-dienes, that can effectively compete with acetone for the coordination sites⁴⁶. It has been shown that the use of a coordinating solvent such as acetone actually inhibits the arenechromium tricarbonyl catalyzed hydrogenation of 1,4-dienes by not allowing the 1,4-diene, a weaker binding ligand, to enter the coordination sphere⁴⁶. The use of a noncoordinating solvent like heptane alleviates such problems. Thus, in the case of using substrates that might be weakly coordinating such as the unreacted coal liquid, the use of a noncoordinating solvent is warranted.

Therefore, a new series of reactions was carried out to address these issues. The severity of the reaction conditions was increased and heptane was employed as a noncoordinating solvent. The glass micro-hydrogenation reactor system used in the previous experiments was not suitable for the increased pressures and temperatures desired. A new system was employed using a modified Parr 71mL high pressure and high temperature reactor. Glass liners allowed the volume to be reduced to 30mL. The reactor was heated and agitated in a fluidized sand bath.

Reactions employing the more severe conditions were carried out with the coal liquid arenechromium tricarbonyl complexes as catalysts for the deuteration of both methyl-2,4-hexadienoate and unreacted coal liquid. The coal liquid arenechromium tricarbonyl complex successfully catalyzed the deuteration of methyl-2,4-hexadienoate to methyl hex-3-enoate-2,5-d₂ at 200°C under 330 psi of dideuterium. The reaction was run for two hours and an approximate

FIGURE 18.
DEUTERATION REACTIONS AND RESULTS UNDER MILD CONDITIONS.

yield of 80-85% was obtained. The deuterium NMR spectrum of the product, methyl hex-3-enoate-2,5-d₂, is shown in Figure 19. In addition, the ¹H NMR spectrum of the product mixture is shown in which integration comparisons of reactant and product peaks allows the yield determination. One can conclude that since the deuteration reaction of methyl-2,4-hexadienoate to methyl hex-3-enoate-2,5-d₂ occurred only under more severe conditions that predominantly monocyclic arenechromium tricarbonyl complexes were formed in the Wilsonville coal liquid. This is not necessarily surprising even though polycyclic aromatics are present because they are harder to complex and in the presence of monocyclic aromatics, the monocyclic arenechromium tricarbonyl product will certainly be favored.

The coal liquid arenechromium tricarbonyl complex failed to catalyze any significant level of dideuterium uptake in the reactions employing the unreacted coal liquid as the deuteration substrate. Apparently, this particular coal liquid does not contain any molecules that are reducible using arenechromium tricarbonyl complexes as the catalyst in either coordinating or noncoordinating solvents employing either mild or more severe conditions. A summary of the results obtained under the more severe conditions is shown in Figure 20.

The active species of the arene-Cr(CO)₃ complex that catalyzes hydrogenation is postulated to be the "Cr(CO)₃" loosely coordinated to the arene ligand. To further investigate this aspect of the chemistry, a variation upon the use of arene-Cr(CO)₃ complexes as hydrogenation catalysts was approached in which chromium hexacarbonyl, Cr(CO)₆, was used "as is" in reactions with coal liquid and with methyl 2,4-hexadienoate as shown in Figure 20. Perhaps, Cr(CO)₆ under severe hydrogenation reaction conditions can generate the "Cr(CO)₃" complex in situ. This process could eliminate the time consuming process of synthesizing the coal arene-Cr(CO)₃ complex. Chromium hexacarbonyl and methyl 2,4-hexadienoate were reacted in heptane at 200°C under 500 psi D₂. Therefore, if any active form of the catalyst was generated, the diene would be deuterated to methyl hex-3-enoate-2,5-d₂. The desired product was formed in good yield. Figure 21 shows the deuterium NMR spectrum displaying the characteristic resonances of the deuterated monoene. Further work is necessary to determine whether or not the "tricarbonyl complex" was formed in

¹H (TOP) AND ²H (BOTTOM) NMR SPECTRA OF COAL LIQUID-Cr(CO)₃ CATALYZED DEUTERIATION OF METHYL 2,4-HEXADIENOATE REACTION PRODUCT.

FIGURE 20.
DEUTERATION REACTIONS AND RESULTS UNDER MODERATE CONDITIONS.

**¹H NMR SPECTRA OF Cr(CO)₆ CATALYZED DEUTERATION OF
METHYL 2,4-HEXADIENOATE REACTION PRODUCT.**

FIGURE 22.
ATTEMPTED COMPLEXATION REACTIONS OF POCAHONTAS COAL WITH Cr(CO)₆.

FIGURE 23.
INFRARED SPECTRA OF POCAHONTAS COAL BEFORE (TOP)
AND AFTER REACTION WITH $\text{Cr}(\text{CO})_6$ (BOTTOM).

situ yielding the active species, or if another pathway was responsible. The reaction with $\text{Cr}(\text{CO})_6$ and the coal liquid failed to produce any deuterated products.

The synthesis of the coal liquid arenechromium tricarbonyl complex using the Wilsonville run 250D coal liquid and triaminechromium tricarbonyl was very efficient. However, the application of this chemistry to less soluble coal liquids and solid coals requires the use of chromium hexacarbonyl due to experimental considerations. The use of triaminechromium tricarbonyl requires filtration of the soluble reaction product to remove the unreacted and insoluble triaminechromium tricarbonyl. This scheme is certainly unsatisfactory for solid coal products. The use of chromium hexacarbonyl requires longer reaction times and excesses of the reagent. However, the reaction is quite suitable for solid coals because, although chromium hexacarbonyl is also a solid that is insoluble in common organic solvents, it can easily be sublimed away as the solvent is removed. It is also a simple reaction in that coal and chromium hexacarbonyl are the only reactants required.

Initial attempts to complex solid, highly aromatic Pocahontas coal (APCS #5) under rigorous conditions, Figure 22, resulted in no complexation which was determined by the infrared spectrum, Figure 23. Although insolubility of the reactants could certainly be the problem, it was thought that perhaps there were substances in solid coal not found in refined coal liquids that might inhibit the reaction. In that respect, O-methylated Illinois No. 6 coal was extracted with dioxane for twenty four hours in order to provide a very coal-like material. This extract was refluxed in dioxane with chromium hexacarbonyl for 5 days after which the solvent and excess chromium hexacarbonyl were removed under vacuum. The infrared spectrum of the product, along with that of O-methylated Illinois No. 6 coal for comparison, Figure 24, shows the characteristic carbonyl frequency of arene complexed $\text{Cr}(\text{CO})_3$ groups indicating successful complexation with aromatic molecules in the coal extract. The infrared spectrum of chromium hexacarbonyl, Figure 11, can be referred to for comparison. Therefore, the reaction with solid Illinois No. 6 coal was attempted. This reaction, carried out for 7 days, proved to be successful as indicated by the $\text{Cr}(\text{CO})_3$ carbonyl stretching frequency of the product shown in Figure 25. The degree of complexation is

FIGURE 24.

INFRARED SPECTRA OF O-METHYL ILLINOIS NO. 6 COAL (TOP) AND [DIOXANE
EXTRACT OF O-METHYL ILLINOIS NO. 6 COAL]- $\text{Cr}(\text{CO})_3$ (BOTTOM).

FIGURE 25.
INFRARED SPECTRUM OF ILLINOIS NO. 6 COAL-Cr(CO)₃.

undetermined, but it is promising that the reaction can be carried out with the associated solubility problems. No attempts to hydrogenate this material were made.

CONCLUSIONS

Organometallic complexes have been successfully synthesized with coal molecules to form arenechromium tricarbonyl complexes. Results showed that, despite the difficulties of carrying out this type of sensitive chemistry on coal materials, high yields of coal-arenechromium tricarbonyl complexes were achieved. This has been achieved using coal liquids, solid coal extracts, and solid coal. Arenechromium tricarbonyl complexes have exhibited diverse chemical properties including the ability to homogeneously catalyze the hydrogenation of dienes under very mild conditions. Research was directed at trying to use such complexes to hydrogenate molecules in coal products. These coal complexes exhibited catalytic activity towards the hydrogenation of dienes under moderate conditions. Unfortunately, neither pure aromatic chromium tricarbonyl complexes nor coal-arenechromium tricarbonyl complexes are apparently able to catalyze the hydrogenation of the coal materials under the mild conditions used in this study. However, it is encouraging that the coal-arenechromium tricarbonyl complexes in fact performed the same hydrogenation chemistry as the pure complexes found in the literature.

While supported and dispersed heterogeneous catalysts are predominately used in the hydrogenation and upgrading of coal products, many problems associated with the use of solid catalysts in such systems certainly exist such as catalyst poisoning as well as deactivation due to "coke" deposits. The results of this study give support to the idea that homogeneous catalytic systems may be feasible for use in the refining of coal products. This study has shown that such chemistry is possible with coal molecules. The use of these monomolecular reagents can increase the effective concentration of the catalyst by many orders of magnitude when compared to supported and dispersed solid catalysts. The many advantages of homogeneous systems warrant further investigation of their use in coal systems, and may provide an economical and chemically efficient method of refining fossil fuel materials.

REFERENCES

1. Stock, L. M., *Coal Sci.*, 1982, **1**, 161-282.
2. Davison, R. M. *Coal Sci.*, 1982, **1**, 84-160.
3. Sternberg, H. W. and Delle Donne, C.L. *Prepr. Pap.- Am. Chem. Soc., Div. Fuel Chem.*, 1968, **12(4)**, 13.; Sternberg, H. W.; Delle Donne, C.L. and Pantages, P., *Prepr. Pap.- Am. Chem. Soc., Div. Fuel Chem.*, 1970, **14(1)**, 87.
4. Mallya N. and Stock L. M., *Fuel*, 1986, **65**, 736.
5. Ignasiak, B.; Carson, D.; Gawlak, M., *Fuel*, 1979, **58**, 833.
6. Gawlak, M.; Carson, B.; Ignasiak, T.; Strausz, O.P. *Preprints of Papers, International Conference on Coal Science*, Maastricht, The Netherlands, 1987; pp 57.
7. Lazarov, L.; Marinov, S.P.; Stefanova, M.; Angelova, G. *Preprints of Papers, International Conference on Coal Science*, Maastricht, The Netherlands, 1987; pp 745.
8. Lazarov, L.; Marinov, S.P. *Fuel*, 1987, **66**, 185.
9. Chambers, R. R., Jr.; Hagaman, E. W.; Woody, M. C. Polynuclear Aromatic Hydrocarbons; Ebert, L., Ed., *Advances in Chemistry*, 1987, **217**; Chapter-15, pp 255.
10. Miyake, M. and Stock, L. M., *Energy and Fuels*, 1988, **2**, 815.
11. Liotta, R.; Rose, K. and Hippo, E., *J. Org. Chem.*, 1981, **46**, 277.
12. Ettinger, M.; Nardin, R.; Mahasay, S. R.; Stock, L. M., *J. Org. Chem.*, 1986, **51**, 2840.
13. Bates, R. B.; Siahaan, T. J.; Suvannachut, K.; Vasey S. K.; Yager K. M., *J. Org. Chem.*, 1987, **52**, 4605.
14. Chatterjee, K. and Miyake, M., unpublished results.
15. Chatterjee, K.; Miyake, M.; Stock, L. M. *Energy and Fuels*, 1990, **4**, 242.
16. Chambers, R. R., Jr.; Hagaman, E. W.; Woody, M. C. *Preprints of Papers, International Conference on Coal Science*, Maastricht, The Netherlands, 1987; pp 741.
17. Schlosser, M. and Strunk, S., *Tetrahedron Letters*, 1984, **25**, 741.
18. Lochmann, L.; Pospisil, J. and Lim, D., *Tetrahedron Letters*, 1966, 257;
Lochmann, L. and Lim, D., *J. Organomet. Chem.*, 1971, **28**, 153.
19. Bates, R. B., personal communication.

20. Meaercker, A., *Angew. Chem. Int. Ed. Engl.*, 1987, **26**, 972.
21. Botto, R. E.; Choi, C. Y.; Muntean, J. V.; Stock, L. M. *Energy and Fuels*, 1987, **1**, 270.
22. Stock, L. M.; Willis, R. S. *J. Org. Chem.*, 1985, **50**, 3566.
23. Our sincere thanks to Mr. Art Kostka of Argonne National Laboratory, who helped us to obtain these spectra.
24. Cram, D. J. in *Fundamentals of Carbanion Chemistry*; Academic Press, New York, 1965, pp. 8-20, 41.
25. Streitwieser, A., Jr.; Juaristi, E.; Neberzahl, L. L. in *Comprehensive Carbon Chemistry*; Part A: Buncl, E. & Drust, T., Eds. Elsevier, New York, 1980, pp. 323-381.
26. Siskin, M. and Aczel, T., *Fuel*, 1983, **62**, 1321.
27. Wachowska, H. *Fuel*, 1979, **58**, 99.
28. Gilman, H.; Young, R. V., *J. Am. Chem. Soc.*, 1934, **56**, 1415; *ibid*, 1935, **57**, 1121.
29. Alemany, L. B. and Stock, L. M., *Fuel*, 1982, **61**, 1088.
30. Eliel, E. L.; Allinger, N. L.; Angyal, S. J.; Morrison, G. A., *Conformational Analysis*, Interscience Publishers, New York, 1965.
31. Streitwieser, A., Jr., personal communication; Cram, D. J., personal communication; Grovenstein, E. Jr., personal communication.
32. Kuznetsov, P.N.; Sukhova, G.I.; Bimer, J.; Salbut, P.D.; Korniyets, E.D.; Belskaya, N.A.; Ivanchenko, N. M., *Fuel*, 1991, **70**, 1031.
33. *Chemistry of Coal Utilization*, H. H. Lowry Ed., Supplementary Volume, John Wiley and Sons, New York, 1963.
34. Wender, I.; Heredy, L. A.; Neuworth, M. B. and Dryden, I. G. C. in *Chemistry of Coal Utilization*, Second Supplementary Volume (Ed. M. A. Elliot), John Wiley and Sons, New York; 1981, Chapter 4 424.
35. Ouchi, K.; Hirano, Y.; Makabe, M. and Itoh, H., *Fuel*, 1980, **59**, 751.
36. Larsen, J. W. and Wei, Y.-C., *Energy and Fuels*, 1988, **2**, 344.
37. Semmelheck, M. F. *J. Organomet. Chem.*, 1976, **1**, 371.
38. Jaouen, G. in *Transition Metal Organometallics in Organic Synthesis*, Vol. 2, H. Alper (ed.), Academic Press 1978.
39. Cais, M., Fraenkel, D., Weidenbaum, K. *Coord. Chem. Rev.*, 1975, **16**, 27.
40. Mahaffy, C. A., Pauson, P. L. *Inorg. Synth.* 1979, **19**, 154.

41. Razuvaev, G. A., Artemov, A. N., Aladjin, A. A., Sirotkin, N. I. *J. Organometal. Chem.*, 1976, **111**, 131.
42. Moser, G. A., Rausch, M. D. *Synth. React. Inorg. Met.-Org. Chem.*, 1955, **280**, 255.
43. Kundig, E. P., Desobry, V., Simmons, D. P., Wenger, E. *J. Am. Chem. Soc.*, 1989, **111**, 1304.
44. Cais, M., Rejoan, A. *Inorg. Chim. Acta*, 1970, **4**, 509.
45. Muir, K. W., Ferguson, G., Sim, G. A., *Inorg. Chem.*, 1966, **5**, 1851.
46. Tucker, J.R., Riley, D.P. *J. Organomet. Chem.*, 1985, **279**, 49.

APPENDIX I

LIST OF PUBLICATIONS

K. Chatterjee and L. M. Stock, Coal Solubilization Through C-Alkylation, Preprints of Papers, Am. Chem. Soc. Fuel Chem. Div., 35(1), 46-50 (1990).

K. Chatterjee, M. Miyake and L. M. Stock, Coal Solubilization. Promotion of C-Alkylation by n-Butyllithium and Potassium t-Butoxide, Energy and Fuels, 4, 242-248 (1990).

M. Ettinger and L. M. Stock, Arenechromium Complexes in Coal. Preprints of Papers. Am. Chem. Soc. Fuel Chem. Div., 36(2), 838-844 (1991).

K. Chatterjee and L. M. Stock, Toward Organic Desulfurization. The Treatment of Bituminous Coals with Strong Bases. Energy and Fuels, 5, 704-707 (1991).

K. Chatterjee, S. Jiralerspong and L. M. Stock, Coal Depolymerization Using Metal Alkoxides in Refluxing Alcohols, Preprints of Papers, Am. Chem. Div., (00(0)), 000-000. Accepted for publication.

END

**DATE
FILMED**

8 / 25 / 92