NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

ANNUAL TECHNICAL REPORT FOR DOE GRANT NUMBER DE-FG03-88ER45378
ENTITLED "APPLICATIONS OF MESOSCOPIC PHYSICS"

Shechao Feng

September 1991 - June 1992

DOE/ER/45378--T1

DE92 015610

In this annual technical report, I shall summarize the various research activities that I and my group have pursued this past year here at UCLA, in the area of the "applications of mesoscopic physics to novel correlations and fluctuations of speckle patterns: imaging and tomography with multiply scattered classical waves", and related topics, during the grant period from September 1991 to June 1992.

Last year we had the great fortune of hosting the visit of Prof. Boris Spivak from Leningrad, who then became a full professor at the University of Washington. Some of the work in the report reflects our joint work here at UCLA.

I will list my research results in order of the various publications which have recently appeared.

1. "Acoustical nondestructive evaluation of heterogeneous materials in the multiple scattering regime": This was a joint work between myself and Dr. Didier Sornette from the CNRS at the University of Nice in France. We were interested to apply the theoretical techniques in the field of mesoscopic fluctuations to the problem of how to reconstruct the position of a defect in a metal or concrete like material, using appropriate averaging and correlations of acoustic speckle patterns, in the multiple scattering limit. We were able to show that indeed such image reconstruction is possible, if one can have sufficient frequency averages of the outgoing speckle pattern, and the resulting image of a new crack is of the

MASTER
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Ed

6/10/92

form of a diffusion pole, which provides information of both the transverse position as well as its depth from the measuring surface. We feel this work has potential large impact in the field of acoustical nondestructive evaluations.

2. "Classical and quantum superdiffusion in a time-dependent random potential": This was a joint work between myself and my former post-doc Leo Golubovic, who left recently to become a post-doc at Cal Tech. Due to the funding decreases from DOE and other sources, I can no longer support a post-doc at present or in the near future. In this work, we set out to consider the problem of wandering of a non-relativistic particle in a time-dependent random potential in d spatial dimensions. The motivation of this problem was that this is a limiting case of the wave scattering problem if the scattering is sufficiently anisotropic, such that one of the spatial dimensions can be regarded as a "time" dimension; then the scattering problem from random potentials maps onto the problem of a quantum mechanical particle wandering through a time-dependent random potential. Its root-mean-square displacement from the initial position increases superdiffusively with time t as $t^{9/8}$ for $d > 1$, and as $t^{6/5}$ in $d = 1$. Its kinetic energy increases as $t^{1/2}$ for $d > 1$, and as $t^{2/5}$ in $d = 1$. These scaling behaviors hold for both the classical and the corresponding quantum-mechanical problem in *continuous* space-time and differ from those of lattice models, where the particle wandering is described by simple diffusion. These new scaling laws are strikingly new, and have the potential of motivating many experimental studies to verify such behaviors. This paper is the most important one that I have written this past year.

3. "Negative Magnetoresistance in Variable Range Hopping Conduction": This is a joint work between myself, my students, and Prof. Spivak. In this work we considered the

orbital mechanism of negative magnetoresistance of variable range hopping conductivity by careful numerical simulations, and qualitative arguments. We find that at small magnetic fields the positive magnetic field corrections to the localization length in two-dimensional samples are universal in the sense that they are independent of the details of the scattering potential (impurity concentration, amplitude of scattering and proximity to the metal-insulator transition). Our result can be summarized by the simple scaling law $L = \gamma \frac{r}{L_H}$, where $L = \log(G(H)/G(0))$ is the magetoconductance in a hopping system, r is essentially the Mott hopping length, which we assume to be long in the low temperature regime, and L_H is the magnetic length. This result holds in the regime $r \gg L_H$ only. The coefficient γ is found to be universal. Implications of these results to transport properties on the dielectric side of the superconductor-insulator transition were also considered in this work.

4. "Reproducible Conductance Fluctuations in Macroscopic Anderson Insulators":

This is a joint work between myself and Dr. Jean-Louis Pichard from Saclay in France. Here we studied theoretically the large magneto-conductance fluctuations in macroscopic Anderson insulators, i.e. variable range (Mott) hopping systems that we spoke about in the last section, which have recently been observed. Based on the hypothesis of large ergodic quantum fluctuations of the localization length when a flux quantum is applied through a localization domain, we showed that the normalized magneto-conductance fluctuations $\delta G_{tot}/G_{tot}$ is $\sqrt{\frac{\xi_p^2}{\Omega_{eff}}}$ for ergodic field changes $\Delta B \sim \phi_0/\xi^2$, where ξ_p is a percolation correlation length which approaches macroscopic length scales at very low temperatures, and Ω_{eff} is an effective volume for the classical self-averaging of the fluctuations. This work was also published as a Physical Review Letter.

5. "Feasibility of far-infrared lasers using multiple semiconductor quantum wells":

This is a joint work between myself and Prof. Qing Hu from MIT. In this work we studied the rather applied but technologically very important problem of whether it is feasible to build a new far-infrared (THz) laser device using semiconductor multiple quantum wells. If our ideas can be realized in the laboratory, this could mean a new generation of small size semiconductor lasers whose wavelengths, in the far-infrared range, can be adjusted at will by the design of the quantum wells. The technological advance due to this device could imply high efficiency space communications in the Tera-Hertz range. Our main ideas are based on selective injection of electrons into an upper subband, and selective removal from a lower subband of a quantum well, using quantum-well energy filters. Photon confinement is achieved between doped injector and collector semiconductor contacts which also serve as good photon reflectors. The lasing threshold current density is estimated to be in the range of $J_{th} \sim 130 A/cm^2$ at 5 THz. Thus the feasibility of a far-infrared laser based on this design seems quite promising.

The above summarizes the various research activities that I have pursued this past year. I hope they are satisfactory to the requirement of the guidelines of the DOE granting agency. On the next page, I shall list all the papers that I have published from the research projects that I have performed during this granting period.

PUBLICATIONS FROM RESEARCH ACTIVITIES BETWEEN 9/91 TO 6/92

1. "Acoustical nondestructive evaluation of heterogeneous materials in the multiple scattering regime", Shechao Feng and D. Sornette, J. Acoust. Soc. Am. 90, 1742 (1991).
2. "Classical and quantum superdiffusion in a time-dependent random potential", L. Golubovic and Shechao Feng, Phys. Rev. Lett. 67, 2115 (1991).
3. "Negative Magnetoresistance in Variable Range Hopping Conduction", H-L Zhao, B. Z. Spivak, M. P. Gelfand, and Shechao Feng, Phys. Rev. B 44, 10760 (1991).
4. "Reproducible Conductance Fluctuations in Macroscopic Anderson Insulators", Shechao Feng and J-L Pichard, Phys. Rev. Lett. 67, 753 (1991).
5. "Feasibility of far-infrared lasers using multiple semiconductor quantum wells", Q. Hu and Shechao Feng, Appl. Phys. Lett. 59, 2923 (1991).

**DATE
FILMED**

8 / 17 / 92