MOLECULAR BIOLOGICAL ENHANCEMENT OF COAL DESULFURIZATION
Contract # DE-AC22-89PC89903

Cloning and Expression of the Sulfoxide/Sulfone/Sulfonate/Sulfate
Genes in *Pseudomonads* and *Thiobacillae*

Eleventh Quarterly Report

Due: April 20, 1992

Submitted: April 28, 1992

DOE/PC/89903--T8

DE92 015666

Steven Krawiec

Department of Molecular Biology
Center for Molecular Bioscience and Biotechnology
Lehigh University

-
- I. INTRODUCTION
 - A. N1-36
 - B. Pulsed-field gel electrophoresis (PFGE) of chromosomal DNAs from *Thiobacillae* spp.
 - C. Fresh isolates with the DbtS⁺ phenotype
 - II. DESULFURIZATION WITH N1-36
 - A. Production of OH-BP and o,o'-biphenol by N1-36
 - B. Regulation of the DbtS⁺ phenotype: oxidation of DBT and desulfurization of OH-BP
 - C. Nutritional characteristics
 - 1. Ethanol vs. glucose as carbon source
 - 2. Monohydroxybiphenyl (OH-BP) as carbon source
 - D. Growth of N1-36 in continuous culture
 - E. N1-36: aka *Rhodococcus erythropolis*
 - 1. Cell and colony morphology
 - 2. Biolog analysis
 - 3. Fatty-acid analysis
 - 4. General conclusion regarding identity
 - F. Acknowledgement
 - III. PULSED-FIELD GEL ELECTROPHORESIS OF CHROMOSOMAL DNAs OF *THIOBACILLUS* spp.
 - A. Qualitative results
 - B. Reliable estimation of sizes of standards
 - C. Some technical advances
 - D. Estimation of generation times
 - E. Acknowledgement
 - IV. FRESH ISOLATES WITH THE PRESUMPTIVE CAPACITY TO DESULFURIZE DBT
 - A. Isolation, abundance, and screening
 - B. Presumptive identities
 - C. Desulfurization
 - D. Acknowledgement
 - V. CONCLUSIONS
 - VI. MILESTONE/SCHEDULE STATUS REPORT
 - VII. COST MANAGEMENT REPORT

MASTER

U.S./DOE Patent Clearance is not required prior to the publication of this document.

I. INTRODUCTION

A. N1-36

N1-36, now presumptively identified as *Rhodococcus erythropolis*, has been grown recurrently in fed batch cultures for durations as long as 210 hours. The turbidity, number of colony-forming-units, pH, and concentrations of dissolved O_2 , glucose, dibenzothiophene (DBT), dibenzosulfone (DBTO₂), and monohydroxybiphenyl (OH-BP) were measured. In the present quarter, the abundance of products produced under conditions judged to be especially favorable were quantified. As expected, OH-BP was produced; what was unexpected was the formation of a product (seen with HPLC) which co-migrates with o,o'-biphenol.

The effect of $SO_4^{=}$ on the oxidation of DBT and the desulfurization of DBTO₂ was assessed. The data indicate that the former process is unaffected while the latter process does not occur in the presence of $SO_4^{=}$.

Also during this quarter, growth of N1-36 in continuous culture was initiated. The observations are presently limited in their extent but serve as useful qualitative guides to further experimentation.

B. Pulsed-field gel electrophoresis (PFGE) of chromosomal DNAs from *Thiobacillae* spp.

Physical characterization of the chromosomes of *T. acidophilus* 27807 and *T. novellus* 8093 using PFGE continued. The chromosome of the former is reliably digested with *Xba*I and *Spe*I while that of the latter is cut with *Spe*I and possibly *Dpn*I. Several technical improvements were introduced into the procedures.

C. Fresh isolates with the *DbtS*⁺ phenotype

Acquisition of gram negative, sexually-competent bacteria with the *DbtS*⁺ phenotype continues to be a major goal. Fresh isolates with these attributes were acquired and have been preliminarily characterized. The principal concern will be whether the phenotype is stable when the isolates are recurrently subcultured in standard bacteriological media.

II. DESULFURIZATION WITH N1-36

A. Production of OH-BP and o,o'-biphenol (!) by N1-36

In December 1991 (during the previous quarter of contracted research), two batch cultures with N1-36 were run for 210 hours in New Brunswick Scientific Bio-Flo III or C-30 "fermenters" (batches 6 and 7, respectively). In the former instance, the sole sulfur source was DBTO₂; in the latter DBT. In both cases, the pH was maintained at 6.0. With "batch 6", 0.3% glucose was added at 26 hours; with "batch 7", glucose was added at 27 hours. The data recorded from these experiments were previously presented. HPLC analysis of production of OH-BP was performed during

U.S./DOE Patent Clearance is not required prior to the publication of this document.

the present quarter (specifically in January of 1992). The data for the disappearance of substrate and appearance of product(s) appear on page 4.

In both instances, the concentration of DBT or DBTO₂ diminishes rapidly. The concentration of OH-BP peaks at a time approximately equivalent to the transition from late exponential phase to stationary phase. As was previously noted, the production of OH-BP is greater when DBTO₂ is the substrate as compared with DBT; for the former, the amount of OH-BP was nearly 50 μ M whereas with the latter the concentration of product was approximately 33 μ M.

An unexpected product eluted from the C-18 HPLC column with a retention time of 1.83 min. When these samples were spiked with o,o'-biphenol, the product and the standard co-eluted as a single symmetrical peak.

B. Regulation of the DbtS⁺ phenotype: oxidation of DBT and desulfurization of OH-BP

Aliquots from three day old cultures of N1-36 were inoculated (5%) into seven media with different individual or combinations of sulfur-sources. The concentration of inorganic S was 200 μ M and the concentrations of organic S were 100 μ M. The media were:

1) MgSO ₄	2) DBTO ₂	4) DBT	6) BP [benzothiophene]
	3) DBTO ₂ + MgSO ₄	5) DBT + MgSO ₄	7) BP + MgSO ₄

Turbidity at 660 nm. was measured at 48.5 hr and 70 hr. (Please see data on page 5.) Also at these times, the ability to desulfurize both DBT and DBTO₂ was measured with aliquots of the cultures. One volume of culture from each of the seven media was challenged with four volumes of buffer containing either DBT or DBTO₂. The data for the disappearance of substrate (DBT or DBTO₂) and appearance of product appear on pages 5 and 6.

The data are highly suggestive (and the implications are very important). The data indicate that: i) DBT and DBTO₂ disappear under all conditions tested, ii) DBT is oxidized to DBTO₂ in the presence of MgSO₄, and iii) the product OH-BP forms only in the absence of MgSO₄.

Note that previous data had suggested that the disappearance of substrate occurred in medium which had supported growth but from which the cells had been removed. Appearance of OH-BP required the presence of cells. Collectively, these data suggest that the first aspects of uptake of the substrate may be non-enzymatic, that oxidation of the sulfur moiety of DBT is not suppressed by SO₄⁼, but that desulfurization is repressed by SO₄⁼.

The data from uninoculated control and the culture lacking DBT or DBTO₂ indicate that the measurements are somewhat noisy. An alternative interpretation is that some physical process is

U.S./DOE Patent Clearance is not required prior to the publication of this document.

N1-36 Batch-6

N1-36 Batch-7

U.S./DOE Patent Clearance is not required prior to the publication of this document.

EXTENT OF GROWTH AS MEASURED BY TURBIDITY AT 660 nm.*

<u>MEDIUM</u>	<u>48.5 hr</u>	<u>70 hr</u>
MgSO ₄	0.268	0.446
DBTO ₂	0.231	0.426
DBTO ₂ + MgSO ₄	0.099	0.284
DBT	0.245	0.416
DBT + MgSO ₄	0.226	0.390
BT	0.076	0.083
BT + MgSO ₄	0.239	0.383

*Samples were diluted 1:10 prior to measuring optical density.

THE LOSS OF SUBSTRATE (DBT or DBTO₂) AND APPEARANCE OF PRODUCT (OH-BP)*
IN 48.5 HR CULTURES AFTER GROWTH IN THE PRESENCE OF DBT or DBTO₂

<u>MEDIUM</u>	<u>ELAPSED TIME</u>	<u>DBT</u>		<u>DBTO₂</u>	
		<u>[SUBSTRATE]</u>	<u>[PRODUCT]</u>	<u>[SUBSTRATE]</u>	<u>[PRODUCT]</u>
MgSO ₄	20'	76647 ^S	- [▲]	35011	-
	120'	91880	-	29643	-
	220'	84237	-	28120	-
DBTO ₂	20'	59921	4040	37648	4882
	120'	55653	5447	27801	6155
	220'	+	5904	26033	5504
DBTO ₂ + MgSO ₄	20'	+	(4326)	38959	-
	120'	56995	(5332)	30030	-
	220'	62625	(6125)	28261	-
DBT	20'	77872	6049	35559	6268
	120'	62496	6832	29758	6505
	220'	27794	6553	27582	6449
DBT + MgSO ₄	20'	96183	-	31692	-
	120'	70948	-	25770	-
	220'	64747	-	31380	-
BT	20'	62166	-	35018	-
	120'	56370	-	26862	-
	220'	50682	-	25182	-
BT + MgSO ₄	20'	62353	-	34264	-
	120'	54470	-	28281	-
	220'	73756	-	27394	-

*Product abundance presented in parentheses is DBTO₂ rather than OH-BP.

^SUnits are integrated area under HPLC peak.

[▲]Peak present; integrator did not give a value.

⁺No OH-BP detected.

U.S./DOE Patent Clearance is not required prior to the publication of this document.

THE LOSS OF SUBSTRATE (DBTO₂) AND APPEARANCE OF PRODUCT (OH-BP)
IN 70 HR CULTURES AFTER GROWTH IN THE PRESENCE OF DBTO₂

<u>MEDIUM</u>	<u>ELAPSED TIME</u>	<u>DBTO₂</u>	
		<u>SUBSTRATE</u>	<u>PRODUCT</u>
MgSO ₄	1'	41597 ^S	- [▲]
	221'	39965	-
DBTO ₂	1'	41362	11509
	221'	35934	10853
DBTO ₂ + MgSO ₄	1'	47838	-
	221'	41439	-
DBT	1'	36092	10054
	221'	37531	8521
DBT + MgSO ₄	1'	40770	-
	221'	37389	-

^SUnits are integrated area under HPLC peak.

[▲]No OH-Bp detected.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

U.S./DOE Patent Clearance is not required prior to the publication of this document.

removing substrate during the course of the assay. But the variation in measurement and the extent to which OH-BP might be physically removed from the samples are less than the reductions observed in the presence of DBT. Another concern is that the amount of product does not consistently increase. However, results pertaining to the abundance of OH-BP in stationary phase indicate that OH-BP disappears with time; a similar effect may explain why the abundance of product at 220 min or 221 min is less than the amount seen at earlier times; this interpretation is being investigated.

C. Nutritional characteristics

1. Ethanol vs. glucose as carbon source

N1-36 was inoculated into media with no carbon source, 0.1% EtOH, 1.0% EtOH, 0.3% glucose, 0.1% EtOH + 0.3% glucose, and 1.0% EtOH + 0.3% glucose. Growth was monitored for 7 days. The data show that glucose alone is a poor carbon source and that any medium with ethanol supports vigorous growth. (Data not shown.)

2. Monohydroxybiphenyl (OH-BP) as carbon source

N1-36 was inoculated into media with no carbon source, 50 μM OH-BP, 0.1% EtOH + 50 μM OH-BP, 1.0% EtOH + 50 μM OH-BP. Growth was monitored for 7 days. The data show that OH-BP alone does not support growth. Again, any medium with ethanol supports vigorous growth. (Data not shown.)

The following HPLC data indicate that the OH-BP introduced into the medium was not consumed:

[OH-BP] in medium at 7 days

<u>Medium</u>	<u>Inoculated</u>	<u>Uninoculated</u>
OH-BP	28.38 μM	32.87 μM
OH-BP + 0.1% EtOH	32.36 μM	31.17 μM
OH-BP + 1.0% EtOH	31.72 μM	31.78 μM

D. Growth of N1-36 in continuous culture

N1-36 was grown in continuous culture in a New Brunswick Bio-Flo III apparatus. The working volume of the culture was approximately 2.1 liters. The dilution rates (D) were 0.05 hr^{-1} (two runs), 0.10 hr^{-1} and 0.20 hr^{-1} . Approximately 10 liters of medium were used at each dilution rate. Hence, the lengths of the runs were ≈ 96 hr, ≈ 48 hr, and ≈ 24 hr, respectively. The growth-limiting nutrient was sulfur; this element was present in the medium in the substrate reservoir in the form of DBTO_2 at a concentration of 50 μM .

U.S./DOE Patent Clearance is not required prior to the publication of this document.

The cell concentration was estimated from optical density at 660 nm. The concentrations of DBTO₂ and monohydroxybiphenyl (OH-BP) in the culture are to be measured (and values will be reported in a subsequent report). The substrate will be measured by HPLC whereas the product will be measured both by HPLC and Multi-Excitation Fluorescence Spectroscopy. (Presently, the bulb on the HPLC detector is blown.)

A number of problems emerged. In an initial run, the DBTO₂ precipitated from the medium; stirring and mild heating of the substrate reservoir appeared to alleviate this problem. More specifically, visible precipitation was not evident in a four day period. Another problem was accumulation of bubbles at the slowest growth rate. This problem was reduced with anti-foam. However, with the resolution of this difficulty, a final problem, viz., contamination, terminated experimentation.

The observations which can be reported at this time are very limited and solely qualitative. Steady-state growth appeared to occur at the two lower growth rates but not at the higher rate. (However, the data are too sparse to offer a definitive statement.) Nonetheless, the results would not be surprising since a dilution rate of 0.2 hr⁻¹ might well exceed the maximum instantaneous growth rate ($\mu_{\max}$) of N1-36 and steady-state growth only occurs when $D \leq \mu_{\max}$. The relation of the instantaneous growth rate and the generation time (G) is $\mu = \ln 2/G$; hence, if G were greater than 3.5 hr, washout would occur.

E. N1-36: aka *Rhodococcus erythropolis*

1. Cell and colony morphology

All the isolates which have the ability to desulfurize DBT or DBTO₂ are Gram-positive, non-endospore forming, non-acid-fast rods. The cells exist singly, or paired as in the letter "V," or, occasionally, connected in groups of 3 or 4 to form the shapes of the letters "N" and "W." The colonies are circular, convex or pulvinate, entire, smooth and butyrous.

2. Biolog analysis

Ninety-six-well Biolog GP microtiter plates, which have 95 wells with distinct substrates plus a redox indicator dye, were used to generate nutritional profiles of the isolates. Data were obtained from the Microlog I computer program pertaining to the percent similarity, distance, and goodness of identification for the unknowns (when compared with species with established identities). The data are summarized in Table 1 on page 9. The identification achieved for all three isolates was *Rhodococcus erythropolis*.

U.S./DOE Patent Clearance is not required prior to the publication of this document.

<u>ISOLATE</u>	<u>Species identity</u>	<u>Similarity</u> *	<u>Distance</u> **
N1-36	<i>Rhodococcus erythropolis</i>	0.830	2.795
N1-36	<i>Rhodococcus erythropolis</i>	0.607	6.35
N1-43	<i>Rhodococcus erythropolis</i>	0.799	3.505
N1-43	<i>Rhodococcus erythropolis</i>	0.716	4.57
Q1a-22	<i>Rhodococcus erythropolis</i>	0.667	4.967
Q1a-22	<i>Rhodococcus erythropolis</i>	0.747	3.89

- * Similarity coefficients estimate relatedness of the experimental pattern with an averaged pattern observed among strains of the identified species.
- ** Distance measures how far the data are from being a perfect match with the average of a given species. Distance is measured in units of differences in numbers of test wells observed on a Biolog plate.

These observations are highly suggestive. Control experiments were run with *Bacillus cereus*. The Biolog program did not correctly identify this organism; the identifications offered in rank order of similarity were *B. sphaericus*, *B. badius*, and *B. brevis*. Additional efforts for species identification will be undertaken.

3. Fatty-acid analysis

The fatty acids of N1-36, N1-43, and Q1a-22 grown on trypticase soy agar were extracted and analyzed by gas chromatography; the digitized chromatographic spectrum was compared with a database of $\geq 60,000$ spectra; no identification of the organism was achieved. (The extraction, chromatography, and comparison was performed by Microbial ID, Inc., Newark, DE.) Notably, the database does contain profiles for *R. rhodochrous* and *R. erythropolis*.

4. General conclusion regarding identity

The species and genus of the isolates remains uncertain. The identification based on the nutritional profile strongly indicate that the identity is *R. erythropolis*. This genus identification is the same as that achieved by Kilbane et al. for an isolate obtained in a similar manner and having a similar phenotype; however, the species-specific identity of the Kilbane isolate differs; it is *R. rhodochrous*. As noted earlier, this identification was not confirmed by gas chromatographic profiles of whole cell fatty acids despite the fact that profiles for both *R. erythropolis* and *R. rhodochrous* were in the database.

F. Acknowledgement

The data regarding N1-36 are observations obtained by Ms. Ping Wang, a Ph. D. student in the Department of Molecular Biology at Lehigh University.

III. PULSED-FIELD GEL ELECTROPHORESIS OF CHROMOSOMAL DNAs OF *THIOBACILLUS* spp.

A. General considerations

A qualitative summary is that the enzyme *SpeI* gives reliable digestions of DNAs from both *T. acidophilus* 27807 and *T. novellus* 8093; with this enzyme, nine bands are obtained with the former organism and ten bands with the latter. For the former organism, reproducible patterns are also achieved with *XbaI* (see upper panel on page 11). The aesthetic quality of the gels is improving notably. Estimates of the sizes of fragments were previously established and appear reliable. Technologies for quantifying the amount of DNA in the bands are being assessed. Precautions against inadequate resolution of fragments which appear to co-migrate are being evaluated. Work persists to find additional enzymes which yield reproducible results. A credible estimate of chromosomal size requires digestions by at least two enzymes; the sums of the sizes of the fragments produced by each of the enzymes must give equivalent or nearly equivalent values.

B. Reliable estimation of sizes of standards

Pulsed field gel electrophoresis (PFGE) was developed for the estimation of size of large nucleic acids. The data now acquired on 20 cm. gels with chromosomes of *Saccharomyces cerevisiae* and "lambda ladders" are fully compatible (please see figure on page 11. The multiple lines are replicate results.)

C. Some technical advances

A variety of technical conditions were experimentally evaluated. For example, the necessity of having freshly grown bacteria was assessed and found not to be essential. In protocols for PFGE, recently harvested bacteria are embedded in agarose plugs; the bacteria are then lysed and the released DNA is subjected to restriction digestion; finally, the restriction fragments are electrophoresed. We observed that agarose plugs in which *T. acidophilus* 27807 had been embedded last August were an adequate source of materials for digestion. Additional refinements in procedure had to do with new materials, e. g., Sea Kem GTG ("genetic technology grade") agarose as well as a new TBE ("Tris-borate-EDTA") buffer.

D. Estimation of generation times

Knowledge of the generation time of the *Thiobacillus* spp. used in the contracted research will be important when attempting to determine that all rounds of chromosome replication associated with a cell generation are complete; failure to consider this matter may mean over-representation of restriction fragments near the origin of replication or under-representation of fragments near the terminus. The generation times of *Pseudomonas aeruginosa* PAO and *Thiobacillus novellus* 8093 were reported to be,

U.S./DOE Patent Clearance is not required prior to the publication of this document.

LANE A: lambda concatemers

LANE B: *T. acidophilus* cut with XbaI

LANE C: uncut plasmid

LANE D: plasmid cut with XbaI

LANE E: plasmid cut with SpeI

LANE F: 5 kb ladder

Collective Data of DNA Standards

U.S./DOE Patent Clearance is not required prior to the publication of this document.

respectively, ≈ 1.5 hr and ≈ 11 hr. These two values were confirmed this month and data were collected which establish that the generation time for *T. acidophilus* 27807 is ≈ 12 hr.

D. Acknowledgement

The data on pulsed-field gel electrophoresis were generated by Ms. Diane Dutt, a technician and graduate student in the Department of Molecular Biology at Lehigh.

III. Fresh isolates with the presumptive capacity to desulfurize DBT

A. Isolation, abundance, and screening

In late February, soil samples were collected at Petrilla Fuel Company in Bethlehem; the sampling sites had previously been the source of organisms which had seemed to hold great promise.

The abundance of bacteria in the samples was assessed on five solid media; the media were distinguished on the basis of identity and strength of carbon source. The media were on one containing salicylic acid (SAL), a second containing benzoic acid (BEN), and a third containing glucose (GLU); the fourth medium was Luria broth (LB) and the fifth medium was 1/10 strength Luria broth supplemented with glucose. The numerical results are summarized in the following table.

COLONY FORMING UNITS PER GRAM OF SOIL SAMPLE

<u>SAMPLE SITE</u>	<u>MEDIUM</u>				
	<u>SAL</u>	<u>BEN</u>	<u>GLU</u>	<u>LB</u>	<u>1/10 LB + GLU</u>
A	6.3×10^4	2.5×10^4	3.8×10^5	1.6×10^5	7.4×10^5
C	2.1×10^3	1.1×10^3	3.7×10^5	6.3×10^5	8.4×10^5
L	TFTC*	1.2×10^E	2.2×10^4	3.5×10^4	8.7×10^4

* TFTC = too few to count

The "fluorescent spray plate assay" was used to identify isolates for further characterization. Eight isolates were obtained from site A; six were from plates growing on benzoic acid and two from plates growing on salicylic acid. A ninth isolate from a plate containing glucose was not characterized. Further, two isolates from site L which had been grown on plates containing salicylic acid were selected.

B. Presumptive identities

Preliminary identification of the organisms was undertaken. All the strains were gram negative; hence, identification using Biolog GN plates was appropriate. Seven of the isolates gave presumptive identifications as *Pseudomonas* spp. and three were presumptively *Klebsiella pneumoniae* subgroup A. These identifi-

U.S./DOE Patent Clearance is not required prior to the publication of this document.

cations are tentative since two of the identifications were labeled (by the computer program) as being "borderline" while five were "poor" and three were "good."

C. Desulfurization

A colorimetric assay was used to detect the presence of phenolic moieties produced from DBTO₂. The assay was used in a standard format with large sample volumes; the abundance of chromogen was measured by spectrophotometry. The assay was also used in a qualitative manner with organisms grown in the presence of DBTO₂ in wells of microtiter plate; color development was detected eyeballometrically. The data indicate that at least three and possibly more strains produce phenolic moieties. (The data are not included because: [i] they are preliminary and [ii] they were collected after the eleventh quarter of contracted research concluded.)

D. Acknowledgement

Mr. Todd Davies, a graduate student, isolated the fresh strains, quantified their abundance, screened for the Dbts⁺ phenotype, performed the Biolog analysis, and subsequently performed the phenol colorimetric assay. He did so during a six-week rotation in my laboratory; he is eager to gain experience and plans to obtain more definitive results during the summer.

V. CONCLUSION

The Dbts⁺ phenotype in N1-36 appears to be stable; hence, kinetic analysis of the production of OH-BP from DBT and DBTO₂ is proceeding. Such physiological characterization is important if successful genetic manipulation is to be achieved.

Characterization of the chromosomal DNA from *Thiobacillus* spp. is being achieved through PFGE. Knowledge of the chromosomal organization (as well as accessory elements) will be important if the genes responsible for the Dbts⁺ phenotype are to be cloned in these sulfur oxidizing organisms.

The presence of the Dbts⁺ in sexually-competent, well-characterized, gram-negative bacteria remains an alluring phenomenon. The recurrent isolation of such strains allows investigation of factors which contribute to the instability of the phenotype in the laboratory. Knowledge of these factors may allow maintenance and control of the phenotype.

U.S./DOE Patent Clearance is not required prior to the publication of this document.

VI.

MILESTONE SCHEDULE/STATUS REPORT

Selection and preliminary characterization of fresh *Pseudomonas* isolates which stably express the Dbts⁺ phenotype:

28-----29-----30-----31-----32-----33-----34-----35-----36

Fresh isolates available and awaiting characterization.

Modeling R' formation and transfer in *Pseudomonas* spp.:

28-----29-----30-----31-----32-----33

This aspect of the work is nearly complete. A manuscript is being drafted.

Kinetics and site of monohydroxybiphenyl production by N1-36:

28-----29-----30-----31-----32-----33-----34-----35-----36

Excellent progress.

Genus and species identification of N1-36 and other gram positive isolates:

29-----30-----31-----32

Either it is *Rhodococcus* or a new genus.

Pulsed-field gel electrophoresis of *Thiobacillus* spp. chromosomes:

28-----29-----30-----31-----32-----33-----34-----35-----36

Good progress. One more enzyme needed for *T. novellus*.

Preparation of final report:

34-----35-----36

U.S./DOE Patent Clearance is not required prior to the publication of this document.

VII.

COST MANAGEMENT REPORT

DATE RUN	04/01/92	LEHIGH UNIVERSITY	REPORT PAGE 387		
TIME RUN	C7:52:09	FINANCIAL RECORDS SYSTEM	PAGE 1		
ACCT: 5-35074		RESEARCH ACCOUNT SUMMARY IN WHOLE DOLLARS AS OF 03/31/92	ACCOUNT PAGE 1		
DEPT: MOLECULAR BIOLOGY		START 06/30/89	KRAMIEC S		
		EXP 06/29/92	MOLECLR BIOL-IAOCCA		
			BUDGET		
			AVAILABLE		
			58		
			5,043-		
			420		
			320		
			6,655-		
			4,245-		
			2,471-		
			209		
			862-		
			1,949		
			15,434		
			235-		
			228-		
			2,360		
			16,156		
			11,910		
			11,910		
DESCRIPTION	PROJ-TO-DATE	BUDGET	BALANCE BEFORE	DOE	OPEN
FACULTY - SUMMER	50,197	50,139	58	COMMITMENTS	COMMITMENTS
SUPPORT STAFF	39,035	40,123	1,088-		3,955
RES ASST STIPEND	28,320	25,200	3,120		2,700
OTHER	1,520	1,200	320		
* TOTAL SALARY/WAGE	119,072	116,662	2,410		6,655
E B REIMB-SPONS RES	28,335	27,807	528		528
** TOTAL COMPENSATION	147,407	144,469	2,938		7,183
SUPPLIES	27,635	30,306	2,471-		
COMMUNICATION/POSTAGE	765	556	209		
DOMESTIC TRAVEL	7,068	7,930	862-		
PUBLICATION COSTS	2,430	481	1,949		
EQUIPMENT (1)	27,160	11,726	15,434		
OUTSIDE SERVICES		235	235-		
EQUIPMENT USAGE		228	228-		
OTHER	4,930	316CR	2,570		
* TOT OTHER DIRECT COST	70,188	1,739	16,156		
*** TOTAL DIRECT COSTS	217,595	198,502	19,093		7,183
INDIRECT COSTS	116,167	113,539	2,628		2,628
**** GRAND TOTAL	333,762	312,041	21,721		9,810

(1) - NOT SUBJECT TO INDIRECT COST.

FOR THIS REPORT, SOME FRS EXPENSE CODES HAVE BEEN CONSOLIDATED TO PROVIDE SUMMARY INFORMATION. TRANSLATION OF THESE CODES TO THE FRS FOUND ON THE R96 SCREEN ON THE DEPARTMENT CENTER/INSTITUTE OFFICE. FRS CODES ARE AVAILABLE TO THE DEPARTMENT CENTER/INSTITUTE OFFICE. FOR MORE INFORMATION, PLEASE CONTACT THE R96 AND/OR THE R93 FRS SCREENS. YOU WILL NEED FURTHER ASSISTANCE. PLEASE CONTACT YOUR PROGRAM ADMINISTRATOR IN THE OFFICE OF RESEARCH AND SPONSORED PROGRAMS.

NOTE: YOU WILL CONTINUE TO RECEIVE THE PAYROLL MANAGEMENT REPORT FOR DETAILED SALARY INFORMATION.

**DATE
FILMED
8/3/92**