KRALL ASSOCIATES

DOE/ER/53280--6

DE92 018557

ANNUAL PERFORMANCE REPORT FOR "A PROGRAM OF FRC THEORY RESEARCH"

Prepared by
Nicholas A. Krall

Prepared for
U.S. Department of Energy
Washington, D.C.

under
Grant No. DE-FG03-88ER53280

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

May 15, 1992

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

1. INTRODUCTION

The original activities carried out under this Grant were specific to the Fusion scheme known as the Field Reversed Configuration (FRC). With the decisions of the Department of Energy to deemphasize experimental activity in the areas known as "alternate concepts," we have generalized our studies. Since the research we have performed on the FRC led us into the area of high beta ($8\pi nT/B^2$) and large orbits (gyroradius comparable to the density or magnetic field scale length), we have chosen to emphasize those areas of plasma physics. The relevance to the mainline fusion program will emerge as our understanding of these areas develops. This statement may not be obvious, but it reflects my observations, developed over the last thirty-two years of fusion research, of how progress is made in this complex field. During the present contract period, we have undertaken three specific studies on large orbit and high β physics which have reached a point of near completion.

One of these studies has demonstrated that in large orbit systems collisional relaxation to a Maxwellian distribution proceeds at a surprisingly slow rate, because of the competing effects of collisions which tend to make a local Maxwellian, which in a large orbit system can make remote regions more non-Maxwellian. This work is described in Section 2, and in complete detail in Appendix A.

A second study reopens the problem of confinement of plasma by a cusp magnetic field. This area of research was in vogue before high speed computing was available to aid the study. The inherent high β and large orbit nature of this problem makes the problem nearly intractable without numerical tools. We

have made both analytic and numerical advances in this area. Section 3 and Appendix B present the status of this work.

Thirdly, we note that the separatrix in the FRC can lead the plasma into an x-point on the cylindrical axis of that device. The magnetic topography at that point is essentially a spindle cusp. Confinement in that region can affect the confinement in the entire device by continuity of flow (the importance of the transport properties of the separatrix region has been stressed for quite awhile now by Dr. Loren Steinhauer of STI Optronics). Thus our studies on cusp confinement can be seen to reflect directly on the properties of the FRC. We have begun to explore this phenomenon, in collaboration with Dr. Steinhauer and expect further interaction with Dr. Akio Ishida, Niigata University, Japan, as well. Section 4 and Appendix C summarize the work to date on this problem.

In the later portions of this Grant, we intend to complete these studies, and begin studies of applications of our work to fusion devices. One possible application is to the effect of finite β on the ion temperature gradient drift instability. This mode is nearly neutrally stable, and the recent Sherwood meeting in Santa Fe, NM included a number of papers on weak effects which could affect this mode. Finite β is an effect so far overlooked. We intend to pursue this facet of eta-i stability in the remainder of the Grant period. Despite the waning of the U.S. effort in FRC experimental research, the world program in this area continues. We intend to begin a collaboration with STI Optronics and Japan on this magnetic geometry, to help the U.S. stay current, and contribute to progress in new areas of fusion research. Section 5 describes a possible program plan.

2. COLLISIONAL EFFECTS IN LARGE ORBIT SYSTEMS

In this study we have shown that for a moderately long time scale, two-body collisions can actually help in maintaining a non-Maxwellian ion velocity distribution in a plasma. This surprising effect occurs when the ions have orbits large enough to transit spatial regions with different physical parameters in a single collision time. Although this is a generic effect expected in all systems where the equilibrium properties vary strongly over a particle orbit, the effect in the present study is applied to ions confined in a spherically convergent ion focus, where collisions at the edge of a spherical potential well help maintain convergent non-Maxwellian flow in the rest of the device.

It is well known that on a long enough time scale, two-body collisions will drive a plasma to a Maxwellian distribution. On shorter time scales this need not be the case. There are systems characterized by large ion orbits in a system with spatially varying density and kinetic energy in which two-body collisions which thermalize the plasma in one region can thereby help maintain a non-Maxwellian flow in other, more spatially extensive, regions. An example of such a system is the spherically convergent ion focus, and we use that geometry as a specific geometry in which to demonstrate the effect.

The idea of this type of device is to create a spherical potential well of sufficient depth to accelerate ions from low energy at the periphery of the well to fusion energies within a focus at the center of the sphere. The potential well which accelerates the ions is maintained at a constant value sufficient to confine the ions within the device, returning them again and again at high velocity to the central focus. Essential to the success of the scheme is that the ions maintain their nonthermal velocity distribution, with primarily

radial flow, long enough to produce fusion in the dense focus at the center of the sphere.

If the ion source is at the periphery and ions return to their birth point (turning points), then an isotropic source at R will clearly produce an anisotropic (non-Maxwellian) distribution at $r < R$, since v_r increases as $\sqrt{2e\phi(r)/m_i}$, while from conservation of angular momentum $v_\perp$ increases as $v_\perp = v_{\perp R} R/r$, with $2e\phi_{\max}/m_i = v_{\perp R}^2 R^2/r_c^2$ determining r_c , the closest approach of an ion to the center, where $v_{\perp R}$ is the velocity perpendicular to $\hat{i}_r$ of an individual ion at $r = R$, and where $\phi = 0$ at the location of the ion source. If the ions must converge to a high density, small radius focus, to form a reactor relevant configuration, ion-ion collisional effects can destroy the configuration, even if they do not cause ions to leave the device.

There are two distinct effects that collisions have on ion convergence. First, as ions born at low energy pass through the dense core of the device, they will be nearly monoenergetic, at the energy of the well. Collisional processes will tend to thermalize the velocity distribution in the core. An ion which is upscattered in energy will have a turning point at a radius larger than the radius at which it was born, and can eventually be lost to the dense core by escape from the device or scattering from fields arising from the mechanism used to produce the potential. This could involve grids for the case of externally produced potentials, or magnetic fields which confine electrons for the case of potentials produced by injecting electrons, and confining them magnetically at the surface of the device.

The second collisional effect involves isotropization of the ion velocity distribution in the bulk of lower density plasma outside the core. In this large scale region, ion-ion collisions can transfer momentum from the primarily radial flow, which produces good spherical convergence, to a more isotropic

distribution of velocities. This deflection by collisions leads to an increase of local azimuthal velocity in this bulk region, which clearly decreases the convergence of the ions to the core.

In addition to the core and the bulk plasma which surrounds it, there is a third region with distinctive collisional properties, namely the edge plasma, where the bulk of the ions have their turning points. In this region ion velocities are low, and collisions much more frequent than in the interior. In this region the ion distribution due to previous collisions in the interior tends to become more anisotropic, while collisions in the edge itself make the edge distribution more isotropic. An isotropic distribution at the edge will produce exactly the strong radial convergence required for the SCIF. Thus due to the large ion orbits, collisional processes in the edge will tend to drive the distribution in the interior to stay anisotropic and convergent. In this, edge collisions compete with local collisions in the interior which attempt to produce a local Maxwellian. If edge collisions dominate, even though large angle scattering will eventually thermalize the entire system, the more rapid small angle scattering (which competes with the fusion reacting time scale) will force the distribution to stay non-Maxwellian in the interior. This effect is simply a matter of competitive rates of thermalization in the various regions.

Appendix A describes this work in detail. Firstly, we calculate the ion collision frequencies in the various spatial regions, for a variety of plasma conditions. Then we carry out a Fokker-Planck-like calculation of the ion distribution which results from the combined effect of collisions in all three regions, and show that the edge effects dominate, producing a Maxwellian edge and a non-Maxwellian convergent interior. Lastly, we apply the results to estimating loss rates in a reactor based on a spherically convergent ion focus.

3. THE THEORY OF CUSP CONFINEMENT

The use of magnetic point cusps to confine plasma is a long standing problem in fusion research. Despite the long history of this problem, and the fairly large number of examples where some portion of a plasma device has cusp geometry (divertors, plasma guns, magnetic guide fields, etc.) we have found that there is no definitive calculation of cusp confinement, either theoretically or experimentally, in the parameter range where plasma density and temperature are sufficiently high that the magnetic field is excluded by diamagnetic effects from most of the volume. The reason for this gap in confinement lore is that high- β cusp confinement is a collective process,^{1,2} with plasma currents acting to exclude the B-field. Collective processes are traditionally difficult to pin down; further, when cusp confinement was an active area of research, tools for numerical simulation of this multidimensional, high density system were not available. Experimentally, the situation was not much better, with plasma formation techniques and available diagnostics leading to a clouded and often contradictory set of cusp confinement measurements. There are several reviews which document this situation.^{3,4}

Because of the occurrence of cusp geometry in a variety of devices, and because of this gap in plasma confinement theory, we have embarked on a program to measure cusp confinement by numerical simulation. Because of the expense in computer time involved in simulating an inherently 3-dimensional, high density, large volume, and non-neutral plasma, we are approaching the problem in scaled stages, choosing plasma parameters which preserve the essence of the cusp problem while minimizing the numerical difficulties.

From a simplified point of view, the question of cusp confinement involves being able to distinguish between several competing confinement ideas. These include the following:

- o Single particle confinement: Even in the absence of diamagnetic exclusion of the B-field from the plasma, particles are confined by mirror reflection as the cusp field increases near the outside of the device. Particles are "scattered" into the loss cone by passing through the low field center of the cusp, where their magnetic moment is no longer invariant. The mirror ratio is the ratio of the maximum cusp field to the field at which the particles again become adiabatic.
- o Ergodic sampling of a field distribution with a few well defined escape holes. When the plasma excludes the B-field, it is possible that a geometry will develop in which there is a virtually field-free interior region, and a high field exterior region, separated by a sharp interface, of the order of an electron gyroradius. In that case, the particles free stream across the interior, reflecting specularly off the interface, until by chance they strike a small area near the corner of the cusp, Figure 1, whereupon they are lost. This is the classic picture of cusp confinement. The question is what is the size of the loss hole? Possibilities include the following:
 - a. A hole of area πr_e^2 , where r_e is the electron gyroradius. This was suggested by Berkowitz.¹

- b. A hole of area $2\pi R r_e$, where R is the characteristic size of the device. This was calculated by Taylor² as an upper limit to the loss rate in a spindle cusp, where there was a symmetry axis which led to a line cusp of length $2\pi R$.
- c. A hole of area πr_i^2 , where r_i is the ion gyroradius. This scaling arises from considerations of the stability of a very thin plasma-field interface.⁵
- d. A hybrid area, $\pi r_i r_e$, estimated from at least 2 experiments.^{6,7}

In our first effort, we have simulated confinement in a spindle cusp. This allows a two-dimensional treatment of the problem, simplifying the computation. We study a neutral plasma, with $T = 120$ keV and $B_{\max} = 3$ kG. This also simplifies the computation by allowing high β studies at acceptably low densities. Essential to the study of course is that r_e and r_i be small compared to the system size R . Therefore we chose $r = 25$ cm. The ion mass is unrealistically taken to be a small multiple of the electron mass m_e , again for numerical convenience, while still maintaining the dimensionless variables of the system in a range appropriate to studying cusp confinement.

In Appendix B, we document the predictions of the various confinement ideas listed above for the parameters of the simulation. We also make a stab at estimating the effects of ambipolarity on the confinement. This work is ongoing, and we expect results by the end of the present contract period. Results to date confirm the low β aspects of the theory, the so-called mirror limit of cusp confinement.

4. CONFINEMENT AT THE FRC SEPARATRIX

Although this was not a direct motivator for our study of cusp confinement, it turns out that there is a direct application of that work to confinement in the FRC. The field reversed configuration (FRC)^{8,9} includes a region of closed field lines bounded by a region of open field lines (Figure 2). Transport of particles and energy from the device includes a continuity condition, which says that the essentially axial loss of plasma along magnetic field lines must be matched by the essentially radial plasma flow across magnetic field lines. Early attempts at modelling FRC loss rates^{10,11} assumed that the plasma density would drop to nearly zero in a distance of the order of the ion gyroradius outside the separatrix, because of the very rapid loss along the open field lines. It was observed however that this assumption was not in agreement with the data, which showed a substantially broader sheath.¹²

A closer examination of the magnetic geometry (Figure 3) shows that the field lines at the separatrix lead the plasma into an end configuration which is essentially a spindle cusp. The confinement properties of the FRC as a whole, as well as the structure of the transition between open and closed field lines, will be influenced by an interplay between transport from the interior of the FRC and transport of plasma from a cusp. We have explored this feature of FRC boundary layer physics, and obtained a prescription for determining the sheath width, and the confinement time in the sheath. This work is detailed in Appendix C. The work is ongoing, in that we intend to use continuity of particle and energy flow to determine the full implications of this boundary layer phenomenon on global confinement in the FRC.

5. FUTURE WORK, INCLUDING PLANNED COLLABORATIVE PROGRAM WITH STI OPTRONICS AND JAPAN ON FRC

In the succeeding performance period, we have a number of tasks in mind. Firstly, the work on cusp confinement and its implications for FRC confinement must be completed. Next, the application of our high β studies to the eta-i mode as cited in the Introduction should be pursued, and might be a significant contribution to the mainline program. A further exciting possibility is a collaboration with the Japanese program on FRC research. With the current decline in DOE expenditures on FRC experiments, it would be valuable to maintain contact with (and also contribute to the growth and success of) this exciting field. A collaboration with Niigata University in Japan has been discussed, with Dr. Loren Steinhauer of STI Optronics, Professor Akio Ishida of Niigata, and myself as the "gang of three" who would spearhead the collaboration. In the following Grant period, I intend to pursue this collaboration. A possible work agenda was drafted by Dr. Steinhauer, as follows:

CONFINEMENT IN FIELD-REVERSED CONFIGURATIONS

"Gang of Three" Strategy Sheet

I. PARTICLE CONFINEMENT

A. Dispersion Relation

- (1) Clearly define and defend the foundational assumptions made in deriving the d.r.; identify and discuss the neglected effects; identify what changes (if any) in extrapolating to a reactor.

- (2) Derive the d.r.
 - (3) Determine the unstable modes and their frequencies including the range of mode numbers (axial, azimuthal) for instability; identify limits on mode numbers, e.g., (a) when the basic assumptions break down, or (b) limits set by geometry or other conditions; identify if new mode types appear in going toward the reactor regime.
- B. Mode Saturation
- (1) Determine the saturation levels for various mechanisms relevant to each axial and azimuthal mode number.
 - (2) Compare the various saturation mechanisms and identify the regimes or conditions over which each is most restrictive.
- C. Transport Rate
- (1) Sum over all allowed unstable modes with the appropriate saturation level to determine the total local transport rate.
- D. Global Confinement
- (1) Identify and characterize other regulative mechanisms such as "profile consistency" which may affect the FRC structure.
 - (2) Derive a global confinement model which accepts an arbitrary local transport (expressed as a formula).
 - (3) Solve the global confinement model with the appropriate transport.
- E. Comparison with Experimental Data Base
- (1) Develop a systematic and broad-based confinement data base. Collect data for both global confinement times and edge layer properties.

- (2) Compare the theory predictions with the data base.
- (3) Predict the confinement as conditions approach the reactor regime.

II. FLUX CONFINEMENT (later)

III. ENERGY CONFINEMENT (later)

Some of the work has already been done or begun by either myself or the other members of the gang. This would be a very cost-effective way of maintaining a critical mass of activity in this area without the DOE having to fund a large program on their own.

REFERENCES

1. J. Berkowitz, K. O. Friedrichs, H. Goertzel, H. Grad, J. Kileen, and E. Rubin, Proceedings of the 2nd Intl. Conf. on Peaceful Use of Atomic Energy, Geneva, 31, 171 (1958).
2. J. B. Taylor (private communication), Culham Report CLM-R58 (1966).
3. I. Spalding in Advances in Plasma Physics, eds. A. Simon and W. B. Thompson (Wiley, New York, 1971), Vol. 4, p. 79.
4. M. G. Haines, Nucl. Fusion 17, 811 (1977).
5. S. Hamasaki, R. C. Davidson, N. A. Krall, and P. C. Liewer, Nucl. Fusion 15, 27 (1975).
6. A. Kitsunezaki, M. Tanimoto, and T. Sakiguchi, Phys. Fluids 17, 1895 (1974).
7. N. Hershkowitz, K. N. Leung, and T. Komesser, Phys. Rev. Lett. 35, 277 (1975).
8. W. T. Armstrong, R. K. Linford, J. Lipson, D. A. Platts, and E. G. Sherwood, Phys. Fluids 24, 2068 (1981).
9. J. Lipson, W. T. Armstrong, J. C. Cochrane, K. F. McKenna, E. G. Sherwood, M. Tuszewski, S. Hamasaki, Appl. Phys. Lett. 39, 43 (1981).
10. S. Hamasaki and N. A. Krall in Proceedings of IEEE 1979 International Conference on Plasma Science, Montreal (IEEE, New York, 1979), p. 143.
11. M. Tuszewski and R. K. Linford, Phys. Fluids 25, 765 (1982).
12. L. Steinhauer (private communication).

APPENDIX A.
THE EFFECT OF COLLISIONS IN MAINTAINING A NON-MAXWELLIAN PLASMA DISTRIBUTION
IN A SPHERICALLY CONVERGENT ION FOCUS

I. INTRODUCTION

In this study we show that for a moderately long time scale, two-body collisions can actually help in maintaining a non-Maxwellian ion velocity distribution in a plasma. This effect occurs when the ions have orbits large enough to transit spatial regions with different physical parameters in a single collision time. The effect is applied to ions confined in a spherically convergent ion focus, where collisions at the edge of a spherical potential well help maintain convergent non-Maxwellian flow in the rest of the device.

It is well known that on a long enough time scale, two-body collisions will drive a plasma to a Maxwellian distribution. On shorter time scales this need not be the case. In this study we show that there are systems characterized by large ion orbits and a spatially varying density and kinetic energy in which two-body collisions which thermalize the plasma in one region can thereby help maintain a non-Maxwellian flow in other, more spatially extensive, regions. An example of such a system is the spherically convergent ion focus device, and we use that geometry both as an important application of this unusual effect of collisions, and as a specific geometry in which to demonstrate the effect.

The idea of this device is to create a potential well of sufficient depth to accelerate ions from low energy at the periphery to fusion energies within

a focus at the center of the sphere. The potential well which accelerates the ions is maintained at a constant value sufficient to confine the ions within the device, returning them again and again at high velocity to the central focus. Essential to the success of the scheme is that the ions maintain their nonthermal velocity distribution, with primarily radial flow, long enough to produce fusion in the dense focus at the center of the sphere.

If the ion source is at the periphery and ions return to their birth point (turning points), then an isotropic source at R will clearly produce an anisotropic (non-Maxwellian) distribution at $r < R$, since v_r increases as $\sqrt{2e\phi(r)/m_i}$, while from conservation of angular momentum $v_\perp$ increases as $v_\perp = v_{\perp R} R/r$, with $2e\phi_{\max}/m_i = v_{\perp R}^2 R^2/r_c^2$ determining r_c , the closest approach of an ion to the center, where $v_{\perp R}$ is the velocity perpendicular to $\hat{i}_r$ of an individual ion at $r = R$, and where $\phi = 0$ at the location of the ion source.

If the ions must converge to a high density, small radius focus, to form a reactor relevant configuration, ion-ion collisional effects can destroy the configuration, even if they do not cause ions to leave the device. There are two distinct effects that collisions have on ion convergence. First, as ions born at low energy pass through the dense core of the device, they will be nearly monoenergetic, at the energy of the well. Collisional processes will tend to thermalize the velocity distribution in the core. An ion which is upscattered in energy will have a turning point at a radius larger than the radius at which it was born and can eventually be lost to the dense core by escape from the device or by deflection from fields associated with the creation of the potential well. This can be electrostatic grids in the case of an externally produced potential, or magnetic fields, if the potential is produced by electrons injected from guns and confined by B-fields which increase with radius.

The second collisional effect involves isotropization of the ion velocity distribution in the bulk of lower density plasma outside the core. In this large scale region, ion-ion collisions can transfer momentum from the primarily radial flow, which produces good spherical convergence, to a more isotropic distribution of velocities. This deflection by collisions leads to an increase of local azimuthal velocity in this bulk region, which clearly decreases the convergence of the ions to the core.

In addition to the core and the bulk plasma which surrounds it, there is a third region with distinctive collisional properties, namely the edge plasma, where the bulk of the ions have their turning points. In this region ion velocities are low, and collisions much more frequent than in the interior. In this region the ion distribution due to previous collisions in the interior tends to become more anisotropic, while collisions in the edge itself make the edge distribution more isotropic. An isotropic distribution at the edge will produce strong radial convergence. Thus due to the large ion orbits, collisional processes in the edge will tend to drive the distribution in the interior to stay anisotropic and convergent. In this, edge collisions compete with local collisions in the interior which attempt to produce a local Maxwellian. If edge collisions dominate, even though large angle scattering will eventually thermalize the entire system, the more rapid small angle scattering (which competes with the fusion reacting time scale) will force the distribution to stay non-Maxwellian in the interior. This effect is simply a matter of competitive rates of thermalization in the various regions.

In Section II, we calculate the ion collision frequencies in various spatial regions, for both experimental and reactor grade plasmas based on spherically convergent ion focii, with the potential produced by an excess of

electrons.^{1,2} In Section III, we give a Fokker-Planck-like calculation of the ion distribution which results from the combined effect of collisions in all three regions, and show that the edge effects dominate, producing a Maxwellian edge and a non-Maxwellian convergent interior. In Section IV we apply the results to estimating loss rates in a reactor based on potentials produced by injected electrons.

II. ESTIMATES OF ION COLLISION RATES

In this section we estimate the various ion collision rates, as discussed in Section I. Because collisions depend on energy and density, we first present the radial density and energy profiles expected in an idealized version of the spherically convergent ion focus, and discuss the collisional rates in the different regions, for both experimental and reactor-grade parameters, using the parameter set in Table I.

A. ION ENERGY AND DENSITY PROFILES

As a first approximation for the motion of a test ion in this type of device, we assume the ion undergoes conservative motion in a spherically symmetric electrostatic potential well. The ion is born with an energy E_0 at a location R . The total energy of the particle is conserved:

$$E_i(r) = \frac{1}{2} m_i (v_r^2 + v_\perp^2) - e\phi(r) \quad . \quad (1)$$

Here E_i is the total energy, m_i is the ion mass, v_r and $v_\perp$ are the radial and azimuthal ion velocities, $v_\perp^2 = v_\phi^2 + v_\psi^2$ in a spherical coordinate system, and $\phi(r)$ is the electrostatic potential energy.

We assume the electrostatic potential has the following spatial form:

$$\phi(r) = \phi_{\max} \left[1 - \left(\frac{r}{R} \right)^p \right] \quad , \quad (2)$$

where $\phi_{\max}$ is the value of the potential at $r = 0$, the center of the device, and $\phi = 0$ at $r = R$, where R is the location of the ion source. The actual form of $\phi(r)$ is not relevant to the qualitative results we obtain.

The test ion radial energy profile is then approximately

$$E_r = e\phi_{\max} \left[1 - \left(\frac{r}{R} \right)^p \right] + E_0 \quad . \quad (3)$$

The ion azimuthal energy profile follows from conservation of angular momentum, $rv_{\perp}(r) = \text{constant}$, leading to

$$E_{\perp} = \frac{1}{2} m_i v_{\perp R}^2 \left(\frac{R}{r} \right)^2 \quad ,$$

where $v_{\perp R}$ is the ion's azimuthal velocity at birth.

We assume that the ion source is active for a time long compared with the time it takes for an ion to flow from $r = R$ to $r \approx 0$ and back again, and that the source produces a low energy ion cloud at $r = R$. The ion distribution everywhere, and the radial ion density profile, are then determined by the potential $\phi(r)$ and the ion distribution function f_i at the birth point, which can be written in terms of the constants of the motion (E and $rv_{\perp}$), since the Vlasov equation for the steady state is $f_i(E, rv_{\perp}) = \text{constant}$. There are three quite different regions of the device, characterized by the relative values of the ion's birth energy, and azimuthal and radial energies. Within a very small radial distance inward from the ion birth point, the potential change is negligible, the ion energy is of the order of it's birth energy, and the density is given by $n = n_{\text{edge}}$. The potential increases inward radially, and since the

maximum well depth is in the keV range, compared with birth energies in the several eV range, it follows that $2e\phi \gg E_0$ a short radial distance from the edge, while $m_i v_{\perp R}^2 (R/r)^2 < 2e\phi$. These conditions hold for a large radial region of the device which we refer to as the bulk. Eventually, however, at a smaller radius, $r \ll R$, $m_i v_{\perp R}^2 (R/r_c)^2 = 2e\phi_{\max}$ and the ions can come no closer to $r = 0$. This defines the dense core region. This inward motion clearly evolves a Maxwellian source at $r = R$ into a radially counterstreaming bi-Maxwellian f_i at $r < R$ proportional to

$$e^{-\frac{m_i}{2T_r} (v_r \pm v_d)^2} e^{-\frac{m_i v_{\perp}^2}{2T_{\perp}(r)}},$$

where the radial drift responds to the radial electric field, $v_d = (2e\phi/m_i)^{1/2}$, and the transverse "temperature" $T_{\perp} = m_i v_{\perp 0}^2 (R^2/r^2)$ increases inwardly to provide conservation of angular momentum, with $m_i v_{\perp 0}^2$ an average perpendicular energy of ions born at $r = R$. If the ion source is distributed rather than sharply localized, the same arguments presented here can be used with reference to azimuthal "slices" of the ion source.

Integrating f_i over velocity yields the density in the various radial regions of the device. As a model for the ion distribution at the birth point, we assume that at birth the ion distribution function is uniform in energy up to some small energy E_0 , and uniform in angular momentum up to some small azimuthal velocity $v_{\perp}$, and take the potential at the birth point to be $\phi = 0$. This distribution is described in terms of the constants of the motion ($E = (1/2)m_i(v_r^2 + v_{\perp}^2) - e\phi$, $rv_{\perp}$) by the function

$$f_i = 3(m_i/E_0)^{3/2} n_{\text{edge}} \quad , \quad 0 < E < E_0 \quad , \quad 0 < rv_{\perp} < R v_{\perp 0}$$

$$= 0 \quad \text{else} \quad .$$

A Maxwellian gives similar results for $n(r)$, but the flat distribution is easier to work with analytically. The Vlasov equation for steady state is $f(E, rv_{\perp}) = \text{constant}$, so the density and distribution everywhere is determined by the potential and the value of f at the ion birth point. There are three regions. For a very small distance from the birth point, the potential is negligible, and the density is given by $n = n_{\text{edge}}$. In the bulk of the plasma, interior from the edge, where $E_0 \ll 2e\phi$, and $m_i v_{\perp 0}^2 (R/r)^2 < 2e\phi$, the density is given by integrating f over $v_{\perp} dv_{\perp} dv_r$ with the result (r_c is defined in Eq. (5) below)

$$n_B = \frac{3}{4} \left(\frac{E_0}{e\phi} \right)^{1/2} \left(\frac{R^2}{r^2} \right) n_{\text{edge}} \quad , \quad r_c < r < R \quad . \quad (4)$$

Assuming that the potential reaches its full value at a moderately large distance from $r = 0$, we see that a mean electron density throughout the bulk of the device will go as $1/r^2$. Moving inward, the $(R/r)^2$ factor eventually becomes substantial, so that the density becomes much larger than n_{edge} . Eventually, the radius is so small that $m_i v_{\perp 0}^2 (R/r)^2 > 2e\phi_{\text{max}}$. Inside that radius the density is changing fairly slowly, on the scale of the changing potential. This defines the radius of the dense core r_c as the radius at which $2e\phi = m_i v_{\perp 0}^2 (R/r)^2$, because outside that radius $n \sim 1/r^2$, while inside that radius n is nearly constant, found by integrating f_i over velocity,

$$n_c \sim \left(\frac{E_0}{e\phi_{\max}} \right)^{1/2} \left(\frac{R}{r_c} \right)^2 n_{\text{edge}} , \quad r < r_c ,$$

$$r_c \approx (m_i v_{\perp 0}^2 / 2e\phi_{\max})^{1/2} R . \quad (5)$$

The size of the core and the central density are seen to depend on the angular momentum at the outer turning points of the ion motion.

B. COLLISION FREQUENCY PROFILE

Using the ion distribution function discussed in the last section, where most of the ions have turning points $v_r = 0$ at the same radius $r = R$ at the periphery of the central potential $\phi(r)$, and taking the radial and azimuthal energy moments of the distribution in the three regions of the device, we find that the core and the edge are isotropic, while the bulk of the plasma is highly anisotropic, with $E_r \gg E_{\perp}$.

In the core, defined by Eq. (5), the ion density is relatively flat; using the above estimates for the ion energy and density, we write the ion-ion collision time in the core, τ_{iic} , as (energies are in eV)

$$\tau_{iic} = \frac{2 \times 10^7 (e\phi_{\max})^{3/2}}{n_c \ln \Lambda_c} , \quad (6)$$

where the Coulomb logarithm in the core is

$$\ln \Lambda_c \approx 7 + 2.3 \log_{10} \left[\frac{(e\phi_{\max})^{3/2} \text{ (eV)}}{(n_c/10^{14} \text{ cm}^{-3})^{1/2}} \right]$$

Equation (6) gives $\tau_{iic} \approx 1 \text{ s}$ and $30 \times 10^{-6} \text{ s}$, respectively, for the EXP and REACTOR parameters in Table I.

In the bulk, the ion density decreases radially outward basically as $1/r^2$, with an additional weak radial dependence from ϕ , which tends to make the density distribution flatter than $1/r^2$ beyond about the $R/2$ point of the device. The ion-ion collision time in the bulk, τ_{iib} , is

$$\tau_{iib} \approx \frac{4}{3} \left(\frac{e\phi_{\max}}{E_0} \right) \left(\frac{r}{R} \right)^2 \left[1 - \left(\frac{r}{R} \right)^p \right]^2 \left(\frac{\ln \Lambda_c}{\ln \Lambda_B} \right) \tau_{iic} ,$$

where

$$\ln \Lambda_B \approx 7 + 2.3 \log_{10} \left[\frac{\left\{ e\phi_{\max} \text{ (eV)} \left[1 - \left(\frac{r}{R} \right)^p \right] \right\}^{3/2}}{(n_B/10^{14} \text{ cm}^{-3})^{1/2}} \right]$$

The edge region is defined as that region in which the ion radial velocity slows down to a value of the order of it's birth speed, that is, $E_r = e\phi_{\max} [1 - (r_e/R)^p] \sim E_0$. The radial extent of the edge region, $\Delta r_e = R - r_e$, is then given by $\Delta r_e \approx (1/p)(E_0/e\phi_{\max})R$. The ion-ion collision time in the edge, τ_{iie} , is given by

$$\tau_{iie} \approx \left(\frac{E_0}{e\phi_{\max}} \right) \left(\frac{\ln \Lambda_c}{\ln \Lambda_e} \right) \tau_{iic} ,$$

where

$$\ln \Lambda_e \approx 7 + 2.3 \log_{10} \left[\frac{E_o^{3/2} \text{ (eV)}}{(n_e/10^{14} \text{ cm}^{-3})^{1/2}} \right] .$$

The collision time τ is not sufficient for comparing the effectiveness of collisions in the three spatial regions, because the time an ion spends in each region is drastically different. A better figure of merit is the ratio of the ion transit time t_{tr} through the region Δr to the collision time τ_{ii} in the region. This essentially gives the number of collisions, N_{coll} , that a test ion undergoes in a particular region of the device per transit.

The number of collisions in the core per pass is then

$$(N_{coll})_c \sim \frac{2R}{\tau_{iic}} \left(\frac{E_o}{e\phi_{max}} \right)^{1/2} \left(\frac{m_i}{2e\phi_{max}} \right)^{1/2} \approx 5 \times 10^{-8}, \text{ EXP}$$

$$\approx 3 \times 10^{-4}, \text{ REACTOR} .$$

Thus one collisional upscatter time corresponds to a large number of ion transits.

The ratio of the number of collisions in the edge region to the number of collisions in the core per transit is (take $p = 3$)

$$\frac{(N_{coll})_{edge}}{(N_{coll})_c} \sim \frac{1}{p} \left(\frac{e\phi_{max}}{E_o} \right) \left(\frac{\ln \Lambda_e}{\ln \Lambda_c} \right) \approx 4 \times 10^2, \text{ EXP}$$

$$\approx 2 \times 10^3, \text{ REACTOR} ,$$

which corresponds to a small fraction of an edge collision per transit for experimental parameters, but to approximately one collision per transit for reactor parameters.

As representative values for the bulk region, we take $r = R/2$. Then the ratio of the number of collisions in the bulk region to the number of collisions in the core per pass is approximately

$$\frac{(N_{\text{coll}})_B}{(N_{\text{coll}})_C} \sim \frac{2R}{r_c} \left(\frac{E_o}{e\phi_{\text{max}}} \right) \left(\frac{\ln \Lambda_B}{\ln \Lambda_C} \right) \approx 5 \times 10^{-2}, \text{ EXP}$$

$$\approx 2 \times 10^{-2}, \text{ REACTOR} .$$

From the above, we see that the number of collisions per pass is by far the highest at the edge region, where the low energy ion distribution is relatively isotropic. The number of collisions is lowest in the bulk of the device, where the radial ion energy is high, and where the energy distribution is highly anisotropic, with $E_r \gg E_{\perp}$.

III. CALCULATION OF THE ION DISTRIBUTION PRODUCED BY COLLISIONS

In the previous section we calculated the relative rates of collisions in the various spatial regions of a SCIF, and found that collisions in the edge regions dominate. More importantly, we found that the collision rate in the interior of the device is very small compared with the transit frequency, so that for many transits the ions return to their original turning points at $r = R$. That is, the definition of "edge region" is preserved for thousands of interior plasma collision times. In this section we give a Fokker-Planck-like calculation of temperature relaxation of the edge distribution due to ion-ion collisions throughout the device. Since the collision frequencies in the bulk and core are much lower than the transit frequency, the edge distribution determines the evolution of the ion density and energy structure in all three regions. The edge distribution evolves (a) due to collisions at the edge, (b) due to the influx of energy-scattered ions (due to collisions in the core) and (c) due to energy-deflected ions (due to collisions in the bulk) coming into the edge region in every transit.

The dominant effect of edge collisions is to relax any temperature anisotropy which develops in this region. The source of ion anisotropy in the edge is the influx of excess $v_{\perp}$ into the edge region due to the conversion of radial to azimuthal energy via ion-ion collisions in the highly anisotropic bulk plasma. The goal of the present calculation is to quantify the extent to which isotropizing collisions in the low ion energy edge can remove or reduce this increase of $v_{\perp}$. The anisotropy at the edge is the primary effect which spreads out the distance of closest approach to $r = 0$ (this is obvious from energy conservation), which in turn could lead to an isotropic interior.

A. ION DISTRIBUTIONS IN THE THREE REGIONS

In Sections III.A and III.B, we assume that the effect of the competition between edge collisions and bulk collisions will be that the edge region will ultimately tend toward a bi-Maxwellian, characterized by T_r and $T_\perp$, and calculate the steady state difference of T_r and $T_\perp$ in the edge region.

Thus for the edge region, we take

$$f_{i,e} = K e^{-\frac{m_i v_r^2}{2T_{r0}}} e^{-\frac{m_i v_\perp^2}{2T_{\perp 0}}} , \quad (7)$$

with K being a normalization constant such that

$$n_{\text{edge}} = \int d^3 v f_{i,e} .$$

Since the ion collision frequency in the interior is low, the ions maintain a dominant set of turning points at $r = R$, so that the interior distribution continues to be the evolution of an edge Maxwellian under the influence of a strong central potential $\phi(r)$. Therefore, we take the ion distribution in the bulk region to be the sum of two radially counter-drifting bi-Maxwellians. The radial drift, $v_D = \sqrt{2e\phi(r)/m_i}$, is due to the electrostatic field. The azimuthal temperature $T_\perp$ increases with decreasing radius, reflecting conservation of angular momentum, $T_\perp(r) = T_{\perp 0} (R/r)^2$. Thus in the bulk

$$f_{i,B} = K e^{-\frac{m_i (v_r \pm v_D)^2}{2T_{r0}}} e^{-\frac{m_i v_\perp^2}{2T_\perp(r)}} . \quad (8)$$

We note that the bulk distribution reduces to the edge distribution at $r = R$, since $\phi(R) = 0$.

We assume that the core is defined as that region in which the azimuthal and radial ion energies are the same, that is, $T_r = T_\perp = e\phi_{\max}$. Thus we model the core distribution as an isotropic Maxwellian at the temperature $T_c = e\phi_{\max}$, since ϕ is virtually constant over the small core region,

$$f_{i,c} = K e^{-\frac{m_i v^2}{2T_c}} \quad (9)$$

For simplicity, we use the density and energy profiles in the three regions of the device derived in Section II, which can be shown to be similar to those produced by a bi-Maxwellian.

B. CALCULATION OF THE STEADY-STATE ION DISTRIBUTION

We model the evolution of $T_\perp$ in the edge region by:

$$\frac{dT_{\perp 0}}{dt} = \left. \frac{dT_{\perp 0}}{dt} \right|_B + \left. \frac{dT_{\perp 0}}{dt} \right|_{ISO} \quad (10)$$

Here, $dT_{\perp 0}/dt|_B$ is the source term for the additional $v_\perp$ coming in to the edge region on each ion transit, due to the conversion of radial to azimuthal energy by collisions in the bulk, and $dT_{\perp 0}/dt|_{ISO}$ is the isotropization rate due to collisions in the edge region. There is a similar equation for T_{r0} . However, we can use conservation of energy to write

$$\frac{dT_{\perp 0}}{dt} = - \frac{d}{dt} \frac{T_{r0}}{2} , \quad (11)$$

and to obtain an equation for the evolution of the edge temperature difference,

$$\Delta T = T_{\perp 0} - T_{r0},$$

$$\frac{d}{dt} \Delta T = 3 \left. \frac{dT_{\perp 0}}{dt} \right|_B + 3 \left. \frac{dT_{\perp 0}}{dt} \right|_{ISO} . \quad (12)$$

There is a similar driver in which T_{r0} is changed by energy scattering in the core, with $T_{\perp 0}$ in the edge changed because energy conservation slightly alters the turning point. We emphasize here the $T_{\perp}$ scattering because $T_{\perp}$ acquired during a transit is carried out to the turning point decreased only by r/R (of order 1/2), while v_r always goes to 0 at the turning point because of the structure of $\phi(r)$.

In the following calculation, we assume that the edge temperature difference is small, $\Delta T = T_{\perp 0} - T_{r0} \ll T$, because the conversion of radial to azimuthal velocity of an ion in each transit through the bulk is small compared with the rate of isotropizing edge collisions, from the estimates in Section II.

The isotropization time for a bi-Maxwellian distribution, in the limit of small temperature differences, is given by:⁵

$$\left. \frac{dT_{\perp 0}}{dt} \right|_{ISO} = - \frac{\Delta T}{\tau_e} , \quad (13)$$

where $\tau_e = (15/4) (m_i^{1/2} T_{\text{edge}}^{3/2} / \sqrt{\pi} e^4 n_{\text{edge}} \ln \Lambda_e)$, and $T_{\text{edge}} = T_{r0}$ to this order of approximation.

To calculate $(dT_{\perp 0}/dt)$ due to the bulk region, where the ion energy distribution is highly anisotropic, we write the Fokker-Planck collision term using the Landau form for the collision operator. We take the energy moment of the kinetic equation

$$\frac{\partial f_i}{\partial t} = - \frac{\partial j_\alpha}{\partial v_\alpha} \quad , \quad (14)$$

where j is the three-dimensional particle flux density in velocity space, representing particle collisions:

$$j_\alpha = a_\alpha f_i - d_{\alpha\beta} \frac{\partial f_i}{\partial v_\beta} \quad , \quad (15)$$

where

$$a_\alpha = \frac{1}{\tau} \left[\langle \Delta v_\alpha \rangle - \frac{1}{2} \frac{\partial}{\partial v_\beta} \langle \Delta v_\alpha \Delta v_\beta \rangle \right] \quad ,$$

$$d_{\alpha\beta} = \frac{1}{2\tau} \langle \Delta v_\alpha \Delta v_\beta \rangle \quad ,$$

as given for example in Reference 6.

Using conservation of energy, we obtain for the increase of $T_{\perp 0}$ due to the bulk,⁷

$$\frac{dT_{\perp 0}}{dt} = - \frac{d}{dt} \frac{T_{ro}}{2} = - \frac{1}{n_B} \int \frac{m_i v_r^2}{2} \frac{\partial f_i}{\partial t} d^3 v = - \frac{m_i}{n_B} \int v_r j_r d^3 v \quad . \quad (16)$$

We compute j_r by writing the flux in this case in Landau form⁷

$$j_r = \frac{1n\Lambda}{8\pi} \left(\frac{4\pi e^2}{m_i} \right)^2 \int v_{rj} \left(f_i \frac{\partial f'_i}{\partial v'_j} - f'_i \frac{\partial f_i}{\partial v_j} \right) d^3 \underline{v}' , \quad (17)$$

with

$$v_{rj} = \frac{\delta_{rj}}{u} - \frac{u_r u_j}{u^3} .$$

Here the primed, unprimed quantities refer to the field, test particles, respectively, and $u = v - v'$. We assume that the field and test particles counter-drift radially:

$$f_i = K e^{-\frac{m_i (v_r - v_D)^2}{2T_{ro}}} e^{-\frac{m_i v_\perp^2}{2T_\perp}} , \quad (18)$$

$$f'_i = K e^{-\frac{m_i (v'_r + v_D)^2}{2T_{ro}}} e^{-\frac{m_i v'^2_\perp}{2T_\perp}} . \quad (19)$$

Then

$$\frac{\partial f_i}{\partial v_r} = -\frac{m_i}{T_{ro}} (v_r - v_D) f_i ,$$

$$\frac{\partial f'_i}{\partial v'_r} = -\frac{m_i}{T_{ro}} (v'_r + v_D) f'_i ,$$

and we have

$$\frac{dT_{\perp}}{dt} = - A \int_{-\infty}^{\infty} dv_r dv'_r \int_0^{\infty} dv_{\perp}^2 dv'_{\perp}^2 v_r$$

$$\cdot \left[\frac{u_{\perp}^2}{u^3} \frac{m_i}{T_{ro}} (u_r - 2v_D) - \frac{u_r u_{\perp}^2}{u^3} \frac{m_i}{T_{\perp}} \right] f_i f'_i \quad ,$$

where A is given by

$$A = \frac{m_i}{n_B} \pi^2 \frac{\ln \Lambda}{8\pi} \left(\frac{4\pi e^2}{m_i} \right)^2 .$$

To do this integral, we make the approximation that $u = |\underline{u}| \approx |v_r - v'_r| - 2 v_D$, so that u can be taken out of the integral. We rewrite the integration variables as

$$\bar{v}_r = v_r - v_D \quad ,$$

$$\bar{v}'_r = v'_r + v_D \quad ,$$

to obtain

$$\frac{dT_{\perp}}{dt} = - AK^2 \int \frac{(v_r + v_D)}{(2v_D)^3} \left[\frac{m_i u_{\perp}^2}{T_{ro}} (v_r - 2v_D) - \frac{m_i u_{\perp}^2}{T_{\perp}} u_r \right] \\ \cdot e^{-\frac{m_i \bar{v}_r^2}{2T_{ro}}} e^{-\frac{m_i \bar{v}_r'^2}{2T_{ro}}} e^{-\frac{m_i v_{\perp}^2}{2T_{\perp}}} e^{-\frac{m_i v_{\perp}'^2}{2T_{\perp}}} d\bar{v}_r d\bar{v}_r' dv_{\perp}^2 dv_{\perp}'^2. \quad (20)$$

Performing the $\bar{v}_r, \bar{v}_r'$ integrations in Eq. (20) gives

$$\frac{dT_{\perp}}{dt} = - \frac{AK^2}{(2v_D)^3} \int_0^{\infty} u_{\perp}^2 \left(\frac{2\pi T_{ro}}{m_i} \right) \left[1 - \frac{T_{ro}}{T_{\perp}} - \frac{2m_i v_D^2}{T_{\perp}} \right] \\ \cdot e^{-\frac{m_i v_{\perp}^2}{2T_{\perp}}} e^{-\frac{m_i v_{\perp}'^2}{2T_{\perp}}} dv_{\perp}^2 dv_{\perp}'^2.$$

The $v_{\perp}, v_{\perp}'$ integrations can then be done to obtain

$$\frac{dT_{\perp}}{dt} = - \frac{AK^2}{(2v_D)^3} \left(\frac{2\pi T_{ro}}{m_i} \right) \left(\frac{2T_{\perp}}{m_i} \right)^3 \left[1 - \frac{T_{ro}}{T_{\perp}} - \frac{2m_i v_D^2}{T_{\perp}} \right] \left(2 - \frac{\pi}{2} \right). \quad (21)$$

We rewrite this as

$$\frac{dT_{\perp}}{dt} = \frac{(2m_i v_D^2 + T_{ro} - T_{\perp})}{\tau_B}, \quad (22)$$

where the isotropization time τ_B is given by

$$\tau_B = \frac{m_i^2 (2v_D)^3}{2\pi e^4 (\ln \Lambda_B) n_B (4 - \pi)} \quad (23)$$

In order to use this as a source term for $(dT/dt)_B$ in the edge region, it is necessary to take into account the fact that an ion in its transit through the device spends more time in the bulk than in the edge. Thus, we multiply Eq. (22) by the ratio of the transit time in the bulk to the transit time at the edge (t_B/t_e). This is equivalent to writing Eq. (22) in units of the edge transit time. We also need to take the following effect into account, which is a consequence of conservation of angular momentum. An ion which gains excess $(1/2)m_i v_{\perp}^2$ in the bulk at radius r , will bring an excess azimuthal energy into the edge of magnitude $\approx (1/2)m_i v_{\perp}^2 (r/R)^2$. We thus have, for use in Eq. (10),

$$\left. \frac{dT}{dt} \right|_B \approx \frac{2m_i v_D^2}{\tau_B} \left(\frac{r}{R} \right)^2 \frac{t_B}{t_e} \quad (24)$$

Equation (12) then becomes

$$\frac{d}{dt} \Delta T \approx \frac{6m_i v_D^2}{\tau_B} \left(\frac{r}{R} \right)^2 \frac{t_B}{t_e} - \frac{3\Delta T}{\tau_e} \quad (25)$$

We use the estimates for the transit time in the bulk and the edge regions from Section II. The ratio of the transit time in the bulk to the transit time in the edge is of the order of $t_B/t_e \sim p \sqrt{e\phi_{\max}/T_{ro}}$, while the ratio of the

isotropization time in the bulk, τ_B , to the isotropization time in the edge, τ_e , is

$$\tau_B/\tau_e \sim \left(\frac{e\phi_{\max}}{T_{ro}} \right)^{3/2} \frac{n_{\text{edge}}}{n_B} \quad (26)$$

Thus, as long as $n_{\text{edge}}/n_B > p$, with p defined in Eq. (2) typically of order 2-3, we expect that the temperature difference in the edge region would remain small, $\Delta T < T_{ro}$. This condition depends on the form of the density profile in the device, which is determined from the edge ion distribution and conservation of energy and angular momentum. Using the flattop ion distribution discussed in Section II, we have that $n_{\text{edge}}/n_B \sim (e\phi_{\max}/E_0)^{1/2} \gg 1$.

The equilibrium solution of Eq. (25) yields $\Delta T = (T_{\perp} - T_r)_{\text{edge}} \sim T_{ro} (T_{ro}/e\phi_{\max})^{1/2} \ll T_{ro}$. Thus the quasisteady solution to this nonlocal collision relaxation process is an ion distribution which is nearly Maxwellian at the edge. Since the ions have turning points in the same vicinity as their birth point, the interior distribution continues to evolve from the edge distribution, conserving energy and angular momentum, so that the non-Maxwellian distributions Eqs. (19)-(20) with $T_{\perp} = T_{\perp 0} (R/r)^2$ continue to describe the bulk plasma. The interior, like the edge, is driven toward isotropy by local collisions, but driven toward anisotropy by influx of ions from the Maxwellian edge. It is because the edge collision frequency per transit is many orders of magnitude larger than the bulk rate that the anisotropic influx so dominates the interior.

IV. APPLICATION TO ION FOCUS LOSS RATES

For the experimental parameters in Table I, the rate of collisions per transit is so small as to be negligible. For reactor parameters, however, the collision rates must be compared to the fusion energy multiplication rate. Therefore in this section we consider in detail the loss produced by collisions for reactor parameters in an ion focus where the potential is produced by magnetically confined electrons.

The previous results show that edge collisions maintain a non-Maxwellian interior for a time larger than the small angle scattering time which in most cases produces a local Maxwellian. Applying this result to ion loss rates, we note that there are two effects. First, large angle scattering will prevent ions from returning to the vicinity of their birth location, thus removing the healing effect of edge collisions which assumes a fixed range of turning points. Second, the slightly anisotropic edge distribution found in Section III will produce some degradation in the core on the time scale of the small angle scattering time in the core.

A. ION LOSS BY LARGE ANGLE SCATTERING

The frequency of 90° scattering is obtained by simply setting $\log \Lambda = 1$ in the collision rates in Section III. To estimate the loss rate we must estimate how large a change in velocity Δv is required to cause an ion to be lost to the core. In this section we consider in detail such losses.

1. Ion Loss Due to Radial Upscatter in the Core

Upscatter in radial velocity in the core by collisions can affect ion convergence in the device. Consider a collision which causes one ion to gain energy and another ion to lose energy. The ion which gains energy in the core can make a larger radial excursion to a radius beyond it's birth radius, with the possibility that the magnetic fields at the new turning point will be sufficient to deflect the ion in $v_{\perp}$, leading to a degradation of the focus.

We estimate the value of δv which could lead to ion "loss," in the sense of a broadened core radius r_c , due to increased deflection by the higher B-field as the upscattered ion makes a larger radial excursion. To do this, we estimate the new radial location r'_0 at which the upscattered ion's radial energy decreases to its birth energy. Neglecting scattering, we assume the ion radial energy profile is, generalizing Eq. (3) to $r_0 \neq R$,

$$E_r = E_0 + e\phi_{\max} \left[\left(\frac{r_0}{R} \right)^p - \left(\frac{r}{R} \right)^p \right] , \quad (27)$$

Due to upscatter in the core, the ion receives a kick in energy ΔE_r ,

$$\Delta E_r \sim e\phi_{\max} \left(\frac{r_0}{R} \right)^p f(2 + f) . \quad (28)$$

where $f = \delta v_r / v_r$.

Then $E_r + \Delta E_r = E_0$ when

$$\frac{r}{R} = \frac{r'_0}{R} \sim (1 + f)^{2/p} \left(\frac{r_0}{R} \right) . \quad (29)$$

If this new turning point r'_0 takes the ion into a region of high magnetic field, $B = B_0(r/R)^m$, the perpendicular deflection of the ion by this magnetic field, $\Delta v_\perp$, changes the core radius by

$$\frac{r_c}{R} = \frac{\Delta v_\perp}{v_\perp} \quad (30)$$

The perpendicular deflection of the ion by the magnetic field can be estimated from straightforward consideration of ion orbits in orthonormal magnetic and electric fields. Assuming that the ions were originally born at a location r_0 at which magnetic deflection would give a contribution to $\Delta v_{\perp 0}$ (and to r_c) of the same order of magnitude as the $\Delta v_\perp$ arising from the thermal spread $T_{\perp 0}$ in the ion source, it is easy to show that the spread Δv due to a larger radial excursion to r'_0 , at which the magnetic field is larger ($B \sim r^m$), is

$$\frac{\Delta v(r'_0)}{\Delta v(r_0)} \sim \frac{4m}{p} \frac{\delta v_r}{v_r} \approx \frac{\Delta r_c}{r_c} \quad (31)$$

Thus ion loss, in the sense that $\Delta r_c/r_c \sim 1$, requires $\delta v_r/v_r$ comparable to unity, which means a single large angle scattering. From the results in Section III, and the reactor parameters in Table I, this gives (setting $\log \Lambda = 1$)

$$\tau_{\text{loss}} \sim 85 \text{ ms} \quad (32)$$

which is large compared to fusion reaction times of order 10 ms, from References 1 and 2.

2. Ion Loss Due to Perpendicular Deflection in the Bulk

Perpendicular deflection due to scattering in the bulk of the device could lead to isotropization of the anisotropic ion distribution. From conservation of angular momentum (without scattering), $rv_{\perp}(r) = \text{constant}$. An increase in the azimuthal ion velocity $\Delta v_{\perp}$ at its birth point is related to an increase in the core convergence radius, Δr_c , by

$$\frac{\Delta v_{\perp}(r_0)}{v_{\perp}(r_0)} = \frac{\Delta r_c}{r_c} \quad (33)$$

Since $v_r \gg v_{\perp}$ through much of the ion orbit, even a small decrease in $\Delta v_r/v_r$ could produce enough $\Delta v_{\perp}/v_{\perp} \sim (\Delta v_r/v_r)(v_r/v_{\perp})$ to degrade the ion focus. However, a small Δv_r would still result in a turning point at $r \approx r_0$, where collisions would restore f_i as calculated in Section III. In order to change r_0 significantly, $\delta v_r/v_r \approx 1$ is required, even for collisions in the bulk. The result must take into account the lower density n_B/n_c in the bulk, and the extended dwell time R/r_c in the bulk. This gives an estimate of the loss time due to collisions in the bulk of

$$\tau_{\text{loss},B} \approx \tau_{\text{loss},\text{core}}(n_c/n_B)(r_c/R) \approx \tau_{\text{loss},\text{core}}(e\phi/E_0)^{1/2} \approx 10 \text{ s} \quad , \quad (34)$$

for reactor parameters. This is negligible compared with τ_{fusion} .

B. Loss Due to the Edge Anisotropy

In Section III we calculated the anisotropy which develops at the edge due to small angle scattering in the rest of the device. In this section we estimate the degradation of the core which results from this anisotropy.

From Section III we showed that the increase in $T_{\perp 0}$ at $r = R$ was of order $E_0(E_0/e\phi)^{1/2}$, and that the increase in core size was $\Delta r_c/r_c \approx \Delta v_{\perp}/v_{\perp}$. Therefore it follows that

$$\frac{\Delta r_c}{r_c} \approx \left(\frac{E_0}{e\phi}\right)^{1/4} \leq 10^{-1} \quad , \quad (35)$$

with this small Δr_c taking place on a time scale of a few ion transits. We conclude that small angle scattering, when the Maxwellization of the plasma near the ion turning points is taken into account, is a very small contribution to core degradation, even though this is the fastest collisional effect.

APPENDIX A REFERENCES

1. R. W. Bussard, Fusion Tech. 19, 273 (1991).
2. N. A. Krall, Fusion Tech. (1992, in press).
5. D. V. Sivukhin, in Reviews of Plasma Physics, Vol. 4, p. 93 (ed. by M. A. Leontovich), Consultants Bureau, New York (1966).
6. V. I. Kogan, in Plasma Physics and the Problem of Controlled Thermonuclear Reactions, Vol. 1, Pergamon Press, New York (1959).
7. B. A. Trubnikov, in Reviews of Plasma Physics, Vol. 1, p. 105, Consultants Bureau, New York (1965).

Table I. Representative ion focus parameters.

	EXP	REACTOR
$e\phi_{\max}$ (depth of the potential well)	10 keV	100 keV
E_0 (ion birth energy)	5 eV	5 eV
n_c (ion core density)	10^{12} cm^{-3}	10^{18} cm^{-3}
R (radius of device)	1 m	2 m
B (magnetic field at R , $m = 3$)	0.2 T	1 T

APPENDIX B.

CONFINEMENT OF A PLASMA BY A CUSP MAGNETIC FIELD

I. INTRODUCTION

The use of magnetic point cusps to confine plasma has been a widely explored topic in fusion physics. It remains a relevant topic because of the number of devices which have a region in which the magnetic topography is cusp-like. These include magnetic divertors, plasma guns, plasma guide fields, and regions such as the x-point in the field reversed configuration. Despite the wide range of existing studies and applications, we feel that there is no definitive calculation of cusp confinement, either theoretically or experimentally, in the parameter range where plasma density and temperature are sufficiently high that the magnetic field is excluded by diamagnetic effects from most of the volume. The reason for this gap in confinement lore is that high- β cusp confinement is a collective process,^{1,2} with plasma currents acting to exclude the B-field. Collective processes are traditionally difficult to pin down; further, when cusp confinement was an active area of research, tools for numerical simulation of this multi-dimensional, high density system were not available. Experimentally, the situation was not much better, with plasma formation techniques and available diagnostics leading to a clouded and often contradictory set of cusp confinement measurements. There are several reviews which document this situation.^{3,4}

Because of the importance of cusp confinement to a number of applications, we have embarked on a program to measure confinement in this geometry using analytic estimates and numerical simulation. Because of the expense in computer time involved in simulating an inherently 3-dimensional, high density, large volume, and non-neutral plasma, we are approaching the problem in well defined stages, and choosing plasma parameters which preserve the essence of the cusp problem while minimizing the numerical difficulties.

From a simplified point of view, the question of cusp confinement involves being able to distinguish between several competing confinement ideas. These include the following:

- o Single particle confinement: Even in the absence of diamagnetic exclusion of the B-field from the plasma, particles are confined by mirror reflection as the cusp field increases near the outside of the device. Particles are "scattered" into the loss cone by passing through the low field center of the cusp, where their magnetic moment is no longer invariant. The mirror ratio is the ratio of the maximum cusp field to the field at which the particles again become adiabatic.
- o Ergodic sampling of a field distribution with a few well defined escape holes. When the plasma excludes the B-field, it is possible that a geometry will develop in which there is a virtually field-free interior region, and a high field exterior region, separated by a sharp interface, of the order of an electron gyroradius. In that case, the particles free stream across the interior, reflecting specularly off the interface, until by chance they strike a small

area near the corner of the cusp, Figure 1, whereupon they are lost. This is the classic picture of cusp confinement. The question is what is the size of the loss hole? Possibilities include the following:

- a. A hole of area πr_e^2 , where r_e is the electron gyroradius. This was suggested by Berkowitz.¹
- b. A hole of area $2\pi R r_e$, where R is the characteristic size of the device. This was calculated by Taylor² as an upper limit to the loss rate in a spindle cusp, where there was symmetry axis which led to a line cusp of length $2\pi R$.
- c. A hole of area πr_i^2 , where r_i is the ion gyroradius. This scaling arises from considerations of the stability of a very thin plasma-field interface.⁵
- d. A hybrid area, $\pi r_i r_e$, estimated from at least 2 experiments.^{6,7}

In our first effort, we are simulating confinement in a spindle cusp. This allowed a two-dimensional treatment of the problem, simplifying the computation. We study a neutral plasma, with $T = 120$ keV and $B_{\max} = 3$ kG. This also simplifies the computation by allowing high β studies at acceptably low densities. Essential to the study of course is that r_e and r_i be small compared to the system size R . Therefore we chose $r = 25$ cm. The ion mass is unrealistically taken to be a small multiple of the electron mass m_e , again for numerical convenience, while still maintaining the dimensionless variables of the system in a range appropriate to studying cusp confinement.

In this report we document the predictions of the various confinement ideas listed above for the parameters of the simulation. We also make a stab at estimating the effects of ambipolarity on the confinement. Later reports will present confinement times found by the simulation. At present, at least the low β parameter range has been simulated, agreeing well with the discussion in Section II.

II. LOSS MODELS FOR THE SPINDLE CUSP

In this section, we summarize calculations we and others have carried out on the various loss models for cusp confinement.¹⁻¹⁰

A. MIRROR LOSS, LOW β

At low density, or in the absence of diamagnetic effects, the particles are confined by reflection from the increasing B-field in the cusp. The mirror ratio is determined by the magnetic field at the radius at which the particles become adiabatic, compared with the magnetic field at the surface of the cusp. The 'collision time' for scattering into the loss cone is, in a cusp, the transit time of the particles across the cusp. This is because in a transit, the particles pass through a low field interior region, and lose their memory of their adiabatic invariant. For a particle of mass m and energy E , the confinement time has been calculated to be⁸

$$\tau_M = 3 \times 10^{-5} B_G^{3/4} R_{cm}^{7/4} m_g^{1/8} / E_{eV}^{7/8} \text{ s} \quad , \quad (1)$$

where the units are indicated as subscripts. In Eq. (1) a factor $R/(2E/m)^{1/2}$ is the transit time across the cusp. The rest is the effect of particle reflection from or transmission through the cusp field.

B. HIGH β CUSP LOSS

At higher density the vacuum field is so distorted that the analysis of mirror loss based on the vacuum fields⁸⁻⁹ is irrelevant. The loss can be envisioned as a particle ergodically sampling the surface which separates the $B = 0$ region from the high B exterior. The surface has a discrete set of "holes" of area πr_0^2 , and the sampling rate is v_T/R , with v_T the thermal speed. The size of the hole, r_0 , is questionable. It has been suggested¹ that $r_0 = r_e$, the electron gyroradius, assuming that the sheath width is r_e . The loss is determined by the relative area of the magnetic surface and the area of the loss holes. If the loss area is in fact πr_e^2 , we have

$$\tau_C = \frac{5 \times 10^{-23} R_{cm}^3 B_G^2}{E_{eV}^{3/2} m_g^{1/2}}, \quad (2)$$

where $R/(2E/m)^{1/2}$ is the transit time contribution to the loss.

C. SPINDLE CUSP LOSS

A direct calculation of the high β regime loss area has been carried out by Taylor for the special case of confinement in a spindle cusp, Figure 2. Using a model which retains two invariants of the particle motion he obtains $r_0^2 = r_e R$ as the size of the loss hole. These are (at least) two reasons why this calculation can fail in a general geometry. First, the spindle cusp has an axis of symmetry leading to a line cusp of length $2\pi R$. This is specific to the

spindle. Second, Taylor included only two invariants in his calculation. Other invariants would extend the confinement time. Using $\pi r_e R$ as the loss area gives

$$\tau_{\text{JBT}} = 5 \times 10^{-9} R_{\text{cm}}^2 B_G / E_{\text{eV}} \quad . \quad (3)$$

III. GENERALIZATION FOR $\beta = 8\pi nE/B^2 \ll 1$

Even when $8\pi nT \ll B_{\max}^2$, there can still be a $\beta = 1$ interface well inside the plasma. In these cases, R in Eqs. (2)-(5) should be interpreted as r_β , the location of the $\beta = 1$ surface, and B as the magnetic field at that radius. For $B = B_0(r/R)^m$ we have¹⁰

$$r_\beta = (8\pi nE/B_0^2)^{1/2m} R, \quad (4)$$

$$B(r_\beta) = (8\pi nE)^{1/2}. \quad (5)$$

The point is that high β confinement might dominate even when the $\beta = 1$ surface is fairly far inside the plasma. We use Eqs. (4)-(5) to generalize Eqs. (2)-(3) for $8\pi nT \ll B_0^2$, with the result

$$\tau_C = \frac{5 \times 10^{-23} R^3 B_0^2 \left(8\pi nT/B_0^2\right)^{1 + \frac{3}{2m}}}{E_e^{3/2} m^{1/2}}, \quad \frac{8\pi nT}{B_0^2} < 1, \quad (6)$$

$$\tau_{JBT} = 5 \times 10^{-9} \frac{R^2 B}{E_e} \left(\frac{8\pi nT}{B_0^2}\right)^{\frac{1}{2} + \frac{1}{m}}, \quad \frac{8\pi nT}{B_0^2} < 1. \quad (7)$$

IV. GENERALIZATION FOR AMBIPOLARITY

After a brief transient, electrons and ions must leave at the same rate, and an electric field will develop during the initial transient to ensure that this is the case. If we assume that the electrons and ions see the same size loss hole, electrons will be lost first, because they sample the $\beta = 1$ surface at a faster rate, i.e., during the transient phase

$$\tau_C^{i,e} \sim \frac{R \sqrt{M_{i,e}}}{\sqrt{E}} K ,$$

$$\tau_{JBT}^{i,e} \sim \frac{R \sqrt{M_{i,e}}}{\sqrt{E}} .$$

The effect will be that an electric potential ϕ_A develops to retard the electrons in the vicinity of the loss holes; a potential of order $e\phi_A = E_0$ can reduce the electron flow to include only those with $1/2 mv^2 > E_0 - e\phi_A$. This will have only a slight effect on the ions, since their transit time is essentially unchanged, as is the escape hole they see. This means that the ambipolar loss rates for C(usp) and JBT confinement are determined by the electron gyroradius and the ion transit time,

$$\tau_C = \frac{5 \times 10^{-23} R^3 B_0^2}{E^{3/2} m_e^{1/2}} \left(\frac{M}{m}\right)^{1/2} \left[\frac{8\pi n E}{B_0^2}\right]^{1 + \frac{3}{2m}} , \quad \frac{8\pi n E}{B_0^2} < 1 , \quad (8)$$

$$\tau_{JBT} = \frac{5 \times 10^{-9} R B_0}{E} \left(\frac{M}{m}\right)^{\frac{1}{2}} \left[\left(\frac{8\pi n E}{B_0^2}\right)^{\frac{1}{2}} + \frac{1}{m} \right], \quad \frac{8\pi n E}{B_0^2} < 1 \quad (9)$$

Ambipolar mirror confinement is somewhat different, in that instead of a loss hole there is a loss cone, different for each species since the "mirror ratio" depends on the gyroradius, which determines the r, B at which the particles become adiabatic. The ambipolar mirror loss is determined by equating $m_e^{1/8}/(E - \Delta E)^{7/8} = m_i^{1/8}/(E + \Delta E)^{7/8}$, which again makes $\tau^e = \tau^i$, and gives

$$\tau_M = 3 \times 10^{-5} \frac{B_G^{3/4} R_{c,0}^{7/4} m_e^{1/8}}{E_{eV}^{7/8}} \left[\frac{1}{2} + \frac{1}{2} \left(\frac{M}{m}\right)^{1/7} \right]^{7/8} \quad (10)$$

V. CONCLUSIONS

The results are given in Eqs. (8)-(10), namely (assuming $8\pi nE < B_0^2$)

$$\tau_M = \left[1 \times 10^{-8} B_G^{3/4} R_{cm}^{7/4} / E_{eV}^{7/8} \right] \left[\frac{1}{2} + \frac{1}{2} \left(\frac{M}{m} \right)^{1/7} \right]^{7/8} s \quad (11)$$

$$\tau_C = \left[1.7 \times 10^{-9} B_G^2 R_{cm}^3 / E_{eV}^{3/2} \right] \left[8\pi n_{cm^{-3}} E / B_0^2 \right]^{1 + \frac{3}{2m}} \left(\frac{M}{m} \right)^{1/2} s \quad (12)$$

$$\tau_{JBT} = \left[5 \times 10^{-9} B_G R_{cm}^2 / E_{eV} \right] \left[8\pi n_{cm^{-3}} E / B_0^2 \right]^{\frac{1}{2} + \frac{1}{m}} \left(\frac{M}{m} \right)^{1/2} s \quad (13)$$

where M/m is the ion/electron mass ratio and B is the field in the cusp, with $B \sim B_0(r/R)^m$. For the parameters presently used in the simulation, $E = 120 \times 10^3$ eV, $B = 3 \times 10^3$ G, $R = 25$ cm (we take $R = 18$ cm in evaluating Eqs. (11)-(13) because the B-field is not spherical. This is arbitrary, but $R < 25$ cm is certainly required), $M/m = 5, 50$, we get the results listed in Table I.

One question, among many, is at what $\beta = 8\pi nE/B_0^2$ the magnetic field surfaces are so distorted that cusp confinement itself is lost. A density of $n = 5 \times 10^{11} \text{ cm}^{-3}$ corresponds to $\beta = 0.3$, which seems a prudent place to stop believing the formulas. With that caveat, Table I shows that the simulation can easily show $M \rightarrow C$, but to see $M \rightarrow JBT$ requires large M/m and careful analysis. Equations (11)-(13) show how to scale future simulations to make $M \rightarrow JBT$ stand out, if it happens at all.

APPENDIX B REFERENCES

1. J. Berkowitz, K. O. Friedrichs, H. Goertzel, H. Grad, J. Kileen, and E. Rubin, Proceedings of the 2nd Intl. Conf. on Peaceful Use of Atomic Energy, Geneva, 31, 171 (1958).
2. J. B. Taylor (private communication), Culham Report CLM-R58 (1966).
3. I. Spalding in Advances in Plasma Physics, eds. A. Simon and W. B. Thompson (Wiley, New York, 1971), Vol. 4, p. 79.
4. M. G. Haines, Nucl. Fusion 17, 811 (1977).
5. S. Hamasaki, R. C. Davidson, N. A. Krall, and P. C. Liewer, Nucl. Fusion 15, 27 (1975).
6. A. Kitsunezaki, M. Tanimoto, and T. Sakiguchi, Phys. Fluids 17, 1895 (1974).
7. N. Hershkowitz, K. N. Leung, and T. Romesser, Phys. Rev. Lett. 35, 277 (1975).
8. R. W. Bussard and N. A. Krall (private communication), Krall Associates Report KA-90-54; EMC2 Report EMC2-0191-02 (1991).
9. R. W. Bussard (private communication), EMC2 Report EMC2-0391-02.
10. N. A. Krall (private communication), Krall Associates Report KA-91-11.

Table I. Predicted confinement time in a 25 cm sphere with a cusp field of 3 kG and plasma energy of 120 kV per particle obtained by adding τ_M to either a C or JBT model for cusp loss. For a loss area of $\pi r_i r_e$, multiply τ_C by $\sqrt{m/M}$; for a loss area of πr_i^2 , multiply τ_C by m/M ; in all cases, however, $\tau \geq \tau_M$, since τ_M is not affected by the sheath size.

$n(\text{cm}^3)$	$M/m = 5; \tau_M = 35 \text{ ns}$		$M/m = 50; \tau_M = 50 \text{ ns}$	
	$\tau_C(\text{ns})$	$\tau_{\text{JBT}}(\text{ns})$	τ_C	τ_{JBT}
2.5×10^{10}	42	35	70	55
5×10^{10}	50	35	100	62
7.5×10^{10}	60	40	135	68
10^{11}	85	42	200	75
2.5×10^{11}	240	50	710	100
5×10^{11}	630	60	1400	125

Figure 1. Geometry of a set of point cusps.

Figure 2. Geometry of a spindle cusp.

APPENDIX C.
CONFINEMENT AT THE SEPARATRIX
IN A FIELD REVERSED CONFIGURATION

I. INTRODUCTION

The field reversed configuration (FRC)^{1,2} includes a region of closed field lines bounded by a region of open field lines (Fig. 1). Transport of particles and energy from the device includes a continuity condition, which says that the essentially axial loss of plasma along magnetic field lines must be matched by the essentially radial plasma flow across magnetic field lines. Early attempts at modelling FRC loss rates^{3,4} assumed that the plasma density would drop to nearly zero in a distance of the order of the ion gyroradius outside the separatrix, because of the very rapid loss along the open field lines. It was observed however that this assumption was not in agreement with the data, which showed a substantially broader sheath.⁵

A closer examination of the magnetic geometry (Fig. 2) shows that the field lines at the separatrix lead the plasma into an end configuration which is essentially a spindle cusp. The confinement properties of the FRC as a whole, as well as the structure of the transition between open and closed field lines, will be influenced by an interplay between transport from the interior of the FRC and transport of plasma from a cusp. In this note we begin an examination of that interplay.

II. CONFINEMENT OF PLASMA BY A CUSP

Although the idea of plasma confinement by a cusp magnetic field has been explored almost from the origin of the U.S. fusion program,⁶ there is at present no experiment, theory, or numerical simulation which gives a definitive answer as to the confinement time to be expected of a plasma in a cusped magnetic field.⁷ The theoretical difficulty is that the problem is at least two-dimensional and high- β ; the confined plasma is predicted to alter the magnetic geometry in a way that improves confinement, and this effect is predicted to depend on spatial loss areas, which requires at least a 2-D treatment. Simplifying assumptions have led to at least three^{6,8,9} disagreeing results for confinement time vs. magnetic field strength and plasma temperature. The experimental picture has been clouded by various plasma formation schemes used, and diagnostic difficulties present in 1960-1970, when the bulk of the work was done. Similarly, the need to have enough plasma that $8\pi nT \sim B^2$, and to study at least a 2-D abstraction of the problem, placed this question outside the numerical capabilities available in 1960-1970.

Recently, a variant of the electrostatic confinement fusion scheme^{10,11,12} required a knowledge of confinement of plasma by a cusp, leading to some new analytic but more especially simulation work on this problem. The results confirmed that there are indeed two regimes of cusp confinement. First, at low $\beta = 8\pi nT/B_{\max}^2$, the confinement is produced by plasma mirroring in the increasingly strong B-field at the cusp throats. The mirror ratio is $B_{\max}/B_{\text{adiabatic}}$, where B_A is the field at the location r_0 where $d \ln B(r_0)/dr \approx a_L(r_0)^{-1}$, with a_L the ion gyroradius, and the "collision time" in mirror theory is replaced by an ion transit time. This is based on the notion that a charged

particle in a cusp will lose its adiabatic memory when traversing the low field ($B(0) = 0$) interior of a cusp, reattaching itself to a field line at the location r_0 specified above, but with an isotropic velocity distribution, as if it had suffered a collision. This low β confinement produces a prediction for a spindle cusp

$$\tau_c \sim \frac{R}{(T/2M)^{1/2}} \left(\frac{R}{2a_L}\right)^{1/2} \sim \frac{R^{3/2}}{T^{3/4}} B_{\max}^{1/2}, \quad (1)$$

where $a_L = (Tmc^2/2e^2B_{\max}^2)^{1/2}$ is the ion gyroradius at $B = B_{\max}$, and $B \sim B_{\max}(r/R)^2$ has been used to model the magnetic field of a spindle cusp.

In contrast, when the plasma pressure is large enough to seriously distort the magnetic field, the equilibrium that results has a large interior region of essentially $B = 0$, bounded by a relatively sharp layer in which B rises to $B = B_{\text{vacuum}} \approx B_{\max}$. In this layer, B -lines coincide with mod- $B = \text{constant}$ lines, except for a small area at the exit of the cusp field, referred to as the "hole in the cusp." The particle motion consists of specular reflection off this layer followed by free streaming across the cusp interior to the next encounter with the boundary. This motion continues until by random walk the particle encounters the loss hole and escapes. This gives by geometric considerations,

$$\tau_c = \frac{R}{(T/2M)^{1/2}} \frac{\pi r_h^2}{4\pi R^2}, \quad (2)$$

where r_h is the radius of the loss hole, which fairly obviously is comparable to the width of the sheath which bounds the plasma. The calculation of r_h is

the crucial and unknown part of the loss calculation. Early work⁸ opined that $r_h = a_e$, the electron gyroradius, implying excellent confinement. Subsequent work found $r_h = a_e^{1/2} R^{1/2}$ under certain restrictive assumptions.⁸ More recently, $r_h = a_e^{0.8} R^{0.2}$ was obtained⁹ in a calculation with fewer assumptions than Ref. 8. A calculation which included stability considerations gave¹³ $r_h \sim a_i$, the ion gyroradius. Work is in progress on a computer simulation of confinement in a spindle cusp. Results are still preliminary but do show that τ is correctly given by Eq. (1) in the low β limit, and is substantially improved as β increases.¹⁴

We can use these ideas to estimate the enhancement of the sheath width outside the FRC closed field-line interior due to the spindle cusp. This does not require complete knowledge of τ_c , only that $\tau_c \gg R/(T/2M)^{1/2}$ which is the loss time along the truly open field lines well outside the separatrix. Both Eqs. (1) and (2) predict $\tau_c \gg R/(T/2M)^{1/2}$. The calculation of the sheath width L_s is then a calculation of the size of the entry port into the cusp field. The same calculation used for obtaining Eq. (1) can be used to find L_s . Particles leaving the closed field region of the FRC will enter the cusp if they reside on a field line which passes through the cusp at a sufficiently interior point that the particles' adiabatic invariants (e.g., the magnetic moment) are broken, and it is trapped in the cusp. But the discussion preceding Eq. (1) shows how to find the size of the adiabatic region; taking $B = B_0(x/L)^2$ and $(d \ln B/dx)_{x_0}^{-1} = a_L$ gives $x_0 = (a_L L^2)^{1/3}$ as the distance from the separatrix to the truly open field lines. The "sheath," which is the distance from $n = 0$ inward to a location where transport is determined by FRC stability considerations, extends from $x_s - x_0$ to $x_s + x_0 + \text{few } a_{Li}$. Note that the cusp entry extends inside the separatrix as well as outside. Note also that in the relation

$$x_s = (a_L L^2)^{1/3} ,$$

L is measured from the cusp center, $B = 0$. To make the result practical, a mapping of the flux line at $x = 1$ back to its location in the racetrack region of the FRC is necessary.

APPENDIX C REFERENCES

1. W. T. Armstrong, R. K. Linford, J. Lipson, D. A. Platts, and E. G. Sherwood, *Phys. Fluids* 24, 2068 (1981).
2. J. Lipson, W. T. Armstrong, J. C. Cochrane, K. F. McKenna, E. G. Sherwood, M. Tuszewski, S. Hamasaki, *Appl. Phys. Lett.* 39, 43 (1981).
3. S. Hamasaki and N. A. Krall in Proceedings of IEEE 1979 International Conference on Plasma Science, Montreal (IEEE, New York, 1979), p. 143.
4. M. Tuszewski and R. K. Linford, *Phys. Fluids* 25, 765 (1982).
5. L. Steinhauer (private communication).
6. J. Berkowitz, K. O. Friedrichs, H. Goertzel, H. Grad, J. Kileen, and E. Rubin, *Proceedings of the 2nd Intl. Conf. on Peaceful Use of Atomic Energy*, Geneva, 31, 171 (1958).
7. I. Spalding in Advances in Plasma Physics, eds. A. Simon and W. B. Thompson (Wiley, New York, 1971), Vol. 4, p. 79.
8. J. B. Taylor (private communication), Culham Report CLM-R58 (1966).
9. W. Grossmann, Jr., *Phys. Fluids* 9, 2478 (1966).
10. R. L. Hirsch, *J. of Applied Phys.* 38, 4522 (1967).
11. R. W. Bussard, *Fusion Tech.* 19, 273 (1991).
12. N. A. Krall, *Fusion Tech.* (1992, in press).
13. N. A. Krall, S. Hamasaki, R. C. Davidson, and P. C. Liewer, *Nucl. Fusion* 14, 27 (1974).
14. J. Watrous (private communication).

Figure 1. Magnetic geometry of the Field Reversed Configuration (FRC).

Figure 2. Magnetic field lines near the x-point in the FRC.

END

**DATE
FILMED
9 / 22 / 92**