DOE/ER/25015--31

DE92 040652

015 A008

COMPUTATION AND GRAPHICS
IN MATHEMATICAL RESEARCH

Progress Report
for Period June 1, 1991 - May 31, 1992

D.A. Hoffman and J. Spruck

University of Massachusetts
Amherst, MA 01003

January 1992
Resubmitted August 13, 1992

Prepared for

THE U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-86ER25015

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed or represents that its use would not infringe privately-owned rights.

TABLE OF CONTENTS

Cover Sheet	i
Table of Contents	ii
Part 1 - Description of the GANG Project and Results from Prior Research	1
1. The Center for Geometry, Analysis, Numerics and Graphics	1
2. Description of GANG Laboratory	3
3. Software Development at GANG	6
4. Mathematical and Scientific Research Activities	7
References	25
Personnel Wholly or Partially Supported by this Grant	34e
Research Visitors	34e
Appendix:	35
Geometry, Computation and Graphics Series List	36

Part 1 – Description of the GANG Project and Results from Prior Research

1 The Center for Geometry, Analysis, Numerics and Graphics

In 1986, David Hoffman and Joel Spruck began to develop a type of computational laboratory, based on interactive scientific computer graphics and high speed computation, that could be used to study geometric interface problems in the physical sciences. The laboratory is now nationally and internationally recognized as an innovative and highly successful model of the use of sophisticated computer graphics in scientific research.

The goal in creating this computational laboratory was to develop an environment to support pure and computational investigations in mathematics and to promote interdisciplinary research that requires a combination of theoretical and experimental methodology. We have had collaborations on this campus with researchers in Polymer Science, Physics and Computer Science and have developed cooperative relationships with other centers in Paris and Bonn. Research in polymer physics continues with scientists in the Program for Polymer Physics at MIT. The Center runs a visitor's program, allowing researchers to spend between one week to several months working on projects in our laboratory. Visitors come from the United States and Europe, including those who are interested in emulating all or part of our environment at their home institutions.

The primary thrust of our Center is interdisciplinary research. A natural byproduct of our research is the development of computational algorithms to simulate mathematical and physical processes. This has resulted in the development of software to simulate various physical phenomena, from constrained plasma flow to the electron microscope imaging of the microstructure of compound materials. Post-doctoral students have had degrees in Chemical Engineering and in Mathematical Physics as well as in pure Mathematics. A secondary but important activity is graduate and undergraduate training through involvement in research projects. The laboratory has a research report series that describes the completed projects of affiliated individuals. Publications from our research have appeared or will appear in publications as diverse as *Nature*, the *Communications of the ACM*, computational physics journals, and the *Journal for Colloid and Interface Science*, as well as numerous mathematics journals. In sum, the activities of the Center involve more than a single department and

the support requirements and scientific concerns are well beyond those of standard academic mathematics.

Historically, the Center's research environment has been focused on its facility at the University of Massachusetts. This facility uses 3D graphics computers, coupled by a high-speed network and directly attached to professional-quality video equipment, to provide an environment for rapid computation and visualization. However, a necessary and increasingly important activity of the laboratory is to extend this environment, in the areas of graphics and distributed computing, in order to support and enhance scientific collaboration with researchers at other locations (on- or off-campus). Toward this end, the Center developed the GANGVideo distributed video software system and the GANG Message Service distributed computing toolset.

GANGVideo, an assortment of video devices (including a scan converter for the real-time output of computer graphic displays), can be accessed and controlled from remote location for interactive viewing or creation of videotapes. GANGVideo supports both interactive and frame-by-frame video animation on several types of video storage media. Further, the Center operates a broadcast video network, running in parallel with the Center's data networks, over which (remotely-located) researchers can access stored video materials or view ongoing computations. The graphical front-end to GANGVideo can be used to make "multi-media mail" from both "live" computing sessions and stored materials without video editing. Currently, videotape is used as the delivery media, but GANGVideo is compatible with future "all-digital" distribution systems.

The GANG Message Service (GMS), now under development at the Center, is a toolset for the development of distributed and collaborative scientific applications. GMS supports the high-speed, application-level exchange of complex scientific data between processes running in a distributed, multi-processor environment, using a simple filesystem-type programming interface. The Center's MESH, IVIEW, and Visual Programming System (VPS) software packages (among others) are being adapted to this environment. The MODERATOR, originally an adjunct to GANGVideo, is also being re-implemented with GMS. The MODERATOR is a software tool for scientific collaboration which uses the GANGtv broadcast network to deliver imagery and graphics in real-time to a collaborating group, and the data networks to deliver user input from members of this group to a running application. All this will be under control of a human moderator who arbitrates control of the application. This system is an early prototype of the real-time collaborative applications imagined for the high-speed digital networks of the future, and serves as a baseline for development of

such applications at the Center.

Through its publications, through its active visitors program, and through the use of innovative computing techniques, the Center is able to extend its expertise to the broader scientific community. This model for scientific interaction has, to date, proven successful. We believe this model is adaptable to the emerging capabilities for real-time interaction over high-bandwidth communication networks. Center research and technical staff are now developing the tools and techniques for this environment.

2 Description of GANG Laboratory

The Center for Geometry, Analysis, Numerics, and Graphics (G.A.N.G.) at the University of Massachusetts maintains a computation and visual analysis facility in the Lederle Graduate Research Center. This facility serves principal investigators David Hoffman and Joel Spruck, along with collaborators, visiting scientists and graduate students (typically, about 20 people). This facility offers computation, computer graphic, videographic and software development services to these researchers.

2.1 Computation and Graphics

The computation facility centers around a Silicon Graphics 4D/240VGX graphics computer. This machine has 4 20MIP processors, 64MB of memory, 3.4GB of disk storage and a 120MIPS graphics engine, and serves as the Center's core development machine. The close coupling of high-speed processing with high-speed graphics has a great effect on the Center's focus in differential geometry: researchers are able to compute, and simultaneously visualize in near real-time, problems which required many minutes of time on earlier computers. Animation sequences are now produced as the problems are being calculated. The Center supports two additional real-time 3D graphics machines - the Silicon Graphics 4D/35TG, networked to the 4D/240GTX with the Fiber Distributed Data Interface (FDDI), and a Silicon Graphics Indigo. Two Sun 3/60 color workstations are used for still-frame color graphics and X11 clients to the server machines. Three Macintosh II workstations, are also in use, and connected to the network by an AppleTalk-Ethernet bridge for access to the SGI and Sun servers.

The Center's Sun 4/260 is the main file server for the Center's client (dataless) workstations. This system has 3.0GB of disk storage, 32MB of memory, and a high-resolution monochrome console. Two Sun SPARCstation-1s provide monochrome access points to users. The Center uses two NeXT workstations for experimentation

with the Mach operating system, for video software development, for inexpensive on-line storage of the Center's picture library using removable optical disk, and for text processing. Three Sun 3/50 monochrome workstations are used for access from offices, and for text processing.

The Center's computer systems are connected to other campus research laboratories and computing facilities (such as the Engineering Computer Services, Math Research Computing Facility, and University Computing Services) via the campus backbone Ethernet. The University has upgrading this backbone to an FDDI trunk expected to become operational during Spring 1992. A T1 speed (1.4Mb/s) link connects the campus to the NSFNet (via NEARNet), with slow-speed external connections to the CSNet and BITNet.

See attached diagram.

2.2 GANGVideo Distributed Video Services

The Center takes a distributed, interactive approach to the use of broadcast video and production of scientific video animations. The GANGVideo system is easy to use and accessible from remote locations. It meets frame and sequence archiving requirements.¹ The Center maintains a data base of images and sequences stored on laser disks, accessible through the GANGVideo software.

In Winter 1990, the Center added broadcast capabilities and a video cable system throughout Center workspaces, offices and presentation areas. Using GANGtv, and the GANGVideo software services, the video output of the Center's computers, as well as archived video materials on any media, can be viewed and controlled from remote locations.

See attached diagram.

2.3 GANG Software

Center staff have developed a number of software tools that are used by the Center's associated researchers for mathematical research and scientific visualization.

¹Using high-resolution video switching and distribution, all of the Center's graphics computers are connected to a Lyon Lamb Real Time Converter (RTC) which supports full-screen and arbitrary-window output in a variety of video formats. Recorded output can be directed to the Center's 3/4" Beta-SP tape recorder (Sony BVU-950), or 1/2" VHS tape (Mitsubishi HR8000A). A Sony LVR5000 Laser Disk Recorder is available for permanent frame storage. Up to 50 minutes of video can be recorded on each LVR disk (write-once), either as still frames or sequences. Media life is expected to exceed 30 years, making the laser disk a far better archiving media than tape.

MESH integrates with user-written surface-generation software to compute the triangulation of minimal surfaces and surfaces of constant mean curvature. SOLSURF computes the zero-set of arbitrary functions of three variables. DISP creates a number of visual representations from N-dimensional arrays of data. The Visual Programming System (VPS) is a suite of programs for creating and visualizing surface descriptions. VPS uses a mouse-driven, visual, hierarchical programming system to create both still pictures and animated sequences. Both supplied and user-written surface primitives can be used. All of the Center's geometry computation programs use the IVIEW 3D geometry viewer, allowing real-time manipulation of geometry data as it is computed. All programs work in concert with the GANGVideo software suite.

GANGVideo is a distributed video control environment that allows network-connected workstations to access and control video production equipment for traditional frame-by-frame animation, or interactive viewing and recording. Using GANGVideo and GANGtv, researchers can browse archived video materials, view the "live" output of graphics computers, and create "video mail" from these sources (with voice annotation) while working at the Center facility, or from remote workstations. Nearly all functions of the video service are under computer control. There are three interfaces to GANGVideo: the graphical user-interface, a command-line interface, and a set of function libraries callable by user-written programs. All interfaces are network transparent.

The GANG Message Service (GMS) is an application-level toolset for distributed computing. GMS allows the intercommunication of arbitrarily complex datasets among computing processes in both distributed (networked) and multi-processor environments. GMS has a very simple file-system style programming interface and includes support tools for prototyping and debugging applications. GMS is now being integrated into the Center's software system in our attempt to create a flexible, transparent distributed computing environment. Center staff are also experimenting with GMS as a distributed scientific collaboration environment, using GANGVideo and GANGtv to control the distribution of real-time imagery among collaborators.

Center software is available via anonymous FTP from Center computers. Center software generally carries either the Berkeley-style copyright, or the Free Software Foundation's "copyleft".

3 Software Development at GANG

Center staff have made progress in developing software tools for both computing and scientific visualization. These tools are now utilized in research on a daily basis at the Center. The following is a summary of the past year's efforts.

The Center's DISP general-purpose array graphics software is now available via anonymous ftp from Center computers. Versions for the Sun3 and Sun4/4C (both under Sunview and OpenWindows), and SGI Personal Iris and Power Series, are available. MESH, the adaptive mesh generator, is available on a more limited basis. An interactive graphic version is available for SGI computers. A non-graphic version is available for Sun4/4C. (MESH is described in [68].)

As the Center's real-time graphics software has evolved, particularly on the Silicon Graphics workstations, many components have become similar in function - user interface, data I/O, geometry viewing transformations, etc.. Jim Hoffman has generalized the geometry viewing functions into a "companion-module", called IVIEW, which is spawned as a subprocess to all the polygon computing applications, including those listed above, the ANNEAL mesh generator, the TREE optical density mapper (both in development), and the VRUN viewer for the Visual Programming System (VPS). Functions formerly specific to a certain program are now "generic" including: hyperbolic transformations, world and model transformation, frame sequence or "flipbook" animation, both "slicing" and "clipping" planes, and 3D stereographic viewing. IVIEW uses shared memory so that polygon data structures can be viewed as they are constructed. To support the Center's recent research in constant mean curvature surface evolution, Jim Hoffman has adapted the Brakke Surface Evolver to use IVIEW. (See the "Video Mail" tape which accompanies this report.) The use of IVIEW in distributed and collaborative environments is discussed further elsewhere in this proposal.

The Center's GANGVideo distributed video software has stabilized and is now used on a daily basis. InterNet-connected computers can control most aspects of video encoding, switching, playback and recording automatically using GANGVideo. Currently, three types of interface are available for GANGVideo: a graphical user interface (using the public domain FORMS library) for interactive use and "live" recording, a suite of command line interfaces for each specific piece of equipment (typically used in shell and VPS scripts), and a set of C-callable library routines. The GANGtv facilities are available through any of the Center's software systems using these interfaces. Current work on GANGVideo includes: a "frame daemon"

that manages laser disk storage, a "configuration daemon" to simplify setup of the video switching and broadcast network, and a more robust database interface for the laser disk video archive. See [94] for a technical summary of GANGVideo.

In anticipation of the wide deployment of high-performance all-digital data/video/voice networks, the Center has adopted a "parallel networks" approach to simulating these high-speed network environments using FDDI and the GANGtv broadcast video network. The Center is using this testbed to investigate new software tools, and enhancements to its existing tools, that promote scientific collaboration.

For example, the MODERATOR implements a simple form of "workgroup computing" that allows several persons to control and view a graphic computation process "simultaneously" under the supervision of a human moderator at the graphics computer console. The MODERATOR runs as a sub-process to the main computational code, accepting network connections from users who are able to view the computation "live" via GANGtv cable television and who control the computation process via mouse or keyboard on their own workstations. The MODERATOR inspired work on the GANG Message Service, the alternative model for distributed processing described below, and now awaits a re-implementation using this new software.

David Oliver, GANG technical director, has designed and is implementing the GANG Message Service (GMS). These are application-independent interprocess communication tools that are network transparent at both the user and programmer levels. GMS implements a "message backplane architecture" that can support both simple communication (such as the byte-stream protocols of the GANGVideo software) and high level communication of complex data structures between parallel or distributed processes. GMS is more fully described in the Technical Summary in [93].

MESH is the first software application being redesigned to use the GANG Message Service. This redesign allows MESH computation modules to be written in any computer language. This new version of MESH is now being debugged. Jim Hoffman has begun work on a GMS interface to the IVIEW geometry viewer that will allow IVIEW to run as an entirely separate module and to be run simultaneously by several viewers (on distributed computers) of the same computation.

4 Mathematical and Scientific Research Activities

During the past several years, David Hoffman and Joel Spruck have worked on a variety of pure and applied problems in Differential Geometry, Calculus of Variations

and Mathematical Physics, with an emphasis on developing new theory, whenever possible, that would lead to rigorous and effective numerical algorithms. We describe in these sections some of the major results obtained.

4.1 Research results under current support: Joel Spruck

1. Multivortex solutions in the electroweak theory

The most successful non-abelian gauge field theory is the electroweak theory of Glashow, Salam and Weinberg (GSW) which unifies the electromagnetic and weak interactions in a single framework and has been taken as the starting point of the grand unified theories of forces in nature. The recent work of Ambjorn and Olesen suggests that when a critical phase is realized, the GSW theory may allow the existence of multivortices in periodic cells. Such solutions are important because they give rise to field condensates, anti-screen the magnetic excitation, and are relevant to the quark confinement problem. The mathematical features of the model are quite interesting: The underlying equations (after reduction of the self-dual equations) are a system of two coupled nonlinear elliptic equations:

$$(4.1) \quad \begin{aligned} -\Delta u &= g^2 e^w + 4g^2 e^u - 4\pi \sum_{j=1}^k n_j \delta(z - z_j) \\ \Delta w &= \frac{g^2}{2\cos^2 \theta} (e^w - \varphi_0^2) + 2g^2 e^u \end{aligned}$$

Here g, θ, φ_0 are physical parameters, the z_j gives the prescribed location of the vortices and Ω is the fundamental domain of a lattice structure or $\Omega = \mathbb{R}^2$. The structure of these equations resembles the prescribed Gaussian curvature equation on a two-dimensional Riemannian manifold (the two-dimensional Yamabe problem).

My former student Yisong Yang and I have jointly worked on this problem and have obtained the following results through the use of a multiconstrained variational approach.

(a) We have shown that under the periodic boundary condition (periodicity up to gauge transformations), the GSW system has an N -vortex solution realizing an arbitrary distribution of vortex-lines if N satisfies

$$(4.2) \quad \frac{4\pi N}{|\Omega|} - g^2 \varphi_0^2 < \frac{8\pi \sin^2 \theta}{|\Omega|}$$

as well as the necessary condition

$$(4.3) \quad g^2 \varphi_0^2 < \frac{4\pi N}{|\Omega|} < \frac{g^2 \varphi_0^2}{\cos^2 \theta}$$

When $N = 1, 2$, our condition (4.3) for the existence is both necessary and sufficient.

(b) We have also studied the system (4.1) over the full plane $\mathbb{R}^2$ using the theory of weighted Sobolev spaces. In this case there is no restriction to the total number of vortices. We have shown that for any distribution of vortices and local vorticities, there is a two-parameter family of gauge-distinct solutions of the equations. Sharp asymptotic decay estimates at infinity are also found for these solutions.

These results will appear as two papers: **On Multivortices in the Electroweak Theory I: Existence of Periodic Solutions, II: Existence of Bogomol'nyi Solutions in $\mathbb{R}^2$** , to appear in *Communications in Math. Physics* 1992.

2. The Chern-Simons Higgs solitons

Unlike magnetic monopoles, vortices (two-dimensional static solutions in the Yang-Mills-Higgs (YMH) theory) cannot carry electric charge as well as magnetic charge. However, charged solitons are important in a wide variety of problems such as proton decay, the quantized Hall effect, high temperature superconductivity and quantum cosmology. Therefore, an effort has been made to modify the YMH model in order to accommodate these classical solutions. The recent work of Witten and Jackiw and Weinberg along with others has indicated that the Chern-Simons Higgs (CSH) theory may provide an answer. In particular, Jackiw and Weinberg showed that, when the Higgs self-interaction is of a special ϕ^6 type, self-duality will be present. In other words, the theory now allows a reduction into a Bogomol'nyi system of partial differential equations and hence multivortices carrying both electric and magnetic charges (dyons) may exist. The most interesting feature of the self-dual CSH model is that both the symmetric vacuum and the asymmetric vacuum are the ground state solutions. Therefore multivortex solutions can approach either the symmetric state or asymmetric state at infinity. The former are called topological while the latter, non-topological. The existence of these types of solutions reduces to the study of the following (scalar) nonlinear elliptic equation in $\mathbb{R}^2$:

$$\Delta u = e^u(e^u - 1) + 4\pi \sum_{i=1}^m \delta(z - z_i)$$

subject to two kinds of boundary conditions at infinity with entirely different levels of difficulty. Yisong Yang and I have obtained the following results.

(a) Topological solitons: u tends to 0 at infinity. These solutions are easier to handle than non-topological ones and there is already an earlier study by R. Wang on their existence via a non-constructive variational method. However, since there is no way to find a solution explicitly, it will be important to obtain a solution constructively. We have successfully constructed a globally convergent iterative method

to find the maximal solutions of the system. These solutions are most physical (or most superconducting) and yield quantized charge, flux, energy, and momentum. Our method has been used effectively to get interesting numerical solutions on a computer.

These results have been summarized in a paper submitted to *J. Computational Phys.* in a paper entitled **Topological solutions in Self-dual Chern-Simons Theory.**

(b) Non-topological solitons: u tends to $-\infty$ at infinity. These solutions have the interesting properties that they may give rise to fractal values of charge, flux, etc. Due to the difficulty in the boundary condition at infinity, we have thus far been unsuccessful in proving the existence of multivortex solutions by well-known PDE or variational methods. For example, the lack of a suitable function space formulation prohibits a variational solution; the difficulty in getting a good sub-solution makes it hopeless to prove the convergence of iterative sequences. However, assuming that the vortices are all superimposed at one point, we have been able to obtain an existence proof of non-topological radially symmetric CSH solitons with an arbitrary vorticity strength. Again, there is no variational formulation of the problem. Our approach is by using a shooting argument to get the desired behavior both at infinity and the origin. The correct shooting data are elusive but they can be captured as the zero set of a function. Actually, we have found a continuous family of gauge-distinct solutions of the model which may provide interesting implications to some theoretical issues of relevant problems.

These results have been submitted to *Commun. Math. Phys.* in a paper entitled **On the existence of Non-Topological Chern-Simons Solitons.**

3. Boundary Value Problems for Surfaces of Constant Gauss Curvature

Given a disjoint collection $\Gamma = (\Gamma_1, \dots, \Gamma_m)$ of smooth Jordan curves in R^3 , a fundamental question is to decide if there exist surfaces of constant Gauss curvature K (K -surfaces) with boundary Γ . More generally, we consider $\Gamma \subset R^{n+1}$ with each Γ_i a closed codimension 2 embedded submanifold, and seek a K -hypersurface of constant Gauss-Kronecker curvature K . Of course, the range of allowable K depends on Γ . An elementary necessary condition for $\Gamma \subset R^3$ to span a surface S of positive Gauss curvature $K(S) > 0$ is that each Γ_i have no inflection points. In particular, a single planar Jordan curve bounded by S must be strictly convex and S must meet the plane transversally. However, Rosenberg has shown that the condition that a single Jordan curve have no inflection points is in general not sufficient to bound an immersed

surface S with $K(S) > 0$; there are topological obstructions.

Thus far, the known existence results find K -hypersurfaces as graphs over a smooth domain $\Omega \subset R^n$ using the theory of Monge-Ampère equations. In 1984, Caffarelli, Nirenberg and Spruck proved that if Ω is strictly convex and ϕ is a C^∞ function on $\partial\Omega$, then $\Gamma = \text{graph } \phi$ bounds a K -hypersurface M for $K > 0$ sufficiently small. Moreover, M is a graph over Ω . This result was extended by Hoffman and Spruck (with H. Rosenberg) who considered domains $\Omega \subset R^n$ with boundary components $\partial\Omega = (\gamma_1, \dots, \gamma_m)$ where each γ_i is strictly convex or concave with respect to Ω . Assuming there is a C^∞ strictly locally convex graph $M = \text{graph } \underline{u}$ $x \in \bar{\Omega}$ with $K(M) > K$, then there is a K -graph $x_{n+1} = u(x)$ with $u = \underline{u}$ on $\partial\Omega$. This last result was motivated by the geometric problem $\Gamma = (C_1, C_2)$ where C_1, C_2 are strictly convex curves in parallel planes. In the special case that there is a translation which moves one of the curves inside the other, the theorem implies there is a K -graph (with respect to the translation direction) for K sufficiently small. This work will appear in *Comm. Pure and Applied Math.* in a paper entitled **Boundary value problems for surfaces of constant Gauss curvature.**

In recent work, Spruck and his Ph.D. student Bo Guan have again used the theory of Monge-Ampère equations to look for K -hypersurfaces but in a much more geometric setting. To motivate this approach consider the problem of finding K -hypersurfaces spanning "strictly extreme" Γ :

$$(4.4) \begin{cases} \text{Let } M \text{ be a smooth strictly convex closed hypersurface in } R^{n+1} \\ \text{and let } D \text{ be a smooth subdomain of } M \text{ with } \partial D = \Gamma = (\Gamma_1, \dots, \Gamma_m). \\ \text{For } 0 < K < K(M) = \inf_{P \in M} K(P) \text{ find a } K\text{-hypersurface spanning } \Gamma. \end{cases}$$

The idea is to represent M as a radial graph over a domain Ω on the unit sphere $S^n \subset R^{n+1}$ (with respect to a suitable choice of origin) and to deform M to a K -graph over Ω . For $0 < K < K(M)$, M serves as a strict subsolution for the resulting problem and the deformation takes place inside M . Thus we arrive at the following formulation of our boundary value problem on S^n :

Let Ω be a smooth domain on $S^n \subset R^{n+1}$. We seek a smooth strictly locally convex hypersurface which can be represented as

$$(4.5) \quad X(x) = \rho(x)x, \quad \rho > 0, \quad x \in \Omega,$$

with Gauss curvature

$$(4.6) \quad K[X(x)] = \psi(x), \quad x \in \Omega,$$

and boundary value

$$(4.7) \quad X(x) = \phi(x)x \quad \text{on } \partial\Omega,$$

where $\psi \in C^\infty(\bar{\Omega})$, $\phi \in C^\infty(\partial\Omega)$, $\psi, \phi > 0$.

We assume that

$$(4.8) \quad \Omega \text{ does not contain any hemisphere,}$$

and that there exists a smooth strictly locally convex radial graph $\bar{X}(x) = \bar{\rho}(x)x$ over Ω satisfying

$$(4.9) \quad \begin{aligned} K[\bar{X}(x)] &\geq \psi(x) + \delta_0 && \text{in } \Omega, \\ \bar{\rho} &= \phi && \text{on } \partial\Omega, \end{aligned}$$

for some $\delta_0 > 0$.

The main result of Spruck and Guan may be stated as follows.

Theorem 4.1. *Under conditions (4.8)-(4.9), there exists a smooth strictly locally convex radial graph $X(x) = \rho(x)x$ with $\rho \leq \bar{\rho}$ which satisfies (4.6)-(4.7). Moreover any such solution satisfies the a priori estimate*

$$(4.10) \quad \|\rho\|_{C^{2,\alpha}(\bar{\Omega})} \leq C, \quad \rho \geq c_0 > 0, \quad \kappa_i \geq \epsilon_0 \quad i = 1, \dots, n,$$

where the κ_i are the principal curvatures of X and C, c_0, ϵ_0 are uniform constants.

In general, solutions to (4.6)-(4.7) are not unique and the solution we find is closest to the subsolution. As an immediate corollary of this result we can solve Problem (4.4):

Corollary 4.2. *Let M be strictly convex hypersurface in R^{n+1} , and let $\Gamma = (\Gamma_1, \dots, \Gamma_m)$ be strictly extreme, that is $\Gamma = \partial D$ for a smooth subdomain $D \subset M$. Then for $0 < K < K(M)$, Γ bounds an embedded K -hypersurface which is contained inside M .*

In the same way we can also prove

Corollary 4.3. *Let M be a globally strictly convex hypersurface with boundary in R^{n+1} . Then for $0 < K < K(M)$, ∂M bounds an embedded K -hypersurface which is contained "inside" M .*

It is also of interest to consider a polyhedral version of Problem (4.4).

$$(4.11) \quad \left\{ \begin{array}{l} \text{Let } \Sigma \text{ be the boundary of a convex polyhedron in } R^{n+1} \text{ and let} \\ \Gamma = (\Gamma_1, \dots, \Gamma_m) \subset \Sigma \text{ be a collection of smooth strictly convex} \\ \text{codimension 2 surfaces such that each face of } \Sigma \text{ contains at most} \\ \text{one } \Gamma_i \text{ strictly in its interior. For } K > 0 \text{ sufficiently small,} \\ \text{find a } K\text{-hypersurface spanning } \Gamma. \end{array} \right.$$

Corollary 4.4. *For K sufficiently small, there is an embedded K -hypersurface which solves Problem (4.11). In particular, any two strictly convex curves in parallel planes bound some K -surface.*

One of the surprising and important features of Theorem 4.1 with respect to the theory of Monge-Ampère equations is that there are no geometric assumptions on $\partial\Omega$. That is, all of the analytic and geometric obstructions to proving existence are now embodied in the assumption of the existence of a strict subsolution of (4.9). This has led us to a careful reexamination of the local regularity estimates of Caffarelli, Nirenberg and Spruck. In particular, obtain a global result which contains an optimal smooth existence result for graphs over domains in R^n of prescribed Gauss curvature.

Theorem 4.5. *Let $\Omega \subset R^n$ be a smooth bounded domain with boundary components $\partial\Omega = (\gamma_1, \dots, \gamma_m)$. Assume there is a smooth strictly locally convex function $\underline{u}$ in $\bar{\Omega}$ satisfying*

$$(4.12) \quad \begin{aligned} \det(\underline{u}_{ij}) &\geq \psi(x, \underline{u}, \nabla \underline{u}) + \delta_0 \text{ in } \Omega \\ \underline{u} &= \phi \text{ on } \partial\Omega \end{aligned}$$

where ϕ, ψ are smooth, $\psi_0 = \inf_{\Omega} \psi > 0$ and $\psi^{\frac{1}{n}}(x, u, p)$ is convex in p . Then, there is a smooth locally convex solution $u \in C^\infty(\bar{\Omega})$ to the Monge-Ampère boundary value problem:

$$(4.13) \quad \begin{aligned} \det(u_{ij}) &= \psi(x, u, \nabla u) \text{ in } \Omega, \\ u &= \phi \text{ on } \partial\Omega. \end{aligned}$$

If $\psi_u \leq 0$ the solution is unique.

The choice $\psi(x, u, p) = K(x, u)(1 + |p|^2)^{\frac{n+2}{2}}$, $K(x, u) \geq K_0 > 0$, corresponds to prescribing the curvature function $K(x, u)$.

4. The Motion of Level Sets by Mean Curvature

Over the last several years, Spruck in collaboration with L. C. Evans has intensively investigated a new approach for defining the evolution of a hypersurface in R^n moving according to its mean curvature. In this PDE approach, given an initial surface Γ_0 , select some continuous function $g: R^n \rightarrow R$ so that

$$(4.14) \quad \Gamma_0 = \{x \in R^n | g(x) = 0\}.$$

Consider then the parabolic PDE

$$(4.15) \quad u_t = (\delta_{ij} - \frac{u_{x_i} u_{x_j}}{|Du|^2}) u_{x_i x_j} \quad \text{in } R^n \times [0, \infty)$$

$$(4.16) \quad u = g \quad \text{on } \mathbb{R}^n \times \{t = 0\},$$

for the unknown $u = u(x, t)$ ($x \in \mathbb{R}^n$, $t \geq 0$). Now the PDE (4.15) says that each *level set of u evolves according to its mean curvature*, at least in regions where u is smooth and its spacial gradient Du does not vanish. Consequently, focusing our attention on the set $\{u = 0\}$, it seems reasonable to *define*

$$(4.17) \quad \Gamma_t \equiv \{x \in \mathbb{R}^n \mid u(x, t) = 0\}$$

for each time $t > 0$ as our generalized mean curvature flow.

Using an appropriate notion of weak solution inspired by the so-called “viscosity solution” approach, we are able to prove the existence and uniqueness of a global in time solution to our mean curvature evolution PDE and to show that Γ_t is *independent* of the choice of g . This approach allows the definition of mean curvature flow even after the onset of singularities. Many surprising and interesting properties of the motion follow from a strong comparison principle for solutions. This work appeared in a fundamental initial paper: **Motion of Level Sets by Mean Curvature I**, *J. Diff. Geometry* 33 (1991) 635–681.

A second paper of Spruck and Evans gives a new and surprising proof of classical short time existence (as well as some other properties) by using the distance function to the motion. The motion is now the zero set of the distance function.

This work will appear in **Motion of Level Sets by Mean Curvature II**, to appear in *Trans. A.M.S.*

Spruck and Evans have completed a third paper studying the generalized mean curvature flow whose purpose is to effect a partial reconciliation of the level set point of view with Brakke’s geometric measure theory approach. The point of this paper is to develop tools to analyze the structure of the singularities of the generalized mean curvature flow. Along the way, several interesting results are proved.

For example, it is shown that $H^{n-1}(\Gamma_t^*) < \infty$ for each time $t > 0$ provided Γ_0 is compact, H^{n-1} -rectifiable with $H^{n-1}(\Gamma_0) < \infty$. Here $\Gamma_t^* = \partial\Gamma_t$, the topological boundary of Γ_t .

Another result of interest is that if some part of our generalized mean curvature evolution can be written as a graph, then that portion is in fact a smooth hypersurface. This is a first regularity assertion for our generalized flow. As a corollary, we show that an initial surface consisting of the boundary of an arbitrary convex set becomes smooth and convex.

This work will appear in *J. Geometric Analysis* in a paper entitled **Motion of Level Sets by Mean Curvature III**.

Recently, Spruck and Evans completed a fourth paper of the series which demonstrates that almost every level set of u is a density one varifold moving by mean curvature in the sense of Brakke. The proof uses some new estimates on the fundamental “graph” approximations to the motion utilized extensively in paper III along with ideas of compensated compactness. This work appears in the paper **Motion of Level Sets by Mean Curvature IV**.

5. Variational Problems in Plasma Physics

Spruck, in collaboration with Eydeland and Turkington at UMass, completed a first paper on variational problems with many constraints that models semilinear problems

$$-\Delta u = \Lambda(u)$$

where the profile function $\Lambda(u)$ is determined implicitly from a minimization problem subject to a family of nonlinear constraints. The motivation for this problem comes from a classic paper of Kruskal and Kulsrud which gives the natural formulation of the magnetohydrodynamic (MHD) equilibrium problem governing configurations of a plasma in a toroidal geometry. This formulation can be succinctly stated as a variational principle: an equilibrium configuration minimizes its total (potential) energy subject to two families of constraints, which determine the flux and mass within the magnetic surfaces in the plasma. These constraints are derived from the families of conserved quantities associated with ideal MHD evolution.

In this paper we study the case of an incompressible plasma (whose magnetic field has no toroidal component) in a cylindrical geometry. In this case, there is only one constraint family and the constraints are convex. We are able to formulate a natural finite constraint approximation to the original variational problem and develop a globally convergent algorithm for its solution. This finite constraint problem gives exact solutions to the physical problem and is only an approximation in the sense that the full set of constraints are not used.

Our constrained variational principle and algorithms furnish us with a foundation for many investigations in plasma physics, especially those concerned with the design of magnetic confinement devices and the analysis of their performance (see below).

This paper appeared as **Multiconstrained Variational Problems of Semilinear Eigenvalue Type: New Formulation and Algorithms**, *Mathematics of Computation* 55 (1990) 509–535.

Continuing this work, Spruck (with Eydeland, Lifshits and Turkington) showed that the algorithm of the previous paper extends to cover the case of a general ax-

isymmetric compressible plasma. In the usual cylindrical coordinates the toroidal region $D = \{x = (r, \phi, z) \in \mathbb{R}^3 : (r, z) \in \Omega, 0 \leq \phi < 2\pi\}$ is determined by its cross-section Ω , a domain in the half-plane $\mathbb{R}_+^2 = \{(r, z) : r > 0\}$. In D the plasma-vacuum system is governed by the ideal magnetohydrodynamic equilibrium equations

$$(4.18) \quad \mathbf{J} \times \mathbf{B} = \nabla p, \quad \nabla \times \mathbf{B} = \mathbf{J}, \quad \nabla \cdot \mathbf{B} = 0.$$

The plasma lies within the free-boundary surface S on which the natural interface conditions hold—namely, $p = 0$, $\mathbf{n} \cdot \mathbf{B} = 0$ on S and $\mathbf{n} \times \mathbf{B}$ is continuous across S , where $\mathbf{n}$ is the unit normal on S . On the fixed boundary ∂D , which is taken to be a perfectly conducting shell, the normal component of $\mathbf{B}$ vanishes.

Under the assumption of axisymmetry with respect to the toroidal angle ϕ , the fields in (4.18) admit the representation

$$(4.19) \quad \mathbf{B} = \nabla \Psi \times \nabla \phi + f \nabla \phi$$

$$(4.20) \quad \mathbf{J} = \nabla f \times \nabla \phi + (L\Psi) \nabla \phi$$

in terms of the scalar unknowns $\Psi = \Psi(r, z)$, $f = f(r, z)$ and the linear elliptic partial differential operator

$$(4.21) \quad L := -r \frac{\partial}{\partial r} \left(\frac{1}{r} \frac{\partial}{\partial r} \right) - \frac{\partial^2}{\partial z^2}.$$

The basis for our computational method is a variational principle which makes use of the functionals

$$(4.22) \quad E(\psi, f, g) = \frac{1}{2} \int_{\Omega} \left[r^{-1} |\nabla \psi|^2 + r^{-1} f^2 + r g^2 \right] dr dz$$

$$(4.23) \quad F_0(f) = \int_{\Omega} r^{-1} f dr dz$$

$$(4.24) \quad F_{\sigma}(\psi, f) = \int_{\Omega} r^{-1} f(\psi + \bar{\psi} - \sigma)_+ dr dz \quad (\sigma \geq \sigma_0)$$

$$(4.25) \quad G_{\sigma}(\psi, g) = \int_{\Omega} r g^{2/\gamma} (\psi + \bar{\psi} - \sigma)_+ dr dz \quad (\sigma \geq \sigma_0),$$

where the notation $s_+ = \max(s, 0)$ is used. Here the pressure p is replaced by the unknown

$$(4.26) \quad g := \left(\frac{2p}{\gamma - 1} \right)^{\frac{1}{2}},$$

as this substitution is convenient for technical reasons.

The basic variational problem governing the equilibrium configurations we study can now be stated as

$$(P_{\infty}) \left\{ \begin{array}{l} E(\psi, f, g) \rightarrow \min \text{ subject to} \\ F_0(f) = F_0^*, F_{\sigma}(\psi, f) = F_{\sigma}^*, G_{\sigma}(\psi, g) = G_{\sigma}^* \quad (\sigma \geq \sigma_0), \end{array} \right.$$

where $F_0^*, F_\sigma^*, G_\sigma^*$ denote given constraint values. This minimization problem is novel in the sense that it imposes two (continuously) *infinite* families of constraints (as σ runs through the magnetic surfaces within the plasma). Consequently, the calculation of the variational equation (a Lagrange multiplier rule) satisfied by a minimizer (ψ, f, g) is not straightforward, since certain regularity properties are required to justify such a calculation. In order to arrive at a variational problem that is tractable to analysis and computation we therefore formulate a natural discretization of the constraint families in (P_∞) . The resulting multiconstrained variational problem, which we call (P_n) is solved by a globally convergent iterative algorithm. An important feature of our formulation is that *solutions to (P_n) give exact solutions* to the equilibrium equations (4.18) (and not approximations!) Also, many of the ideas introduced have application to other optimization problems with nonconvex constraints.

This work will appear as **Multiconstrained Variational Problems in Magnetohydrodynamics I: Equilibrium**, preprint.

Continuing this work, Spruck (with A. Lifshits, Eydeland and Turkington), has applied the previous work to study the slow evolution of a plasma that describes flux conserving and adiabatically compressed tokamaks. The method allows us to easily study magnetic island formation and annihilation. This work will appear as **Multiconstrained Variational Problems in Magnetohydrodynamics II: Slow Evolution**, preprint.

6. Asymptotic Symmetry and applications

Spruck, in collaboration with Caffarelli and Gidas, completed his work on the classification of isolated singularities of the conformally invariant scalar field equation:

$$\Delta u + u^{\frac{n+2}{n-2}} = 0$$

$$u \geq 0.$$

In doing so, a powerful new variant of the Alexandrov reflection technique that is measure theoretic and localizable was developed. This work appeared in **Asymptotic Symmetry and the Local Behavior of Semilinear Elliptic Equations with Critical Sobolev Growth**, *C.P.A.M.* 42 (1989) 271-297.

Spruck with Caffarelli continued the above mentioned work to study variational problems with critical Sobolev growth. In particular, they proved that the Dirichlet problem

$$\Delta u + \lambda u^{\frac{n+2}{n-2}} = 0 \quad \text{on } \Omega$$

$$u = \phi \quad \text{on } \partial\Omega$$

for nonnegative ϕ , $\phi \not\equiv 0$, is well-posed. This result is of particular interest because it clarifies the reasons for blow up. This work recently appeared in *Variational Problems with Critical Sobolev Growth and Positive Dirichlet Data*, *Indiana University Math. J.* 39 (1990) 1-18.

4.2 Research results under current support: David Hoffman

During the past several years, Hoffman and collaborators have studied surface interfaces, satisfying various volume constraints, that are also critical points for physically meaningful energies. This work has three interlocking components: extending mathematical theory of existence and behavior of interfaces; development and use of new algorithms for computing and displaying new equilibria; application of the theoretical and computational mathematics to real problems in materials science.

1. Existence and limiting behavior of embedded minimal surfaces of finite topology

The fundamental paper *Embedded minimal surfaces of finite topology*, *Annals of Mathematics* 131 (1990) 1-34 written with W. Meeks, established the existence of an infinite sequence of embedded minimal surfaces of genus $k \geq 1$ with three topological ends, generalizing the genus-one example of Costa and gave a uniform proof of embeddedness. It is proved in that paper that these surfaces are the most symmetric examples of genus- k with three ends. A symmetry extension theorem was presented that gives circumstances under which symmetries of the Gauss mapping of a minimal surface can extend to the surface in R^3 . Included at the beginning of this paper is a synopsis of the known results about embedded minimal surfaces of finite total curvature and finite topology, as well as an estimate (in terms of the total winding at infinity) of the maximal order of self-intersection that a minimal surface can have.

In subsequent work Meeks and Hoffman investigated the behavior, as $k \rightarrow \infty$ of these surfaces. This was published in *Limits of minimal surfaces and Scherk's fifth surface*, which appeared in the *Archive for Rational Mechanics and Analysis*, Volume 111 No. 2, 1990 181-195.

Further progress has been made in finding new examples. Meeks and Hoffman discovered that each of the three-ended embedded minimal surfaces, discussed above,

could be deformed through a one-parameter family. The original undeformed surfaces have two catenoid ends and one planar end between them. The deformed surfaces have three catenoid ends. This is carried out in **One-parameter families of embedded minimal surfaces of finite total curvature** (with H. Karcher and W. H. Meeks), which is in the final stages of preparation and will appear soon. The difficulty in giving a mathematical proof of the existence of these embedded surfaces had to do with the fact that one must solve the period problem while keeping the logarithmic growths of the catenoid ends strictly ordered. This additional constraint made the problem much more delicate. This paper also contains the first results describing the structure of the space of embedded minimal surfaces of finite total curvature. After a natural geometric normalization, we prove that the space of all complete embedded minimal surfaces of genus less than or equal to k and with r or fewer ends is compact, but that degeneration (drop in genus and drop in number of ends) can occur at the boundary of these spaces.

2. Global theoretical results concerning embedded minimal surfaces of finite topology

The theoretical requirements of the above paper also had an influence on the contents of **Complete embedded minimal surfaces of finite total curvature** (Survey article for Springer Encyclopaedia of Mathematics Volume on minimal submanifolds, R. Osserman ed.), which I am writing with Hermann Karcher and which is in the final stages of preparation. The necessary theoretical framework for the results of **One-parameter families of embedded minimal surfaces of finite total curvature** were added to this survey, after being suitably generalized.

In **The asymptotic behavior of properly embedded minimal surfaces of finite topology** *Journal of the AMS* 2 No. 4 (October, 1989) 667–681, which is another paper written with Meeks, we investigate annular ends of properly embedded minimal surfaces. It is proved there that there can be, at most, two annular ends with infinite total curvature. Subsequent work of Y. Fang (who did his dissertation under the supervision of Hoffman) and Meeks [89] established that these ends had certain growth properties and could be isolated in the following sense. If a properly embedded minimal surface of finite topology has two ends of infinite total curvature, then each such end must be contained in its own halfspace and the remaining ends of the surface are all flat and located in the slab between the ends.

These two papers used a very general result proved by Hoffman and Meeks in **The strong halfspace theorem for minimal surfaces**, *Inventiones Math.* 101, (1990)

373–377. The result states that any two properly immersed minimal surfaces in R^3 must intersect, unless they are parallel flat planes; in particular, a properly immersed minimal surface cannot lie in a halfspace of R^3 .

Meeks and Hoffman wrote an invited survey article for the special Geometry issue of the *American Mathematical Monthly*. This article, Minimal surfaces based on the catenoid, *American Mathematical Monthly*, 97 No. 8 (1990) 702–730, also touches on the subject of singly and doubly-periodic, embedded minimal surfaces, which we will discuss below. Another version of this article was translated into French: La Caténoïde par les deux bouts, *Quadrature, Bimestriel* No. 5, (Août Septembre 1990) 31–48 (with W. H. Meeks).

3. Singly-periodic embedded minimal surfaces, I

An early survey of results and conjectures is given in our paper New examples of singly periodic minimal surfaces and their qualitative behavior, which was published in *Contemporary Mathematics* 101, 1989, 97–106. In the paper The structure of singly-periodic minimal surfaces, *Inventiones Math.* 99, (1990) 455–481, which is joint work with M. Callahan and W. Meeks, we prove the existence of singly-periodic embedded minimal surfaces with an infinite number of flat ends and any odd genus ≥ 3 . These are the first examples found, since those of Riemann, of properly embedded minimal surfaces with an infinite number of ends and an infinite symmetry group. We show how to construct them from three different points of view: Weierstrass representation; conjugate surface construction; and as desingularized limit of a sequence of surfaces whose curvature is blowing up at a point. Following this paper, the same authors wrote The structure of singly-periodic minimal surfaces, *Inventiones Math.* 99, (1990) 455–481, in which they showed that many of the properties of the surfaces constructed in the first paper were forced. That is, they characterized to a great extent the singly-periodic embedded minimal surfaces with an infinite number of topological ends. In this work the notion of the “limit tangent plane” is introduced for the first time in the literature and its fundamental properties are proved. It has been used a great deal in the past two years. As a consequence of the theory developed in this paper, we were able to show that every embedded doubly-periodic surface that has finite topology in the quotient must necessarily have one topological end. This is a very strong restriction.

In this same paper we also proved that the surfaces constructed in “The structure of singly-periodic minimal surfaces” could be deformed through a family of embedded singly-periodic minimal surfaces that are invariant under a screw motion: that is, the

composition of a translation and a rotation about the axis of translation. The proof of the existence of these surfaces was nonconstructive, a generalization of the method of desingularizing the limit of a sequence of surfaces whose curvature is blowing up at a point. The conjugate surface construction will not work because, after reducing by its cyclic group of screw motions to a compact surface, there are not enough symmetries left to produce a fundamental domain bounded by lines or planar geodesics. The standard Weierstrass representation will not work either because the Gauss map is not well-defined on the quotient surface; only the logarithmic derivative descends. This means that the period problem for these screw-motion-invariant surfaces must be solved in terms of a Gauss map, which itself is computed by integration and has its own period problem of a different sort. We were able to do this by a combination of theoretical and computational techniques. The results will appear in **Smooth deformation families of singly periodic minimal surfaces invariant under screw motions**, written with M. Callahan and H. Karcher. The successful results of the computation can be seen in the video entitled "A Family of Minimal Surfaces with Screw-Motion Symmetry and Infinitely Many Ends" on the videotape "Deformation Families of Embedded Minimal Surfaces, Part 1," made by M. Callahan in August, 1991 at GANG.

4. Singly periodic embedded minimal surfaces, II

M. Wohlgemuth and I have been able to add handles to the singly-periodic examples described above. This appears in **New examples of singly periodic minimal surfaces**, which is in preparation. Illustrations of these surfaces can be seen in the article "Minimal surfaces based on the catenoid," mentioned above. Their construction is also depicted in the video lecture accompanying our article **Construction of minimal surfaces with and without conformal representation**, which appeared in "Computing Optimal Geometries," Jean Taylor, ed. AMS Selected Lectures in Mathematics. This article describes computational techniques for computing minimal surface interfaces, including solving the Plateau Problem by direct area minimization methods. It also contains an animation of the deformation of the Karcher-Hoffman surface $\mathcal{W}_k$, for $k = 1$, which is a desingularization of $2k + 1$ coaxial helicoids. This surface is also depicted on the cover of the special Geometry Issue of the *MAA Monthly*, which was mentioned above. The generalized Weierstrass Representation for these surfaces will appear in **One parameter families of singly-periodic minimal surfaces derived from exotic triply-periodic examples**, an article which is in preparation with H. Karcher. The exotic examples referred to

are the triply-periodic surfaces discovered by the crystallographers Fischer and Koch [38, 39, 40, 41].

Because the Riemann examples have been crucial in development of the theory of singly-periodic examples, we have focused attention on them. We have found that while they are in many ways the simplest examples (they are characterized, along with the catenoid, by the fact that they are fibred by circles in parallel planes) they are much more rigid than the recently discovered examples. In **Embedded minimal annuli bounded by a pair of straight lines in parallel planes**, a joint work with H. Karcher and H. Rosenberg about to appear in *Comm. Math. Helv.*, we prove that if the lines referred to in the title of the paper are parallel, then the surface must be part of a Riemann example. In a sequel, which is now in preparation, entitled **The classification of complete embedded periodic minimal surfaces of total curvature -8π in the quotient**, we show that if the lines are not parallel, there is no solution. That implies that the examples of Riemann can not be modified to be invariant under a screw motion. (This has also been proved independently by Pérez and Ros [98].) Also in this paper, we partially classify singly periodic embedded minimal surfaces with one topological end and low total curvature, extending some results of Meeks-Rosenberg and of Wei [112]. This paper is joint work with H. Rosenberg and E. Toubiana. We also have recently completed project with W. Rossman, a graduate student at UMass who works with GANG researchers. It begins with the observation that the Riemann examples, which form a one-parameter family, come in pairs that are not congruent but are isometric. (The conjugate surface to a Riemann example is another, different Riemann example.) We then show that the possible limit surfaces (in the topology of convergence in compact subsets of R^3) in this family consist of Riemann examples, the catenoid and the helicoid, in addition to the degenerate limits of a flat (perhaps multiply covered) plane. This will appear in **Limits of Riemann's examples**, (with W. Rossman).

Recent results of Pascal Romon, a student from U. Paris VII who is now visiting GANG, indicate that if one considers embedded minimal annuli bounded by parallel lines in parallel planes, then even if additional flat ends are allowed, the surface must be a Riemann example. Specifically, he has shown that an annular embedded minimal surface, bounded by parallel lines and containing one intermediate flat end, must be the Riemann example. The technique should generalize [101]. If however, one allows higher topology then recent results of Wei at GANG show that there are such examples. They have the property that they divide space into two regions whose asymptotic volume-ratio is not $1/2$. This property is important for applications to

compound polymers and Wei's examples are the first of this type to be discovered. We are currently working with Wei to find a general existence proof for these important examples. One of them can be seen in the "Video Mail" tape produced at the GANG Center.

5. The morphology of microstructures in compound polymers

Polymer physicists are interested in the properties of substances with very large molecular units, consisting of two or more different long polymers covalently bonded together. These are the so-called "block copolymers". These bonded polymer molecules are often naturally repelling. When many of these block copolymers are brought together in a condensed phase, they move so as to isolate the different molecular components. If the material is allowed to anneal to a stable state, the resulting configuration is called an *equilibrium domain morphology*. It is determined by the region between the two separated phases, which we model as a surface, and refer to as the *interface*.

The interface geometry is governed by forces dominated by the necessity to reduce surface energy. Because minimization of surface area is the most powerful driving force one expects, to a good approximation, that surfaces of constant mean curvature or (as a special case) minimal surfaces should appear. Basic work in the 80's by Scriven [102], Anderson [8], Anderson, Davis, Nitsche and Scriven [10], Anderson, Henke, Hoffman and Thomas [6] and Bellare and Davis et al. [13] established that these were appropriate models and also found many new periodic surfaces of constant mean curvature. A novel feature of this research was and is that the impetus for mathematical advances in the existence theory for these surfaces came first from practical scientific concerns and then from computer simulation. The interaction between pure mathematics, experimentation² and computer simulation is reflected in the aforementioned work as well as our papers, which will be cited below.

Our work on this subject began with E. L. Thomas, a polymer physicist who is now at MIT, and David Anderson who was the first GANG postdoctoral student in 1986. It has continued with J. Bellare and then with others at MIT and elsewhere. In *Natural minimal surfaces in mathematics and science*, a video lecture with animated computer graphics, December, 1990 produced by *Science Television*, we give an overview to date (August 1990), of our research on equilibrium domain morpholo-

²One of the ways that these polymer structures are investigated is by means of transmission electron microscope images of very thin slices of material. Typically, one phase is doped to be more opaque than the other. Then, by varying the angle of the beam, one can get a series of gray scale images, which then must be interpreted to reveal the microstructure of the material.

gies and tied it in with the developing mathematical theory. In 1990, the volume **Computer Graphics and Geometric Analysis: Proceedings of the Conference on Differential Geometry, Calculus of Variations and Computer Graphics**, which we edited with P. Concus and R. Finn, appeared. The conference at MSRI, which we helped to organize, was perhaps the first to have featured lectures by mathematicians and polymer physicists and computer graphics experts.

The paper **Algorithms for the computer simulation of 2D projections from structures determined by dividing surfaces**, written with D. Anderson, E. Thomas, J. Hoffman et al., will appear in the *Journal of Colloid and Interface Science*. It gives details of the algorithms we use to simulate various exploratory techniques (including electron microscopy) on the computer and compare the results with laboratory data. The investigation of grain boundaries, treated in this paper, led to the consideration of the Scherk family of doubly-periodic minimal surfaces as possible interfaces. With new discoveries, by Karcher, Wei and Hoffman, of other possible models for twist boundaries in grains, it became necessary to know precise information (numerical and analytic) about the Scherk family. This appears in **Analytic data for the Scherk family of doubly-periodic minimal surfaces** which is a GANG Tech report. Work on this subject continues with E. L. Thomas and S. Gido. In particular, we present new models for twist boundaries, using recent results about the limiting behavior of periodic surfaces, as well as an analysis of dislocations from the point of view of minimal surfaces invariant under a screw motion. This work, which is now being completed, and will be submitted to the journal *Microscopy*, is in the paper **Block copolymer grain boundary morphology, I: lamellar twist boundary**.

Other work of Hoffman relevant to this topic is contained in the paper **Some basic facts, old and new, about triply periodic embedded minimal surfaces**, *Colloques de Physique, Colloque No. 7 Supplement au Journal de Physique*, Tome 51. December 1990, 197-209. This volume is the Proceedings of a CNRS conference on surface interfaces, which was attended by mathematicians, polymer physicists and chemists. I gave two talks at the Conference: one was the lead-off talk, a survey of the relevant geometry and math; the other was a talk upon which this paper was based. Unusual for a conference paper, it actually contains a new result: a theorem that explains the blow up of a computation, done by Leibler and Maggs, to compute deformation families of triply-periodic embedded minimal surface interfaces of genus 3. I was asked this question at the conference, and was able to show that there was forced symmetry-breaking in the family they were trying to approximate. The

article also surveys results published elsewhere, and it discusses the use of minimal surfaces as physical models. (They get a mixed review from me). I was also on the Scientific Advisory Committee for this conference.

Other publications containing mathematics relevant to the study of equilibrium domain morphologies is contained in [23, 25, 26, 61, 62].

References

- [1] aPE Version 2.0 Reference Manual. Technical report, Ohio Supercomputer Graphics Project, 1990.
- [2] AVS 3.0 Developer's Guide. Technical report, Stardent Computer Corporation, 1991.
- [3] M. Accetta, R. Baron, D. Golub, R. Rashid, A. Tevanian, and M. Young. Mach: A New Kernel Foundation for UNIX Development. Technical report, School of Computer Science, Carnegie Mellon University, Pittsburg, PA, 1986.
- [4] W. Allard. On the first variation of a varifold. *Ann. of Math.*, 95:417-491, 1972.
- [5] J. Ambjørn and P. Olesen. On Electroweak Magnetism. *Nuclear Physics*, B315:606-614, 1989.
- [6] D. Anderson, C. Henke, D. Hoffman, and E. L. Thomas. Periodic area-minimizing surfaces in block copolymers. *Nature*, 334(6184):598-601, 1988. August 18 issue.
- [7] D. Anderson and E. L. Thomas. Microdomain morphology of star copolymers in the strong-segregation limit. *Macromolecules*, 21:3221-3230, 1988.
- [8] D. M. Anderson. *Studies in the Microstructure of Microemulsions*. PhD thesis, University of Minnesota, June 1986.
- [9] D. M. Anderson, J. Bellare, J. T. Hoffman, D. Hoffman, and E. L. Thomas. Algorithms for the computer simulation of 2D projections from structures determined by dividing surfaces. to appear in the *Journal of Colloid and Interface Science*.
- [10] D. M. Anderson, H. T. Davis, J. C. C. Nitsche, and L. E. Scriven. Periodic surfaces of prescribed mean curvature. *Advances in Chem. Phys.*

- [11] D. M. Anderson, D. C. Martin, and E. L. Thomas. Maximum entropy data restoration using both real and Fourier space analysis. *Acta Crystallographica*, A45, pages 686-698, 1989.
- [12] S. Angenent. Shrinking Doughnuts. Proceedings of the Conference on Elliptic and Parabolic Equations. Gregynog, Wales, 1989.
- [13] J. R. Bellare, H. T. Davis, W. G. Miller, and L. E. Scriven. Polarized optical microscopy of anisotropic media: imaging theory and simulation. *J. Colloid Interface Sci.*, 136(2):305-326, 1990.
- [14] K. A. Brakke. *The motion of a surface by its mean curvature*. Princeton University Press, Princeton, New Jersey, 1978.
- [15] K. A. Brakke. *Surface Evolver Manual Version 1.65*. Geometry Center, University of Minnesota, 1991. Technical Report.
- [16] F. Brito and R. Sá Earp. Geometric configurations of constant mean curvature surfaces with planar boundary. to appear in *Anais da Academia Brasileira de Ciências*.
- [17] F. Brito and R. Sá Earp. On the configuration of certain prescribed mean curvature surfaces with planar boundary. preprint.
- [18] F. Brito, R. Sá Earp, W. H. Meeks III, and H. Rosenberg. A halfspace containment property for constant mean curvature surfaces bounded by a convex planar curve. to appear in the *Indiana Journal of Math.*
- [19] R. Bryant. A duality theorem for Willmore surfaces. *Journal of Differential Geometry*, 20:23-53, 1984.
- [20] R. Bryant. Surfaces in conformal geometry. In R. O. Wells, editor, *The Mathematical Heritage of Hermann Weyl*, volume 48 of *Proceedings of Symposia in Pure Math.* AMS, 1988.
- [21] L. A. Caffarelli, J. J. Kohn, L. Nirenberg, and J. Spruck. The Dirichlet Problem for Nonlinear Second-Order Elliptic Equations IV, Starshaped Compact Weingarten Hypersurfaces. *Current topics in P.D.E.*, pages 19-27, 1986.
- [22] L. A. Caffarelli, L. Nirenberg, and J. Spruck. The Dirichlet Problem for Nonlinear Second-Order Elliptic Equations I. Monge-Ampere Equations. *Comm. in Pure and Applied Math*, 37:369-402, 1984.

- [23] M. Callahan, D. Hoffman, and J. Hoffman. Computer graphics tools for the study of minimal surfaces. *Communications of the ACM*, 31(6):648-661, 1988.
- [24] M. Callahan, D. Hoffman, and W. H. Meeks III. Embedded minimal surfaces with four ends. in preparation.
- [25] M. Callahan, D. Hoffman, and W. H. Meeks III. Embedded minimal surfaces with an infinite number of ends. *Inventiones Math.*, 96:459-505, 1989.
- [26] M. Callahan, D. Hoffman, and W. H. Meeks III. The structure of singly-periodic minimal surfaces. *Inventiones Math.*, 99:455-481, 1990.
- [27] M. Callahan and H. Karcher. Smooth deformation families of singly periodic minimal surfaces invariant under screw motions. *GANG preprint*.
- [28] Y-G. Chen, Y. Giga, and S. Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. preprint, 1989.
- [29] H. I. Choi and R. Schoen. The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature. *Invent. Math.*, 81:387-394, 1985.
- [30] R. Draves. A Revised IPC Interface. In *Proceedings of USENIX Mach Workshop*. USENIX Association, San Diego CA, 1990.
- [31] R.P. Draves, M. B. Jones, and M. R. Thompson. MiG - The Mach Interface Generator. Technical report, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, 1989.
- [32] R. Droms and R. Haber. Workshop Report of the SIGGraph / SIGComm Workshop on Computer Graphics and Networking. *Computer Graphics Quarterly*, vol 25, no 4, 1991.
- [33] J. Grudin (Editor). Collaborative computing. *Communications of the ACM*, vol 34 no 12, December 1991.
- [34] L. C. Evans and J. Spruck. Motion of Level Sets by Mean Curvature II. to appear in *Trans. A.M.S.*
- [35] L. C. Evans and J. Spruck. Motion of Level Sets by Mean Curvature III. to appear in *J. Geometric Analysis*.

- [36] L. C. Evans and J. Spruck. Motion of Level Sets by Mean Curvature IV. preprint.
- [37] L. C. Evans and J. Spruck. Motion of Level Sets by Mean Curvature I. *J. Diff. Geometry*, 33:635-681, 1991.
- [38] W. Fischer and E. Koch. On 3-periodic minimal surfaces. *Zeitschrift für Kristallographie*, 179:31-52, 1987.
- [39] W. Fischer and E. Koch. On 3-periodic minimal surfaces with non-cubic symmetry. *Zeitschrift für Kristallographie*, 183:129-152, 1988.
- [40] W. Fischer and E. Koch. Topological characterization of 3-periodic minimal balance surfaces. *Zeitschrift für Kristallographie*, 185:293, 1988.
- [41] W. Fischer and E. Koch. New surface patches for minimal balance surfaces. I. Branched catenoids. *Acta Cryst.*, 45:166-169, 1989.
- [42] M. Gage and R. S. Hamilton. The heat equation shrinking convex plane curves. *J. Differential Geometry*, 23:69-96, 1986.
- [43] M. Gerald-Yamosaki. Distributed Library. Technical report, NASA NAS, Moffet Field, CA, 1990.
- [44] M. Gerald-Yamosaki. Cooperative Visualization of Computational Fluid Dynamics. Technical report, NASA NAS, Moffet Field, CA, 1991.
- [45] S. Gido, J. Gunther, D. Hoffman, and E. L. Thomas. Block copolymer grain boundary morphology I. Lamellar twist boundary. to be submitted to *Macromolecules*.
- [46] S. Gido and E. L. Thomas. Block copolymer grain boundary morphology II. preprint.
- [47] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer-Verlag, New York, 2nd edition, 1983.
- [48] M. Grayson. The heat equation shrinks embedded plane curves to round points. *J. Differential Geometry*, 26:285-314, 1987.
- [49] S. Greenberg. Sharing Views and Interactions with Singer-user Applications. In F. H. Lochovsky and R. B. Allen, editors, *ACM SIGOIS Bulletin*, volume

vol 11 no 2,3 of *Proceedings of the Conference on Office Information Systems*. Association for Computing Machinery, New York, NY, 1990.

- [50] Bo Guan and J. Spruck. Boundary value problems on S^n for surface of constant Gauss curvature. preprint.
- [51] D. Hoffman and H. Karcher. Singly periodic minimal surfaces derived from crystallographic examples. in preparation.
- [52] D. Hoffman, H. Karcher, and W. H. Meeks III. One-parameter families of embedded complete minimal surfaces of finite topology. preprint.
- [53] D. Hoffman, H. Karcher, and H. Rosenberg. Embedded minimal annuli in $\mathbb{R}^3$ bounded by a pair of straight lines. preprint.
- [54] D. Hoffman and W. H. Meeks III. A complete embedded minimal surface in $\mathbb{R}^3$ with genus one and three ends. *Journal of Differential Geometry*, 21:109–127, 1985.
- [55] D. Hoffman and W. H. Meeks III. Complete embedded minimal surfaces of finite total curvature. *Bulletin of the AMS*, 12:134–136, 1985.
- [56] D. Hoffman and W. H. Meeks III. Properties of properly embedded minimal surfaces of finite total curvature. *Bulletin of the AMS*, 17(2):296–300, 1987.
- [57] D. Hoffman and W. H. Meeks III. A variational approach to the existence of complete embedded minimal surfaces. *Duke Journal of Math.*, 57(3):877–894, 1988.
- [58] D. Hoffman and W. H. Meeks III. The asymptotic behavior of properly embedded minimal surfaces of finite topology. *Journal of the AMS*, 2(4):667–681, 1989.
- [59] D. Hoffman and W. H. Meeks III. Embedded minimal surfaces of finite topology. *Annals of Mathematics*, 131:1–34, 1990.
- [60] D. Hoffman and W. H. Meeks III. Les surfaces minimales: la caténoïde par les deux bouts. *Quadrature*, 5:31–47, 1990.
- [61] D. Hoffman and W. H. Meeks III. Limits of minimal surfaces and Scherk's Fifth Surface. *Arch. Rat. Mech. Anal.*, 111(2):181–195, 1990.

- [62] D. Hoffman and W. H. Meeks III. Minimal surfaces based on the catenoid. *Amer. Math. Monthly, Special Geometry Issue*, 97(8):702-730, 1990.
- [63] D. Hoffman and W. H. Meeks III. The strong halfspace theorem for minimal surfaces. *Inventiones Math.*, 101:373-377, 1990.
- [64] D. Hoffman, H. Rosenberg, and J. Spruck. Boundary Value Problems for Surfaces of constant Gauss curvature. to appear in *Comm. Pure and Applied Math.*
- [65] D. Hoffman, E. Toubiana, and H. Rosenberg. The classification of complete embedded periodic minimal surfaces of total curvature -8π in the quotient. in preparation.
- [66] D. Hoffman and M. Wohlgemuth. New embedded periodic minimal surfaces of Riemann-type. in preparation.
- [67] D. Hoffman and M. Wohlgemuth. Limiting behavior of classical periodic minimal surfaces. in preparation, 1989.
- [68] J. T. Hoffman. MESH manual. *GANG preprint series II*, #35.
- [69] J. T. Hoffman. A visual programming system and image synthesis application. *GANG preprint series II*, #19; to appear in *Leonardo*.
- [70] J.T. Hoffman. The Visual Programming System. Technical report, Center for Geometry, Analysis, Numerics and Graphics, University of Massachusetts, Amherst MA, 1991.
- [71] G. Huisken. Flow by mean curvature of convex surfaces into spheres. *J. Differential Geometry*, 20:237-266, 1984.
- [72] A. Jaffe and C. Taubes. Vortices and Monopoles. Birkhauser, Boston, 1980.
- [73] S. M. Kaplan, A. M. Carroll, and K. J. MacGregor. Supporting Collaborative Processes with ConversationBuilder. In Peter deJong, editor, *ACM SIGOIS Bulletin*, volume 12 no 2,3 of *Proceedings of the Conference on Organizational Computing Systems*. Association for Computing Machinery, New York, NY, 1991.
- [74] H. Karcher. Comparison theorems for minimal surfaces. preprint.
- [75] H. Karcher. Embedded minimal surfaces derived from Scherk's examples. *Manuscripta Math.*, 62:83-114, 1988.

- [76] H. Karcher, U. Pinkall, and I. Sterling. New minimal surfaces in S^3 . *Journal of Differential Geometry*, 28(2):169–186, 1988.
- [77] N. Korevaar and L. Simon. Boundary regularity for solutions of prescribed mean curvature problems. in preparation.
- [78] M. D. Kruskal and R. M. Kulsrud. Equilibrium of a magnetically confined plasma in a toroid. *Physics of Fluids*, 1:265–274, 1958.
- [79] R. Kusner. Conformal geometry and complete minimal surfaces. *Bulletin of the AMS*, 17(2):291–295, 1987.
- [80] R. Kusner. Comparison surfaces for the Willmore problem. *Pacific J. Math.*, 138:317–345, 1989.
- [81] H. B. Lawson. Complete minimal surfaces in S^3 . *Annals of Math.*, 92:335–374, 1970.
- [82] J. J. Lee. XSketch: A Multi-user Sketching Tool for X11. In F. H. Lochovsky and R. B. Allen, editors, *ACM SIGOIS Bulletin*, volume vol 11 no 2,3 of *Proceedings of the Conference on Office Information Systems*. Association for Computing Machinery, New York, NY, 1990.
- [83] Daniel Lenoski, James Laudon, Kourosh Gharacholroo, Wolf-Dietrich Weber, Anoop Gupta, and John L. Hennessey. Overview and Status of the Stanford DASH multiprocessor. In *Proceedings of the International Symposium on Shared Memory Multiprocessing*, pages 102–108, Tokyo, Japan, April 1991.
- [84] K. Leopere. Mach 3 Kernel Principles. Technical report, Open Software Foundation and Carnegie Mellon University, 1991.
- [85] H. Lewy. On the boundary behavior of minimal surfaces. *Proceedings of the National Academy*, 37:103–110, 1951.
- [86] W. H. Meeks III and B. White. The space of minimal annuli bounded by an extremal pair of planar curves. preprint.
- [87] W. H. Meeks III and B. White. Minimal surfaces bounded by a pair of convex planar curves. *Bulletin of the AMS.*, 24(1):179–184, 1991. Research Announcement.

- [88] W. H. Meeks III and B. White. Minimal surfaces bounded by convex curves in parallel planes. *Commentari Mathematici Helvetici*, 66:263–278, 1991.
- [89] W. H. Meeks III and S. T. Yau. The classical Plateau problem and the topology of three-dimensional manifolds. *Topology*, 21(4):409–442, 1982.
- [90] G.M. Neilson. Visualization in Scientific and Engineering Computing. *IEEE Computer*, vol 24, no 9, 1991.
- [91] V. I. Oliker. The problem of embedding S^n into R^{n+1} with prescribed Gauss curvature and its solution by variational methods. *Trans. A.M.S.*, 295:291–303, 1986.
- [92] D. M. Oliver. Scientific Visualization: Work in Progress at GANG. Technical report, Center for Geometry, Analysis, Numerics and Graphics, University of Massachusetts, Amherst MA, 1990.
- [93] D. M. Oliver. The GANG Message Service. Technical report, Center for Geometry, Analysis, Numerics and Graphics, University of Massachusetts, Amherst MA, 1991.
- [94] D. M. Oliver. The GANGVideo Distributed Video Services. Technical report, Center for Geometry, Analysis, Numerics and Graphics, University of Massachusetts, Amherst MA, 1991.
- [95] D. M. Oliver. Workshop Report of the SIGGraph'91 Workshop on Computer Graphics in the Network Environment. *Computer Graphics Quarterly (to appear)*, vol 26, no 2, 1992.
- [96] S. Osher and J. A. Sethian. Fronts propagating with curvature dependent speed: algorithms based on Hamilton-Jacobi formulations. *J. Computational Physics*, 79:12–49, 1988.
- [97] M. Overmars. Forms Library Version 2.0. Technical report, Department of Computer Science, Utrecht University, 1991.
- [98] J. Pérez and A. Ros. Some uniqueness and nonexistence theorems for embedded minimal surfaces. preprint.
- [99] D. Phillips, J. Torborg, C. Molor, M. Pique, and D. Greenberg. Distributed Graphics: Where to Draw the Line? In *Panel Proceedings ACM SIGGraph'89*. Association for Computing Machinery, Chicago IL, 1989.

- [100] S. P. Reiss. Connecting Tools Using Message Passing in the Field Environment. *IEEE Software*, 1990.
- [101] P. Romon. Unicity of the double covering of the riemann example. *GANG preprint, to appear*.
- [102] L. E. Scriven. Equilibrium bicontinuous structures. *Nature*, 263:123-125, 1976.
- [103] J. A. Sethian. Recent numerical algorithms for hypersurfaces moving with curvature-dependent speed: Hamilton-Jacobi equations and conservation laws. *J. Diff. Geometry*, 31:131-161, 1990.
- [104] L. Simon. Existence of Willmore surfaces. In *Proceedings of the Australian National University*, 1986.
- [105] J. Spruck and Y. Yang. The Existence of Nontopological Solitons in Chern-Simons Theory. preprint.
- [106] J. Spruck and Y. Yang. On Multivortices in the Electroweak Theory I: Existence of Periodic Vortices. to appear in *Comm. in Math. Physics*.
- [107] J. Spruck and Y. Yang. On Multivortices in the Electroweak Theory II: Existence of Bogomol'nyi solutions in R^2 . to appear in *Comm. in Math. Physics*.
- [108] J. Spruck and Y. Yang. Topological Solutions in the Self-Dual Chern-Simons Theory. preprint.
- [109] J. Terstriep. DICE: The Distributed Interactive Collaboration Environment. Technical report, National Center for Supercomputing Applications, Urbana-Champaign IL, 1991.
- [110] C. Upson, B. Brown, S. Dyer, D. Kamins, and J. Rasure. Future Directions in Visualization Software Environments. In *Panel Proceedings ACM SIGGraph'91*. Association for Computing Machinery, Chicago IL, 1991.
- [111] T. Vogel. Unbounded parametric surfaces of prescribed mean curvature. *Indiana Univ. Math. J.*, 31:281-288, 1982.
- [112] F. Wei. *The existence and topology of properly embedded minimal surfaces in R^3* . PhD thesis, University of Massachusetts at Amherst, 1991.
- [113] H. Wente. An existence theorem for surfaces in equilibrium satisfying a volume constraint. *Arch. Rational Mech. Anal.*, 50:139-158, 1973.

Figure 1

34a

Figure G 1

Ray Tracing Simulation
80° Scherk Twist Boundary

Scherk's Surface 90°

Possible helical deformation

Scherk's First Surface
90° twist boundary

$A \neq B$

Karcher's Surface
Possible 90° twist boundary

Figure 6.2

SCIENCE SCOPE

Material Science Initiative to Take a Step Forward

■ Next month, a key White House committee is expected to urge the federal government to launch a major multiyear national initiative in materials research with the potential to nearly double the estimated \$1.4 billion already spent each year.

Approval by the committee, the Federal Coordination Council for Science, Engineering, and Technology (FCCSET), would be an important first step toward the inclusion of a materials initiative in the 1993

budget early next year. Earlier FCCSET-backed projects include this year's much-heralded high-performance computing and science education initiatives.

A computer simulation of certain polymer surfaces.

The Bush Administration is considering pumping more funds into materials science after a series of reports over the past 2 years made an impressive case that the skills and wares of

materials researchers—the latter consisting of everything from Silly Putty to super-alloy turbine blades to ceramic-polymer composites in tennis rackets and aircraft—are critical to national well-being.

The embryonic initiative has at least one well-placed fan: presidential science adviser D. Allan Bromley, who has long talked about coordinating materials research. And researchers such as Bill Appleton, a materials research director at Oak

Ridge, couldn't be happier. The message that "materials are extremely important to everybody" finally seems to be getting across in Washington, Appleton says.

This short article from *SCIENCE* (5 April 1991) contains an image of Scherk's doubly-periodic minimal surface, produced in the GANG for the purposes of numerical simulation of an idealised grain boundary in a copolymer blend, annealed in a thin sheet. The interaction of theoretical mathematics, algorithm development, numerical simulation and computer graphics with applied physical science is encapsulated in this image (which is not properly credited as to its scientific meaning or provenance). Our proposed research has as one of its main purposes the facilitation of this crucial scientific communication, which is cross-disciplinary and involves simultaneous use of several modes of communication and thinking.

THE AMERICAN MATHEMATICAL MONTHLY

75

Volume 97, Number 8

October 1990

GEOMETRY

The cover illustration is a minimal surface discovered by David Hoffman (Amherst) and Hermann Karcher (Bonn). It is singly-periodic and behaves at infinity much like the union of helicoids. The image was made by James Riordan and Jim Hoffman (GANG, University of Massachusetts, Amherst)

Personnel Wholly or Partially Supported by this Grant

Ph.D. students

Bo Guan - Motion of Interfaces and curvature (working with Spruck) (May 28, 1989 - December 28, 1991)

Guoghang Sun - Motion of interfaces by curvature functions (working with Spruck) June 1, 1989 - December 28, 1991)

Edward Thayer - Scientific programming graphics development involving problems in differential geometry, and Ph.D research in minimal surface and beginning his Ph.D work. (working with D. Hoffman) (May 27, 1990 - May 25, 1991)

Wayne Rossman - Ph.D student involving problems in differential geometry in euclidean and hyperbolic space. (Working with D. Hoffman and W. Meeks, III) January 20, 1991 - May 25, 1991 and January 12, 1992 - May 23, 1992)

Scientific and Graphics Programmers and Researchers

James T. Hoffman - Graphics programming for all scientific projects, development of graphics interface for visual analysis of numerical simulations, development of algorithms for the analysis of data, graphics software development.

Pascal Romon - Graphics and scientific programming, assisted Jim Hoffman and worked intensively with David Hoffman on research projects.

Postdoctoral Research Associate

Fusheng Wei (6/1/91 - 5/31/92)

Research Visitors supported by this Grant (past 12 months).

Tom Ilmanen (Institute for Advanced Study)	November 15, 1991
Harold Rosenberg (University of Paris VII)	Oct. 1 - 11, 1991
Jean Taylor (Rutgers University)	September 27, 1991
Peter Laurence (Courant Institute)	May 22 - 23, 1991
Frank Baginski (George Washington University)	December 7 - 10, 1990
Meinhard Wohlgemuth (Bonn Germany)	July 29 - August 12, 1991
Yi Fang (Courant Institute)	August 14 - 16, 1991
Vitaly Volpert (Institute of Chemical Physics USSR Academics of Sciences)	April 10, 1991
Jonathan Goodman (Courant Institute)	March 26 & 27, 1991
Alexander Bobenko (Bonn Germany)	March 14 & 15, 1991
Brian Wetton (Courant Institute)	March 5 & 6, 1991
Silvio Levy (Princeton University)	February 28 - March 2, 1991
Eric Boix (Ecole Polytechnique, Paris)	January 3 - 20, 1991
David Chopp (University of California, Berkeley)	February 7 - 9, 1991
Pascal Romon (ENS and University of Paris VII)	February 24 - March 3, 1991
Charles Frohman (SUNY at Buffalo)	January 10 - 11, 1991

Hermann Karcher (University of Bonn, Germany)	March 8 - 27, 1991
Harold Ziegler (Ruhr-University, Bochum)	November 7 & 8, 1990
Michael Callahan (Harvard University)	November 1 - 3, 1990
Yisong Yang (University of New Mexico)	November 18 - 21, 1990
Laurent Capdequi (Ecole Polytechnique, Paris)	July 27 - August 29, 1991

Partial List of Other Research Visitors who visited this DOE supported facility. July 1, 1991 - December 31, 1991

Bill Goldman (University of Maryland)
 Conrad Pohlthier (University of Bonn, Germany)
 Nicholas Korevaar (University of Utah)
 Daniel Ruberman (Brandis University)
 Frank Morgan (Williams College)
 Michael Anderson (Stony Brook)
 Todd Kapitula (Brown University)
 Lucio Rodriguez (Inst. for Pure-Appl. Math. Rio de Janeiro, Brazil)
 James Riordan (University of Minnesota)
 Ivan Sterling (Berlin, University of Toledo)
 Eric Boix (Ecole Polytechnique)

In July, GANG also hosted a four hour workshop with live demonstrations and lectures, for 65 participants in the National Science Foundation (NSF) sponsored Regional Geometry Institute, "Geometry in the Machine Age."

Appendix

1. Geometry, Computation and Graphics Preprint List

Center for
Geometry, Analysis, Numerics, and Graphics

G.A.N.G.
University of Massachusetts at Amherst

APPENDICES

The "Geometry, Computation and Graphics" series II includes preprints and technical reports in Geometric Analysis, Numerical Computation, Scientific Graphics and Applications. (May 1990 - present)

1. M. Wohlgemuth.
Higher Genus Minimal Surfaces by Growing Handles out of a Catenoid.
2. A. Lifschitz.
Essential Spectrum and Local Stability Condition in Hydrodynamics.
3. L.C. Evans and J. Spruck.
Motion of Level Sets by Mean Curvature III.
4. D. Hoffman.
Some Basic Facts, Old and New, About Triply Periodic Embedded Minimal Surfaces.
5. D. Hoffman, H. Karcher and H. Rosenberg.
Embedded Minimal Annuli in R^3 Bounded by a Pair of Straight Lines.
6. B. Turkington, A. Eydeland, A. Lifschitz and J. Spruck.
Multiconstrained Variational Problems in Magnetohydrodynamics, I: Equilibrium.
7. A. Lifschitz, B. Turkington, J. Spruck, and A. Eydeland.
Multiconstrained Variational Problems in Magnetohydrodynamics, II: Slow Evolution.
8. D. Hoffman, H. Rosenberg, and Joel Spruck.
Boundary Value Problems for Surfaces of Constant Gauss Curvature.
9. J. Spruck and Y. Yang.
On Multivortices in the Electroweak Theory I: Existence of Periodic Solutions.
10. J. Spruck and Y. Yang.
On Multivortices in the Electroweak Theory II: Existence of Bogomol'nyi Solutions in R^2 .
11. D.M. Anderson, J. Bellare, J.T. Hoffman, D. Hoffman, J. Gunther, and E.L. Thomas.
Algorithms for the Computer Simulation of 2D Projection from Structures Determined by Dividing Surfaces.
12. D.Hoffman.
Computing Minimal Surfaces With and Without Conformal Representation.

13. B. Guan and J. Spruck.
Interior Gradient Estimates for Solutions of Prescribed Curvature Equations of Parabolic Type.
14. Y. Fang.
Complete Minimal Surfaces of Finite Total Curvature.
15. B.C. Morris.
Variational Study of Interstellar Magnetic Gas Clouds: Theory, Modeling, and Computation.
16. N. Korevaar and R. Kusner.
The Structure of Constant Mean Curvature Embeddings in Euclidean Three Space.
17. R. Kusner.
A Maximum Principle at Infinity and the Topology of Complete Embedded Surfaces with Constant Mean Curvature.
18. N.J. Korevaar, R. Kusner, W.H. Meeks, III and B. Solomon.
Constant Mean Curvature Surfaces in Hyperbolic Space.
19. J.T. Hoffman.
A Visual Programming System and Image Synthesis Application.
20. J. Spruck and Y. Yang.
Topological Solutions in the Self-Dual Chern-Simons Theory
21. B. Guan and J. Spruck.
Interior Gradient Estimates for Solutions of Prescribed Curvature Equations of Parabolic Type.
22. F. Wei.
The Existence and Topology of Properly Embedded Minimal Surfaces in R^3 .
23. N. Whitaker.
Numerical Solution of the Hele-Shaw Equations Using the Point Vortex Method.
24. W. Meeks, III, R. Earp, F. Brito, and H. Rosenberg.
Structure Theorems for Constant Mean Curvature Surfaces Bounded by a Convex Planar Curve.
25. W. Meeks, III and B. White.
Minimal surfaces bounded by convex curves in parallel planes.
26. W. Meeks, III, and C. Frohman.
The Topological Uniqueness of Complete One-ended Minimal Surfaces and Heegaard Surfaces in R^3 .
27. W. Meeks, III and H. Rosenberg.
The Geometry of Periodic Minimal Surfaces.

28. W. Meeks, III and B. White.
The Space of Minimal Annuli Bounded by an Extremal Pair of Planar Curves.
29. W. Meeks, III and C. Frohman.
The Ordering Theorem for the ends of Properly Embedded Minimal Surfaces.
30. W. Meeks, III and H. Rosenberg.
The Geometry and Conformal Structure of Properly Embedded Minimal Surfaces of Finite Topology in R^3 .
31. F. Wei.
The Existence and Uniqueness of Properly Embedded Doubly Periodic Minimal Surfaces.
32. H. Rosenberg.
Hypersurfaces of Constant Curvature in Space Forms.
33. B. Guan and J. Spruck.
Boundary Value Problems on S^n for Surfaces of Constant Gauss Curvature.
34. F.E. Baginski.
The Computation of One-parameter Families of Bifurcating Elastic Surfaces.
35. J. Hoffman.
MESH Manual.
36. D.M. Oliver.
The GANG Video Distributed Video Services.
37. D.M. Oliver.
Scientific Visualization: Work in Progress at GANG.
38. D.M. Oliver.
The GANG Message Service.

To Appear:

D.Hoffman.
"Analytic and Quantitative Data for the Scherk Family of Minimal Surfaces".

M. Callahan, D. Hoffman, H. Karcher.
"Singly Periodic Minimal Surfaces in Variet Under a Sorew Motion".

END

**DATE
FILMED**

11 / 2 / 92