ABSTRACT

Recent research findings in our laboratory have determined that organized large scale vortex structures can play a dominate role in the particle dispersion process in free shear flows. The effect of these large scale vortices on the particle dispersion process is especially significant if the time scales associated with the aerodynamic response time of the particles and the large scale vortices are of the same order. Our recent experimental results demonstrate that if this time scale condition exists the particle dispersion process will be dominated by the large scale vortices. Moreover, the particle dispersion rate can be significantly affected if forcing techniques are imposed to control the large scale structure interactions.

Our previous research has focused on the relatively simple case of the plane mixing layer which has a time average vorticity in only one direction and consequently generates vortex structures of like sign. The proposed research will extend our present research thrust to a more complex free shear flow environment, the plane wake. The specific flow to be investigated will be the plane wake produced by a splitter plate with a variable thickness trailing edge. This flow is of interest from both scientific and technological viewpoints. The plane wake flow has time average vorticity in two directions and thus produces large scale vortex structures of differing signs. These vortex structures may potentially engage in various interactions depending on the signs of the interacting vortices. In addition, the maximum turbulence levels in this flow are significantly influenced by the scale ratio between the initiating boundary layers and the splitter plate trailing edge thickness. Thus, the particle dispersion process in this flow may be significantly more complex since both the length scale parameter and the time scale parameter already studied will be operative

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

870

in this flow.

We propose to investigate this two-phase flow using both experimental and numerical approaches. The experimental techniques will involve image analysis of flow and particle visualization studies supported by three-dimensional point laser velocimetry measurements. The numerical technique will employ discrete vortex techniques for modeling the flow coupled with the use of Lagrangian particle trajectory analysis.

The outcome of this proposed research will be an improved understanding of the particle dispersion process in a more complex flow environment which has close ties to technologically important flows. Improved physical models as well as the potential for developing control or design strategies for affecting particle dispersion are also probable benefits of this research program.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

I. INTRODUCTION

Over the last five years our research group at Washington State University has been involved in a program aimed at developing a new physical concept concerning the turbulent dispersion of fine particles. This physical concept assumes that large scale vortex structures can play a dominate role in the particle dispersion process in free shear flows. This research program has been pursued using both time dependent numerical techniques and focused laboratory experiments. Our initial research efforts have primarily concentrated on particle dispersion in plane mixing layers. We now propose to extend this work to a more complex flow situation; namely, plane wakes. A summary of our recent findings concerning the particle dispersion process in mixing layers follows. A detailed discussion of our proposed research begins in Section II.

For the past 1.5 years our laboratory experimental program has been supported by the Department of Energy under grant number DE-FG06-86ER13567. This research program has focused on the particle dispersion process in plane mixing layers. The experiments have involved solid spherical particles in an air flow. The high density ratio between the particles and the air substantially simplifies the force interaction analysis. This feature eases the computational requirements for the supporting numerical simulation efforts (Please see section III B. for details). In addition, our experiments have primarily been conducted with low particle concentration levels such that the particles are assumed to have a negligible effect on the flow field (one way coupling).

Plane mixing layers possess a simple time average geometry and are dominated by time average vorticity aligned in the spanwise direction with only one sign. This flow had been extensively studied by numerous previous investigators and has been found

to produce quasi-deterministic large scale vortex structures. A review of the work on organized vortex structures in mixing layers can be found in Ho and Huerre (1984).

The results of our present experimental investigation demonstrate that large scale vortex structures play an important role in the particle dispersion process (Kamalu, et al., 1987, 1988 and Wen, et al., 1987). This finding agrees with our previous conjectures (Crowe, et al., 1983, 1985) and our previous numerical simulations (Crowe, et al., 1985, Gore, et al., 1985, Chung and Troutt, 1988). A review article by Crowe, Chung and Troutt (1988) summarizing these findings and their connection to previous work has been submitted for publication.

In our proposed physical model it was suggested that if the aerodynamic response time of the dispersing particles, τ_A , and the characteristic time, τ_f , of the organized vortex structures are of the same order then enhanced particle dispersion might occur. The aerodynamic response time, τ_A , is defined as the time necessary for a solid particle released from rest to achieve $(1 - 1/e)$ or 63 percent of the velocity of a surrounding steady fluid flow. For Stokes flow it can be shown that $\tau_A = \frac{\rho_p d_p^2}{18\mu}$, where ρ_p is the particle density, d_p is the particle diameter, and μ is the fluid viscosity. The characteristic time, τ_f , of the large scale vortex structures is estimated as $\tau_f = \frac{\delta}{\Delta U}$, where δ is a measure of the local mixing layer thickness and ΔU is the velocity difference across the layer. A ratio of these two time scales is defined as the Stokes Number, $St = \tau_A/\tau_f$.

A schematic illustrating this physical concept is shown in figure 1. Particles with very small time scale ratios are shown to follow closely the fluid particle mixing and to distribute themselves approximately homogenously throughout the mixing layer. Conversely, particles with very large time scale ratios have insufficient time to respond to the

large scale structures and are essentially unaffected by the turbulence. However, particles with intermediate time scale ratios are expected to be strongly affected by the large scale vortex structures. These particles are too large to closely follow the small scale turbulent fluctuations and therefore respond predominately to the large scale vortices. These particles may be centrifuged to the outer edges of the large scale vortices or even beyond the boundaries of the mixing layer.

Support for this physical model concerning the particle dispersion process in plane mixing layers has been provided by both our numerical and experimental studies. Figure 2 shows the predictions of particle spatial distribution using two-dimensional discrete vortex simulations for dispersion in a plane mixing layer, from Chein and Chung (1988). The particle dispersion patterns appear to qualitatively support the physical model. Recently acquired experimental results in the form of visualizations of the flow and particle dispersion patterns provide even further support of the concept.

Figure 3 shows smoke-wire flow visualizations of a mixing layer under both natural and subharmonically forced conditions. The smoke particles used to visualize the flow are nominally 1-5 micron in diameter and can be assumed to have very small time scale ratios. The smoke particles thus follow the flow fluctuations closely and disperse with the fluid particles. The high speed stream is on the top half of the pictures. The velocity ratio for this flow is $U_H/U_L = 2.5$. Acoustic forcing at the first subharmonic of the initial vortex passage frequency stimulates the first vortex pairing interaction thus increasing the initial spreading rate of the mixture layer. The forcing also increases the organization of the resulting downstream vortex structures.

Instantaneous dispersion patterns for 40 micron diameter spherical glass particles

are shown in figures 4a and b for both naturally evolving and subharmonically forced mixing layers. Light illumination for both the smoke and particle visualizations are from diffuse flash sources. These particles have time scale ratios with values of order unity. The experimental patterns confirm our speculations concerning the centrifuging of the particles to the outer edges of the vortex structures. This centrifuging effect becomes even more apparent for the subharmonically forced situation where the resulting paired vortex structures are highly regular.

Laser velocimetry time average velocity profiles for the flow and the 40 micron particles are compared in figures 5 and 6. The results for the streamwise velocity component, presented in figure 5, show that particle velocities gradually approach the fluid velocities until no major qualitative or quantitative differences exist between the velocity profiles for the particles and fluid.

A significantly different situation however is observed for the time average cross-stream component of the flow and particle velocity shown in figure 6. These measurements reveal an interesting qualitative difference between the flow and particle cross stream velocity profiles. The cross-stream velocities of the fluid tend to be negative (downward) on the high speed side of the flow and positive (upward) on the low speed side indicating, as expected, a net entrainment of fluid particles into the mixing layer. The cross-stream solid particle velocities, however, display exactly the opposite trend, indicating that the solid particles are being predominately flung outward from the center of the mixing layer. This time average result strongly supports our physical model and our interpretations of the visualization results. A similar result has also been observed by Kobayashi, et al. (1987) for a vertically oriented mixing layer flow.

Other recent experimental efforts by our group concerning particle dispersion in the planar mixing layer have involved evaluating the downstream development of particle concentration levels using interdata times from laser velocimetry measurements. The results of the laser velocimetry measurements for both naturally evolving and forced mixing layer flows have also been enlightening concerning the particle dispersion process. Specific details concerning these results can be found in Kamalu, et al. (1988) which is included as Appendix A.

Further on-going experiments in our laboratory on mixing layer flows are focusing on the instantaneous three-dimensional nature of the particle dispersion process, and the effects of controlled forcing on the particle dispersion process.

II. SUBJECT OF PROPOSED INVESTIGATION

II. A. General Features of the Plane Wake Flow

The subject of the proposed investigation involves a direct extension of our previous efforts concerning particle dispersion in plane mixing layer flows to the more complex flow environment associated with plane wakes. Plane wakes are categorized as free shear flows like mixing layers and jets so the dynamics of these flows can be expected to be strongly influenced by large scale vortical flow structures. In fact, recently published results on plane wake flows (Wynanski, et al., 1986) demonstrate that organized vortical structures play an important role in their development.

Plane wake flows are significantly more complex than mixing layer flows because the time average vorticity in the plane wake velocity profile contains both positive and negative values. This time average feature of the plane wake vorticity or velocity profiles

END

**DATE
FILMED**

11 / 16 / 93