DOE/ER/53176--8

DE92 040854

Theoretical Plasma Physics

Progress Report

November 1, 1991 - October 31, 1992

Allen H. Boozer and George M. Vahala

College of William and Mary
Williamsburg, VA 23185

May 1992

Prepared for the U.S. Department of Energy under Grant Number
DE-FG05-84ER53176

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Received by OSTI

SEP 11 1992

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Theoretical Plasma Physics

During the past year, we have worked in the areas of classical and anomalous transport, three dimensional equilibria, divertor physics, as well as diagnostic techniques using waves. Although much work has been done on classical transport, the validity of the guiding center drift equations, which are the basis of much of the theory, has received little attention. We have an ongoing research program to understand the limitations of the drift approximation. We have published our work on three dimensional equilibria which shows that quasi-helical symmetry is broken in third order in the inverse aspect ratio. We have also published our work on the modification of the current profile due to tearing modes. This work is relevant to the maintenance of a steady-state tokamak by the bootstrap current. Divertor physics is a primary area that required development for ITER. One of the few methods by which the physics of the divertor can be modified or controlled is magnetic perturbations. We have been studying the effect of magnetic perturbations on the divertor scrapeoff layer in collaboration with Hampton University. The evolution of magnetic field embedded in a moving plasma is a dynamics problem of potential importance in which we are working. Renormalization techniques gave important insights first in the theory of phase transitions. The applications of these techniques has extended to many areas of physics, including turbulence in fluids and plasmas, an area in which we have a continuing research program. Essentially no diagnostics for magnetic fluctuations inside a fusion-grade plasma exist. We have a collaborative program with Old Dominion University and the Princeton Plasma Physics Laboratory to develop such a diagnostic based on the conversion of electromagnetic waves from the ordinary to the extraordinary mode.

Table of Contents

I.	The Drift and the Exact Particle Hamiltonians	3
II.	Magnetic Field Strength in Three Dimensional Equilibria	7
III.	Modification of the Current Profile by Tearing Modes	9
IV.	Divertor Calculations	10
V.	Magnetic Fields in Moving Plasmas	13
VI.	Renormalization Group Theory Techniques in Turbulence	15
VII.	Magnetic and Density Fluctuations in Tokamak Plasmas	17
	References	19
	List of 1991-1992 Publications	
	Allen H. Boozer	23
	George M. Vahala	24

I. The Drift and the Exact Particle Hamiltonians

The guiding center drift equations, and not the exact equations of motion, are the basis for most of the theory of particle confinement in asymmetric devices, such as stellarators and rippled tokamaks. There are three reasons: the drift equations can be integrated much faster, the constraints associated with magnetic moment conservation are more obvious in the drift equations, and the drift equations require far less information about the magnetic configuration. Remarkably, the conditions for the trajectories of the drift equations to adequately approximate the true particle trajectories have received only limited consideration.

The relation between the drift and the exact particle trajectories will be of great importance to the success of the ITER tokamak. The confinement of the thermonuclear alphas is critical to ignition, and the studies of alpha confinement in the presence of toroidal ripple have all been made in the drift approximation. Of all of the particles of interest to fusion, the drift approximation is most questionable for fusion alphas due to their large gyroradii, $\rho_\alpha = 0.27 \text{ meters/B}_{\text{tesla}}$, and the large number of toroidal transits that they make before slowing down. For example, the conservation of the magnetic moment, which is a necessary condition for the validity of the drift approximation, is sensitive to the variations in field strength along the field lines. The shortest wavelength variation of this type is due to the toroidal ripple which has a wavelength in a reactor of about 2 meters. Fusion alphas have a slowing down time that is of order ω_c/ϵ^4 with ω_c the cyclotron frequency and $\epsilon \equiv \rho_\alpha/\lambda$ with λ the shortest scale over which the magnetic field varies, $\epsilon \approx 0.14$. Unfortunately, the standard derivation of the drift equations, which is due to Alfven,¹ is only valid for a time ω_c/ϵ .

The distinction between the exact and the drift trajectories is also important for understanding the transport properties of quasi-helical stellarators.^{2,3} The magnetic field strength in a quasi-helical stellarator depends on only two variables and has the form $B(\psi, \theta - N\phi)$ with ψ the toroidal flux, θ a poloidal magnetic angle, ϕ a toroidal magnetic angle, and N the number of periods of the stellarator. The properties of the drift

trajectories are determined by the field strength alone,⁴ and the quasi-helical form for the field strength implies the existence of a conserved canonical momentum, P_h . The exact equations of motion depend not only on the field strength but also on the full metric for the magnetic coordinates. In a quasi-helical stellarator the metric tensor depends in a complicated way on all three magnetic coordinates, ψ , θ , and ϕ , so the canonical momentum P_h is not an invariant of the exact Hamiltonian. We have shown that for curl-free magnetic field, the lowest order in ϵ [the gyroradius divided by the system size] correction to the drift equations already depends in a generic way on the metric, so one can use perturbed drift equations to study the invariance properties of P_h . Although quasi-helical symmetry is broken by terms of order ϵ to the drift equation, our preliminary results indicate that the primary effect is an oscillation in the value of the canonical momentum P_h rather than a total loss of this invariant. The importance of quasi-helical stellarators that have a helical invariant P_h is that their neoclassical transport and high energy particle confinement properties resemble those of an axisymmetric tokamak,⁵ but they have the stellarator advantages of intrinsic steady state (no need for current drive) and the absence of disruptions. Despite the potential importance of quasi-helical symmetry to the fusion program, we believe that we are the only group in the world working on the fundamental theory of its transport properties.

The procedure that we have used in these studies is to write the exact Hamiltonian H_e for the particle trajectories, which has six canonical coordinates, as the sum of a guiding center Hamiltonian H_g , which depends on only four canonical coordinates, and a perturbation $\tilde{H}_e$, which is chosen to be as small as possible,

$$H_e(\mu, \theta_g, \mathbf{P}, \mathbf{Q}) = H_g(\mu, \mathbf{P}, \mathbf{Q}) + \tilde{H}_e(\mu, \theta_g, \mathbf{P}, \mathbf{Q}).$$

The four canonical variables of the guiding center Hamiltonian are $(\mathbf{P}, \mathbf{Q})$, the magnetic moment is $\mu = \pi v_\perp^2 / 2B$, and the canonically conjugate gyrophase is θ_g . Kruskal has shown⁶ that the breaking of magnetic moment conservation is slower than ϵ^N / ω_c with N any integer. Since $d\mu/dt = -\partial \tilde{H}_e / \partial \theta_g$, the conservation properties of the magnetic moment imply that

the perturbation $\tilde{H}_\epsilon$ can be chosen so it scales as $\exp(-c_0/\epsilon)$ with c_0 a constant.

The Hamiltonian H_d that is equivalent to the standard drift equations of Alfvén differs from the guiding center Hamiltonian H_g . By the guiding center Hamiltonian, we mean the most accurate Hamiltonian that has only four canonical variables. One can write

$$H_g(\mu, \mathbf{P}, \mathbf{Q}) = H_d(\mu, \mathbf{P}, \mathbf{Q}) + \tilde{H}_g(\mu, \mathbf{P}, \mathbf{Q}).$$

The perturbation $\tilde{H}_g$ has a magnitude of order ϵH_d . The importance of the term $\tilde{H}_g$ is made clearer if the one uses the magnetic coordinates (ψ, θ, ϕ) that were introduced by Boozer⁷ to study particle drift motion. He found⁴ that in these coordinates the standard drift equations, which are represented by the Hamiltonian H_d , depend on the complexity of the magnetic field structure only through the magnetic field strength $B(\psi, \theta, \phi)$. We find that both the exact Hamiltonian H_ϵ and the lowest order in ϵ expression for $\tilde{H}_g$ depend on the metric tensor g_{ij} of the transformation from ordinary spatial coordinates $\mathbf{x}$ to magnetic coordinates (ψ, θ, ϕ) .

To understand these results, consider a curl-free magnetic field that is written in the forms $\mathbf{B} = \nabla\psi \times \nabla\theta + \nabla\phi \times \nabla\chi(\psi)$ and $\mathbf{B} = \mu_0 G \nabla\phi$ given by Boozer.⁷ One can show that ψ is the toroidal flux, $-\chi$ is the poloidal flux, θ is the poloidal angle, ϕ is the toroidal angle, and G is the current, in ampere-turns, flowing in the external coils. The rotational transform is $\iota = d\chi/d\psi$. It is useful to define $\hat{\mathbf{e}}_\theta \equiv \nabla\theta - \iota \nabla\phi$, so $\mathbf{B} = \nabla\psi \times \hat{\mathbf{e}}_\theta$. The metric tensor g_{ij} can be given in terms of $g^s \equiv \hat{\mathbf{e}}_\theta \cdot \hat{\mathbf{e}}_\theta$, $g^\psi \equiv \nabla\psi \cdot \nabla\psi$, $g^c \equiv \nabla\psi \cdot \hat{\mathbf{e}}_\theta$, the field strength B^2 , and the external current G . The components g^ψ , g^s , and g^c are not independent of the field strength; they satisfy the identity $B^2 = g^\psi g^s - g^{c2}$. It is a general result of canonical transformation theory that if the position vector $\mathbf{x}$ is given as a function of coordinates ξ^i , the canonical coordinates $\mathbf{x}, \mathbf{p}$ of a Hamiltonian can be canonically transformed to coordinates (ξ^i, P_i) by letting $P_i = \mathbf{p} \cdot \partial \mathbf{x} / \partial \xi^i$. The canonical momentum $\mathbf{p}$ has the form $\mathbf{p} = m\mathbf{v} + e\mathbf{A}$ with e the particle charge. The linear momentum $m\mathbf{v}$ can be written in a covariant representation

$$m\mathbf{v} = v_{\parallel}\mathbf{B} + s\hat{\mathbf{e}}_s - p_s\nabla\psi,$$

and the theory of coordinate transformations implies $\nabla\xi^i\cdot\partial\mathbf{x}/\partial\xi^j=\delta_{ij}$. One can then show that the exact Hamiltonian, $H_e=mv^2/2$, has canonical coordinates (s,Θ,φ) with $\Theta=\theta+p_s/e$ and canonical momenta $P_{\Theta}=e(\psi+s)$, $P_{\varphi}=(\mu_0 G/B)mv_{\parallel}-e\chi$, and p_s . In these canonical coordinates

$$H_e(P_{\Theta},\Theta;P_{\varphi},\varphi;p_s,s) = \frac{1}{2m} \frac{B^2}{(\mu_0 G)^2} [P_{\varphi}+e\chi+s]^2 + \frac{1}{2m} [e^2 g^s s^2 - 2eg^c s p_s + g^v p_s^2].$$

If the magnetic field is spatially constant, $\Psi\equiv P_{\Theta}/e$ and Θ are constants of the motion and are the coordinates of the center of the circle about which a particle located at (ψ,θ) gyrates, the coordinate φ advances due to the motion along the field lines, and the canonical coordinates p_s and s oscillate at the gyrofrequency.

To obtain Hamiltonians that are in the forms that were discussed earlier, we use a generating function to effect a transformation from the (p_s,s) coordinate pair to the (μ,θ_g) pair. This canonical transformation makes changes of order ε^2 to the $(P_{\Theta},\Theta;P_{\varphi},\varphi)$ canonical coordinates, which become the $(\mathbf{P},\mathbf{Q})$ coordinates. If one is retaining only the lowest order in ε expression for $\tilde{H}_g$, one need not make a distinction between $(\mathbf{P},\mathbf{Q})$ and $(P_{\Theta},\Theta;P_{\varphi},\varphi)$. The expressions that we obtain for H_d and the lowest order expression for $\tilde{H}_g$ are

$$H_d = \frac{1}{2m} \frac{B^2}{(\mu_0 G)^2} (P_{\varphi} + e\chi)^2 + \mu B,$$

which is the standard expression for the drift Hamiltonian, and $\tilde{H}_g = (P_{\varphi} + e\chi) \Delta$ with

$$\Delta = - \frac{\mu B}{2e(\mu_0 B)^2} \left\{ \frac{dt}{d\psi} g^{\psi} + \left[\frac{\partial}{\partial\Theta} + \frac{\partial}{\partial\varphi} \right] g^c - \frac{g^c}{g^s} \left[\frac{\partial}{\partial\Theta} + \frac{\partial}{\partial\varphi} \right] g^s \right\}.$$

The drift Hamiltonian H_d is independent of the metric, which has the important consequence that quasi-helical symmetry is possible. The exact Hamiltonian depends on all components of the metric as does the guiding center Hamiltonian H_g . Even the lowest order expression for the perturbation $\tilde{H}_g$ depends on all components of the metric. The dominant contributions to the breaking of the helical invariant P_h of the drift Hamiltonian by the exact Hamiltonian can, therefore, be studied using the lowest order expression for the perturbation $\tilde{H}_g$.

An article "The Exact and Drift Hamiltonian," which discusses this work will appear in the August 1992 issue of the Physics of Fluids B. The authors are Qun Yao, a graduate student in physics, and Allen Boozer.

II. Magnetic Field Strength in Three Dimensional Equilibria

Our work on three-dimensional equilibria reached a state of completion during the past year with the publication of two papers in the Physics of Fluids B.^{9,3} The authors were David Garren, who completed a doctorate in physics and accepted a postdoctoral position in plasma physics at the Naval Research Laboratory, and Allen Boozer. In these papers we derived a general expression for the magnetic field strength in an equilibrium plasma. The expression is an expansion in the inverse aspect ratio ϵ about the magnetic axis $r_0(\psi)$ of the equilibrium. The general form is

$$B(\psi, \theta, \varphi) = B_0(\varphi) [1 + \kappa(\varphi)x + F(\theta, \varphi)\psi + C(\theta, \varphi)\psi^{3/2} + \dots].$$

The field strength along the magnetic axis is $B_0(\varphi)$ and $\kappa(\varphi)$ is the curvature of axis. The distance $x(\psi, \theta, \varphi) = \psi^{1/2} \rho(\varphi) \cos[\theta - \alpha(\varphi)]$ is the approximate distance to the magnetic surface in locally Cartesian coordinates that use $(\hat{\kappa}, \hat{z}, \hat{b})$ as the $(\hat{x}, \hat{y}, \hat{z})$ unit vectors with $\hat{b}$ the tangent and $\kappa \hat{\kappa}$ the curvature vector of the axis and $\hat{z} = \hat{\kappa} \times \hat{b}$. The functions $\rho(\varphi)$ and $\alpha(\varphi)$ can be chosen freely except that ρ must be positive definite. The function $F(\theta, \varphi)$ has the form

$$F(\theta, \varphi) = W(\varphi) + \tilde{F}(\varphi) \cos[2\theta - \alpha_2(\varphi)].$$

The three functions $W(\varphi)$, $\tilde{F}(\varphi)$, and $\alpha_2(\varphi)$ can be chosen freely. A large positive $W(\varphi)$ will give a magnetic well. The function $C(\theta, \varphi)$ has the general form

$$C(\theta, \varphi) = C_1(\varphi) \cos[\theta - \alpha_1(\varphi)] + C_3(\varphi) \cos[3\theta - \alpha_3(\varphi)].$$

The two functions $\alpha_1(\varphi)$ and $\alpha_2(\varphi)$ can be chosen freely, but only one of the functions $C_1(\varphi)$ and $C_3(\varphi)$ can. The various powers of the toroidal flux ψ in the expression for $B(\psi, \theta, \varphi)$ can be taken as an expansion in the inverse aspect ratio ϵ . The toroidal flux ψ is $B_0 \pi r^2$ with r an average radius of a magnetic surface, so the inverse aspect ratio, $\epsilon \equiv r/R$ with R the major radius, obeys $\epsilon \propto \psi^{1/2}$.

The general equation for the magnetic field strength $B(\psi, \theta, \varphi)$ demonstrates that quasi-helical symmetry,² $B(\psi, \theta - N\varphi)$, is broken by terms of third order in the inverse aspect ratio ϵ . To obtain quasi-helical symmetry through second order, one makes the choices of free functions $\alpha(\varphi) = N\varphi$, $\rho(\varphi) \propto 1/\kappa(\varphi)$, $\alpha_2(\varphi) = 2N\varphi$, $W(\varphi)$ constant, and $\tilde{F}(\varphi)$ constant. The third order, $\psi^{3/2}$, terms break the quasi-helical symmetry, though we have shown that one can achieve exact quasi-helical symmetry on a single ψ surface. For example one can obtain exact quasi-helical symmetry on the surface ψ_0 through ϵ^3 order by letting $\rho(\varphi) = \rho_0(\varphi) + \rho_1(\varphi)$ with $\rho_0(\varphi) \propto 1/\kappa(\varphi)$ and $\rho_1(\varphi) = -\psi_0 C_1(\varphi)/\kappa(\varphi)$. The other third order functions are chosen to have the forms $\alpha_1(\varphi) = N\varphi$, $\alpha_2(\varphi) = 3N\varphi$, and $C_3(\varphi)$ constant.

Our work is related to earlier expansions around the magnetic axis developed by Mercier,⁹ Lortz and Nührenberg,¹⁰⁻¹² and Solov'ev and Shafranov.¹³ These authors solved for the magnetic field as a function of the ordinary spatial position $\mathbf{x}$. We solve for the position and the field strength as a function of the magnetic coordinates ψ, θ, φ , which greatly simplifies the determination of the properties of equilibria. A related inverse method, though not in the appropriate magnetic coordinates for transport studies, was developed by Lortz and Nührenberg.¹⁰

The methods that we have developed can be used to clarify the types of three dimensional equilibria that are available to the fusion program. One would like an equilibrium to have perfect confinement of drift orbits, a magnetic well, high local shear in regions of bad curvature, a small bootstrap current, and a small Shafranov shift. The class of equilibria that have these features can be defined using our methods.

III. Modification of the Current Profile by Tearing Modes

Investigations of the steady-state maintenance of the tokamak current by a combination of the bootstrap current and tearing modes has been brought to a state of completion by the publication of a paper in the Physics of Fluids B.¹⁴ The paper was by Rick Weening, who has completed a doctorate in physics and taken a postdoctoral position in fusion plasma physics in Lausanne, and Allen Boozer.

Our work was based on a code that solves the diffusion equation for magnetic flux that is associated with the Ohm's law

$$\mathbf{E} \cdot \mathbf{B} = \eta \mathbf{j} \cdot \mathbf{B} - \nabla \cdot [\lambda \nabla (\mathbf{j} \cdot \mathbf{B} / B^2)].$$

This is the general differential form for Ohm's law in the presence of unresolved MHD activity.¹⁵ In the code λ is made large in regions of the plasma that are occupied by magnetic islands. These regions are determined by solving for the Δ' of tearing mode theory¹⁶ and allowing the regions of large λ to increase in width at a rate proportional to $\eta \Delta'$ as would be expected in the so-called Rutherford regime of tearing mode growth.¹⁷ That is, if Δ' is positive, the width of a region in which λ is large becomes wider; if Δ' is negative the width becomes narrower until it reaches zero width.

The code has been used to study the accessibility of various tokamak current profiles and the potential for the bootstrap current to make the tokamak self-maintaining. The paper of Cheng, Furth, and Boozer¹⁸ mapped the regions of internal inductance ℓ_i and q space in which tokamaks can be

stable to tearing modes. This map has been used to understand disruptions in JET.¹⁹ The code allows us to investigate the time evolution of tokamak discharges through ℓ - q space and address a number of issues associated with tokamak disruptions.

A bootstrap current as large as 80% of the total toroidal current has been reported by the JT60 group.²⁰ The bootstrap current creates no poloidal flux and by itself cannot maintain the tokamak current,²¹ but the tearing mode relaxation of the hollow current profiles associated with a strong bootstrap effect does create poloidal flux.²² Our code has found current profiles that can be maintained by tokamak plasma through the combination of the bootstrap effect plus tearing modes. In other words, the code is calculating current profiles that are consistent with self-sustainment of the tokamak current.²²

IV. Divertor Calculations

The prediction and preferably the control of the width of the scrapeoff layer of a divertor is a central issue in the design of ignition tokamaks such as ITER. Divertor simulations generally assume that the plasma is bounded by a sharp separatrix, which is correct only if the toroidal symmetry is perfect. Toroidal asymmetries in the magnetic field cause the last confining magnetic surface to lie inside the ideal separatrix and create a layer of open magnetic field lines that lies between the last confining magnetic surface and the ideal separatrix. The width of this layer, which we call the stochastic layer, is approximately proportional, near the X-point, to the square root of the toroidally asymmetric perturbation.²³ It is important to know the effects of naturally occurring perturbations, such as toroidal ripple or MHD modes, on the design of divertors. The introduction of magnetic perturbations may allow an optimization of the width of the divertor scrapeoff layer and a control of the heat deposition on the divertor plates. Our work in this area is an ongoing collaboration of Allen Boozer with Alkesh Punjabi and Arun Verma of Hampton University.

In the simulations that assume a sharp separatrix, the width of the scrapeoff layer is determined by the assumed diffusion coefficient, which

is usually of order $1 \text{ m}^2/\text{s}$. The field lines of the scrapeoff layer strike the divertor plates in a stripe with a width that is simply determined by axisymmetric magnetics.

If the magnetic perturbations are sufficiently large, the scrapeoff layer will be dominated the stochastic layer. The field lines of the stochastic layer strike the divertor plate in a stripe that is in proportion much narrower (about a factor of 6) than is the divertor stripe filled by the field lines of an axisymmetric scrapeoff layer. The greater width of the scrapeoff layer in comparison to the divertor stripe has important implications for the design of divertors.

The basic features of magnetic field line motion near a separatrix are generic and can be efficiently investigated using an area preserving map. Pomphrey and Reiman²⁴ have made related calculations using wires to model a divertor configuration instead of a map. The simplest divertor map has a single positive parameter k ,

$$\begin{aligned}x_{n+1} &= x_n - k y_n (1-y_n) \\ y_{n+1} &= y_n + k x_{n+1},\end{aligned}$$

an O-point at $x=0$, $y=0$, and an X-point at $x=0$, $y=1$. The map provides a good representation of the magnetic field lines of a single-null tokamak divertor. Toroidal asymmetries can be simulated using k or by additional terms.²⁵ Whether toroidal asymmetries are simulated using k or other representations does not make a qualitative difference. The smaller k , the closer the last confining magnetic surface passes to the X-point. For $k=0.6$, the closest approach is approximately $x=0$, $y=0.9971$, and for $k=0.9$, the closest approach is approximately $x=0$, $y=0.854$. Each iteration of the map corresponds to a toroidal angular advance ζ_0 . Near the O-point the rotational transform of the field lines per iteration $1/q_I$ is

$$\sin(2\pi/q_I) = k (1-k^2/4)^{1/2}.$$

One obtains a transform of roughly unity near the axis if 10 iterations of the map constitute a toroidal circuit, $\zeta_0 = 2\pi/10$.

The width of the stripe on the divertor plates is uniquely determined

by the stripe formed by field lines as they cross the plane $y=1$, which passes through the X-point. The toroidal location ζ at which field lines cross the x - ζ plane is determined using a continuous analogue to the discrete map. Letting $\psi = \zeta/\zeta_0$, $0 \leq \psi \leq 1$, the (area preserving) continuous analogue is

$$\begin{aligned} x(\psi) &= x_n - k y_n (1-y_n) \psi \\ y(\psi) &= y_n + k x(\psi) \psi. \end{aligned}$$

The behavior of field line trajectories that cross the last confining surface at a random point can be assessed by adding a small area-expanding step to each iteration of the discrete map, $x_{\text{new}} = (1 + R\delta)x_{\text{old}}$ with R a random number between 0 and 1 and δ the expansion coefficient. For $k=0.6$, we find that for $\delta \ll 10^{-13}$ the strike points form helical stripes that are independent of δ . For larger values of δ the strike points form a single stripe, which has a width that scales as $\delta^{1/4}$.

The open magnetic field lines of the stochastic layer, which lies between the last confining surface and the unperturbed separatrix, enter the region from one divertor plate and leave striking another divertor plate after an integer number of complete poloidal circuits N_p . The discrete stripes on the divertor plates correspond to the number of poloidal circuits made by the open field lines. The width of the stripes scales as $1/N_p^2$. The open field lines that approach the last confining surface make many poloidal circuits before striking a divertor plate in a stripe that is much narrower than the stochastic layer ($\approx 15\%$) but nonetheless of non-zero width. An open field line that makes many poloidal circuits in going from one divertor plate to another in essence densely fills the stochastic layer except for the regions in the stochastic layer that are occupied by magnetic islands.

The complicated filling of the stochastic region by the open field lines gives an associated complexity to the ambipolar electric potential that exists along any open magnetic field line that is embedded in a plasma. Although we have not calculated the particle transport from the

EXB drifts associated with the ambipolar potential, we expect a diffusive effect that would smear the stripe structures.

V. Magnetic Fields in Moving Plasmas

Fusion plasmas have a very low electrical resistivity, and as a first approximation they and the associated magnetic fields evolve ideally. A magnetic field $\mathbf{B}(\mathbf{x},t)$ evolves ideally if its time dependence is given by

$$\partial \mathbf{B} / \partial t = \nabla \times (\mathbf{u} \times \mathbf{B}).$$

The velocity of the magnetic field lines is $\mathbf{u}(\mathbf{x},t)$.

Using methods that date back to Lundquist,²⁶ one can show that the vector potential of an ideally evolving field, obeys

$$\mathbf{A} = \sum A_i(\xi) \nabla \xi_i.$$

The ξ coordinates are defined by the transformation equations $\mathbf{x}(\xi,t)$ with

$$(\partial \mathbf{x} / \partial t)_\xi = \mathbf{u}(\mathbf{x},t)$$

and the arbitrarily specified initial condition $\mathbf{x}(\xi,0)$. The choice $\mathbf{x}(\xi,0) = \xi$ makes ξ the Lagrangian coordinates of the magnetic field. By taking a curl of the vector potential to obtain the magnetic field $\mathbf{B}$ and a curl of the magnetic field to obtain the current $\mathbf{j}$, Allen Boozer demonstrated, in a paper that will appear in the *Astrophysical Journal*,²⁷ that the scalar quantities B^2 , j^2 , and $\mathbf{j} \cdot \mathbf{B}$ can be evaluated given the initial magnetic field $\mathbf{B}(\mathbf{x},0)$ and the metric tensor

$$g_{ij} = (\partial \mathbf{x} / \partial \xi_i) \cdot (\partial \mathbf{x} / \partial \xi_j).$$

For example, the magnetic field strength is given by

$$B^2 = \sum b^i(\xi) g_{ij} b^j(\xi) / J^2$$

with J the Jacobian of the transformation, $J^2 = \|g_{ij}\|$ and

$$b_i(\xi) = \sum \epsilon^{ijk} (\partial A_k / \partial \xi_j).$$

One can show that if the flow is divergence free, $\nabla \cdot u = 0$, the Jacobian is independent of time. The quantities $b_i(\xi)$ are given by the initial magnetic field and the initial transformation equations. Indeed with the choice $x(\xi, 0) = \xi$, the b_i are just the three Cartesian components of the initial magnetic field.

The metric tensor g_{ij} is symmetric and positive-definite, so it has three positive eigenvalues, which can be ordered by size, $\lambda_1 \geq \lambda_2 \geq \lambda_3$. The product of the three eigenvalues is the square of the Jacobian, $J^2 = \lambda_1 \lambda_2 \lambda_3$. If the flow $u(x, t)$ is chaotic, as most natural flows are, neighboring points separate exponentially. The characteristic rate of exponential separation is called the Liapunov exponent ω . The eigenvalues of the metric tensor are simply related to the Liapunov exponent as time goes to infinity. If a flow is divergence free, the relation between the Liapunov exponent and the eigenvalues as $t \rightarrow \infty$ is

$$[\ln(\lambda_1)]/t = 2\omega, \quad [\ln(\lambda_2)]/t = 0, \quad \text{and} \quad [\ln(\lambda_3)]/t = -2\omega.$$

The relation between the eigenvalues of the metric tensor and B^2 implies that the field strength characteristically rises in a chaotic flow at the exponential rate $\exp(2\omega t)$. The dissipation, which is proportional of j^2 , rises much faster, at the exponential rate $\exp(6\omega t)$, so resistive dissipation of the field will rapidly become a dominate process.²⁷

The most obvious application of these results is to astrophysical rather than fusion plasmas. Nonetheless, the technique of following the evolution of a magnetic field (or more simply a passive scalar) using the metric tensor is potentially an important technique in plasma physics, which may have applications of fusion relevance.

VI. Renormalization Group Theory Techniques in Turbulence

Over the past few years²⁸⁻³⁶ George Vahala and coworkers have been engaged in applying renormalization group theory (RNG) and subgrid modeling to problems in fluid and plasma turbulence. There are basically two different formulations of RNG: the ϵ -RNG approach and the recursive RNG approach. The ϵ -RNG approach, in the form developed by Yakhot & Orszag³⁷, has gained considerable popularity in the turbulence community. It has also been applied by Longscope and Sudan³⁸ to the reduced MHD (Strauss³⁹) equations. We, on the otherhand, while applying the ϵ -RNG to some selected plasma turbulence problems^{28,29,31}, have been basically developing the recursive RNG approach^{30,32-36} - an approach initially introduced by Rose⁴⁰ to consider the problem of the advection of a passive scalar. Both methods involve the successive elimination of high k -mode bands and the estimation of the effects of these removed bands on the resolvable scales. The ϵ -RNG method passes to a differential limit of infinitesimal bands, ignores all higher order nonlinearities introduced by the RNG procedure and calculates the transport coefficients in the limit of $k \rightarrow 0$. Detailed critiques on these approximations have appeared in the literature^{35,41,42}. The recursive RNG method does not need to take a differential limit, treats higher order nonlinearities generated, and permits the determination of the transport coefficients over the entire resolvable scales. Closure in the ϵ -RNG theory is achieved by letting $\epsilon \ll 1$ approach the limit $\epsilon \rightarrow 4$. Closure in the recursive RNG is achieved by only including the first level of new nonlinearities generated by the RNG procedure. In ϵ -RNG, the transport coefficients are determined readily from the solution of ordinary differential equations while in recursive RNG the coefficients are determined, at each resolvable wavenumber k , from the somewhat difficult determination of a fixed point from a difference-integro recursion relation.

One of the important distinctions between ϵ -RNG and recursive RNG in the treatment of nonlinear interactions and their effects on the transport coefficients is that

- (a) ϵ -RNG only treats non-local interactions, while
- (b) recursive RNG treats both non-local and local interactions.

We now briefly summarize some of our major recursive RNG results³⁶ for 3D Navier-Stokes incompressible turbulence. The elimination of high k ("subgrid") bands leads to a cubic nonlinearity in the renormalized equation of motion. Higher order nonlinearities are shown to be higher order in the nonlinear vertex interaction perturbation (which eventually is set to unity!) and are neglected. This closure scheme is reminiscent of the original RNG theory of Wilson on the 2D Ising spin model in equilibrium statistical mechanics. To recover the same physics Wilson requires that he obtain the same partition function after decimating the number of degrees of freedom. As a result of eliminating some degrees of freedom, the new Hamiltonian now involves not only the Ising nearest-neighbor interactions but also new interactions arising from diagonal and from 4-spin interactions. Wilson drops the 4-spin interactions, since it can be shown that the strength of these interactions is very weak. However he keeps the new RNG-induced diagonal interactions and shows that they play a crucial role in the RNG recursion relations from which the fixed points are determined.

In the case of turbulence, we are far from equilibrium. However to achieve closure we drop fourth and higher order nonlinearities (by perturbation arguments) and find that the RNG-induced triple nonlinearities play a crucial role in determining the structure of the transport coefficients. In particular, in the energy transport equation, we³⁶ have shown that the inclusion of the triple nonlinearities leads to the important effect of backscatter of energy from the small scales to large scales. This effect has been seen in direct numerical simulations and large eddy simulations, but is not present in ϵ -RNG theories. The non-local interactions lead to a constant small- k eddy viscosity, as is expected from the definition of transport coefficients. However, local interactions are now shown to play a crucial role in treating the interaction of the highest resolvable- k with the nearby subgrid (eliminated) k -band. This interaction leads to a sharp cusp or rise in the eddy viscosity near this cut-off wavenumber. Again, this effect has been verified quantitatively by analysis of both direct numerical simulations and large eddy simulations and constitutes a strong validation of our recursive RNG theory. This is especially relevant since fluid turbulence simulations are at a good level of sophistication and these simulation methods are now being implemented in plasma physics. This work was performed in collaboration

with Ye Zhou (a former William & Mary graduate student who is now at Center for Turbulence Research, Stanford University).

VII. Magnetic and Density Fluctuations in Tokamak Plasmas

Recently, George Vahala, Linda Vahala, and Norton Bretz⁴³⁻⁴⁵ have proposed a diagnostic for detection of magnetic fluctuations in tokamak plasmas by using an $O \rightarrow X$ mode conversion process. This is of particular relevance to fusion plasmas because of the interconnection between magnetic fluctuations and transport and because of possible high levels of magnetic fluctuations induced by α -driven modes. We have also shown how the O-mode cutoff layer can be used as a filter to separate an incident O-mode from a scattered X-mode (or vice versa). In perpendicular propagation, due to wave polarization considerations, the mode conversion $O \rightarrow X$ can only be achieved by magnetic fluctuations. This fact allows us to distinguish between scattering from magnetic fluctuations rather than from density fluctuations, even though it is expected that the level of magnetic fluctuations could be a factor 10^4 lower than that for density fluctuations. In particular, we⁴⁵ have considered both the effects of finite microwave beam widths and polarization limitations of receiver/transmitter antennas. More work, however, needs to be done on the effect of fluctuations on the cutoff layers and on reflectometry diagnostic on fluctuation measurements and these studies are underway.

It is well known⁴⁶ that density fluctuations play an important role in the propagation of lower hybrid (LH) waves in a toroidal plasma. The effect of these fluctuations on the LH waves was obtained by a Monte Carlo solution of a wave kinetic equation derived assuming weak turbulence theory. However, in considering LH heating and current drive, the computer models have concentrated on the confinement properties of the suprathermal tail electrons created by these LH waves and the transport properties of the background electrons. Since it has been found to be computationally prohibitive to incorporate the Monte Carlo scattering solver into these transport codes, the effects of scattering on LH waves has not been pursued further. In conjunction with Linda Vahala

[Old Dominion University] and Paul Bonoli [MIT], we^{47,48} have extended the study of fluctuations on LH waves by also incorporating the effects of magnetic fluctuations. A wave kinetic equation has been derived incorporating the effects of both density and magnetic fluctuations and the Monte Carlo scattering code has been suitably modified. We have found considerable internal magnetic fluctuation effects on the power deposition profiles to the electrons, even when the level of these magnetic fluctuations are a factor of 10^4 lower than those of density fluctuations (e.g., density fluctuations arising from drift wave turbulence).

Moreover, we^{48,49} have been able to correlate these effects with hard X-ray experimental data by examining the hard X-ray spatial profile as detected on the JT-60 tokamak in Japan. These hard X-rays arise from prompt collisions of the suprathermal electrons with the background plasma ions and hence the experimental profiles can be correlated to electron power deposition profiles. We have calculated the corresponding electron power deposition profiles for the JT-60 LH parameters from the Monte Carlo solution of our wave kinetic equation which includes both density and magnetic fluctuations. Excellent agreement was found for the three different profiles which corresponded to different initial LH launch $k_{||}$ if one incorporated edge turbulent density fluctuations. However, for the experimental JT-60 parameters considered, there were no significant effects on the electron power deposition profiles when magnetic fluctuations (with intensity down by a factor of 10^4) were also included.

We have also examined the effects of density and magnetic fluctuations on the electron power deposition for LH wave propagation in the much higher magnetic field/density Alcator C-Mod tokamak. In this case, we found quite significant magnetic fluctuation effects - effects that should be visible in hard X-ray data.

References

1. H. Alfven, Ark. Mat. Astron. Fys. **27A**, 1 (1940).
2. J. Nührenberg and R. Zille, Phys. Lett. **A129**, 113 (1988).
3. D. A. Garren and A. H. Boozer, Phys. Fluids **B3**, 2822 (1991).
4. A. H. Boozer, Phys. Fluids **27**, 2441 (1984).
5. A. H. Boozer, Phys. Fluids **26**, 496 (1983).
6. M. Kruskal, in **Pendiconti del Terzo Congresso Internazionale sui Fenomeni D'Ionizzazione nei Gas tenuto a Venezia** (Societa Italiana di Fisica, Milan, 1957), p. 56.
7. A. H. Boozer, Phys. Fluids **24**, 1999 (1981).
8. D. A. Garren and A. H. Boozer, Phys. Fluids **B3**, 2805 (1991).
9. C. Mercier, Nucl. Fusion **4**, 213 (1964).
10. D. Lortz and J. Nührenberg, Z. Naturforsch. **31a**, 1277 (1976).
11. D. Lortz and J. Nührenberg, Nucl. Fusion **17**, 125 (1977).
12. D. Lortz and J. Nührenberg, Z. Naturforsch. **34a**, 167 (1979).
13. L. S. Solov'ev and V. D. Shafranov, in Review of Plasma Physics, edited by M. A. Leontovich (Consultants Bureau, New York, 1970), Vol. 5, p. 1.
14. R. H. Weening and A. H. Boozer, Phys. Fluids **B4**, 159 (1992).
15. A. H. Boozer, J. Plasma Phys. **35**, 133 (1986).

16. H. P. Furth, P. H. Rutherford, and H. Selberg, Phys. Fluids 16, 1054 (1973).
17. P. H. Rutherford, Phys. FLuids 16, 1903 (1973); A. H. Boozer, Phys. Fluids 27, 2055 (1984).
18. C. Z. Cheng, H. P. Furth, and A. H. Boozer, Plasma Phys. and Cont. Fusion 29, 351 (1987).
19. J. A. Wesson, R. D. Gill, M. Hugon, F. C. Schüller, J. A. Snipes, D. J. Ward, D. V. Bartlett, D. J. Campbell, P. A. Duperrex, A. W. Edwards, R. S. Granetz, N. A. O. Gottardi, T. C. Hinder, E. Lazzaro, P. J. Lomas, N. Lopes Cardozo, K. F. Mast, M. F. F. Nave, N. A. Solomon, P. Smeulders, P. R. Thomas, B. J. D. Tubbing, M. F. Turner, and A. Weller, Nucl. Fusion 29, 641 (1989).
20. S. Ishida, Y. Koide, T. Ozeki, M. Kikuchi, S. Tsuji, H. Shirai, O. Naito, and M. Azumi, Phys. Rev. Lett. **68**, 1531 (1992).
21. R. J. Bickerton, J. W. Connor, and J. B. Taylor, Nat. Phys. Sci. **229**, 110 (1971).
22. A. H. Boozer, Phys. Fluids **29**, 4123 (1986).
23. A. Boozer and A. Rechester, Phys. Fluids **21**, 682 (1978).
24. N. Pomphrey and A. Reiman, Princeton Plasma Physics Lab. Report PPPL-2801 (1992).
25. T. J. Martin and J. B. Taylor, Plasma Phys. Contr. Fus. **26**, 321 (1984).
26. S. Lundquist, Phys. Rev. **83**, 307 (1951).
27. A. H. Boozer, to be published Astrophys. J. (July 1992).
28. Y. Zhou and G. Vahala, Phys. Lett. **124A**, 355 (1987).

29. Y. Zhou and G. Vahala, J. Plasma Physics **39**, 511 (1988).
30. Y. Zhou, G. Vahala, and M. Hossain, Phys. Rev **A37**, 2590 (1988).
31. Y. Zhou and G. Vahala, J. Plasma Phys. **41**, 67 (1989).
32. Y. Zhou, G. Vahala, and M. Hossain, Phys. Rev. **A40**, 5865 (1989).
33. Y. Zhou and G. Vahala, Phys. Lett. **A147**, 43 (1990).
34. Y. Zhou and G. Vahala, J. Plasma Physics **45**, 239 (1991).
35. Y. Zhou and G. Vahala, Phys. Rev. A (to be published, 1992).
36. Y. Zhou and G. Vahala, "Reformulation of Recursive Renormalization Group Based Subgrid Modeling of Turbulence", submitted to Phys. Fluids **A**.
37. V. Yakhot and S. A. Orszag, J. Sci. Comput. **1**, 3 (1986).
38. D. W. Longcope and R. N. Sudan, Phys. Fluids **B3**, 1945 (1991).
39. H. R. Strauss, Phys. Fluids **19**, 134 (1976).
40. H. A. Rose, J. Fluid Mech. **81**, 719 (1977).
41. L. M. Smith and W. C. Reynolds, Phys. Fluids **A4**, 364 (1992).
42. S. H. Lam, Phys. Fluids **A4**, (April, 1992).
43. L. Vahala, G. Vahala and N. Bretz, Rev. Sci. Instrum., **61**, 3022 (1990).
44. L. Vahala, G. Vahala and N. Bretz, Proc. 9th Topical Conf. on Radio Frequency Power in Plasmas. (AIP Conference Proceeding, Charleston S.C. August 1991).
45. L. Vahala, G. Vahala and N. Bretz, Phys. Fluids **B4**, 619 (1992).

46. P. T. Bonoli, Ph. D Thesis (1981); P. T. Bonoli and E. Ott, Phys. Rev. Lett. **46**, 424 (1981); Phys. Fluids **25**, 359 (1982).
47. G. Vahala, L. Vahala and P. T. Bonoli, Rev. Sci. Instrum (to be published, 1992).
48. G. Vahala, L. Vahala and P. T. Bonoli, "Effect of Magnetic and Density Fluctuations on the Propagation of Lower Hybrid Waves in Tokamaks". Phys. Fluids B (submitted).

List of 1991-1992 Publications

Allen Boozer:

1. The Drift Hamiltonian in a Magnetic Field with a Separatrix
Allen H. Boozer
Phys. Fluids B 3, 875-879 (1991).
2. Magnetic Field Strength of Toroidal Plasma Equilibria
D. A. Garren and A. H. Boozer
Phys. Fluids B 3, 2805-2821 (1991).
3. Existence of Quasihelically Symmetric Stellarators
D. A. Garren and A. H. Boozer
Phys. Fluids B 3, 2822-2834 (1991).
4. Electron Transport in Stochastic Fields of RFP ZT-40M
A. Punjabi, A. Verma, M. Kim, and A. H. Boozer
Proc. XX Int. Conf. on Phenomena in Ionized Gases, Barga, Italy, July 8-12, 1991 (V. Pappalardo and M. Vaselli ed., Institute of Atomic and Molecular Physics-CNR, Pisa, Italy, 1991) p. 424-425.
5. Completely Bootstrapped Tokamak
R. H. Weening and A. H. Boozer
Phys. Fluids B 4, 159-170 (1992).
6. Dissipation of Magnetic Energy in the Solar Corona
A. H. Boozer
To be published in Astrophys. J. (July 1992).
7. Onsager Symmetry of Transport in Toroidal Plasmas
A. H. Boozer
To be published in Phys. Fluids B (August 1992).
8. The Exact and Drift Hamiltonian
Q. Yao and A. H. Boozer
Phys. Fluids B (August 1992).

George Vahala:

1. Filamentation in High β Plasmas and Flux Penetration in Type II Superconductors
L. Vahala and G. Vahala, Phys. Essays **4**, 223 (1991).
2. Aspects of Subgrid Modeling and Large Eddy Simulation of MHD Turbulence
Y. Zhou and G. Vahala, J. Plasma Physics **45**, 239 (1991).
3. Detection of Internal Magnetic Fluctuations by Electromagnetic Scattering in Tokamaks
L. Vahala, G. Vahala, and N. Bretz, Proc. 9th Topical Conf. on Radio Frequency Power in Plasmas (AIP Conference Proceeding, Charleston S.C. August 1991).
4. Electromagnetic Wave Scattering from Magnetic Fluctuations in Tokamaks
L. Vahala, G. Vahala, and N. Bretz, Phys. Fluids **B4**, 619 (1992).
5. Effect of Fluctuations on Lower Hybrid Power Deposition and Hard X-Ray Detection
G. Vahala, L. Vahala, and P. T. Bonoli, Rev. Sci. Instrum (to be published, 1992).
6. Local Interactions in Renormalization Methods for Navier-Stokes Turbulence
Y. Zhou and G. Vahala, Phys. Rev. A (to be published, 1992).

END

DATE
FILMED

11 / 16 / 92