AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

BROOKHAVEN NATIONAL LABORATORY

July 1994

BNL-60633

A Relativistic Cascade for Heavy Ion CollisionsT.J. Schlagel^{a,b}, Y. Pang^a, and S.H. Kahana^a,^a Brookhaven National Laboratory, Upton, NY 11973^b State University of New York, Stony Brook, NY 11794

ABSTRACT

Au on Au collisions at the BNL/AGS (11.6 GeV-A/c) are expected to produce a short-lived state of matter at high baryon density. If the baryons reach sufficiently high density, they may produce a quark-gluon plasma (QGP). The signals from a QGP phase may be difficult to distinguish from those of ordinary hadronic matter. We have constructed a relativistic cascade (ARC) for hadrons in an attempt to model the dynamics of ordinary hadronic matter in a heavy ion collision, in the hopes that deviations from the cascade results may indicate new physics. In this contribution I will discuss the formation of high baryon density matter, and its effect on antiproton production.

Invited talk presented at the 5th Conference on
The Intersections of Particle and Nuclear Physics
St. Petersburg, Florida
May 31 - June 6, 1994

This manuscript has been authored under contract numbers DE-FG02-93ER40768 (TJS) and DE-AC02-76CH00016 with the U.S. Department of Energy. Accordingly, the U.S. Government retains a non-exclusive, royalty-free license to publish or reproduce the published form of this contribution, or allow others to do so, for U.S. Government purposes.

MASTER *ds*
DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

A RELATIVISTIC CASCADE FOR HEAVY ION COLLISIONS

T. J. Schlagel^{a,b}, Y. Pang^a, and S. H. Kahana^a

^aDepartment of Physics, Building 510A
Brookhaven National Laboratory, Upton, NY 11973

and

^bState University of New York
Stony Brook, NY 11794

Abstract

Au on Au collisions at the BNL/AGS (11.6 GeV-A/c) are expected to produce a short lived state of matter at high baryon density. If the baryons reach sufficiently high density, they may produce a quark-gluon plasma (QGP). The signals from a QGP phase may be difficult to distinguish from those of ordinary hadronic matter. We have constructed a relativistic cascade (ARC) for hadrons in an attempt to model the dynamics of ordinary hadronic matter in a heavy ion collision, in the hopes that deviations from the cascade results may indicate new physics. In this contribution I will discuss the formation of high baryon density matter, and its effect on antiproton production.

The first ultra-relativistic collisions between truly heavy ions (Au on Au) were performed in 1992 at the BNL/AGS. It is hoped that in these collisions, with sufficiently high energy and baryon number, that a region of compressed hadronic matter will be formed. When the baryon density or temperature rises above a critical value, the quarks and gluons confined in the hadrons are freed, and a state of matter called the quark-gluon plasma (QGP) is formed. The challenge to experimentalists and theorists is to extract information on the QGP phase of matter from the final particles which are actually detected. One approach to studying these collisions is to construct a completely hadronic model for the elementary collisions of particles, and compare the results of simulations to experiment. Deviations from the model might then be potential signals for the formation of QGP.

We have developed an hadronic cascade model for relativistic heavy ion collisions called *A Relativistic Cascade* (ARC)[1, 2, 3]. The basic constituents of ARC are baryons (N , Δ^{1232} , Λ , $\bar{N}$, $\bar{\Delta}^{1232}$) and mesons (π , ρ , K). Scattering by a potential is neglected, so particles travel along straight line trajectories between two-body collisions, greatly simplifying the solution of the dynamics. Since the initial velocity of the nucleons is large ($v_p^{\text{CM}} > 0.9$ at AGS energies) the use of relativistic kinematics is required. Scattering between two particles occurs when $b_{\text{cm}} < \sqrt{\sigma(\sqrt{s})}/\pi$, where b_{cm} is the minimum distance of approach between the particles in their mutual center of mass frame, and $\sigma(\sqrt{s})$ is the energy dependent cross section for the two particles. The two-body scattering cross sections are taken from the CERN-HERA data compilation[4]. Elastic and inelastic

Figure 1: Proton rapidity distribution in central Au on Au collisions at 11.6 GeV-A/c. The solid curve shows the cascade prediction ($b < 2$ fm), while the E866[5] preliminary data are shown by the black circles. The white circles show the data reflected about midrapidity ($y_{nn} = 1.6$).

Figure 2: Rapidity distribution of produced particles. The cascade predictions are shown by the diamonds while the preliminary E866[5] data are the circles.

cross sections are used to determine the branching ratios into different final channels. In the inelastic channel the produced particles are distributed using Lorentz invariant phase space

$$\int W_T W_L \delta^4 \prod_i \frac{d^3 p_i}{E_i} \quad (1)$$

where the delta function is to ensure conservation of energy and momentum for the collision and the additional phase space weights

$$W_T = e^{-p_t^2/\beta^2}, \quad W_L = e^{-\alpha(1 \pm p_L/p_{max})} \quad (2)$$

are added to assign limited p_t for all produced particles and leading particle behavior for the incoming particles. The parameters α and β are adjusted to fit pp data. The elastic channel is handled separately using momentum transfer and energy as the variables.

Comparisons of ARC to data on Si induced reactions demonstrate the cascade gives a good description of these collisions. Predictions for proton and meson rapidity distributions in central Au on Au collisions are shown in Figures 1 and 2 along with the preliminary data from experiment E866[5]. The proton distribution at midrapidity is expected to be measured during the next heavy ion run at the AGS (Fall '94).

Figure 3: Evolution of the local baryon density in central collisions. Pictured are Si on Si (14.6 GeV-A/c), Si on Au (14.6 GeV-A/c), and Au on Au (11.6 GeV-A/c) in the nucleon-nucleon center of mass frame at four different times. The grayscale at the bottom indicates the density in units of nuclear matter density. A color version of this figure can be obtained from the author.

An advantage of the cascade model is that it contains the complete space-time history of the collision, and so can give information on the evolution of the system. We have used this information to analyze the local baryon density, ρ_{local}^B . Figure 3 shows ρ_{local}^B (in units of nuclear matter density, $\rho_0 = 0.16 \text{ fm}^{-3}$) for three systems studied at the AGS, Si on Si, Si on Au, and Au on Au. The evolution of the Si induced reactions is quite similar. By the time $t = 0 \text{ fm/c}$ (when the centers of the nuclei overlap in the absence of interactions), the projectile has completely entered the target, and there is a region roughly the size of the Si with baryon density $> 5\rho_0$. Note that the minimum peak baryon density we should see is $\rho_{\text{local}}^B/\rho_0 = 2\gamma_{\text{ev}} = 5.8$ (γ_{ev} is the gamma of the projectile and target in the equal velocity frame) which comes from the overlap of the target and projectile nuclei. By $t = 2 \text{ fm/c}$, the density has passed its peak value, and the system decompresses. In the Si on Si case there is a two component distribution, but the Si projectile completely disintegrates in Si on Au.

The situation in Au on Au collisions is quite different. Now at $t = 0 \text{ fm/c}$ a large volume of high baryon density matter is formed. The density continues to increase until 2 fm/c , approaching a peak value of $9.5\rho_0$, with 25% of the baryons at $\rho_{\text{local}}^B > 5\rho_0$. After $t = 4 \text{ fm/c}$ the nuclei appear to have lost their original identity, replaced by a single expanding source. The region of high baryon density lasts twice as long as in the Si on Au case.

One might ask, if we are truly creating matter with such high baryon density, then why does the hadronic cascade approach still work? Perhaps the signals from the QGP formation are small and are washed out by the final state interactions of the hadrons. If this is the case, finding evidence for the QGP will require a very detailed comparison with experimental data. It may also be that we have not achieved the ideal environment for forming the high density plasma. In Figure 4 we show the evolution of the rapidity

Figure 4: Evolution of the rapidity distribution of baryons in central Au on Au collisions at 11.6 GeV-A/c in the center of mass frame. The curves show target (solid) and projectile (dashed) baryon distributions as they are followed through the cascade. The final spectra is shown at the bottom.

Figure 5: Rapidity distributions in Au on Au collisions for target and projectile protons separately as a function of beam momentum. The large gap between target and projectile central rapidities at 11.7 GeV-A/c reduces the effect of the high baryon density achieved.

distribution of baryons which have originated from projectile or target separately. The peaks of the two distributions always remain separated, meaning that the target and projectile nucleons have not completely mixed in momentum space, although they have reached a high density in coordinate space. As the beam momentum is lowered the rapidity gap between the two peaks decreases (Figure 5), so the density in momentum space increases. Interestingly, the coordinate space density decreases slowly with energy. Even for a beam momentum of 4 GeV-A/c, the peak baryon density reaches values of $\rho > 7\rho_0$, and the system is even longer lived. Therefore the possibility of forming a high density plasma may improve at somewhat lower energies, though still accessible at the AGS.

Antiproton production and annihilation can be strongly effected by dense matter, and has been suggested as a useful probe[6]. We have previously studied antiprotons in Si induced reactions[2]. Our conclusion from this work is that in order to treat the large cross section processes of annihilation ($\sigma_{p\bar{p}} > 200$ mb at low relative momentum) realistically in a cascade, it is necessary to include the interactions with a third particle prior to annihilation, which can scatter the antiproton along another trajectory, thus reducing the annihilation probability. The solution is to include an extra time for the annihilation to occur,

$$\tau_a = D/2v, \quad (3)$$

where D is the distance of that particles at the time of closest approach, and v their velocity. This extra time has no effect in very dilute systems, when the probability for screening is small.

Recently experiment E878 has measured antiproton production in Au on Au collisions at 10.8 GeV-c/A at $p_t = 0$ [7]. In their comparisons with cascade models, they have shown that ARC overpredicts the rapidity distribution but gives approximately the correct overall shape. In contrast, the model RQMD[8], which has enhanced production but uses the full annihilation cross section, underpredicts the spectrum and shows a large dip at y_{nn} , while the experimental data is flat. We can explore within the contexts of ARC the effects of increased annihilation by making a small change to τ_a . If we assume that the proton and antiproton do not have to be overlapping, but can annihilate at some distance χ , then the annihilation time becomes

$$\tau'_{\text{ann}} = (D - \chi)/2v. \quad (4)$$

We have considered three cases: full screening ($\chi = 0$), half overlap ($\chi = 0.8$ fm) and surface annihilation ($\chi = 1.6$ fm). The results of these calculations are shown in Figures 6 and 7. The transverse momentum spectra near the target and projectile rapidities suffers an overall change in normalization, but the shape remains much the same. Near midrapidity, increasing χ causes the development of a dip at $p_t = 0$, due to the large baryon density and strong annihilation at low relative momentum. This dip is clearly seen in the rapidity distribution measured at $p_t = 0$, but is not evident in the experimental data. Thus it is clear that there is a need for including the screening mechanism for a proper description of antiprotons.

Complicating the extraction of information on antiprotons is the contamination of $\bar{\Lambda}$, which can subsequently decay to an antiproton[9]. The ARC calculation does not include $\bar{\Lambda}$ production, nor does E878 correct for them. So this remains a large source of uncertainty in any comparison.

In conclusion, the hadronic cascade code ARC gives a good description of the global dynamics of the collision. A space-time analysis shows that the densities reached are large, but there is still not complete overlap in momentum space – decreasing the beam energy increases the momentum overlap, but changes the peak baryon density reached only a little. The agreement of the cascade predictions with the current data mean that searching for evidence of a plasma will be a painstaking process. The importance of screening of antiproton annihilation is confirmed in Au on Au collisions, but comparisons with data are complicated by many uncertainties. The situation may become clearer with subsequent Au beam measurements.

REFERENCES

1. Y. Pang, T. J. Schlagel, and S. H. Kahana, Phys. Rev. Lett. **68**, 2743 (1992); T. J. Schlagel, S. H. Kahana, and Y. Pang, Phys. Rev. Lett. **69**, 3290 (1992); S. H. Kahana, Y. Pang, and T. J. Schlagel, *Proceedings of Heavy Ion Collisions at the AGS, HIPAGS '93*, 13–15 January, 1993, MIT, Cambridge, MA, MITLNS-2158.

Figure 6: Transverse momentum distributions of antiprotons in central ($b < 4$ fm) Au on Au collisions at 10.8 GeV-A/c for $\chi = 0$ (circle), $\chi = 0.8$ (diamond) and $\chi = 1.6$ (square). For $\chi = 1.6$ the antiproton spectra develops a hole at $p_t = 0$.

Figure 7: Rapidity distribution of antiprotons at $p_t = 0$. These spectra should be compared to the data of experiment E878, shown in Shiva Kumar's contribution to these proceedings.

2. S. H. Kahana, Y. Pang, T. J. Schlagel, and C. B. Dover, Phys. Rev. C **47**, R1356 (1993).
3. A. J. Baltz *et al.* Phys. Rev. Lett. **69**, 3290 (1992).
4. A. Baldini, V. Flaminio, W. G. Moorhead and D. R. O. Morrison, *Total Cross-Sections for Reactions of High Energy Particles*, Landolt-Börnstein New Series, Volume 12A and Volume 12B, Ed. O. Madelung, (Springer Verlag, Berlin, 1988).
5. H. Hamagami *et al.*, Nucl. Phys. A **566**, 27c (1994).
6. S. Gavin, M. Gyulassy, M. Plümer, and R. Venugopalan, Phys. Lett. B **234**, 175 (1990).
7. M. Bennett, talk given at 1994 Joint April Meeting of The APS and the AAPT, April 1994, and private communication; S. Kumar, these proceedings.
8. A. Jahns *et al.*, Phys. Rev. Lett. **72**, 3464 (1994).
9. P. Rothschild, Ph.D. thesis.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DATE

FILMED

8/25/94

END