AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

**SELECTIVE EXCITATION, RELAXATION, AND
ENERGY CHANNELING IN MOLECULAR SYSTEMS**

Comprehensive Progress Report

April 1990 - July 1993

William C. Rhodes

Institute of Molecular Biophysics

Department of Chemistry

Florida State University

Tallahassee, Florida 32306

August 1993

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-86ER60473

MASTER

EP

SELECTIVE EXCITATION, RELAXATION, AND
ENERGY CHANNELING IN MOLECULAR SYSTEMS

William C. Rhodes
Institute of Molecular Biophysics
Department of Chemistry
Florida State University
Tallahassee, Florida 32306

Grant #: DE-FG05-86ER60473

Department of Energy

TABLE OF CONTENTS

INTRODUCTION.....	2
RESULTS.....	5
I. Quantum Dynamics and Thermodynamics of Conformational	
Transitions in Biological Macromolecules	5
II. Coherent Dynamics	23
III. Reaction Rate Theory	26
IV. Fundamental Subsystem Equations of Motion	27
PUBLICATIONS	30
RELATION TO STATED OBJECTIVE - PLANS FOR CONTINUATION.....	31
GRADUATE STUDENTS TRAINED	31
OTHER FEDERAL SUPPORT	31
PRESENT STATE OF KNOWLEDGE.....	31
ESTIMATE OF UNOBLIGATED BALANCE.....	32

SELECTIVE EXCITATION, RELAXATION, AND ENERGY CHANNELING IN MOLECULAR SYSTEMS

William Rhodes

Comprehensive Progress Report

INTRODUCTION

The research discussed here involves theoretical studies of the response, relaxation, and correlated motion in the time-dependent behavior of rather large molecular systems. By large systems, we mean systems having a large number of degrees of freedom. They range in size from a single large polyatomic molecule to a protein molecule in its natural environment. By response we mean the changes in the electronic and vibrational motion of a molecule (or a portion of a molecule we can call a chromophore) due to interactions with an externally applied field (such as light) or to intermolecular interactions in systems not at equilibrium. Such responses are the associated relaxation and the channeling of energy as the system moves toward a new equilibrium.

The present studies are part of an ongoing development of a research program which historically includes the following topics.

- (1) Electronic spectral properties of helical biological polymers. Focus has been on the relation between conformation and optical properties, such as absorption spectral structure (e.g., hypochromism) and circular dichroism.
- (2) Nonradiative transitions in molecules. Focus has been placed on the role of the exciting light in determining the structure of the prepared excited state. The role of intramolecular

interactions and modulation by the molecular medium have been studied in previous phases of this work. Intermolecular excitation energy transfer by coherent (exciton) or by noncoherent (Förster) mechanisms is included in this topic.

- (3) **Macroscopic system dynamics.** Focus is on the role of the medium in modulating the dynamics of a molecule (a chromophore). Emphasis is placed on the effects of coherent structure, coherence relaxation (memory), and the interplay of coherence and dissipation in determining how kinetics arises from coherent quantum dynamics.
- (4) **Biomolecular dynamics.** Focus is placed on the role of subsystem modulation (e.g., by low-frequency vibrational modes) in the channeling of processes in biological macromolecules and on the role of subsystem cycles in the natural functions of protein molecules. Emphasis is placed on the nature of thermodynamic modes and the microscopic basis of entropy.

The underlying theme of our research is subsystem modulation dynamics, in which the dynamics of a part of a system (as determined by certain substructures or degrees of freedom) are modified by interaction with the remaining degrees of freedom (referred to as the background). Important concepts include coherence, correlation, dissipation, memory, adiabatic motion, symmetry breaking localization and stabilization, collective modes, and resonance structures. The main idea is that quantum mechanical correlations between the components of a system develop with increasing time and these correlations play a major role in determining the balance between *coherent* and *dissipative* forces acting on each component of the system. These forces help establish the *pathways* through which processes are channeled.

The central theme of our research is the *interplay of coherence and dissipation in determining the nature of dynamic structuring and energy flow i.e. molecular transformation mechanisms.*

Our primary goal is two-fold:

- (i) We are working to develop fundamental subsystem equations of motion showing how nonlinear and dissipative dynamics of a particular subsystem arises from correlated interactions with the remainder of the system, *viz.* substituent groups, solvent, lattice modes, etc. One consequence is the formation of *resonance structures* and *resonance networks*, which can function to control and channel processes.
- (ii) We are applying the concepts and principles of quantum dynamics and thermodynamics to develop understanding of control and energy transfer mechanisms involved in the biological functions of protein molecules, such as enzymes, G-proteins, and channel proteins. Transfer/control mechanisms in biological macromolecules are both *global* and *local* in nature. The results associated with goal(i) above are useful in these studies.

In addition to formal theoretical development, our research deals with several phenomenological aspects of molecular systems. These are (a) response/relaxation dynamics associated with molecular excitation by external perturbations. Specifically, these include optical properties and processes in biological polymers and the role of molecular conformation; (b) Reaction rate theory approached via the use of projection operator methods and the Liouville equation; (c) Excitation energy transfer within and between molecules by such mechanisms as exciton transfer, Förster (resonance) transfer, and (Davydov) solitons; (d) Control and energy transformation mechanisms in isothermal protein cycles of enzymes, G-proteins, and channel proteins.

RESULTS

I. Quantum Dynamics and Thermodynamics of Conformational Transitions in Biological Macromolecules

A. The Characterization of Thermal Modes

I. Introduction

We are told by the elementary textbooks that the thermodynamic state of a system is determined by the internal energy dependence on a set of extensive variables including entropy, S , volume, V , molar composition, and any extensive variables conjugate to external forces acting on the system. Consider, for example, the case of one mole of carbon with a certain value of S and V (and no external forces). There may still be ambiguity unless we know whether the carbon is in the form of graphite or diamond. The point is that, in addition to molar composition, the molecular-level structure of the system is important. The spectrum of thermal modes depends on the microscopic structure. For the case of carbon, each structural form is stable for times far longer than is required for thermal equilibration to occur. This is due to *constraints* produced by potential barriers separating the structures.

The lesson of the above carbon example is that absolute thermodynamic equilibrium is a questionable concept; rather, in any real system (except possibly some pure elementary particle systems) thermodynamic equilibrium is constrained in nature and depends on structural parameters of the system. The carbon case is an obvious example, but a similar situation exists for structurally more complex systems such as biological macromolecules. Many protein molecules exist in several conformations, each of which is stable over time periods sufficient for thermal equilibration. Thus, conformational space for these molecules is partitioned into conformational subspaces by energy

barriers which provide thermodynamic constraints. Each conformational subspace has its own spectrum of thermal modes. For solutions, solvent binding adds additional important structural parameters.

Obviously, chemical (and physical) processes in systems, in general, produce changes in the spectrum of thermal modes. Mode types may be transformed: oscillator $\leftrightarrow$ translational; librational $\leftrightarrow$ rotational; etc. Also, mode frequencies may change. An important case is one in which the frequency change of an oscillator mode is so great that the mode is changed from being thermally active to being thermally inactive, or *vice versa*. In the following we refer to the inactive $\rightarrow$ active transformation as *thermal mode thawing*. This is a purely quantum mechanical effect whose theoretical basis is discussed in every elementary text. However, the role of thermal mode thawing and freezing in chemical processes (at constant T), especially large molecule reactions in condensed systems, appears not to be widely appreciated and, thus, is a major emphasis of this work. It is important to realize that mode thawing (freezing) here refers not to the effects of increasing temperature but to the effects of a chemical (or physical) process in changing mode frequencies.

The major focus of this research is on the role of thermal mode transformations in physical and chemical processes. Our primary purpose is (i) to show how the entropy of oscillator thermal modes naturally decomposes into two terms having definite geometrical and physical meanings (relevant to the work of others) and (ii) to show how linear thermodynamic relations can be explained in terms of the transformation of thermal modes associated with chemical processes.

2. Background

In 1971, Benzinger¹ pointed out that the Gibbs free energy change, ΔG , for any process is not affected by subtracting a term from the enthalpy change, ΔH , provided the same term is subtracted from the entropy term, $-T\Delta S$. This alone is not a profound observation, but Benzinger went on to point out that the quantities contributing to ΔH and ΔS for any process (viz., chemical reaction)

naturally produce such a cancelling term. The term in question is the thermal energy component of ΔH , obtained by integrating the change in heat capacity at constant pressure, ΔC_p , over the temperature range $0K$ to T (and adding enthalpies due to phase transitions of components within the temperature range). The remaining component of ΔH is the change in enthalpy at $0K$ which we refer to as the change in zero-point enthalpy, ΔH_0 .

Thus, as Benzinger has pointed out¹, by subtracting the thermal component of the enthalpy change from ΔH and subtracting T^{-1} times this same term from ΔS , one obtains an identical ΔG value in terms of the change in zero-point enthalpy and the change in the remaining component of the entropy, which Benzinger refers to as the "free entropy", ΔS_f . More succinctly, $\Delta G = \Delta H_0 - T\Delta S_f$.

For small molecule reactions in vapor phase the thermal component of ΔH is small compared to the zero-point enthalpy component. As far as these kinds of reactions are concerned, Benzingers' observation produces results having mainly incidental academic value. As Benzinger has pointed out¹, however, this is not true for large molecule reactions in condensed media. Here, ΔC_p can be quite large, producing a thermal component of ΔH which may be much larger than the zero-point component. As an example of this, Benzinger refers to the results of Danforth, et al.,² for the unfolding of ribonuclease.

An important consequence of Benzinger's observation is that, typically, for biological macromolecule reactions, ΔG is determined by the minor component of ΔH together with the "free entropy" term, which is considerably smaller than the total entropy change, ΔS . Unfortunately, Benzinger leaves his discussion on a pessimistic note that the experimental determination of the all-important zero-point enthalpy and the free entropy terms will depend on the difficult measurements of ΔC_p over the entire temperature range, $0K$ to T . While this is true, it does not detract from the conceptual significance of his results (*vide infra*).

The cancellation of major components of ΔH and $T\Delta S$ in large molecule reactions is suggestive of linear thermodynamic relations in organic molecule reactions - a topic which has long been appreciated, as is exemplified in the classic comprehensive work of Leffler and Grunwald.³ The essential feature of linear thermodynamic relations is the existence of a linear relation between two thermodynamic quantities for a family of similar chemical reactions. Two broad categories are as follows:³

- (i) a linear relation between $\Delta G_k^\ddagger$ (activation free energy) and ΔG_k ; and
- (ii) a linear relation between ΔH_k and ΔS_k for families of reactions in which the members (denoted by index, k) vary from one another, for example, by substituent groups or by solvent. The theoretical basis for the linear free energy relation between $\Delta G^\ddagger$ and ΔG goes back to the early work of Leffler.⁴

A good example of linear thermodynamic relations in the recent literature is found in the work of Saltiel and Sun^{5,6} on photoisomerization of stilbene in various n-alkane solvents. The singlet excited state isomerization of stilbene shows a linear free energy relation⁶ between $\Delta G^\ddagger$ and ΔG (category one, above) and a linear relation⁵ between $\Delta H^\ddagger$ and $\Delta S^\ddagger$ (category two, above).

Of great significance to the present work, however, are the thermodynamic properties of reactions involving biological macromolecules^{7,8}. Linear relations, especially in category two above, have been observed in families of protein reactions. Lumry and Gregory⁷ have proposed that Benzinger's compensation thermal component of ΔH is responsible for the linear relation between ΔH and ΔS in protein transformations involving conformation changes. In the course of discussions of these properties, the term compensation entropy⁷ has been applied to the entropy term which cancels the thermal energy component of ΔH . We continue to use this terminology

here. The values of the zero-point enthalpy changes, as well as the changes in compensation and free entropies for the refolding of ribonucleases, have been studied by Chun.⁹ References 7 and 8 contain many references to the thermodynamic properties of biological macromolecular reactions.

3. *Basic Features*

We begin with a formulation of the entropy of a set of quantum oscillators at thermal equilibrium. The thermal energy is regarded as an ideal gas of thermal quanta which, like photons, are not conserved. The thermodynamic systems considered here are assumed to be closed. Although our starting point is the same as that of Planck,¹⁰ we deviate from Planck's formulation and proceed to develop the following points:

- (1) By making a simple transformation, we observe that the entropy, expressed as a function of the number of thermal quanta, consists of two additive components each having definite geometrical and physical meaning.
- (2) We define the *canonical thermal curve* as the curve obtained upon graphing the average number of thermal quanta per mode against the average entropy per mode. The canonical thermal curve is very flat for small values of the entropy, but becomes exponential with increasing entropy (Fig. 1).
- (3) One of the two entropy components is simply related to the thermal energy of the system and, thus, corresponds to the Benzinger compensation entropy¹, S_c . The other entropy term corresponds to Benzinger's free entropy, S_f , which together with the zero-point energy level (i.e., the energy at zero temperature) produces the Helmholtz free energy of the system. The two entropy components have simple geometrical representations in terms of the canonical thermal curve (Fig. 1).

- (4) Upon scaling the canonical thermal curve with the size of the thermal quantum, we obtain a family of equilibrium thermal curves for the energy, E , of the system as a function of the entropy (Fig. 2). We refer to the quadrant of the E - S plane for positive E and S as the E - S field, which for large S corresponds to the *classical region* and for small S produces the *quantum aberration region*. The properties of each region is explored (Fig. 3).
- (5) Chemical and physical processes, in general, may produce changes both in the sizes of mode quanta (i.e., mode frequencies) and the zero-point energy levels of the various thermal modes.
- (6) The concept of *thermal mode thawing* is developed, whereby a given type of mode undergoes a transition between the strongly quantum and the strongly classical regions of the E - S field as a result of a thermodynamic process.
- (7) *Isothermal mode thawing curves* are defined as curves in the E - S plane corresponding to constant temperature. It is shown that these curves are horizontal in the classical region and curve in to the origin in the quantum aberration region. This results in a meaningful structure for the E - S field (Fig. 2).
- (8) Because of the definite quantitative relation between the two entropy components, it is not possible for the Benzinger compensation term, *per se*, to account for observed linear thermodynamic relations.
- (9) We show that linear thermodynamic relations can be attributed to the number of thermal modes undergoing frozen $\leftrightarrow$ thawed or thawed $\rightarrow$ thawed transitions within the thermodynamic process (e.g., chemical reaction). Specifically, systematic variation in the number of such modes among the members of a family of reactions, with other parameters of the process being held constant, is sufficient to produce linearity.

The consequences of the above findings for biomolecular processes and for condensed system processes, in general, are considered.

4. Results

The novel aspects of the thermodynamic relations developed above are based largely on two important features: (i) the observation that the entropy of thermal modes, expressed in terms of thermal quanta according to Planck, is naturally given by the superposition of two terms, each having definite geometrical and physical meanings; and (ii) the concept of thermal mode freezing and thawing (as well as softening) as a fundamental factor in determining thermodynamic relations for families of chemical reactions. Both features depend on the quantum nature of thermal modes.

The natural decomposition of S into two terms provides a theoretical basis to Benzinger's observation¹ that the thermal energy of a system is cancelled by T times an entropy term, S_c . In the "strong" classical limit we have $S_c = g$, the number of thermal modes (dimensionless unit). We have shown, however, that the equality of TS_c with the thermal energy cannot account for observed linearity between ΔH and ΔS . The remaining free entropy term, ΔS_f , is at least as large as ΔS_c .

The concept of thermal mode transformation, in general, and thermal mode thawing, in particular, and their role in governing the relative changes of thermodynamic quantities are the most important elements of this work.

More specifically, we have shown that a sufficient condition to have a linear relation between ΔH and ΔS for a family of reactions is that the number of thermal modes undergoing a frozen $\leftrightarrow$ thawed transformation varies systematically among the family members. It is reasonable that the varying parameter be the number of modes because experimentally the quantity which varies from reaction to reaction is either the size of the solvent molecule or the size of the substituent group among a homologous series. Thus, different (but similar) solvents, for example, contribute a different number of thermal modes of the same type to the system.

Furthermore, it is noteworthy that the quantities which vary in families of reactions do not involve the reaction sites (or centers) directly, rather, they involve secondary, or background, degrees of freedom which are coupled to the primary reaction centers. This indicates the importance of moderate to weak coupling with otherwise passive degrees of freedom in governing the magnitudes of the thermodynamic changes in the system. In other words, *thermal modes which undergo transformations in a reaction are part of the reaction system and are not simply part of the heat bath.*

Overall, the picture which emerges from these studies focuses on the quantum mechanical nature of thermal modes and the fundamental role of these modes in determining the dynamical properties of systems at (or near) thermal equilibrium. In the following, we briefly describe the key concepts related to the ongoing development of this research topic.

1) Local Phase Transitions

The fact that linear thermodynamic relations are associated with changes in secondary structural components of a reaction, such as solvent or substituent groups, suggests that strong transformations in thermal modes belonging to these components play an important role. Lumry and Gregory categorize the structural regions of protein molecules according to a *knot* and *matrix* concept and terminology.^{7,8} The term *knot* refer to a rigid structural region usually containing the active or binding site and the term *matrix* refers to larger, more loosely organized structural regions. We suggest that, during the course of a protein reaction (which may be part of its natural function), associated solvent and background structural regions of the protein may undergo thermodynamic transitions that are like phase transitions in character.¹² Such local (or pseudo-) phase transitions could be tightly coupled to the protein reaction, thus contributing greatly to the overall thermodynamics of the process. The possible collective nature of the *thermodynamic modes* associated with these background transitions needs to be explored.

2) *Irreversible Thermal Curves*

The thermal curves shown in Figs. 1 and 2 are equilibrium curves associated with one kind of thermal mode. Real systems have many kinds of thermal modes having different thermal relaxation times. If these systems are quenched, for example, the actual thermal curve will lie above the equilibrium curve. A point on this curve corresponds to a metastable quenched state, having a higher free energy than the state on the equilibrium thermal curve at the same temperature. This amounts to a time-dependent removal of certain slowly relaxing thermodynamic modes from the pool of thermal modes of the system. These removed modes are thus transformed into effective components having increased work potential (compared to the fully equilibrium state on the equilibrium thermal curve). The possible importance of such thermally generated metastable states is being studied.

3) *Isothermal Subsystem Cycles*

The natural functions of many proteins involve cyclic processes. All parts of the protein molecule and associated molecules, such as solvent, which undergo structural changes during the cycle may be regarded as a subsystem operating at constant temperature. It is convenient and instructive to consider such isothermal subsystem cycles in terms of the energy-entropy plane. We have recently made conjectures about the quantum mechanical nature of such cycles in biological processes.¹³ The possible role of transitions of background modes associated with solvent and protein structural regions, which are coupled to the centers of primary functions (e.g., active sites), is now under study. Observable linear thermodynamic relations provide a valuable tool.

REFERENCES

1. Benzinger, T.H., *Nature* 1971, 229 100-102.
2. Danforth, R., Krakauer, H., and Sturtevant, J.M., *Rev. Sci. Instr.* 1967, 38, 484-487.
3. Leffler, J.E., and Grunwald, E., In *Rates and Equilibria of Organic Reactions*; John Wiley and Sons: New York 1976.
4. Leffler, J.E., *Science* 1953, 117, 340-341.
5. Saltiel, J. and Sun, Y.-P., *J. Phys. Chem.* 1989, 93, 6246-6250.
6. Sun, Y.-P. and Saltiel, J., *J. Phys. Chem.* 1989, 93, 8310-8316.
7. Lumry, R. and Gregory, R.B., In *The Fluctuating Enzyme*; Welch, G. R., Ed.; John Wiley and Sons: New York 1986; pp 1-190.
8. Lumry, R. and Gregory, R.B., *J. Mol. Liquids* 1989, 42 113-144.
9. Chun, P.W., *Internat. J. of Quantum Chem.: Quantum Biol. Symp.* 1988, 15, 247-258.
10. Planck, M., *Verhandlungen der Deutschen Physikalischen Gesellschaft* 1900, 2, 237-245. [Engl. translation by ter Haar, D., *The Old Quantum Theory*; Pergamon Press: Oxford, 1967, pp 82-90].
11. Planck, M., *Annalen der Physik* 1901, 4, 553-563.
12. A caveat arises when one considers first-order phase transitions in the context of conformational changes in complex molecular systems such as proteins and molecule-solvent coupled systems. Clearly, the thermal denaturation of macromolecules, such as those considered in Refs. 1, 2, 7, and 8, involve rather sharp transitions of conformation in a narrow temperature range. Such transitions have some of the features of phase transitions. This raises the question as to whether these systems should be considered as being fixed in composition with a temperature dependent microscopic structure or whether they should be regarded as a temperature dependent equilibrium mixture of different components (molecular species), as is done in Refs. 1, 2, 7, and 8 (and in most other studies). There is merit in both approaches. In this paper we adopt the conventional approach, with the understanding that we are now investigating conformational transitions in macromolecules in terms of "pseudo-phase transitions."
13. Rhodes, W., *J. Mol. Liquids* 1989, 41, 165-180.

FIGURE CAPTIONS

Fig. 1 The canonical thermal curve for a system having one type of thermal mode.

a) Broad entropy range, showing that $\bar{S}_c \approx 1$ for large $\bar{S}$; $\bar{a}$ is a dimensionless Helmholtz potential.

b) Region of small $\bar{S}$, for which $\bar{S}_c < 1$.

Fig. 2 Thermal equilibrium curves for representative values of the thermal quantum size, ϵ . Isothermal mode thawing curves are indicated by dotted curves for three temperature values. The values of $\bar{E}, \bar{S}, \bar{A}$, and T are indicated for the state labelled by the dot. The numerical scale for $\bar{E}$ gives values relative to $\bar{E}_0$.

Fig. 3 The $\bar{E} - \bar{S}$ field showing the classical region, consisting of horizontal isothermal mode thawing curves, and the quantum aberration region, in which the isothermal mode thawing curves are attracted to the origin. A representative value of the Helmholtz potential, $\bar{A}$, is shown. The slope of each line segment is the temperature for the state specified at the midpoint of that segment.

FIGURE 1a

FIGURE 1b

FIGURE 2

FIGURE 3

B. Biological Macromolecule Activation Cycles

1. *Quasi-Phase Transitions in Protein Molecules.*

We have recently investigated the properties of thermal modes in the harmonic limit. The purpose of these studies was to understand the relation between those modes which contribute to thermal equilibrium (thermal modes) and those modes which contribute to raising the work potential (free energy) of the system. The role of thermal mode thawing and thermal mode softening in chemical processes has been explored.

A major result of these studies is the concept of quasi-phase transitions as part of the natural functions of biological macromolecules (esp. proteins). The idea is that bound solvent and bulk regions of a protein molecule undergo "freezing" and "thawing" associated with the chemical and conformational transformations of the macromolecule during its operating cycle. We have made progress on the following aspects.

- (i) Quasi-phase modes. We have formulated a potential energy diagram in terms of a phase coordinate which is prototypic of a phase transition. It shows how energy and entropy change upon transition, so as to maintain constant (Helmholtz) work potential.
- (ii) The phase cell concept. It is clear that phase transitions require a minimum number of modes which is much greater than 1. This results from the collective nature of phase modes and the requirement of an order parameter. The minimum number of coupled modes which will support a phase transition we refer to as a *phase cell*. We have proposed that the phase-like regions in protein molecules consist of one, or a few, phase cells. Progress has been made to develop the statistical mechanics of a system in terms of phase cells and in a manner which takes into account an appropriate order parameter, such as density. It is important to show how the "melting" curve for quasi-phase transitions differ from those for true phase transitions.

- (iii) Harmonic equivalence for thermal modes. We have shown how the actual thermal modes of a system at any given temperature can be replaced by an equivalent set of harmonic oscillator modes (*vide supra*). This simple principle is useful for analyzing thermal curves in the energy-entropy plane. In particular, we find it useful for studying thermal curves of quasi-phase transitions in the energy-entropy plane.

2. *Isothermal Protein Cycles*

In our studies of quantum mechanical mechanisms associated with the thermodynamics of protein conformational transitions we have found it quite useful to use the energy-entropy plane, whereby the ordinate is energy and the abscissa is entropy. Thermodynamic states are represented by points (E_i, S_i) , where fixed values of other extensive variables are understood. The protein (macromolecules) is represented by M and a small molecule (substrate, ligand, activator, etc.) is denoted by X . We are interested in the coupled process $(M, X) \rightarrow M^*, X'$ and the mechanisms of free energy transfer. Two important features suggested by our research are the following.

- (a) Temperature buffering in biological macromolecules. We propose that quasi-phase transitions in protein molecules, concomitant with $M \rightarrow M^*$, provide a thermal "sponge" which acts to absorb or release heat associated with the process $X \rightarrow X'$. In acting as a thermal sponge (heat source/sink), the quasi-phase transitions also act as a temperature buffer. The propensive local temperature changes may not be so important in many laboratory situations or in other nonbiological processes, but large local temperature changes can be traumatic for molecular biological processes.
- (b) Free energy transfer by protein molecules. Protein molecules exist in many metastable conformational states, which we denote as $M(E_i, S_i)$. These states form a grid of points on the E-S plane. The mesh of this grid is made finer by quasi-phase transitions in the protein molecule. At temperature T each point (E_i, S_i) corresponds to a definite Helmholtz work

potential (free energy), A_i . In geometrical terms A_i is the intercept of the energy axis with a line of slope T passing through the point (E_i, S_i) . For a given protein molecule it is possible that A_i is degenerate, as determined by the family of points $(E_i, S_i)_T$ on the line T . Here, degeneracy is defined as the existence of more than one thermodynamic state having a given value, A_i , of the free energy.

We propose the following: (i) In the evolution of biological systems, there has evolved a propensity for free energy (work potential) to be conserved in biochemical reactions. (ii) Protein molecules function as mediators of free energy transfer. If $-\Delta A$ is the free energy change for the biochemical reaction $X \rightarrow X'$, then the coupled protein transformation $M \rightarrow M^*$ has free energy change in the neighborhood of $+\Delta A$. (iii) There is a propensity for the overall entropy change of the couple $(X, M) \rightarrow (X', M^*)$ to be minimal. This arises from the need for local temperature buffering. (iv) Quasi-phase transitions in the protein molecule provide the thermal sponge needed for T buffering/entropy conservation.

The overall result of the above proposition is:

$$(\Delta E_M, \Delta S_M) \approx (-\Delta E_X, -\Delta S_X)$$

The double requirement that both work potential and entropy be conserved seems to place severe demands on the protein molecule, M . However, a dense grid of metastable thermodynamic states, $M^*(E_i, S_i)$, resulting from a multitude of conformations and quasi-phase transitions gives the activated (excited) protein molecule a high degree of thermodynamic flexibility. The transition $M \rightarrow M^*$, which is driven by coupling to the $X \rightarrow X'$ subsystem, is somewhat analogous to a spectroscopic transition from a discrete state to a continuum.

Consider the protein molecule activation

$$M(E_o, S_o) \rightarrow M^*(E_i, S_i)$$

at temperature T. We may regard this isothermal transition as occurring by the virtual path

$$M(E_o, S_o) \rightarrow M^*(E_o + \Delta A, S_o) \rightarrow M^*(E_i, S_i).$$

In those cases where the second step is a quasi-phase transition, the path shown is real. An analogous virtual path for the $X \rightarrow X'$ subsystem can be formulated. Since X is usually a small molecule subsystem (e.g., metabolites), there would be no quasi-phase transition and the virtual path could not be real. This difference between the M and X subsystems underscores the importance of protein molecule versatility. Protein molecules evolve to be thermodynamically adapted to their $X \rightarrow X'$ subsystems!

One of the important concepts to come out of our current work on this subject is that of thermodynamic resonances. A thermodynamic resonance is defined as *a coupling between two subsystems in which free energy is transferred from one subsystem to the other with conservation of both energy and entropy*. We propose that protein molecules evolve in a manner that produces thermodynamic resonances (or near-resonances) by coupling with their "substrate" reactions.

II. Coherent Dynamics

A. Multimode Squeeze Operators and Squeezed States

Multimode squeeze and rotation operators are defined such that they have extremely similar algebraic properties as those of their single-mode counterparts. We have shown that the introduction of N-mode squeeze operators provides a convenient set of parameters to describe the variances of the quadrature amplitudes in multimode Gaussian squeezed states. Some important properties of

these N-mode unitary operators were investigated. It was also shown that the time-evolution operator for a general N-mode quadratic Hamiltonian can be conveniently expressed as an operator product containing an N-mode squeeze operator, and N-mode rotation operator, and an N-mode displacement operator.

B. The Phase Operator

The description of the phase properties of a quantum mechanical harmonic oscillator in terms of a Hermitian phase operator has been a difficult and an unsolved problem for more than sixty years since the first attempt of Dirac in his description of the quantized electromagnetic field in 1927. The difficulty lies in the fact that an appropriate Hermitian phase operator that is the analog of the classical phase variable is hard to define. Following Dirac, many researchers such as Louisell, Susskind and Glogower, Lerner, Carruthers and Nieto, and Barnett and Pegg attacked this problem but all failed to establish a well defined Hermitian phase operator. It is nowadays a common belief that no such phase operators can be constructed, and the phase is considered a classical observable without a corresponding Hermitian operator counterpart. Because of their crucial importance in quantum optical processes, phase properties of harmonic oscillators have recently attracted substantial interest, especially since the experimental observations of squeezed states of light which are characterized by the phase sensitive quantum noise in the two quadrature amplitudes. However, all the work done so far on the phase properties of harmonic oscillators uses phase operators that are not well defined. This implies that previous work does not give a proper description or analysis of the phase properties but can only provide some information about the phase indirectly.

Contrary to the common belief that no Hermitian phase operators can be found to describe the phase properties of a quantum harmonic oscillator, we have succeeded in constructing a well-defined Hermitian phase operator with an appropriate classical limit. The approach is different in nature from those of many previous attempts which were more or less based on the idea of polar decomposition of the annihilation operator. The eigenvalue spectrum and eigenstates of the phase

operator are obtained. We have investigated some important properties of the phase operator and phase states. We find, for example, that the eigenstates of the number operator for positive semi-definite integers, n , are not states with random phase unless $n \rightarrow \infty$.

C. Dynamics of Squeezed States of Multimode Harmonic Oscillators

We have shown that the time evolution of systems governed by multimode quadratic Hamiltonians can be described succinctly and conveniently using newly-defined multimode squeeze and rotation operators. As an interesting application of this formulation, we have formulated the dynamics of time-dependent multimode harmonic oscillators. The most interesting dynamical property of such oscillators is the time-dependent transition amplitudes among the multimode boson number states. These transition amplitudes are important to many physical problems. For example, the evaluation of these transition amplitudes is the crux of time-dependent formulation of molecular spectroscopies. When the Duschinsky effect is brought into consideration, the normal coordinates for the excited-state vibrations are different from those of the ground state. Generally, the normal vibration coordinates undergo a rotation and possibly a displacement upon electronic excitation. A photon-induced transition from the ground state is a case in which a multimode harmonic oscillator undergoes a sudden change in frequencies, masses, and normal coordinates. We have shown that this is one of the simplest cases in which the time evolution operator can be solved exactly and hence the time-dependent transition amplitudes can be analytically evaluated.

III. Reaction Rate Theory

A. Projected Equations of Motion

Most theories of reaction rates are (near) equilibrium thermodynamic theories in that they assume that the reactants are at thermal equilibrium at all times and that they flow or diffuse into products which instantly (rapidly) relax to thermal equilibrium. Examples are Eyring Absolute Rate Theory and Kramers Theory. A hydrodynamic metaphor is a filled fluid reservoir (reactants) separated from a (nearly) empty reservoir by a slightly porous wall. This concept of a reaction is one of (near) equilibrium flow or diffusion.

We have recently begun developing the projected Schrödinger and Liouville equations for the flow from one region of conformational space (reactants) to another region (products). The two regions are coupled only at the junction of the two subspaces (transition state). By a simple, but adept, mathematical manipulation my graduate research student, S. Svirschevsky, has shown that the two subspaces are coupled by the quantum mechanical flux operator. This is a general and promising result which we are now enthusiastically developing. One objective is to relate our results with conventional transition state theory. The flux operator should provide new features of the transition state (structure) for chemical reactions.

B. Theory of Reaction Rates at Avoided Crossings

The energy surfaces of a molecule frequently cross whenever an approximate Hamiltonian is used. When a more exact Hamiltonian is used, however, energy surfaces of the same symmetry tend to repel one another at the crossing point, producing a peak on the lower-energy surface above which is a valley on the upper surface. The kinetics of motion across such avoided crossings were formulated many years ago by Landau and separately by Zener. The probability of an adiabatic

crossing depends on the coupling between the two surfaces and the difference between the slopes of the two diabatic surfaces at the crossing point. Avoided crossings play a role in electron transfer processes and in conformational transitions in molecules.

In formulating a theory for reaction rates at avoided crossings we make use of both Landau-Zener theory and transition rate theory. The preexponential factor of the rate constant contains an integral over the momentum of the reaction coordinate in which the integrand is a product of three terms: (a) the Maxwell-Boltzmann Gaussian; (b) the Landau-Zener (L-Z) adiabatic transition probability; and (c) a linear momentum term. The L-Z term has a relatively large tail which persists at large momentum values. For weak L-Z coupling (i.e., for small coupling of the diabatic surfaces) the tail of the L-Z function strongly overlaps the Maxwell-Boltzmann function. The result is that the large momentum range dominates in its contribution to the rate constant. This corresponds to tunneling of the reaction coordinate through the well formed by the upper energy surface.

We are currently investigating the significance of tunneling in the weak coupling limit. The mass of the reaction coordinate plays a role in that the L-Z adiabatic crossing probability increases with the mass. As the mass increases the velocity for a given momentum of the reaction coordinate decreases, thereby increasing the probability of adiabatic crossover. This suggests that bound solvent may play a special role in enhancing reaction rates in such systems.

IV Fundamental Subsystem Equations of Motion

This area of research is basic to all of the others described above. It concerns the development of equations of motion for a system (primary subsystem) interacting with a background subsystem (surroundings, solvent, substituents, etc.). The purpose is to determine how dissipation arises in the dynamics of a system (W. Rhodes, J. Phys. Chem. 86, 2657 (1982); J. Phys. Chem. 87, 30 (1983)). This is a very important, ongoing area in our program.

A. Adiabatic Modulation Approach to Dissipation

During the 1980's we developed an approach to coherence relaxation (dissipation) whereby the background modes of the medium (e.g., solvent) move on the adiabatic energy surfaces formed by the interaction of those modes with the molecule (primary subsystem) in question. The motion of the wavepackets of the background modes on two different adiabatic surfaces results in a time-dependent coherent relaxation function. This is phase relaxation which breaks the phase between the two states of the molecule to which the two adiabatic energy surfaces belong. In the usual description of dephasing it is assumed that phase is broken by collisions between the molecule in question and solvent molecules. Our approach avoids this assumption and provides a clear picture in terms of overlapping quantum mechanical wavepackets:

The adiabatic modulation formulation lends itself to simple model calculations. The molecule Hamiltonian including external interactions induce transitions between states of the molecule (only nonstationary dynamics are being considered). Each state has its own energy hypersurface formed by the background medium modes. As a model we assume that the two surfaces are identical, harmonic and that they are weakly displaced relative to one another. More specifically, the minimum of one lies inside the zero-point energy contour of the other. The ground state wavefunction of one surface suddenly finds itself on the other potential surface (as a result of a molecule transition) and it begins to move on its new surface. The quantity of importance is the time-dependent overlap of the moving wavepacket with the initial one. A high degree of recurrence in a relatively short time indicates absence of coherence relaxation (dissipation). In a trial calculation we chose a surface of dimension 10 (i.e., 10 background coordinates) and used randomly chosen frequencies over a ten-fold range. The overlap (relaxation) function decayed rapidly to almost zero and over a time range of many oscillator periods the maximum recurrences were of order 10^{-8} .

These initial results are merely suggestive that very weak adiabatic modulation can produce dissipation. In order to establish their validity, we will continue studies with varying parameters and extended time ranges. It is exciting to think that in coherent quantum mechanics there is a simple basis for dissipation.

B. An Entropic Evolution Equation of Motion

This topic was conceived by and is being developed by Sperantza Svirschevsky, a graduate research assistant working with me. An equation of motion for the density operator is derived with use of calculus of variations and Lagrange multipliers. The constraints are that probability and average energy are conserved and that entropy increases with time. Preliminary results give the Lagrange multipliers in terms of familiar physical quantities and reveal constraints on the evolution of the subspaces of state vector space. This is a promising topic of study and its development is just getting under way.

PUBLICATIONS FOR THE PERIOD: 1990-1993

**Laboratory of William Rhodes
Chemistry Department
Florida State University**

1. Localization vs. Delocalization of the Davydov Soliton: Dispersive Collapse Under the Schrödinger Equation, W. Rhodes and A. Nicholls, Phys. Rev. Letters, 64, 1174-77 (1990).
2. Multimode Squeeze Operators and Squeezed States, Xin Ma and W. Rhodes, Phys. Rev. A, 41, 4125 (1990).
3. Phase Properties of the Quantized Single-mode Electromagnetic Field, Xin Ma and W. Rhodes, Phys. Rev. A, 43, 2576 (1991).
4. Phase of Quantum Harmonic Oscillators, Xin Ma and W. Rhodes, Phys. Rev. A (accepted with revision).
5. Dynamics of Time-dependent Multimode Harmonic Oscillators, Xin Ma and W. Rhodes, Phys. Rev. A (submitted).
6. Eigenstates of Unitary k-Boson Operators, Xin Ma and W. Rhodes, Phys. Rev. A (submitted).
7. Quantum Thermodynamics: The Microscopic Basis of Entropy and Linear Thermodynamic Relations, W. Rhodes, J. Phys. Chem., 95, (1991).

RELATION TO STATED OBJECTIVES - PLANS FOR CONTINUATION

It was pointed out in the original proposal that the number of specific objectives exceeded what we would probably achieve, but that objectives in each of the areas of research would be attempted. This has turned out to be the case. Most of the objectives not attained are being included in the renewal proposal.

GRADUATE STUDENTS TRAINED

1. Sperantza Svirschevki - second year graduate student in the Molecular Biophysics Doctoral Program. She is expected to receive her doctorate in 1995.

Research Topic: Subsystem Equations of Motion; Irreversibility; Reaction Rate Theory

OTHER FEDERAL SUPPORT

The Department of Energy provides sole federal support for this research program.

PRESENT STATE OF KNOWLEDGE

The ways in which biological systems organize, coordinate, and control the various levels of molecular structure, energy channeling, and information processing is one of the most fundamental unsolved problems in science. Until a few years ago the available methods put emphasis on determination of structures and biochemical pathways. Recently, however, the development of computational technology and methods of studying fast processes in complex molecular systems

has shifted the emphasis to dynamical and global aspects of molecular level biology. The stage is being set for major breakthroughs in the near future. Quite probably the 21st century will be to biology what the 20th century has been to physics.

One of the central problems of molecular level biology is to understand the relations between energy transformation and control mechanisms. The purpose of our research is to develop concepts and knowledge of principles and mechanisms whereby biological macromolecules operate to control living processes.

Knowledge of how natural systems work is necessary to understand how environmental perturbations, chemical and radiative, affect living organisms at the molecular level.

It is essential that DOE strongly support basic research which is fundamental to its mission; namely, the development of knowledge and understanding of all aspects of energy and its relation to living systems. This includes bioenergetics as well as the effects of environment on life processes. Thus, along with research on energy utilization by society and other practical aspects of energy, this area of basic scientific research is vital to the mission of DOE.

ESTIMATE OF UNOBLIGATED BALANCE

The unobligated balance on October 31, 1993 is estimated to be \$9,000.

**DATE
FILMED**

8 / 30 / 93

END