AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

40/6-8-93 95(2)

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76-CHO-3073

PPPL-2902
UC-420

PPPL-2902

EXPANSION OF PARAMETER SPACE FOR
TOROIDAL ALFVÉN EIGENMODE EXPERIMENTS IN TFTR

BY

K.L. WONG, J.R. WILSON, Z.Y. CHANG, ET AL.

MAY, 1993

PPPL PRINCETON
PLASMA PHYSICS
LABORATORY

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial produce, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

NOTICE

This report has been reproduced from the best available copy.
Available in paper copy and microfiche.

Number of pages in this report: 29

DOE and DOE contractors can obtain copies of this report from:

Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831;
(615) 576-8401.

This report is publicly available from the:

National Technical Information Service
Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22161
(703) 487-4650

Expansion of Parameter Space
for Toroidal Alfvén Eigenmode Experiments in TFTR

K. L. Wong, J. R. Wilson, Z. Y. Chang, E. Fredrickson, G. W. Hammett,
C. Bush, R. Nazikian, C. K. Phillips, J. Snipes*, and G. Taylor

Princeton Plasma Physics Laboratory,
Princeton University, P.O. Box 451, Princeton, New Jersey 08543.

Abstract

Several techniques were used to excite toroidal Alfvén Eigenmodes in the Tokamak Fusion Test Reactor (TFTR) [Proceedings of the Thirteenth International Conference on Plasma Physics and Controlled Nuclear Fusion Research, 1990, Washington, D. C. (IAEA, Vienna, 1990), Vol. I, P. 9] at magnetic fields above 10 kG. These involve pellet injection to raise the plasma density, variation of plasma current to change the energetic ion orbit and the q-profile, and ICRF heating to produce energetic hydrogen ions at velocities comparable to 3.5 MeV alpha particles. These experimental results are presented and relevance to fusion reactors are discussed.

MASTER

I. Introduction

Toroidal Alfvén Eigenmodes¹ (TAE) may become unstable in tokamak fusion reactors. They are driven by the energetic alpha particles produced in d-t fusion reactions. This instability is potentially dangerous because TAE modes can eject the fast particles before they slow down in the plasma, and therefore reduce the alpha-heating efficiency. Until a few years ago, it was generally believed that one could see this instability only in fusion reactors because the Alfvén speed V_A in tokamaks was comparable to velocities of light ions in the MeV energy range. However, by operating at very low magnetic field² (about 10 kG), it is possible to reduce V_A to the magnitude comparable to the injection velocities of existing neutral beams used for plasma heating. Two concurrent simulation experiments were performed with neutral beams tangentially injected into tokamaks at low magnetic field, and TAE modes were indeed excited.^{3,4} As expected, the instability can eject energetic particles from the plasma, but the instability threshold is substantially higher than the initial estimate based on electron Landau damping alone. Stimulated by these experimental results, several important damping mechanisms have been identified recently, namely, ion Landau damping,⁵ collisional damping,⁶ continuum damping⁷ and radiation damping.⁸ TFTR normally operates at magnetic fields above 30 kG. Many diagnostics cannot function at 10 kG, and plasmas at such a low magnetic field have low plasma temperature and high collision frequency. They are not well diagnosed, making it very difficult to extrapolate these results to the fusion reactor regime. In the previous experiment,^{3,9} collisional damping dominates, and recent theory¹⁰ can explain the instability threshold within a factor two. However, for fusion reactor parameters, collisional damping becomes less

important. If one wants a high alpha particle heating efficiency, this instability can impose some constraints on the parameter space in which fusion reactors operate. Therefore, it is important to map out the instability boundary for plasma parameters relevant to fusion reactors. The plasma parameters in tokamaks are closely related to the magnetic field. In the past two years, several experiments were performed in TFTR in order to expand the parameter space for TAE mode studies, i.e., to excite TAE modes at higher magnetic field. We use several different techniques based on intuitive considerations derived from the properties of the instability. The experimental results are presented in this paper. In sect. II, we describe an experiment which employs pellet injection to raise the plasma density. Another method which involves variation of plasma current is described in sect. III. In both of these experiments, TAE modes are driven by 100 keV deuterium neutral beams tangentially injected into circulating particle orbits. Recent theories^{10,11} show that energetic trapped ions can also excite TAE modes. This was observed during ICRF hydrogen minority heating in TFTR¹². The instability amplitude behavior is different from that driven by neutral beams. This feature and a qualitative explanation are presented in sect. IV. A brief summary is given in sect. V.

II. Pellet Injection

In order to excite TAE modes with energetic ions, the ion velocity V_α must be comparable to the Alfvén speed V_A . This calls for higher fast ion velocities for TAE excitation at higher magnetic field. At the same

injection energy, changing the beam ion species from deuterium to hydrogen would raise V_{α} by 40%, but this requires a certain change of neutral beam hardware which is highly undesirable because it interferes with other experiments on TFTR. We can achieve the same condition by a factor of two increase in plasma mass density which reduces V_A by 40%. This can be done with pellet injection. Two lithium pellets are injected into a 630 kA plasma at 15 kG magnetic field with 14 MW of deuterium neutral beams at energies up to 100 keV. Before pellet injection, there is no sign of TAE activity. Immediately after the pellets go in, the density jumps up and the plasma temperature cools down. Figure 1 shows the evolution of the line-integrated electron density measured by an infrared interferometer, and the total plasma kinetic energy determined magnetically. Evidence of TAE activity appears in the Mirnov coil signals 30 ms after pellet injection. It lasts more than 100ms. Figure 2 shows the frequency spectrum of the Mirnov coil signal at three different times. The peak signal has a frequency near the expected TAE frequency, and it drifts towards higher frequencies at later times because the density decreases in time. Beam emission spectroscopy¹³ observes signals at the same frequency, but the analysis requires long integration time, and cannot provide reliable measurements of the mode structure and the mode numbers. Due to the transient nature of the plasma and the lack of high quality data, detailed analysis becomes very difficult. We are not able to do reliable transport modelling of the plasma. With the same technique, we tried to excite TAE modes at 20 kG magnetic field, and the attempt was not successful.

III. Plasma Current Variation

The existence of TAE modes depends on the structure of the Alfvén spectrum which is sensitive to the q -profile. In the previous experiment³, the V_α/V_A ratio was varied by changing the magnetic field, and the plasma current was changed by the same proportion in order to maintain similar q -profiles. It was found that at $I_p=630\text{kA}$, $B=15\text{kG}$, TAE modes disappeared. These data were presented in Fig. 1 of Ref. 3. Recently, we found out that at $B=15\text{kG}$, TAE activity returned when we lowered the plasma current to 420 kA. The results are presented in Fig. 3. Of course, the Alfvén spectrum is changed when we change $q(a)$ from 2.8 to 4.2. At the same time, the excursion of the energetic ion orbit from the flux surface becomes larger when we lower the plasma current. When $q(a)$ is larger, it is more likely that the TAE mode frequency will intersect the Alfvén continuum and the mode will encounter continuum damping. Our observations are consistent with the orbit effect. The finite orbit effect on the driving term of the instability was analyzed by Fu and Cheng¹⁰. The results are schematically shown in Fig. 4. If the orbit size is negligible, the strength of the instability drive rises abruptly when V_α exceeds V_A . The sharp-edge transition is smoothed out by the orbit effect as shown by the dotted line in Fig. 4. When the instability is driven by circulating ions with $V_\alpha > V_A$, finite orbits have a stabilizing effect which reduces the growth rate. In our experiment, $V_\alpha \sim 0.8V_A$ and finite orbits have a destabilizing effect as indicated by the arrow in Fig. 4. It enhances the driving term and makes it easier to excite TAE modes. This is only a qualitative explanation. We do not know the plasma parameters well enough for detailed quantitative analysis.

The effect of q-profile variation on the TAE instability can be explored by current ramping. We find it easier to excite TAE modes in TFTR immediately after the plasma current is ramped up. This is demonstrated in Fig. 5. At 15 kG magnetic field, 630 kA plasma current and 14 MW of beam power, there is no sign of TAE activities in the middle of the flat current waveform. However, by ramping I_p from 350 kA to 630 kA, evidence of TAE activities appear in the Mirnov coil signal shortly after I_p reached 630 kA. It lasts more than 200 ms as shown in Fig. 6. A weaker peak at approximately twice the TAE frequency is also observed. The major difference in these plasmas is in the q-profile. A lower magnetic shear is expected with I_p ramping up. This tends to reduce continuum damping⁷ as well as collisional damping.¹⁰ We cannot measure the q-profiles in these plasmas and it is not clear whether or not the TAE modes intersect the Alfvén continuum. Collisional damping alone can explain the instability threshold, and continuum damping, if present, can be of comparable magnitude. The q-profile evolves on the L/R time scale which is comparable to the time interval in which TAE activities appear.

The techniques discussed so far allow us to excite TAE modes with 100 keV deuterium neutral beams in 630 kA plasmas at 15 kG magnetic field. This is an improvement compared to the 10 kG plasmas since the Michelson interferometer system¹⁴ on TFTR can operate at 15 kG and we can obtain the time history of the electron temperature profile from electron cyclotron emission. However, there is no significant improvement in the plasma parameters, and the data are still not good enough for reliable transport analysis. Therefore, detailed calculation of the instability threshold cannot be carried out. Attempts to drive TAE modes at 20 kG with these techniques were not successful. It appears necessary

to raise the energetic ion velocity in order to excite TAE modes at higher magnetic field.

IV. Energetic Trapped Ions from ICRF

It is well known that very high energy hydrogen ions (0.5-1.0 MeV) are produced in ICRF hydrogen minority heating in TFTR. Recent theory shows that these energetic trapped ions can excite TAE modes via wave-particle drift resonance.^{10,11} MHD activities near the TAE frequency ($\omega = V_A/2qR$) was observed in ICRF experiments, and they were identified as TAE modes¹² from the density dependence of the mode frequency. The TAE mode signals are dominant in the Mirnov coil frequency spectrum as shown in Fig. 7. The main peak has some fine structures which consist of several peaks at slightly different frequencies.¹² They correspond to TAE modes with different mode numbers. The same frequency spectrum is observed by the microwave reflectometer in most of the cases. In these experiments, the toroidal magnetic field at R=2.62m is typically set at 33kG so that the hydrogen cyclotron frequency on the magnetic axis is close to the RF frequency of 47 MHz. The central plasma temperature is about 5 keV. With helium gas for initial break down and density feedback, these plasmas are very reproducible, and most of the diagnostics work well. The threshold RF power is about 3 MW while we typically operate at 5 MW. These TAE modes behave differently from those driven by tangential neutral beams. The beam driven TAE modes have a bursting behavior. When the number of fast particles is above the instability threshold, the mode grows rapidly; a few percent of the fast particles are ejected³ and the mode becomes stable again. The population of the energetic particles is

clamped near the instability threshold.⁹ This phenomenon is very similar to the fishbone¹⁵ activity driven by perpendicular neutral beam injection. The pulsation state can be modelled by two coupled first order differential equations.^{16,17} Ejection of fast particles occurs in a very short time interval (comparable to the bounce period), and the damping time of TAE mode is short compared with the time required to raise the energetic particle population back to the instability threshold. Therefore, a pulsation scenario is expected from the theory of Berk, Breizman and Ye.¹⁸

In TAE modes driven by trapped ions, the amplitude usually varies slowly with a typical time scale of several hundred milliseconds. The mode amplitude can be modulated by sawtooth activities as shown in Fig. 7a, or it can be slowly varying so that it evolves into a quasi steady state as in Fig. 7b or Fig. 7c. We believe that the difference in the mode behavior from the neutral beam injection case^{3,9} is due to the different energetic ion orbits.

In the hydrogen minority ICRF heating¹⁹ experiment, the fundamental cyclotron resonance zone for hydrogen ions is on a vertical plane passing through the plasma core. The hydrogen ions are heated to a perpendicular temperature of about 500 keV. The fast ion population peaks near the plasma core where the temperature is the highest. We expect most of the energetic ions to be magnetically trapped with reflection points near the cyclotron resonance zone. Following G. W. Hammett,²⁰ we can represent their velocity distribution function in the following approximate form:

$$f(W, \xi) = f_0(W) [\exp(-|\xi - \xi_0|/\sigma_\xi) + \exp(-|\xi + \xi_0|/\sigma_\xi)] \quad (1)$$

where $\xi = v_{||} / v$

ξ_* = pitch angle of particles with reflection points at the
cyclotron resonance zone

$$\sigma_{\xi} = \{ Z_{\text{eff}} / [4A \langle Z_i^2 / A_i \rangle (1 + W^{3/2} / W_c^{3/2})] \}^{1/2}$$

$$W_c = 14.8 A T_e \langle Z_i^2 / A_i \rangle^{2/3}$$

= critical energy at which electron drag equals ion drag.

For our experimental parameters, $T_e = 4$ keV, $Z_{\text{eff}} = 3$, $W = (3/2)T_{\perp} = 750$ keV, $A=1$ for hydrogen ions, $\langle Z_i^2 / A_i \rangle = 0.5$, $W_c = 30$ keV, $\sigma_{\xi} = 0.11$. Let ξ_1 be the pitch angle (evaluated at the resonance zone) for barely trapped particles. For $R=2.62\text{m}$, $r=0.3\text{m}$, Eq. (1) yields

$$f(W, \xi_1)/f(W, \xi_*) \sim 4.6 \times 10^{-2} \ll 1 \quad (2)$$

If the energetic ions are governed by neoclassical transport, they are expected to localize in the plasma core with $r \leq 0.3\text{m}$. However, presence of other transport mechanisms can result in a broader radial profile. At $r > 0.3\text{m}$, the ratio $f(W, \xi_1)/f(W, \xi_*)$ is even smaller than that given in Eq. (2). This means that very few of the energetic ions are barely trapped; most of them have their reflection points near the resonance zone, and it is not easy to change these particles into circulating particles in one bounce period. It would require a very large amplitude TAE mode with $\delta B/B > 10^{-2}$ for this to be possible. In this experiment, preliminary estimates based on reflectometer measurements indicate that $\delta B/B$ is of the order of 10^{-4} . Therefore, low amplitude TAE modes can eject the energetic ions only after many bounce periods of interaction; their effect on the energetic ions

is quite similar to small pitch angle scattering due to Coulomb collisions. One only expects sudden changes of the energetic ion orbits in the transition from barely circulating to barely trapped orbits and vice versa. Particle loss mechanisms due to TAE modes driven by energetic circulating ions have been investigated by extensive orbit calculations.^{21,22} It was found that shortly after the TAE modes are turned on, most of the energetic ions were lost during the transition from a barely circulating to a barely trapped orbit. This loss mechanism which happens on a very fast time scale (the bounce period of the ions) is not applicable to the energetic trapped ions for the following reasons. First, there are very few energetic ions near the trapped-circulating boundary in velocity space. Eq. (2) is derived under the assumption of pitch angle scattering by Coulomb collisions. The presence of low amplitude TAE modes would tend to raise σ_{ξ} , but we still expect the majority of the energetic trapped ions to have their reflection points near the cyclotron resonance zone. Second, a barely trapped particle has the largest guiding center orbit excursion from the magnetic surface on which the reflection points lie. Fig. 8 depicts the particle orbits projected on a poloidal plane. Let us neglect the unlikely event of multiple reversal of $v_{||}$ near the banana tips. When a barely trapped particle is changed into a barely circulating particle via pitch angle scattering, there are two possibilities. It can either go into orbit C1 which is almost the same as the outer half of the barely trapped orbit, or it can go into orbit C2 which is almost the same as the inner half of the barely trapped orbit. If the original barely trapped particle is confined, the barely circulating particle is also confined. Therefore, this process does not cause the prompt loss of energetic ions. Finally, let us consider the transition from the barely circulating orbit C2 to a barely trapped orbit which is the dominant

prompt loss mechanism for the energetic beam ions in the previous TAE experiment.³ Unlike neutral beam injection, the barely circulating energetic ions in the ICRF experiment come from barely trapped energetic ions which are confined. On the time scale of the particle transit time, these barely circulating ions can only go back to a trapped orbit very similar to the original confined trapped orbits, i.e., small pitch angle scattering does not provide a fast escape channel for the energetic particles produced by ICRF. Because of the above reasons, the major effect of the low amplitude TAE modes on the energetic ions is to move them from one trapped orbit to a neighbouring one. This aspect is quite similar to the effect due to magnetic field ripples,²³ except that the banana tips need not have the same magnetic field strength because the particle energy and magnetic moment are no longer constants of motion. The TAE modes would enhance the transport of the energetic ions across the magnetic field and relax their density gradient. It is a relatively slow diffusion process which tends to reduce the driving force for the instability. The energetic ions can slowly diffuse out of the machine rather than scatter into a prompt lost orbit and escape suddenly. The driving term for the TAE mode can be written in the following form:

$$\gamma_{\text{drive}} \propto \beta_f (\omega_{*f} / \omega - 1) F(v) \quad (3)$$

where $\beta_f \propto n_f T_f$

$$\omega_{*f} = k_{\theta} K T_f c / (e B L_f)$$

$$L_f = [d/dr (\ln n_f)]^{-1} .$$

Assuming 500 keV hydrogen ion temperature, the neoclassical diffusion coefficient for these trapped ions is about 200 cm²/sec. It is conceivable

that TAE modes can raise this by one to two orders of magnitude. Suppose the fast ions have a diffusion coefficient of $D_f \sim 10^4 \text{ cm}^2/\text{sec}$. The time it takes for a fast particle to diffuse a distance of 10 cm is about 10 ms, and one would expect L_f to change within this time scale which is short compared with the growth time shown in Fig. 7. Let us assume that L_f is a function of the TAE amplitude A whose temporal evolution is governed by the equation

$$dA/dt = (\gamma_{\text{drive}} - \gamma_{\text{damp}}) A = \gamma A \quad (4)$$

If γ can be expressed in terms of A with no explicit time dependence, then Eq. (4) is integrable and the evolution of A can be expressed in terms of the initial condition at $t=0$. From the data shown in Fig. 7, it is apparent that the TAE mode does not grow exponentially. It appears that the driving term and the damping term almost cancel each other, i.e., $\gamma \ll \gamma_{\text{drive}} \sim \gamma_{\text{damp}}$. The amplitude changes on such a slow time scale that the plasma equilibrium can change. Although we cannot measure the evolution of the fast ion density distribution, the data suggest that the plasma parameters evolve in such a way that the TAE modes are marginally unstable and one can pick a slowly varying γ to model the evolution of the mode amplitude. We attribute this to the lack of a rapid loss mechanism for the energetic trapped ions. A fast particle probe outside the plasma edge detects a continuous fast particle flux that correlates with the TAE mode amplitude.¹² This observation is consistent with our interpretation. In order to calculate the variation of the growth rate, it is necessary to know the detailed history of many plasma parameters, including those for

the energetic ions. Some of these are measured in the experiment, the rest can, in principle, be inferred from plasma modelling with a transport code. However, this is not an easy task for plasmas in this experiment. These plasmas involve at least four ion species, namely, helium which is the working gas for initial break down and density feedback, deuterium and carbon released from the limiter, and hydrogen which is the minority species heated by the ICRF waves. It is very difficult to determine the species split. Analysis of such plasmas is in progress.

V. Summary and Discussion

Several techniques were used to expand the operating regime for TAE mode experiments in TFTR. With 100 keV tangentially injected neutral beams, pellet injection and variation of plasma current allow us to excite TAE modes at 15 kG magnetic field. Attempts at 20 kG were not successful, probably due to the limited beam ion velocity. Using the energetic trapped hydrogen ions, TAE modes can be excited at 33 kG magnetic field. Approximately 80% of the RF power goes into the hydrogen ions in these experiments. With 5 MW of RF power, about 4 MW goes into hydrogen ions, which already exceeds the fusion power that is expected in the alpha particles in a $Q=0.3$ d-t plasma in TFTR.²⁴ The velocities of the hydrogen ions are comparable to those alpha particles produced by the d-t reaction. These plasmas are well diagnosed and very reproducible. It offers an excellent opportunity to study TAE modes in a low radioactivity environment. The amplitude of these TAE modes has a behavior different from those driven by neutral beams. We attribute this to the difference in the energetic ion orbits.

Looking towards the forthcoming d-t experiment in TFTR, the 3.5 MeV

alpha particles will have large orbits due to the relatively low plasma current. The typical orbit size is comparable to that of the energetic beam ions in the previous simulation experiment.³ Both the trapped and the circulating alpha particles can contribute to the excitation of TAE modes. For the modes whose amplitudes peak near the magnetic axis, the circulating particles will have a larger contribution primarily because there are more of them. If these modes are excited, it is likely to have a pulsation behavior. If the alpha particle drive is not strong enough to overcome the damping, ICRF produced energetic ions can be used to assist the TAE excitation. We plan to use hydrogen minority heating at 64 MHz and ³He minority heating at 43 MHz for this purpose. This would assure that we can study the alpha-TAE interaction in TFTR even though the alpha particles alone cannot drive the instability. In large fusion reactors where the alpha particle orbit is relatively small, TAE activities, if present, may exhibit slow amplitude variation similar to what we observe in the ICRF experiment.

Finally, we have to point out that in these simulation experiments with fast ions from neutral beams and ICRF heating, the energetic ion velocity distributions are highly anisotropic. This is different from the alpha particles produced by d-t reactions in large tokamaks where prompt orbit loss is negligible.

Acknowledgments:

The authors would like to thank the ICRF group for their support of the TAE experiment. Many helpful discussions with Dr. L. Chen and Dr. G. Y. Fu are gratefully appreciated. This project is supported by the U. S. Department of Energy under contract No. DE-AC02-76-CHO-3073.

* Permanent address: Plasma Fusion Center, Massachusetts Institute of Technology, Cambridge, MA 02139.

References

1. M. N. Rosenbluth and P. H. Rutherford, Phys. Rev. Lett. 34, 1428 (1975); C. Z. Cheng, L. Chen, and M. S. Chance, Ann. Phys. (NY) 161, 21 (1985).
2. K. L. Wong, Bull. Am. Phys. Soc. 33, 2098 (1988).
3. K. L. Wong, R. J. Fonck, S. F. Paul, D. R. Roberts, E. D. Fredrickson, R. Nazikian, H. K. Park, M. Bell, N. L. Bretz, R. Budny, S. Cohen, G. W. Hammett, F. C. Jobes, D. M. Meade, S. S. Medley, D. Mueller, Y. Nagayama, D. K. Owens, and E. J. Synakowski, Phys. Rev. Lett. 66, 1874 (1991).
4. W. W. Heidbrink, E. J. Strait, E. Doyle, G. Sager, and R. T. Snider, Nucl. Fusion 31, 1635 (1991).
5. R. Betti and J. P. Freidberg, Phys. Fluids B 4, 1465 (1992).
6. N. N. Gorelenkov and S. E. Sharapov, Phys. Scr. 45, 163 (1992).
7. M. N. Rosenbluth, H. L. Berk, J. W. Van Dam, and D. M. Lindberg, Phys. Rev. Lett. 68, 596 (1992); F. Zonca and L. Chen, Phys. Rev. Lett. 68, 592 (1992).
8. R. R. Mett, H. L. Berk, D. M. Lindberk and M. N. Rosenbluth, Bull. Am. Phys. Soc. 37, 1434 (1992).
9. K. L. Wong, R. Durst, R. J. Fonck, S. F. Paul, D. R. Roberts, E. D. Fredrickson, R. Nazkian, H. K. Park, M. Bell, N. L. Bretz, R. Budny, C. Z. Cheng, S. Cohen, G. W. Hammett, F. C. Jobes, L. Johnson, D. M. Meade, S. S. Medley, D. Mueller, Y. Nagayama, D. K. Owens, S. Sabbagh, and E. J. Synakowski, Phys. Fluids B 4, 2122 (1992).
10. G. Y. Fu and C. Z. Cheng, Phys. Fluids B 4, 3722 (1992).
11. H. Biglari, F. Zonca, and L. Chen, Phys. Fluids B 4, 2385 (1992).
12. J. R. Wilson, M. G. Bell, H. Biglari, M. Bitter, N. L. Bretz, R. Budny, C. E. Bush, L. Chen, Z. Chang, D. Darrow, P. C. Efthimion, E. Fredrickson, G. Y. Fu, R. Goldfinger, B. Grek, L. R. Grisham, G. Hammett, R. J. Hawryluk, D. Hoffman, J. C. Hosea, A. Janos, D. Jassby, F. C. Jobes, D. W. Johnson, J.

Machuzak, R. Majeski, D. K. Mansfield, E. Mazzucato, K. M. McGuire, S. S. Medley, D. Mueller, M. Murakami, R. Nazikian, D. K. Owens, H. Park, S. Paul, C. K. Phillips, A. T. Ramsey, D. Rasmussen, F. Rimini, J. H. Rogers, A. L. Roquemore, G. Schilling, J. Schivell, G. L. Schmidt, J. Stevens, B. C. Stratton, J. D. Strachan, E. Synakowski, G. Taylor, M. Ulrickson, K. L. Wong, M. Yamada, K. M. Young, M. C. Zarnstorff, and S. J. Zweben, IAEA Fourteenth International Conference on Plasma Physics and Controlled Nuclear Fusion Research, Wurzburg, Germany, 30 Sept-7 Oct, 1992, paper IAEA-CN-56/E-2-2.

13. R. D. Durst, R. J. Fonck, K. L. Wong, C. Z. Cheng, E. D. Fredrickson, and S. F. Paul, Phys. Fluids B 4, 3707 (1992).

14. F. J. Stauffer, D. A. Boyd, R. C. Cutler, M. Diesso, M. P. McCarthy, J. Montague and R. Rocco, Rev. Sci. Instrum. 59, 2139 (1989).

15. K. McGuire, R. Goldston, M. Bell, M. Bitter, K. Bol, K. Brau, D. Buchenauer, T. Crowley, S. Davis, F. Dylla, H. Eubank, H. Fishman, R. Fonck, B. Grek, R. Grimm, R. Hawryluk, H. Hsuan, R. Hulse, R. Izzo, R. Kaita, S. Kaye, H. Kugel, D. Johnson, J. Manickam, D. Manos, D. Mansfield, E. Mazzucato, R. McCann, D. McCune, D. Monticello, R. Motley, D. Mueller, K. Oasa, M. Okabayashi, K. Owens, W. Park, M. Reusch, N. Sauthoff, G. Schmidt, S. Sesnic, J. Strachan, C. Surko, R. Slusher, H. Takahashi, F. Tenney, P. Thomas, H. Towner, J. Valley, and R. White, Phys. Rev. Lett. 50, 891 (1983).

16. L. Chen, R. B. White, and M. N. Rosenbluth, Phys. Rev. Lett. 52, 1122 (1984).

17. W. W. Heidbrink, H. H. Duong, J. Manson, E. Wilfrid, C. Oberman, submitted to Phys. Fluids B.

18. H. L. Berk, B. N. Breizman, and Huanchun Ye, Phys. Rev. Lett. 68, 3563 (1992).

19. T. H. Stix, Nucl. Fusion 15, 737 (1975).
20. G. W. Hammett, Ph. D. Thesis, Princeton University, 1986.
21. C. T. Hsu and D. J. Sigmar, Phys. Fluids B 4, 1492 (1992).
22. D. J. Sigmar, C. T. Hsu, R. White, and C. Z. Cheng, Phys. Fluids B 4, 1506 (1992).
23. R. J. Goldston, R. B. White, and A. H. Rooyer, Phys. Rev. Lett. 47, 647 (1981).
24. R. V. Budny, M. G. Bell, H. Biglari, M. Bitter, C. E. Bush, C. Z. Cheng, E. D. Fredrickson, B. Grek, K. W. Hill, H. Hsuan, A. C. Janos, D. L. Jassby, D. W. Johnson, L. C. Johnson, B. LeBlanc, D. C. McCune, D. R. Mikkelsen, H. K. Park, A. T. Ramsey, S. A. Sabbagh, S. D. Scott, J. F. Schivell, J. D. Strachan, B. C. Stratton, E. J. Synakowski, G. Taylor, M. C. Zarnstroff, S. J. Zweben, Nucl. Fusion, 32, 429 (1992).

Figure Captions

Figure 1: Pellet injection experiment. (a). Line-integrated density measured by an infrared interferometer along a vertical chord at $R=2.47m$. The pellet is injected at 3.8 sec and the frequency spectra of the Mirnov coil signal at t_1 , t_2 , and t_3 are shown in Fig. 2. (b). Total plasma stored energy measured by a diammagnetic loop.

Figure 2: Frequency spectra of the Mirnov coil signal after pellet injection. (a). Spectrum at 3.84 sec. (b). Spectrum at 3.90 sec. (c). Spectrum at 3.95 sec.

Figure 3: Frequency spectra of Mirnov coil signal at various plasma current. (a). $I_p=630$ kA. (b). $I_p=420$ kA. The magnetic field is at 15 kG with 10 MW of neutral beams.

Figure 4: Schematic of the finite orbit effect on excitation of TAE modes by circulating particles based on the result by Fu and Cheng¹⁰. $V_\alpha/V_A \sim 0.8$ in our experiment.

Figure 5: Frequency spectra of Mirnov coil signal for different current waveforms. (a). Flat I_p waveform at 630 kA. (b). 350 to 630 kA ramp up waveform. The magnetic field is at 15 kG with 13 MW of beam power.

Figure 6: Evolution of the frequency spectrum of the Mirnov coil signal during and after current ramp up. The magnetic field is at 15 kG.

Figure 7: Evolution of the frequency spectrum of the Mirnov coil signal during ICRF hydrogen minority heating. The dominant peak was identified as TAE modes. (a) Amplitude modulation by sawtooth activities. The low frequency peaks in the spectra are due to sawteeth crashes (RF power turned on at 3.0 sec). (b) Slow amplitude variation (RF power turned on at 2.5 sec). (c) Growth towards saturation (RF power turned on at 2.5 sec).

Figure 8: Projection of the particle orbits on a poloidal plane. The solid

curve represents the barely trapped orbit, and the two dotted curves represent the two barely circulating orbits with opposite $v_{||}$.

Fig. 1

Fig. 2

Fig. 3

PPPL#93X0025

Fig. 4

PPPL#93X0026

Fig. 5

PPPL#93X0027

Fig. 6

PPPL #93X0020

Fig. 7

PPPL#93X0021

Fig. 8

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA
 Prof. M.H. Brennan, Univ. of Sydney, AUSTRALIA
 Plasma Research Lab., Australian Nat. Univ., AUSTRALIA
 Prof. I.R. Jones, Flinders Univ, AUSTRALIA
 Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA
 Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA
 Prof. M. Goossens, Astronomisch Instituut, BELGIUM
 Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM
 Commission-Europeen, DG. XII-Fusion Prog., BELGIUM
 Prof. R. Boucqué, Rijksuniversiteit Gent, BELGIUM
 Dr. P.H. Sekanaka, Instituto Fisica, BRAZIL
 Instituto Nacional De Pesquisas Especiais-INPE, BRAZIL
 Documents Office, Atomic Energy of Canada Ltd., CANADA
 Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA
 Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA
 Prof. J. Teichmann, Univ. of Montreal, CANADA
 Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA
 Prof. T.W. Johnston, INRS-Energie, CANADA
 Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA
 Dr. C.R. James., Univ. of Alberta, CANADA
 Dr. P. Lukáč, Komenekého Univerzita, CZECHO-SLOVAKIA
 The Librarian, Culham Laboratory, ENGLAND
 Library, R61, Rutherford Appleton Laboratory, ENGLAND
 Mrs. S.A. Hutchinson, JET Library, ENGLAND
 Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS
 P. Mähönen, Univ. of Helsinki, FINLAND
 Prof. M.N. Bussac, Ecole Polytechnique., FRANCE
 C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE
 J. Radet, CEN/CADARACHE - Bat 506, FRANCE
 Prof. E. Economou, Univ. of Crete, GREECE
 Ms. C. Rinni, Univ. of Ioannina, GREECE
 Dr. T. Mui, Academy Bibliographic Ser., HONG KONG
 Preprint Library, Hungarian Academy of Sci., HUNGARY
 Dr. B. DasGupta, Sahe Inst. of Nuclear Physics, INDIA
 Dr. P. Kaw, Inst. for Plasma Research, INDIA
 Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL
 Librarian, International Center for Theo Physics, ITALY
 Miss C. De Palo, Associazione EURATOM-ENEA, ITALY
 Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY
 Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY
 Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
 Prof. I. Kawakami, Hiroshima Univ., JAPAN
 Prof. K. Nishikawa, Hiroshima Univ., JAPAN
 Director, Japan Atomic Energy Research Inst., JAPAN
 Prof. S. Itoh, Kyushu Univ., JAPAN
 Research Info. Ctr., National Instit. for Fusion Science, JAPAN
 Prof. S. Tanaka, Kyoto Univ., JAPAN
 Library, Kyoto Univ., JAPAN
 Prof. N. Inoue, Univ. of Tokyo, JAPAN
 Secretary, Plasma Section, Electrotechnical Lab., JAPAN
 S. Mori, Technical Advisor, JAERI, JAPAN
 Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
 J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
 D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
 Prof. B.S. Uley, Univ. of Waikato, NEW ZEALAND
 Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
 Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
 Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
 Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
 Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
 Dr. O. Petrus, AL I CUZA Univ., ROMANIA
 Dr. J. de Villiers, Fusion Studies, AEC, S. AFRICA
 Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
 Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
 Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
 Dr. L. Stanflo, Univ. of UMEA, SWEDEN
 Library, Royal Inst. of Technology, SWEDEN
 Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
 Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
 Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
 Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
 Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. Apparatus, USSR
 Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
 Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
 Librarian, The Ukr.SSR Academy of Sciences, USSR
 Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
 Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
 Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
 Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
 Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
 Librarian, Max-Planck-Institut, W. GERMANY
 Prof. R.K. Janev, Inst. of Physics, YUGOSLAVIA

**DATE
FILMED**

8 / 18 / 93

END