AIIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

Assessment of CO₂ Projections

Written and edited by I.G. Enting¹,
T.M.L. Wigley² and M. Heimann³

Draft 29/3/93

Model calculations and descriptions contributed by

J.A. Taylor⁴, J. Lloyd⁵ and G. Farquhar⁵

W.R. Emanuel⁶

B.C. Cohen⁷

I.G. Enting¹ and K.R. Lassey⁸

A.A. Keller⁹ and R.A. Goldstein⁹

M. Heimann³

F. Joos¹⁰ and U. Siegenthaler¹⁰

A. Jain¹³ and D. Wuebbles¹³

T.-H. Peng⁶

C. Le Quere¹¹ and J.L. Sarmiento¹¹

J. Alcamo¹² and M. Krol¹²

L.D.D. Harvey¹⁴

J. Viecelli¹³

T.M.L. Wigley²

1. Division of Atmospheric Research, CSIRO

2. Office for Interdisciplinary Earth Studies, UCAR

3. Max Planck Institut für Meteorologie

4. CRES, Australian National University, Canberra, Australia

5. RSBS, Australian National University, Canberra, Australia

6. Oak Ridge National Laboratory, Oak Ridge Tennessee, USA

7. UN/ECE, Geneva

8. National Institute of Water and Atmospheric Research, NZ.

9. Electric Power Research Institute, PO Box 10412, Palo Alto, CA 94303, USA.

10. Physics Institute, University of Bern, CH-3012, Bern, Switzerland.

11. Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, New Jersey, 08544, USA

12. RIVM (National Institute of Public Health and Environmental Protection)
PO Box 1, 3720 BA, Bilthoven, The Netherlands

13. Lawrence Livermore National Laboratory, PO Box 808 L-263, Livermore, CA 94551-9900, USA.

14. Dept Geography, University of Toronto, Toronto, Canada

Assessment of CO₂ Projections

Contents

List of Tables	3
List of Figures	4
Abstract	5
1. Introduction	6
2. Carbon Budget Issues	20
3. Modelling Approaches	30
4. Strategies for Achieving Carbon Balance	40
5. Summary of Models	50
6. Initialisation	60
7. Summary of Results for Forward Calculations	70
8. Summary of Results for Stabilisation Calculations	80
9. Summary of Results for Impulse Responses	90
10. Assessment of Differences and Similarities	100
11. Assessment of Uncertainties	110
12. Additional Calculations	120
13. Extensions	130
14. Conclusions	140
References	200
Appendix A: Instructions for Calculations	300
Appendix B: Data Sets Provided	400
Appendix C: Descriptions of Models	500
Appendix D: Tables of Results	600
Appendix E: Plots of Results	700
Appendix F: Notation	800

Tables

5.1 Models	x
5.2 Characteristics of models	x
6.1 Mean budgets 1980-89	x
6.2 ¹⁴ C results	x
7.1 Forward Projections to 2050	x
7.2 Forward Projections to 2100	x
8.1 Stabilisation cases, peak emissions	x
8.2 Stabilisation cases, emissions in 2200	x
8.3 Air-sea fluxes, S650	x
8.4 Fertilisation fluxes, S650	x
9.1 Equilibrium impulse response	x
9.2 Perturbation impulse response	x
A.1 Land-use emissions, 1850-1990	x
A.2 Industrial emissions 1840-1990	x
A.3 IS92a-f scenarios, Total and deforestation emissions	x
A.4 IS92a-f scenarios, Industrial and deforestation emissions	x
D.1 Calculations contributed	x
D.2 Details of forward initialisations	x
D.3 Ocean fluxes pre-1990	x
D.4 Fertilisation fluxes pre-1990	x
D.5 Emissions S650	x
D.6 Emissions S450	x

Figures

Calculated emissions for S650 case	Cover
4.1 Relations between components of atmospheric budget	x
7.1 Concentrations predicted using IS92a	x
7.2 Concentrations predicted with emissions fixed at 1995 levels	x
8.1 Prescribed CO ₂ concentrations for stabilisation studies	x
8.2 Calculated emissions for S450 case	x
8.3 Calculated emissions for S650 case	x
8.4 Effect of delay in stabilisation at 550	x
9.1 Impulse response to pre-industrial flux	x
9.2 'Ocean-only' response functions	x
9.3 'Biota-only' response functions	x
10.1 Relations between future emissions and current budget	x
B.1 Prescribed fit to past CO ₂	x
B.2 Prescribed fit to past CO ₂ (Recent decades)	x
B.3 Prescribed 'land-use' flux	x
E.1 Calculated emissions for S350 case	x
E.2 Calculated emissions for S550 case	x
E.3 Calculated emissions for S750 case	x
E.4 Calculated emissions for DS450 case	x
E.5 Calculated emissions for DS550 case	x
E.6. Concentrations for 1% compound emission reduction	x
E.7. Concentrations for 2% compound emission reduction	x
E.8 Concentrations predicted using IS92b	x
E.9 Concentrations predicted using IS92c	x
E.10 Concentrations predicted using IS92d	x
E.11 Concentrations predicted using IS92e	x
E.12 Concentrations predicted using IS92f	x
E.13 1995 impulse response relative to S650	x

Abstract

Various projections of the relation between future CO₂ concentrations and future emissions were undertaken as part of the scientific assessment for Working Group 1 of the Intergovernmental Panel on Climate Change. There were three type of calculation: (i) forward projections, calculating the atmospheric CO₂ concentrations resulting from specified emission scenarios, (ii) inverse calculations determining the emission rates that would be required to achieve stabilisation of CO₂ concentrations via specified pathways and (iii) impulse response function calculations required for determining Global Warming Potentials.

The use of a standardised set of conditions allows an intercomparison of models. Sensitivity studies explore other aspects of the uncertainties of such projections. This report documents the specifications, the models that were used and the results that were obtained. Some preliminary interpretations of the results are included.

Assessment of IPCC CO₂ Modelling Study**Issues, file b-issue.tex**

Draft, printed March 30, 1994

Last change 30/3/94

This page is temporary for the purposes of holding production information and will not form part of the final document.

1. Introduction

This report documents a set of calculations which relate future atmospheric CO₂ concentrations to industrial emissions. These calculations were performed at the request of Working Group 1 (WG1) of the Intergovernmental Panel on Climate Change (IPCC). Calculations were contributed by groups from a number of countries (a list is given as Table 5.1). The specifications for the calculations were prepared by Tom Wigley and Ian Enting who were charged with this duty at the IPCC WG1 Bureau and Ozone Assessment 'Experts' Meeting (Bath, UK, 18-19 February, 1993). These instructions are reproduced in this document as Appendix A.

In assessing these calculations it must be emphasised that they are not 'predictions' of a specific future. We prefer the term 'projections' — calculations of the possible consequences of prescribed courses of action. Similarly, these results do not constitute recommendations on the part of the contributors, the editors or the IPCC. They are intended to present ranges of options that may be relevant to dealing with the problems arising from greenhouse gas emissions.

One aspect of the calculations that prevents them being regarded as predictions is the neglect of feedbacks between CO₂ and climate. The uncertainties in such feedbacks are too large to justify defining any standard cases for comparative studies at this time. A small number of calculations with feedbacks were contributed. These are discussed separately in section 12.

The various cases are specified in terms of scenarios: future values of emissions or concentrations. In defining such cases it is important to ensure that the set of definitions is complete both in the mathematical sense of being fully determined and in the physical sense of including all relevant carbon fluxes. However, such completeness must avoid both 'double-counting' of any fluxes and mathematical 'over-specification' which would generally imply inconsistencies in the description. These issues are discussed at greater length in section 2b below.

The main calculations are those concerned with the stabilisation of atmospheric concentrations of CO₂. This emphasis is motivated by the provisions of the Framework Climate Convention. For the present modelling exercise, we have constructed a set of pathways, defining CO₂ concentrations as functions of time, that lead to constant CO₂ concentrations in the 21st or 22nd centuries.

- Validation
- Uncertainties

Previously, projections of future CO₂ had been included in the IPCC reports (IPCC 1990, 1992). The update (IPCC, 1992) included few specific projections of CO₂. However it did introduce new emission scenarios, IS92a-f, and included assessments of the climatic consequences using the STUGE system (Wigley et al., 1991). This integrated system included a carbon cycle model that was an earlier version of the model W whose results are reported here.

Our objective has been to present options based on a credible scientific assessment. Scientific credibility is achieved as a result of critical examination. While the results in this report have been scrutinised during the process of compilation and the overall report has been subject to peer review (not yet applicable for release 1), the ultimate assessment of its credibility must come from making the results freely available.

2. Carbon Budget Issues

2a Budget issues

We write the atmospheric carbon budget (in units of Gt C y^{-1}) as:

$$2.123 \frac{dC}{dt} = Q_{\text{foss}}(t) + D_n(t) - S_{\text{ocean}}(t) - S_{\text{fert}}(t) + S_{\text{resid}}(t) \quad (2.1)$$

where:

$C(t)$ is the atmospheric CO_2 concentration (in ppmv),

$Q_{\text{foss}}(t)$ is the industrial emission rate,

$D_n(t)$ is the net carbon flux from land-use change (including delayed effects from past land-use changes),

$S_{\text{ocean}}(t)$ is the net carbon uptake by the oceans,

$S_{\text{fert}}(t)$ is the net flux associated with CO_2 -enhanced growth,

$S_{\text{resid}}(t)$ is a residual-sink term, associated with short-term climatic variability and any neglected processes.

There are a number of important points to note about the use of this equation:

1. The budget equation (2.1) describes the perturbation about an assumed pre-industrial steady state. All the terms (except possibly $S_{\text{resid}}(t)$) refer either to anthropogenic forcing ($Q_{\text{foss}}(t)$ and $D_n(t)$) or to the natural response to such forcing $\frac{d}{dt}C$, $S_{\text{ocean}}(t)$, $S_{\text{fert}}(t)$. These fluxes are net fluxes superimposed on large natural gross fluxes of carbon. In many cases the models will consider the gross fluxes, particularly when calculating isotopic effects.
2. In practice we need to work with time averages of equation (2.1). At the very least we need to average over the seasonal cycle since these intra-annual changes are not included in this perturbation budget. In practice we may need to average over periods longer than a year in order to define a globally representative rate of change of CO_2 . Multi-year averaging can also remove some of the short-term climatic variability that we have assigned to the term $S_{\text{resid}}(t)$.
3. The standard cases defined in the instructions assume that the residual 'missing-sink' term, $S_{\text{resid}}(t)$, is zero.
 - c data
 - floating start

2b Issues of definition

As noted in the introduction, it is essential to ensure that the specifications of the carbon budget are both complete and consistent. Completeness is required in both in the mathematical sense of being fully determined and in the physical sense of including all relevant carbon fluxes. However such completeness must avoid both 'double-counting' of any fluxes and mathematical 'over-specification' which would generally imply inconsistencies in the description.

The process of mapping fluxes onto compartments with neither double-counting or under-counting is complicated by the different levels of description that are adopted in different

fields of study. In some cases, it is necessary to adapt the preferred model structure in order to match the definitions appropriate data.

Industrial This is intended to include all anthropogenic contributions to the net atmospheric carbon budget, except for those classified as associated with land-use change. In practice, the contributions are fossil fuel usage and cement production.

Fluxes from land-use change This term is taken as being the net flux associated with present and past land use changes. This definition is chosen to match that used by the Woods Hole group (Houghton et al., 1983 and subsequent studies). It includes fluxes associated with deforestation (including decay of forest products), changes in soil carbon, changes associated with changing agricultural regimes and forest regeneration on previously cleared land.

Oceans This term is taken as the net transfer from the atmosphere to the oceans. There is believed to be a natural cycle of carbon from atmosphere to vegetation transferred to the oceans via rivers and then returned to the atmosphere through outgassing (Sarmiento and Sundquist, 1992). This cycle can be ignored in the present global modelling. It becomes important when interpreting ocean CO₂ partial pressure data and (to a lesser extent) when interpreting inversions of spatial distributions of CO₂.

'Fertilisation' This needs to include all changes in terrestrial carbon that are not associated with land-use change. . . .

3. Modelling Approaches

3a. General Issues

The ocean models used in the calculations presented here span a range of forms from response function descriptions to general circulation models. Enting (1987) has described this range in terms of the 'modelling spectrum' concept of Karplus (1977) with models ranging from 'black-box' (purely inductive) to 'white box' (deduced from basic principles). The general issue for all levels of modelling is whether the model parameters can reasonably be regarded as being the same in the future as at present.

One way of grouping modelling approaches is

Extrapolation: This is based on extrapolating the trends in CO₂. This is generally done in statistical terms (e.g. Mannermaa and Karras, 1989). This approach has no capability of relating concentrations to emissions. It is making implicit assumptions about the continuation of current patterns of emissions.

Total response function description: The CO₂ increase is related to emissions through a response function formalism. This approach was used for the present exercise by Cohen (Model C). He derived his response function empirically by statistical fits to historical data. It would, however, be possible to derive this type of total response function from mechanistic models.

Combination of response functions and parameterised models: This description includes numerous possibilities. This approach was used for the present exercise

by Wigley (Model W) and Harvey (Model T). They used response function representations of the ocean and parameterised models of the terrestrial systems. The ocean response functions were derived from mechanistic ocean models.

Parameterised models: These are models with 'lumped' descriptions of actual processes. The way in which they differ from parameterisations in terms of response functions is that the same parameterisation is assumed to apply in more than one situation, making validation and/or calibration possible. For carbon cycle studies, the general assumption is that such parameterised models are adequate for defining relations between the behaviours of the different carbon isotopes, thus allowing the use of isotopic data in model calibration. All of the terrestrial components used in the present studies have been parameterised to some degree. The majority of the ocean models were highly parameterised.

Mechanistic models: These derive the behaviour deductively from fundamental physical and chemical principles. The various ocean general circulation models (Models H, O, Q) fall into this class, although even here some empirical parameterisation is involved.

Other important general issues in the modelling are:

The role of statistics x

- statistics
- Forward vs inverse
- inverse can be done by iterated forward or off-line with response fn.
- DEs
- Note Laplace transform

Some aspects of inverse modelling are noted in note A.6.D below. More detailed discussions are given by Enting and Mansbridge (1987) and Wigley (1991, 1993).

3b. Ocean models

- Response functions
- Parameterised
- OGCMs

The box-diffusion model provides an example of the range of uncertainty in calibrating parameterised models. Table 3.1 lists the range of estimates of the diffusion parameter. The ^3H calibration by Broecker et al. (1980) was used in some of the CO_2 projections reported in the IPCC (1990) assessment (Enting 1991).

Calibration	K	Reference
^{14}C (natural)	3987	Oeschger et al. (1975)
^3H (bomb)	5364	Broecker et al. (1980)
^{14}C (bomb)		Siegenthaler (1983)
Various ^{14}C	6245	Enting and Lassey (1993)
		Model A
	7865	Model B
Various ^{14}C	5859	Model E ₁
Various ^{14}C	x	Model E ₂
		Model P

Table 3.1. Calibrations of the box-diffusion model of ocean carbon uptake. K is the eddy diffusion coefficient in m^2y^{-1} .

Hilda table ?

A more complicated model is the HILDA model. This incorporates a 2-region description of the ocean, an advective circulation and a depth-dependent eddy-diffusion coefficient. Although these features had been incorporated into previous models (both separately and in combination) the HILDA model has been widely studied (Siegenthaler and Joos, 1992) and it is used as the ocean component of several of the calculations reported here.

- MH to write bits on OGCMs
- MH to write bits on penetration depth

Model	Total 1950	Mixed 1950	Total 1973	Mixed 1973	Total $\Delta 50 - 73$	Mixed $\Delta 50 - 73$	z	$\Phi_O z_{obs}/z$
Obs.						157		
E								
H					9.27	189	320	
O					7.18	159		

Table 3.3. Ocean ^{14}C levels. The mixed layer values are in ‰. Depth-integrated totals are in 10^9 atoms cm^{-2} . The penetration depth z is in metres. The observations are from the analysis by Broecker et al. (1985).

3c. Terrestrial models

- regional vs age class disaggregation
- beta on NPP
- deforest, Skee as net
- Layered description

3d. Issues for future work

Since the time that the instructions in Appendix A were drawn up, work on the carbon cycle has continued and some of the issues have become clearer. A number of particular activities have contributed to this. Firstly, there is, of course, the experience of having a number of groups perform the set of calculations. This exercise has revealed a number of issues that were less obvious before we commenced. Secondly, there have been a number of scientific meetings addressing the issues of concern in making projections of CO₂. (The Global Change Institute on the Global Carbon Cycle, Snowmass Village, July 1993; The Fourth International CO₂ Conference, Carqueiranne, September 1993).

In general, these discussions have left us with clearer statements of the problems rather than new solutions. Among the key issues are:

- Can we do better than 'neutral' biosphere? (i.e. unchanging biomass). The assumption of no future biotic change was implicit in many of the calculations produce for the IPCC (1990) report.
- Is a declining 'land-use' source (eg IS92a implying managed biosphere) an appropriate assumption? In the present studies, it has been adopted as an appropriate case to use in stabilisation scenarios, on the basis that any commitment to reduce CO₂ emissions is likely to be implemented by reductions in both industrial and biotic releases.
- Since the 'land-use' component is relatively small, does the precise form matter particularly?
- Are there any global integrals or constraints for terrestrial ecosystems that apply in a way analogous to the ¹⁴C constraint on oceanic CO₂ uptake? For example, are there physiological constraints that can be translated into maximum carbon loadings for particular ecosystems?
- Is terrestrial modelling preferable to scenarios for D_n ?
- Is a more integrated description (sometimes called 'layered modelling') appropriate, e.g. with both industrial and land-use fluxes modelled in terms of specific societal changes?

4. Strategies for Achieving Carbon Balance

The atmospheric carbon balance, with the assumption that CO₂-fertilisation accounts for all the 'missing sink', is expressed as:

$$2.123 \frac{dC}{dt} = Q_{\text{ross}}(t) + D_n(t) - S_{\text{ocean}}(t) - S_{\text{fert}}(t) \quad 4.1$$

This provides an essential constraint on the modelling. In general terms, the specifications in Appendix A invoke two principles:

1. The current (i.e. 1980's mean) carbon budget should balance, i.e. equation (4.1) should be satisfied.
2. There should be continuity in processes from the past to the future: the future balance should be achieved using the same processes as for the past.

Two of the ways in which these requirements translate into procedures for initialising the models for calculations beyond 1990 are:

The standard forward initialisation: Use $Q_{\text{foss}}(t)$, $D_n(t)$ exactly as specified, with $S_{\text{ocean}}(t)$ calculated (possibly using a ^{14}C calibration) and $S_{\text{fert}}(t)$ calculated. The parameterisation of S_{fert} is tuned to give the best possible agreement with the observed record of $C(t)$. The precise details of how this is done are left open.

The standard inverse initialisation: Use specified $Q_{\text{foss}}(t)$ and calculated $S_{\text{fert}}(t)$ and $S_{\text{ocean}}(t)$ and track the prescribed $C(t)$ in order to deduce $D_n(t)$. The model parameters are to be tuned to give $\bar{D}_n = 1.6 \text{ GtC/yr}$ averaged over the 1980s.

Post-1990: The fluxes S_{ocean} and S_{fert} are to be calculated using the same parameters as in the initialisation phase. The land-use flux D_n is prescribed. This leaves the model integration defining a relation between $Q_{\text{foss}}(t)$ and $C(t)$, either of which can be used to determine the other. Calculations of both types are requested in the instructions.

- At least some cases in IPCC 1990 used 'neutral-biosphere' post-1990.
- mix and match Fig 4.1

The prescribed CO_2 concentration for the period 1 Jan 1980 to 31 December 1989 shows an increase of 15.9 ppmv, an average rate of increase of 3.38 GtC/yr. For the same period our standard cases had average sources of $\bar{Q}_{\text{foss}} = 5.45 \text{ GtC/yr}$ and $\bar{D}_n = 1.58 \text{ GtC/yr}$. In terms of the atmospheric budget shown in equation (4.1), this constrains the sum of oceanic and fertilisation fluxes to be $S_{\text{ocean}} + \bar{S}_{\text{fert}} = 3.65$ or more generally $S_{\text{ocean}} + \bar{S}_{\text{fert}} = 2.07 + \bar{D}_n$. This is shown diagrammatically in Figure 4.1. It is possible to take combinations of the oceanic and terrestrial fluxes, whether computed by full models or partial models and combine them in pairs subject to the balance constraint.

5. Summary of Models

5.1 Descriptions

Code	Authors	Institution
A	Taylor, Lloyd and Farquhar	Aust. National University
B	Emanuel	ORNL
C	Cohen	UN Economic Comm. Europe
E	Enting and Lassey	CSIRO (DAR) and NIWA
F	Freidlingstein	
G	Goldstein	EPRI
H	Heimann	MPI, Hamburg
J	Joos and Siegenthaler	Bern
L	Wuebbles and Jain	Livermore
M	Moore	U. New Hampshire
O	Orr and Monfray	Saclay
P	Peng	ORNL
Q	Le Quere	Princeton
R	Alcamo	RIVM
T	Harvey	U. Toronto
V	Viecelli	Livermore
W	Wigley	OIES, UCAR

Table 5.1 Index of models.

The invitation to participate in the modelling exercise for the IPCC drew responses from many groups using a range of different models. For convenience of identification, each model has been given an identifying letter (generally with a mnemonic association with author or institution). These are listed in Table 5.1

Table 5.2 summarises the main characteristics of the various models. They are grouped according to the representation of the oceans, comprising one statistical representation of all sink processes (model C), two with response function representations (T and W); four versions of the box-diffusion model introduced by Oeschger et al. (1975) (A, B, E, P); four versions of the HILDA model (G, J, K, L); one implementation of the 12-box ocean model of Bolin et al. (1983) (model M) and three ocean general circulation models (H, O and Q). These types of model are described in section 3b, above.

The representations of the terrestrial components are more diverse.

Model	Ocean	Land	Init	Fertilisation
F	None			
C	(with land)	(with ocean)		
T	Response function	6-box	Inverse	Logarithmic
W	Response function	4-box	Inverse	Hyperbolic
A ₁	BDM	ANU-BACE	Inverse	
A ₂	BDM	ANU-BACE	Forward	
E ₁ , E ₂	BDM	2-box	Forward	Hyperbolic
P	BDM			
Y	BDM			
G	Hilda	6×6	Forward	
J	Hilda	4-box	Inverse	Logarithmic
L	Hilda	6-box	Inverse	Logarithmic
K	Hilda	?		
M	12-box			
R	?	Image 2.0		
V				
H	HAMOCC-3	OBM3m	Inverse	≈ Hyper.
O	LODYC GCM	None	Inverse	None
Q	GCM	None	Inverse	None

Table 5.2. Characteristics of models.

The models differ in the extent to which they have complied with the specifications in Appendix A. We can group them as:

Full models with standard inverse initialisation: These followed essentially the procedure of Section 4. The models were J, L, T and W. Models A₁ and H used a modified inverse approach, performing forward runs to tune the fertilisation to give a budget that balanced over the period 1960–1990, and then running the model in inverse mode over both past and future to deduce industrial emissions.

Full models with standard forward initialisation: These followed essentially the forward procedure of section 4. The models were A₂, E.

Ocean-only models: As noted above, these can provide useful information for those cases prescribed as inverse calculations. The models were O, Q.

Terrestrial-only models: Similarly these can be useful for the inverse cases. The only case here is model F.

Models with climatic feedbacks: The standard specifications requested that climatic feedbacks NOT be included. We discuss runs with climate feedbacks in section 12. For the moment we note that these cases will not be directly comparable with the no-feedback cases. Runs with climate feedbacks are either excluded from comparisons or are flagged specially. The models were R, T.

Models failing to fit the prescribed stabilisation curves adequately: These were G, V.

Others: The statistical representation (model C) failed to comply with the basic process description. Perhaps a more important failing is that it worked in terms of responses for total CO₂ rather than the perturbation.

In contrast model B is not directly comparable with the other models because its description is more detailed than envisaged in the specified scenarios.

To summarise, the models that can be directly compared for the stabilisation calculations are A₁, E₁, E₂, H, J, L, P and W. Models R and T differ by including climatic feedbacks; they are not directly comparable but are included in various tabulations of results for completeness. For forward calculations this set of 10 models can be augmented by models G and V; these failed to give adequate fits to the prescribed CO₂ concentrations in the inverse calculations.

6. Initialisation

6.1 Budgets

A key specification in the modelling exercise was the requirement for a balanced atmospheric carbon budget consistent with observations. Table 6.1 summarises the way in which various models achieved this balance over the period 1980-89.

	Initialised	Fossil	Ocean	Fertil	Land-use	Biota	$\frac{dC}{dt}$
A ₁	Inverse*	5.65	2.03	1.85	1.56	0.29	1.59
H	Inverse*	5.63	1.47			0.76	1.59
J	Inverse	5.45	2.16			-0.08	1.59
L	Inverse	5.45	2.14		1.60	0.12	
M	Inverse	5.58	1.72			0.48	1.59
T	Inverse	5.41	1.79	1.95	1.68		1.59
W	Inverse	5.45	2.00	1.67	1.60	0.07	1.59
A ₂	Forward	5.49	2.00	1.87	1.29	0.58	1.39
E	Forward	5.48	2.10			0.27	1.50
O	Ocean-only		2.10				
Q	Ocean-only		1.81				
F	Biota-only						

Table 6.1. Mean budgets (or budget components) for 1980-89. The '≡' signs denote values specified as inputs to particular calculations. Fluxes are in GtC/yr and are sinks except for the land-use term. The 'biota' term is fertilisation minus land-use. The mean atmospheric increase is in ppmv/yr. The * denotes a modified inverse initialisation.

6.2 Validation

- ¹⁴C as main data set
- inventory, penetration
- normal is atmospheric forcing, Enting uses bomb forcing to allow decay
- response function indirect

- surface response functions
- some cases are from published descriptions

Table 6.3 shows the cases for which the models have been compared to (or calibrated with) ^{14}C data.

	t	A	D	E	G	H	J	K	L	M	O	P	Q	R	T	V	X	Y
Atmosphere	all	-	-	Y	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mixed layer	all	-	-	Y	-	-	Y	-	Y	-	Y	-	-	-	-	-	-	-
Inventory	all	-	-	-	-	-	Y	-	-	-	Y	-	-	-	-	-	-	-
Mixed layer	1974	-	-	Y	-	-	Y	-	Y	-	Y	-	-	-	-	-	-	-
Inventory	1974	-	-	Y	-	-	Y	-	-	-	Y	-	-	-	-	-	-	-
Penetration z	1974	-	-	Y	-	-	Y	-	-	-	Y	-	-	-	-	-	-	-

Table 6.3. ^{14}C calibration data. 'Y' denotes ^{14}C data contributed. 'P' denotes fit to ^{14}C data described in published description of model.

Figure captions for sections 1 to 6

Figure 4.1 Relation between components of the atmospheric carbon budget averaged over 1980-89. The lines for various mean 'land-use' fluxes specify the combinations of ocean flux and fertilisation flux that are compatible with a balanced atmospheric budget. Labelled points along the axes show fluxes calculated for some of the models. Within the specifications of appendix A, the 'fertilisation' and 'ocean' components of different models can be combined, subject to implying an acceptable 'land-use term'.

Figure 4.1

Assessment of IPCC CO₂ Modelling Study**Results, file b-res.tex**

Draft, printed March 31, 1994

Last change 30/3/94

This page is temporary for the purposes of holding production information and will not form part of the final document.

7. Summary of Results for Forward Calculations

- Table of key results
- List main differences

Model	IS92c	IS92a	IS92e	DEC0%	DEC1%	DEC2%
A ₂		518.6		464.2	442.0	426.2
B ₁		520.9				
B ₂		522.8				
B ₃		512.6				
E	437.7	494.1	541.6	438.9	416.5	400.8
G		509.8		458.3	437.1	422.0
J	442.1	498.7	546.0	443.4	421.0	405.4
L	441.0	497.9	545.5	443.7	421.4	405.8
W	437.4	497.4	547.4	438.6	414.3	397.1
R	432.1	495.1	548.1			
T*	438.1	494.9				

Table 7.1. Projections of CO₂ for 2050 for forward calculations. The two 'feedback' cases, R and T, are included for completeness in the lower section.

Model	IS92c	IS92a	IS92e	DEC0%	DEC1%	DEC2%
A ₂		731.3		537.4	467.5	432.1
B ₁		734.0				
B ₂		731.2				
B ₃		728.4				
E	468.7	683.9	913.1	492.7	425.4	392.9
G		719.8		534.0	464.1	428.2
J	474.2	687.7	910.4	498.2	430.3	397.2
L	473.0	687.0	908.3	502.6	434.2	402.7
W	456.4	676.5	909.2	481.4	411.4	377.3
R	447.6	667.8	899.7			
T*	470.5	672.8				

Table 7.2. Projections of CO₂ to 2100. Again, the feedback cases, R and T, are kept separate.

Tables 7.1 and 7.2 list the projections to 2050 and 2100 for 6 of the scenarios: IS92a (the de facto 'business-as-usual'), IS92c, IS92e (the two extremes) and the three 'science scenarios, DEC0%, DEC1% and DEC2%. The results fall into two main groups, models A₂, B₁, B₂ and G, and the others (E, J, L, T) which give lower projections differing from each other by only a small amount.

8. Summary of Results for Stabilisation calculations

Model	S350	S450	S550	S650	S750	DS450	DS550
A	-1.41 2105	6.75 2001	7.38 2033	9.55 2076	11.68 2075	7.26 2011	8.05 2020
E ₁	-1.26 2111	6.92 2007	8.42 2075	10.71 2075	12.94 2075	7.57 2013	8.69 2039
H		6.57 2000		9.04 2075			
J	-1.12 2109	6.66 2005	8.27 2075	10.59 2075	12.85 2075	7.31 2011	8.43 2039
L	-1.48 2108	6.82 2005	8.02 2075	10.35 2075	12.64 2075	7.45 2012	8.43 2033
M		6.26 2000		9.43 2076			
P		6.10 1990		7.83 2075			
W	0.03 2112	6.78 2012	9.16 2075	11.44 2079	13.68 2077	7.41 2013	9.26 2073
R	-0.26 2100	7.15 2040	8.99 2056	10.77 2058	12.69 2059	7.38 2037	9.37 2043
T	-1.43 2106	6.82 2005	8.13 2075	10.62 2079	13.13 2081	7.45 2012	8.40 2029
O		4.77 2007		8.51 2079			
Q	-0.95 2097	4.37 2002		7.46 2076			
B ₁	-0.12 2076	5.65 1998	7.99 2104	9.98 2104	11.70 2104	6.03 2008	7.62 2104
B ₂	0.52 2074	5.47 2001	8.34 2104	10.33 2104	11.99 2104	5.89 2011	8.01 2104
B ₃	0.38 2098	6.37 2002	7.74 2075	9.66 2077	11.53 2075	6.90 2011	7.76 2066
C		7.10 2000		16.15 2075			
G	0.00 2020	7.10 2000		13.70 2050			
V _A		6.90 2002		16.93 2047			
V _B		7.48 2010					

Table 8.1. Stabilisation cases. Maximum emissions (minimum for S350) and year. The horizontal lines separate groups that are not directly comparable with the top group: feed-back cases (R,T), ocean-only (O,Q), and those that fail to fit the specifications (see section 5).

- List main differences

This section describes the results of 'inverse' calculations which determined the emissions required to achieve stabilisation of atmospheric CO₂ concentrations via specified concentration scenarios. Seven sets of stabilisation scenarios were prescribed, with stabilisation at 350, 450, 550, 650 and 750 ppmv. For the 450 and 550 stabilisation cases two concentration pathways were defined: a standard case and a delayed case in which the concentrations initially followed a higher pathway and were subsequently modified to produce the lower stabilisation level. Figure 8.1 shows the seven prescribed concentration functions. The 'delayed' cases are shown as dashed.

Figures 8.2 and 8.3 show the calculated industrial emissions for various models for the S450 and S650 scenarios. The results from the two models which included climatic feedbacks are shown with short dashes. The results for the other scenarios are plotted in Appendix E. Tables 8.1 and 8.2 list some key characteristics of the results for all seven scenarios and for all models. These tables include (in their lower sections) results that are not strictly comparable in terms of the specification in Appendix A. Two models are 'ocean-only', two include climatic feedbacks, two produced poor fits to the specified concentration histories and one did not 'balance the current carbon budget', i.e. an unidentified residual sink was required for balance.

For the S650 case, fluxes between the atmosphere and oceans are given in Table 8.3 and fertilisation fluxes are given in Table 8.4.

Model	S350	S450	S550	S650	S750	DS450	DS550
A	0.40	1.36	2.14	3.03	4.71	1.35	2.10
E ₁	0.25	1.53	2.57	3.68	5.46	1.51	2.51
H		1.89		3.16			
J	0.34	1.64	2.73	3.88	5.81	1.63	2.67
L	0.31	1.39	2.27	3.33	5.20	1.37	2.22
M		1.55		2.88			
P		1.32		2.65			
W	1.02	2.73	4.16	5.58	7.57	2.71	4.10
T	0.21	1.53	2.85	4.51	7.01	1.51	2.80
O		1.95					
Q	0.37	1.16	1.77	2.39	3.81	1.15	1.75
B ₁	2.50	3.54	4.40	5.18	6.83	3.53	4.37
B ₂	3.07	4.08	4.87	5.62	7.22	4.07	4.84
B ₃	1.80	2.81	3.59	4.35	5.94	2.80	3.57
C		2.03		2.94			
G	0.00	1.49		1.50			
V _A		2.82		0.37			
V _B		5.55					

Table 8.2. Emissions in 2200 for stabilisation cases.

Model	1990	2000	2050	2100	2200	2300
A	2.50	2.97	4.88	5.92	4.45	3.20
B	2.74	3.15	4.26	4.39	3.07	2.30
E ₁	2.30	2.70	3.70	3.80	2.60	2.00
H	1.63	1.87	2.55	2.72	2.18	1.85
J	2.41	2.75	3.68	3.76	2.52	1.84
L	2.38	2.65	3.27	3.22	1.90	1.35
M	1.90	2.18	3.16	3.46	2.70	2.12
O	2.36	2.76	4.18	4.71		
P	2.43	2.78	3.73	3.82	2.63	1.96
Q	2.01	2.35	3.34	3.49	2.36	1.65
W	2.24	2.59	3.98	4.52	3.55	2.57
R	1.63	1.69	2.57	2.85		
T	1.98	2.30	3.20	3.95	3.54	2.80
V	4.22	5.35	10.26	7.08	2.80	

Table 8.3. Air-sea fluxes for S650.

Model	1990	2000	2050	2100	2200	2300
A	-2.04	-2.25	-2.19	-1.66	-0.47	-0.15
F						
T	-2.21	-2.63	-3.31	-3.01	-1.20	-0.50
W	-1.85	-2.12	-2.98	-3.07	-2.03	-1.17

Table 8.4 Fertilisation fluxes for S650.

Figure 8.4 shows the effect (in emission terms) of delaying reductions in CO₂ emissions, or more specifically, choosing a higher target and then revising the target downwards. The higher emissions that are allowed in the 'pre-revision' period are more than offset by the requirements for greater reductions later and indeed for an indefinite period into the future.

Figure 8.4 includes results from one 'ocean-only' model. In this case the results are comparable because the unknown terrestrial fluxes will cancel when taking the emission difference.

9. Summary of Results for Response Functions

- Table of key results
- List main differences

Impulse response functions have been calculated as perturbations from specific cases by adding 10 GtC to the atmosphere, integrating forward and calculating the concentration differences from the case without the 10GtC input.

The equilibrium response adds 10 GtC to a steady state situation. Equilibrium response functions have been calculated using models E, G, H, J, L and W.

The perturbation response functions are calculated as perturbation from the S650 case by the addition of 10 GtC to the atmosphere during 1995 and then integrating forward with the emissions deduced in the S650 case.

The response function, $G_{\text{Pert}}(t)$, can then be evaluated as

$$G(t) = 2.123[C(t) - C_{\text{S650}}(t)]/10.0 \quad (9.1)$$

$G(t)$ is taken to be 1.0 at $t = 0$, defined as mid-1995. Perturbation calculations have been undertaken with models E, H, J, L, R and W.

The calculation of global warming potentials requires the evaluation of the time-integrated CO_2 radiative forcing. With the the radiative forcing, F , of CO_2 (in Wm^{-2}) approximated by

$$F = 6.3 \ln \frac{C(t)}{C(0)}, \quad (9.2)$$

the integrated extra forcing is

$$\int \Delta F = \int_0^T \frac{6.3}{C(\tau)} G(\tau) d\tau \quad (9.3)$$

Again $\tau = 0$ is defined as mid-1995. Integrated forcings have been calculated for the five models listed above.

Model	0	10	25	50	100	200	300	400	500
E	1.0	0.546	0.386	0.289	0.234	0.204	0.187	0.174	0.163
G	1.0	0.639	0.552	0.497	0.435	0.369	0.329	0.299	0.276
H	1.0	0.430	0.283	0.217	0.173	0.141			
J	1.0	0.560	0.410	0.316	0.248	0.204	0.185	0.172	0.163
L	1.0	0.643	0.459	0.354	0.302	0.254	0.228	0.212	0.202
W	1.0	0.629	0.442	0.309	0.225	0.176	0.155	0.143	0.135

Table 9.1 Equilibrium response function, calculated relative to constant pre-industrial concentrations.

Model	0	10	25	50	100	200	300	400	500
E	1.0	0.624	0.493	0.419	0.401	0.425	0.408	0.389	
H	1.0	0.639	0.477	0.368	0.281	0.215			
J	1.0	0.628	0.507	0.435	0.401	0.393	0.376	0.359	0.348
L	1.0	0.695	0.555	0.495	0.477	0.459	0.424	0.399	0.386
R	1.0	0.786	0.616	0.531	0.488				
W	1.0	0.709	0.555	0.430	0.368	0.347	0.320	0.280	0.247

Table 9.2. Perturbation response function, calculated relative to S650.

The most widely quoted ocean response function is

$$G_O(t) = 0.131 + 0.201e^{-t/362.9} + 0.321e^{-t/73.6} + 0.249e^{-t/17.3} + 0.098e^{-t/1.9} \quad (9.x)$$

from the ocean GCM of Maier-Reimer and Hasselmann (1987). Enting and Mansbridge (1987) parameterised the response of the box-diffusion model as

$$G_O(t) = \quad (9.x)$$

Combining response functions

Assume that we have a linear system consisting of atmosphere, ocean and biosphere. We define the atmospheric impulse response function $G_a(\tau)$ as a Greens function

$$n_a(t) = \int_0^\infty G_a(\tau) Q(t - \tau) d\tau \quad (9.x)$$

where $n_a(t)$ represents the excess atmospheric CO_2 content at time t and $Q(t)$ the atmospheric CO_2 source.

We may similarly define partial impulse response functions $G_{aoc}(\tau)$ and $G_{abio}(\tau)$ representing the impulse response function of the atmosphere-ocean, respectively atmosphere-biosphere system. How can $G_a(\tau)$ be reconstructed from known $G_{aoc}(\tau)$ and $G_{abio}(\tau)$?

We define a further Greens function $G_{oc}(\tau)$ relating the net ocean uptake $S_{oc}(t)$ at time t as a convolution integral of previous changes in atmospheric CO_2 concentrations

$$S_{oc}(t) = \int_0^\infty G_{oc}(\tau) \dot{n}_a(t - \tau) d\tau \quad (9.y)$$

$G_{aoc}(\tau)$ is the impulse response of the atmosphere-ocean system, hence the balance equation for the atmosphere is given by

$$\begin{aligned} \dot{h}_{aoc} &= \delta(t) - S_{oc}(t) \\ &= \delta(t) - \int_0^\infty G_{oc}(\tau) \dot{h}_{aoc}(t - \tau) d\tau \end{aligned} \quad (9.z)$$

where $\delta(t)$ indicates the pulse input at time $t = 0$. Integrating (9.z) in time gives

$$G_{aoc}(t) = 1 - \int_0^\infty G_{oc}(\tau) G_{aoc}(t - \tau) d\tau \quad (9.a)$$

An analogue equation holds for the partial biospheric impulse response

$$G_{abio}(t) = 1 - \int_0^\infty G_{bio}(\tau) G_{abio}(t - \tau) d\tau \quad (9.b)$$

The balance equation of the entire atmosphere-ocean-biosphere system in the case of a unit pulse input gives

$$G_a(t) = 1 - \int_0^\infty (G_{oc}(\tau) + G_{bio}(\tau)) G_a(t - \tau) d\tau \quad (9.c)$$

Equation (9.c) relates G_a with G_{oc} and G_{bio} , which themselves are related to the partial impulse response functions G_{aoc} and G_{abio} by equations (9.a) and (9.b).

The convolution integrals can be eliminated by taking Laplace transforms and solving the resulting algebraic equations for G_a . In practice the partial impulse response functions are given in numerical form. By numerical fitting these can be represented as a series of exponential functions (plus possibly a constant offset). The series can be easily transformed and algebraically solved for the Laplace transform of G_a . Since only exponentials are involved, rational functions result which are readily transformed back into a series of exponentials.

Using g or more specifically $g(p)$ to denote the the Laplace transform of a function $G(t)$ we obtain the transformed equations (9.a), (9.b) and (9.c)

$$g_{aoc} = 1/p - g_{oc} g_{aoc} \quad (9.g)$$

$$g_{abio} = 1/p - g_{bio} g_{abio} \quad (9.h)$$

and

$$g_a = 1/p - (g_{oc} + g_{bio})g_a \quad (9.k)$$

Solving equations (9.g), (9.h) and (9.k) for g_a as a function of g_{aoc} and g_{abio} we obtain

$$\begin{aligned} g_a &= \frac{1}{p\left(\frac{1}{pg_{aoc}} + \frac{1}{pg_{abio}} - 1\right)} \\ &= \frac{g_{aoc}g_{abio}}{g_{aoc} + g_{abio} - pg_{aoc}g_{abio}} \end{aligned} \quad (9.m)$$

10. Assessment of Differences and Similarities

The results tabulated in the preceeding sections and the appendices show a relatively wide range of projections. It is, however, possible to discern a few general characteristics.

- The forward projections are very tightly grouped, while the stabilisation cases show a much wider proportional spread. This comes from two well-understood mathematical characteristics of the problems. Firstly, inverse calculations (i.e. the stabilisation calculations) amplify uncertainties much more than forward calculations. Secondly, since all the models are tuned to fit the historical record which has been a period of increasing emissions, they will tend to agree with each other (and be expected to be most accurate) when projecting the consequences of a similar period of future growth in emissions. The differences between models will become apparent when they are used for calculations (whether forward or inverse) involving periods of significant decreases in emissions as required for stabilisation.

11. Assessment of uncertainties

11a. General issues

A very important part of this modelling exercise is the assessment of the uncertainties involved in the CO₂ projections. Some of the key contributions to uncertainties in the model results are:

- Model error
- Budget uncertainty
- Imprecise calibration

There are many aspects of uncertainty analysis and a range of options in addressing them:

1. We can investigate the model-dependence by comparing the results of a set of carefully standardised cases. Much of the present exercise is oriented to this approach.
2. The consequences of the underlying assumptions can also be explored using a set of standardised cases.
3. The sensitivity of the results to variations in model parameters can be explored by systematic exploration of the parameter space (either deterministically or stochastically).

4. A more relevant measure of uncertainty is provided by assessing the sensitivity to those sets of parameter variations that lead to results consistent with observations. This can be assessed using a statistical approach to model calibration. Rather than determine fixed sets of parameters by seeking exact fits to specific data items, models can be calibrated by determining sets of compatible parameter values that give model results within an acceptable range of observed values.

11b Previous studies

There have been a number of previous studies that have either compared a range of models or studies uncertainties within the framework of a single model.

Killough and Emanuel (1981)

Laurmann and Spreiter (1983)

Gardner and Trabalka (1985)

Enting and Pearman (1983, 1986, 1987)

There have been several studies that attempted to analyse the uncertainties in projections of CO₂ in terms of the sensitivities to model parameters: Kandlikar (1992), Filar (1993), Parkinson et al. (1993), Wigley (1993).

11c. Sensitivity to carbon budget

The instructions in Appendix A prescribe a specific set of calculations to address this question. The budget uncertainty is characterised in terms of uncertainty in the land-use component; the specified calculations require the use of 0.6 and 2.6 Gt C y⁻¹ for the 1980's in place of the 1.6 Gt C y⁻¹ specified as the standard case.

These calculations have been performed using models J and W.

11d Calibration sensitivity

Perhaps more important than the extent to which the results depend on the model parameters is the question of the sensitivity to those sets of parameter variations that are consistent with the observational data. To address this question, Gardner and Trabalka (1985) used a Monte Carlo approach, generating sets of randomly distributed parameters and looked at the variability of results for the subset of cases that were consistent with observations.

Enting and Pearman (1983, 1986, 1987) embedded the sensitivity analysis within the model calibration procedure. This approach has been applied to the case with the box-diffusion model used to calculate the IS92a scenario (Enting and Lassey, 1993). They investigated the question of how much variation was possible in the atmospheric concentration for the year 2100, subject to retaining a specified degree of agreement between the model results and the calibration data.

12. Additional Calculations

- Climate sensitivity
- Other initialisations

12a Climate sensitivity

12b. Other

13 Extensions

13a Terrestrial modelling

13b. WEC Energy Use Scenarios

14. Conclusions

Acknowledgements

The publication of this report has been supported by the Australian Department of Environment, Sport and Territories. Rachel Law and Cathy Trudinger gave invaluable assistance with the production.

Figure captions for figures for sections 7 to 14

Figure 7.1 Concentrations predicted using IS92a. The curves with short dashes (models R, T) were calculated with climatic feedbacks included.

Figure 7.2 Concentrations predicted with emissions fixed at 1995 levels

Figure 8.1 Prescribed CO₂ concentrations for stabilisation studies. The dashed curves are the 'delayed' cases DS450 and DS550.

Figure 8.2 Calculated emissions for S450 case. The curves with short dashes (models R, T) were calculated with climatic feedbacks included.

Figure 8.3 Calculated emissions for S650 case. The curves with short dashes (models R, T) were calculated with climatic feedbacks included.

Figure 8.4 Effect of delay in stabilisation at 550. Difference in emissions for DS550 and S550. The curves with short dashes (models R, T) were calculated with climatic feedbacks included.

Figure 9.1 Impulse response to pre-industrial flux

Figure 9.2 'Ocean-only' response functions

Figure 9.3 'Biota-only' response functions

Figure 10.1 Relations between future emissions and current budget

Figure 7.1

Figure 7.2

Figure 8.1

Figure 8.2

Figure 8.3

Figure 8.4 (incomplete)

Assessment of IPCC CO₂ Modelling Study**Appendix A: Instructions, file a-rules.tex**

Draft, printed March 21, 1994

Last change 13/12/93

This page is temporary for the purposes of holding production information and will not form part of the final document.

Appendix A: Instructions for Calculations

This appendix reproduces the instructions that were prepared for the calculations by T. Wigley and I. Enting. Draft versions of these instructions were circulated to potential participants (and other interested parties) by the co-ordinators. A final version was circulated from the IPCC Working Group 1 secretariat at Bracknell. For the purposes of this appendix, the various sections of the instructions have been re-numbered by a prefix 'A.', in order to avoid ambiguity. A number of minor typographical errors have been corrected. Other changes are flagged with footnotes.

Carbon Cycle Model Calculations for the IPCC

Working Group 1 of IPCC is seeking calculations relating future atmospheric CO₂ concentrations to emissions, particularly for cases involving possible stabilization of concentrations. At a joint meeting of the IPCC WG1 Bureau and the Chairs of the International Ozone Assessment Panel (Bath, U.K., 18-19 February 1993) Tom Wigley and Ian Enting were charged with the responsibility of convening and coordinating this activity. We are seeking participation from carbon cycle modelling groups world-wide.

The primary task of WG1 in this context is to provide illustrative emissions pathways that lead to concentration stabilization.

Contents

- A.1: Summary of calculations required.
- A.2: Deliverables.
- A.3: Background.
- A.4: Data sets prescribed.
- A.5: Reporting and model comparison process.
- A.6: Technical notes
 - A.6.A: Land-use emissions prior to 1850.
 - A.6.B: Net flux vs. gross flux from deforestation.
 - A.6.C: Avoiding emission discontinuities.
 - A.6.D: Iterative inverse calculations.
 - A.6.E: Details of IS92a-f.
 - A.6.F: Other aspects of the scenarios.

We can be contacted through:

T.M.L. Wigley
UCAR Office for Interdisciplinary Earth Studies
PO Box 3000 Boulder
Colorado 80307-3000
USA
Phone 1-303-497-1646
Facsimile 1-303-497-1679
wigley@ncar.ucar.edu (internet)
T.WIGLEY (OMNET)

I.G. Enting
Division of Atmospheric Research
CSIRO
Private Bag No. 1, Mordialloc
Victoria 3195
Australia
Phone 61-3-5867696
Facsimile 61-3-5867600
ige@dar.csiro.au (internet)

Acknowledgements The coordinators acknowledge valuable comments from Uli Siegenthaler, Martin Heimann and Mike Oppenheimer.

A.1. Summary of calculations required.

A.1a. General

The calculations required are integrations of global carbon cycle models from pre-industrial times (starting at 1765) to 2100 or preferably to 2200. There are two aspects to the exercise: (a) an assessment of the uncertainty due to uncertainties regarding the current carbon budget; (b) an assessment of the uncertainties arising from differences between models. In order to separate these effects, we propose a set of standard conditions to explore inter-model differences and then a series of sensitivity studies to explore the consequences of current uncertainties in the carbon cycle.

A.1b. Standard integrations

The calculations have been grouped in order of importance since the full set of 'interesting' cases is larger than many groups will want to undertake. The bold face 4 or 5 character abbreviations are used throughout our description to distinguish cases. The annotation 'forward' or 'inverse' refers to the period from 1990 onward. For the inverse calculations we provide specified concentration histories. For the forward calculations the sources are specified. The choice between inverse and forward treatment of the period 1765 to 1990 is left open. A specified concentration history for 1765 to 1990 is provided for use in inverse calculations and for fitting to (as well as possible) in forward calculations. Technical Note A.6.D notes some relevant aspects of inverse calculations.

Essential

- i: Stabilization at 650ppmv. Deduce industrial emissions. *inverse S650*
- ii: Stabilization at 450ppmv. Deduce industrial emissions. *inverse S450*

Highly desirable

- iii, iv: Impulse response function (as input to GWP calculations). Calculated relative to pre-industrial state and as perturbation to S650. *forward Iinit and Ipert*

v: IS92a. *forward* IS92a

vi: IS92a to 2000 then constant industrial emissions. *forward* DEC0%

vii: IS92a to 2000 then industrial emissions decreasing at 1% per year. *forward* DEC1%

viii: Stabilization at 350ppmv. Deduce industrial emissions. *inverse* S350

Desirable

ix: Stabilization at 550ppmv¹ with delayed implementation. Deduce industrial emissions. *inverse* DS550

x: Stabilization at 450ppmv with delayed implementation. Deduce industrial emissions. *inverse* DS450

xi – xii: IS92c, IS92f. (The extreme cases²). *forward* IS92c and IS92f

Possibly useful

xiii: Stabilization at 750ppmv. Deduce industrial emissions. *inverse* S750

xiv: Stabilization at 550ppmv. Deduce industrial emissions. *inverse* S550

xv: IS92a to 2000 then industrial emissions decreasing at 2% per year. *forward* DEC2%

xvi – xviii: IS92b, d, e. *forward* IS92b IS92d IS92e

A.1c. Specifications for standard cases

- From 1990 onwards, all standard cases use specified net flux from land-use change, calculate any fluxes from CO₂ fertilization and, depending on which case is considered, either deduce industrial emissions from specified concentrations (*inverse*) or calculate concentrations from specified industrial emissions (*forward*).
- The future net flux from land-use change is described in Note A.6.E. Forward integrations of the IS92a–f industrial sources should use the IPCC 1992 deforestation scenarios for 2000 to 2100 as given. All other cases should use the modified IS92a (i.e. IS92a* from Table 4 of Note A.6.E) which goes to zero at 2100 and remains zero. All cases should use linear interpolation over 1990 to 2000 to connect the land-use flux used in the initialization phase with which ever case is used after 2000.
- For the effect of land-use change pre-1990, the net flux from Houghton is provided as an option (Note A.6.A). If some alternative is used then the net flux from land-use change should be 1.6 Gt C y⁻¹ averaged over 1980–89 inclusive (except in the sensitivity studies). This is the IPCC 1990 'best guess' value and is close to the estimates by Houghton.

¹Early versions of the instructions referred to a non-existent DS650 case at this point.

²This is an error. The upper extreme is IS92e, not IS92f, and so IS92e is used in the comparisons in other sections of this report

- The atmospheric CO₂ concentration should agree as well as possible with the supplied fit to the observational record. If the period 1765 to 1990 is treated with inverse modelling, then the supplied concentration record should be used.
- The rate of increase of atmospheric CO₂ should be fixed at 1.7 ppmv y⁻¹ for mid-1990 (i.e. $t = 1990.5$), except in sensitivity studies. (See Note A.6.C). For inverse calculations from 1990 onwards (i.e. the stabilization cases), the 1990 concentration needs to be fixed at the starting value of the prescribed time series. We specify 354.17 ppmv for mid-1990.
- The calculation of the impulse response should be as a perturbation from pre-industrial equilibrium with concentration 278 ppmv (with perturbation from S650 as an optional extra). The input should be 10 Gt C over 1 year. For the perturbation case, the 10 Gt should be released during 1995. The integrations should be for 200 years or preferably 500.
- Recommended constants 1 ppmv = 2.123 Gt C in atmosphere. Ocean area = 3.62×10^{14} m².

A.1d. Output

Essential

- i: Fossil carbon emissions (in gigatonnes per year) and CO₂ concentrations (in ppmv) for each case considered, for the period 1765 to 2100.
- ii: Contents of terrestrial biosphere reservoirs (annual data in Gt C) and fluxes between reservoirs (in Gt C y⁻¹), especially atmosphere-to-ocean and atmosphere-to-terrestrial-biosphere.

Desirable

- iii: Mixed layer ¹⁴C, as $\Delta^{14}\text{C}$ in ‰.
- iv: Ocean ¹⁴C inventory for 1/1/74 and preferably over the period 1940–1990. Given as excess ¹⁴C atoms per square meter.
- v: Results as described above extended to 2200 or beyond.

Further details are given in section A.2.

A.1e. Sensitivity studies

Modellers will perform such sensitivity calculations as they judge necessary. Our suggestions of standard cases for intercomparisons are:

- i: Explore sensitivity to fixing the rate of concentration increase to the mid-1990 value of 1.7 ppmv y⁻¹.
- ii: Explore sensitivity to fixing net flux from land-use change at an average of 1.6 Gt C y⁻¹ over 1980–90. We suggest using 0.6 and 2.6 Gt C y⁻¹.

- iii: Compare calculations with two different forms of CO₂-enhanced production:
 - (a): Growth logarithmic $\propto [1 + \beta \ln(C/C_0)]$, (b): Saturation $\propto G_\infty \frac{C-C_c}{C+b}$ with $C_c = 80$ ppmv and the constraint $G_\infty = (C_0 + b)/(C_0 - C_c)$ and b used for tuning.
 - (see Allen et al. *Global Biogeochemical Cycles*, 1, p1).

A.1f. Data provided

- i: $C(t)$ 1765 to 1990. For input to inverse calculations and fitting in forward cases.
- ii: $C(t)$ for 1990 to 2200 for the stabilization cases. (7 cases)
- iii: Fossil carbon emissions for 1844–1990.
- iv: Atmospheric ¹⁴C for 1800 to 1990 for cases with ¹⁴C forced to track atmospheric levels.
- v: Net release from land-use change, $D_n(t)$ from Houghton.

Time series (i) and (ii) are provided in two versions, one for beginning of year values and one for mid-year values. We also provide alternative versions of these time series with mid-1990 gradients of 1.6 and 1.8 ppmv for use in the sensitivity studies.

A.2. Deliverables

The most important outputs are the emissions (for the inverse calculations) and the concentrations (for the forward calculations). The simplest way to provide these data is as annual $E(t)$ and $C(t)$ time series over the full period of the analysis (1765 to the chosen end date). Emissions data must be broken down into industrial and net land-use emissions.

To aid in the interpretation of the results (inter-model differences), other output items are desirable. In particular, 1765-to-end-year annual time series of total terrestrial biomass and flux into the ocean will allow the full budget, and its changes, to be assessed for each model. A breakdown of terrestrial biomass changes into components (above-ground living biomass, litter, soil carbon, etc.) is also desirable, as are gross fluxes, if possible.

We recommend the following file format for ASCII files of time series:

Line 1: Integers N and M

Lines 2 to $N + 1$: Arbitrary identification information.

Lines $N + 2$ to $M + N + 1$ Output records.

The items in the record should be separated by blanks and in floating point format, i.e. suitable for reading by Fortran or Pascal free format reads to real variables. We suggest that the first few items should be

Time As year, with at least one decimal place so that mid-1990 is 1990.5

CO₂ concentration In ppmv, suggest 2 decimal places.

Fossil carbon emissions in Gt C y^{-1} . Suggest at least 2 decimal places.

The ^{14}C data should be reported as time series with the same file structure as above and the records as

Time As above.

Atmospheric ^{14}C As $\Delta^{14}\text{C}$ in ‰.

Mixed layer ^{14}C As $\Delta^{14}\text{C}$ in ‰.

Ocean excess ^{14}C inventory As excess (above pre-industrial) ^{14}C atoms per square meter.

The suggested period for ^{14}C output is 1940 to 1990, expanded to 1765 to 1990 if desired.

Finally, full details should be given of model structure and parameter values used in each run, together with information on how differences between net and gross land-use emissions were accounted for, how the model was initialized, and any other particulars on model structure or modelling strategy that you consider relevant.

A.3. Background

A.3a. The choice of integrations

Working Group 1 of IPCC is seeking calculations of future CO_2 levels, particularly those related to possible stabilization of concentrations. Article 2 of the Climate Convention states that the "ultimate aim" of the Convention is

"...to achieve stabilization of greenhouse gas concentrations ... at a level that would prevent dangerous anthropogenic interference with the climate system ... within a time frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened and to enable economic development to proceed in a sustainable manner"

The primary task of WG1 in this context is to provide illustrative emissions pathways that lead to concentration stabilization.

At the eighth session of the IPCC (Harare, Zimbabwe, 11-13 November 1992) it was agreed that the deliverables under the 1993-95 workplan of WG1 do not include calculations of the climate implications of the 1992 IPCC emissions scenarios developed by Leggett et al. (1992). Nor will these scenarios be revised¹. Thus, IPCC WG1 has no remit to pursue the CO_2 concentration implications of the IPCC92 scenarios. Nevertheless, since such calculations are likely to assist in interpreting the results of stabilization experiments, they are included in the optional part of the present exercise.

In addition, we believe that many modellers will wish to go beyond the basic calculations implied by the requirements of WG1 in other ways. We have listed such additional calculations to try to ensure the maximum degree of comparability between those models for which additional calculations are performed. On the other

¹ However a set of new energy-use scenarios have been calculated by the World Energy Council. These are discussed in section 13a of this report.

hand, groups will differ in their capabilities to produce large numbers of calculations. Therefore we have ranked the calculations in order of importance (section 1b).

The reasons for including the various additional calculations are as follows:

Additional stabilization cases (S350, S550, S750) These give a more extensive range of options.

The 'science' cases (DEC0%, DEC1%, DEC2%) These cases with specified percentage reductions in industrial carbon releases, allow direct comparison of model differences using concentrations from forward calculations. In many contexts, forward calculations are easier to explain.

The 'delayed' cases (DS450, DS650) These provide alternative pathways to stabilization that may be more realistic, given the difficulties of achieving stabilization, and at any rate provide cases for assessing the implications of delay.

IS92a-f These provide projections that can be related to specific energy policies as described in the IPCC 1992 WG1 update.

Impulse response function (Iinit, Ipert) This is required for determining the greenhouse warming potentials (GWP) that characterise the relative importance of the various greenhouse gases.

A.3b. The basis of technical specifications

Our principal requirements are that models should produce an acceptable fit to the record of CO₂ concentrations over the industrial period, and that models should treat processes in the past and future in an equivalent manner. We also require that the components of the model's carbon budget should lie within the recognized uncertainties in the observed carbon budget.

There are two different ways that these requirements may be achieved, using either forward or inverse modelling. In the forward modelling approach, input emissions from industrial sources and land-use changes are specified, and model parameters adjusted to achieve an optimum fit between modelled and observed concentrations. In the inverse approach, the model is run so as to match observed concentrations exactly, producing emissions as output. Model parameters are then adjusted to optimize the fit between modelled emissions and independent estimates of these emissions. Modelling groups are free to choose either of these methods (or, if possible, both) for the period 1765 to 1990.

In choosing the modelling strategy modellers should consider

- The fact that the atmospheric CO₂ history is better known than most other components.
- The important role of ¹⁴C in constraining ocean carbon uptake.

A further requirement is that the calculations performed by the various modelling groups be as easy to compare and interpret as possible. To this end, we have imposed certain constraints on the calculations (see section 1c) in addition to providing standardized input data sets for past concentrations, past industrial emissions and past net emissions from land-use changes: See Note A.6.A.

Use of land-use emissions data as an input raises two problems, specifying emissions prior to 1850, and accounting for differences between net and gross emissions: see Notes A.6.A and A.6.B.

A.4. Data sets prescribed

All input data will be provided as annual time series on Mac or PC floppy disk. Ian Enting will also be able to send the files by e-mail to any group that supplies an address accessible from internet. In addition, the data sets are available through the ftp facility described in section A.5.

Concentration data are smoothed versions of data from ice cores (Neftel et al., 1985; Friedli et al., 1986) and atmospheric observations (MLO data from Keeling, 1991). They begin in 1765, which is the required starting year for all calculations. Industrial emissions are those given by Keeling (1991) and Marland and Boden (1991), linearly extrapolated back to zero in 1844 to avoid a step function start at the beginning of the published record. The additional data for 1990 and revisions for 1985 to 1989 are from Marland (personal communication). Land-use-change emissions are the latest data from R.A. Houghton (1993, personal communication, see also Notes A.6.A and A.6.B). Industrial and land-use emissions data will be supplied on floppy disk as annual values.

Year Decade	0	1	2	3	4	5	6	7	8	9
1850	0.4357	0.4675	0.4824	0.4942	0.5040	0.5122	0.5189	0.5245	0.5296	0.5344
1860	0.5419	0.5336	0.5389	0.5440	0.5491	0.5544	0.5569	0.5604	0.5645	0.5688
1870	0.5718	0.5816	0.5877	0.5915	0.5958	0.6001	0.6049	0.6072	0.6089	0.6102
1880	0.6113	0.6422	0.6563	0.6629	0.6684	0.6728	0.6768	0.6777	0.6793	0.6790
1890	0.6872	0.6844	0.6850	0.6844	0.6833	0.6821	0.6804	0.6795	0.6812	0.6830
1900	0.6862	0.7485	0.7668	0.7817	0.7946	0.8060	0.8170	0.8221	0.8249	0.8270
1910	0.8283	0.7731	0.7572	0.7460	0.7435	0.7338	0.7372	0.7365	0.7360	0.7359
1920	0.7360	0.7846	0.7990	0.8110	0.8166	0.8274	0.8246	0.8243	0.8047	0.7804
1930	0.7747	0.7832	0.7786	0.7736	0.7677	0.7616	0.7674	0.7652	0.7616	0.7571
1940	0.7509	0.7300	0.7231	0.7161	0.7124	0.7094	0.7578	0.7682	0.7740	0.7759
1950	0.7824	0.9463	0.9872	0.9918	1.0885	1.1264	1.1551	1.0907	1.0828	1.0870
1960	1.0854	1.2317	1.2707	1.3092	1.3390	1.3627	1.3327	1.3323	1.3452	1.3500
1970	1.2972	1.2771	1.2459	1.2791	1.2595	1.2554	1.3386	1.3629	1.3628	1.4045
1980	1.4149	1.4351	1.4795	1.5098	1.5674	1.5959	1.6606	1.6942	1.7113	1.7053
1990	1.7129	—	—	—	—	—	—	—	—	—

Table A.1 Revised emissions from land-use change. From Houghton — see papers from IPCC meeting on feedbacks, Woods Hole, October 1992. Annual releases in Gt C.

Year Decade	0	1	2	3	4	5	6	7	8	9
1840	0.000	0.000	0.000	0.000	0.000	0.006	0.012	0.019	0.025	0.031
1850	0.037	0.044	0.050	0.056	0.062	0.068	0.075	0.081	0.087	0.093
1860	0.093	0.099	0.098	0.106	0.115	0.122	0.129	0.138	0.137	0.142
1870	0.145	0.162	0.176	0.188	0.184	0.189	0.192	0.196	0.197	0.208
1880	0.227	0.244	0.263	0.280	0.282	0.276	0.279	0.298	0.322	0.328
1890	0.350	0.365	0.369	0.362	0.377	0.399	0.412	0.432	0.455	0.497
1900	0.525	0.540	0.553	0.606	0.613	0.647	0.696	0.771	0.737	0.769
1910	0.805	0.822	0.866	0.929	0.838	0.831	0.895	0.945	0.932	0.829
1920	0.959	0.828	0.891	1.005	0.989	1.006	1.006	1.097	1.091	1.172
1930	1.077	0.968	0.874	0.919	0.996	1.032	1.146	1.226	1.161	1.233
1940	1.300	1.337	1.334	1.364	1.352	1.204	1.271	1.422	1.517	1.497
1950	1.638	1.775	1.803	1.848	1.871	2.050	2.185	2.278	2.338	2.471
1960	2.586	2.602	2.708	2.855	3.016	3.154	3.314	3.420	3.596	3.809
1970	4.091	4.242	4.409	4.648	4.656	4.629	4.895	5.034	5.082	5.366
1980	5.264	5.129	5.094	5.085	5.243	5.416	5.600	5.698	5.912	6.024
1990	6.097	—	—	—	—	—	—	—	—	—

Table A.2. Industrial emissions, as tabulated in Trends 91, linearly extrapolated pre-1860. For 1985 to 1990 new values are from Marland, personal communication. Annual releases in Gt C.

A.5. Reporting and model comparison process

In order to enhance communication for the exercise we have set up an *anonymous ftp* area at CSIRO¹. This will contain data files, news, utility programs etc.

To access it, the procedure *with user input typed like this* and **machine prompts** in bold is

normal prompt *ftp ftp@dar.csiro.au*

ftp> Name: *ftp* or *anonymous*

password Type your e-mail address

ftp> cd IPCC

ftp> ascii

ftp> get filename using actual filenames. This can be repeated as required.

ftp> quit

The file 'read.me' contains a list of the files currently in the area, together with a brief description. The file 'news' will contain progress reports.

All participants will be kept informed of progress as this exercise proceeds. There will be an opportunity to discuss results at the Global Change Institute in Snowmass, Colorado (July 18-30, 1993), to which many modellers have been invited. However, the main venue for discussing results will be at a special IPCC Working Group 1 meeting on 18 September, 1993, following the Fourth International CO₂ Conference in Carqueiranne, France².

¹ The description below describes the current access mode. The instructions as originally circulated described an earlier form

² Details of discussion ???

On the morning of 18 September, representatives of each modelling group will be given an opportunity to present their calculations. Subsequent discussion will focus on areas of disagreement or uncertainty that have emerged from the presentations, with a view to either resolving those issues which can be resolved immediately, or identifying those issues where disagreement or uncertainty remains and which can only be addressed in the longer term. In order to facilitate discussion at this meeting, results will have to undergo at least some preliminary inter-comparison and interpretation prior to the meeting. We are therefore requesting that results be submitted to either or both of the conveners by the end of July at the latest. Results should be provided in microcomputer-readable form (Mac or PC disk). The eventual plan is to publish the main results as a multi-authored paper in a leading journal. Participants are, however, encouraged to publish complete details of their results individually.

An overall assessment of the modelling exercise will form part of the carbon cycle chapter (lead authors D. Schimel, T. Wigley, D. Alves, M. Heimann, U. Siegenthaler³, I. Enting) of the 1995 IPCC WG1 assessment.

A.6. Technical notes

Technical note A.6.A: Land-use emissions prior to 1850

Since Houghton's published land-use emissions data begin in 1850 (see Table A.1) and since the extrapolated industrial emissions begin in 1844, forward calculations will give constant concentrations over 1765–1843. We can, however, be quite confident that concentration levels increased over this period. While there is weak evidence that some of this increase may have been part of a natural fluctuation, possibly associated with the Little Ice Age, in order to ensure comparability our standard case ignores this possibility.

Since the initial (1850) land-use emissions value is substantially greater than zero, we can also be reasonably sure that land-use emissions were non-zero over 1765–1849. After consultation with Houghton, we have therefore extended the Houghton record back to 1765 by linearly interpolating from 0.2 Gt C y⁻¹ in 1765 to the Houghton value in 1850.

Technical note A.6.B. Net flux vs. gross flux from deforestation

It should be noted that the Houghton data and the IS92a-f 'deforestation' fluxes are net emissions that account for fluxes associated with processes like regrowth. Depending on model structure, required land-use emissions input (or output when run in inverse mode) may be either net or gross emissions. A number of carbon cycle models (including our own) require gross land-use emissions as their required input. With such models, an imposed gross land-use emissions flux leads to additional model-generated fluxes and, hence, to a difference between the model-determined net terrestrial carbon mass change and the input land-use changes, $D_g(t)$. At least part of this difference may be interpreted as effective regrowth. However, being model-dependent, this will not necessarily be consistent with regrowth values that may be

³Uli Siegenthaler has been replaced by D. Raynaud

Model L

Name: LLNL Carbon Cycle Model

Description:

Modellers: D.J. Wuebbles and A.K. Jain

Contact:

Institution: Global Climate Research Division, Lawrence Livermore National Laboratory Livermore, CA 94550 USA

Summary:

Ocean: Surface mixed layer, plus deep ocean treated as a continuous advective-diffusive system.

Terrestrial: Single reservoir.

Initialisation:

Other:

There is an important feedback for the atmospheric CO_2 , namely, the direct interaction of atmospheric CO_2 with the deep ocean which is nearly free of excess CO_2 . We have taken this feedback into account by introducing a polar sea feedback parameter, π_c , defined as the change in the surface concentration in the polar region relative to that in the non-polar region. This parameter is similar to that used by IPCC in their energy balance model to represent the variation of polar sea temperature. The value of π_c would lie between 0.0 and 1.0. For $\pi_c = 0.5$, the model estimated value is 2.08 GtC/yr which is in good agreement with the IPCC estimates of 2.0 ± 0.8 GtC/yr. Therefore, we have used $\pi_c = 0.5$ in our model calculations.

Calibration:

The average oceanic inventory of bomb ^{14}C is determined by integrating the difference between the pre-bomb (1950) and post-bomb (1974) vertical profiles. An average model estimated total CO_2 concentration of 2.31 mol m^{-3} is used to convert model ^{14}C units into ^{14}C inventories. The model estimated inventory is $6.75 \times 10^9 \text{ atom/m}^2$.

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model M

Name:

Description:

Modellers: Moore, Braswell

Contact:

Institutions: U. New Hampshire

Summary:

Ocean:

Terrestrial:

Initialisation:

Other:

Calibration:

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model O

Name: Paris GCM

Description:

Modellers: J.C. Orr† and P. Monfray‡

Contact:

Institutions: †Laboratoire de Modelisation du Climat et de l'Environnement, DSM / CEN Saclay / CEA, L'Orme des Merisiers, Bât. 709, F-91191 Gif-sur-Yvette Cedex, FRANCE

‡Centre de Faibles Radioactivites, Laboratoire CNRS-CEA, F-91198 Gif-sur-Yvette Cedex, FRANCE

Summary:

Ocean: LODYC Ocean GCM.

Terrestrial: None

Initialisation:

Other:

Calibration:

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model P

Name:

Description:

Modeller: T.-H. Peng

Contact:

Institution: Environmental Sciences Division of Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A.

Summary:

Ocean: Box-diffusion model.

Terrestrial:

Initialisation:

Other:

Calibration:

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model Q**Name:****Description:****Modellers:** C. Le Quere and J.L. Sarmiento**Contact:****Institution:** Program in Atmospheric and Oceanic Sciences,
Princeton University, Sayre Hall, Forrestal Campus,
P.O.Box CN710, Princeton, NJ 08544-0710, USA**Summary:****Terrestrial:** None in version 1.**Ocean:****Initialisation:****Other:****Calibration:** The uptake of bomb-produced ^{14}C predicted by Toggweiler et al.'s P model is 16% less than observations. The changes made since publication do not alter the total uptake of carbon by more than a few percent.**Documentation:** Toggweiler et al. (1989).**Cases run:** Q₁ ocean only. Q₂ with vegetation.**Additional information:****Modeller's acknowledgements:**

Model R

Name: IMAGE 2.0

Description:

Modellers: J. Alcamo and M. Krol

Contact:

Institutions: National Institute of Public Health and Environmental Protection P.O.

Box 1, 3720 BA Bilthoven THE NETHERLANDS

Summary:

Ocean:

Terrestrial:

Initialisation:

Other:

Calibration:

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model T**Name:****Description:****Modeller:** L.D. Harvey**Contact:****Institution:** Dept. of Geography, University of Toronto, Totonto, Ontario, Canada.**Summary:****Ocean:****Terrestrial:****Initialisation:****Other:****Calibration:****Documentation:** Harvey (1989). see also Global Biogeochemical Cycles, 1991.**Cases run:****Additional information:****Modeller's acknowledgements:**

Model V

Name:

Description:

Modeller: J.A. Viecelli

Contact:

Institution: Lawrence Livermore National Laboratory

Summary:

Ocean:

Terrestrial:

Initialisation:

Other:

Calibration:

Documentation:

Cases run:

Additional information:

Modeller's acknowledgements:

Model W**Name:****Description:****Modeller:** T. Wigley**Contact:****Institution:** Office for Interdisciplinary Earth Studies, UCAR, Boulder Colorado**Summary:*****Ocean:*** Response function representation.***Terrestrial:******Initialisation:******Other:*****Calibration:****Documentation:** Wigley (1991, 1993).**Cases run:****Additional information:****Modeller's acknowledgements:**

Assessment of IPCC CO₂ Modelling Study**Appendix D: Tables, file a-tables**

Draft, printed March 31, 1994

Last change 13/12/93

This page is temporary for the purposes of holding production information and will not form part of the final document.

		A	B	C	E	F	G	H	J	L	M	O	P	Q	R	T	V	W
IS92a	C	Y	Y	-	Y	-	Y	-	Y	Y	-	-	-	-	Y	Y	-	Y
IS92b	C	-	-	-	Y	-	-	-	Y	Y	-	-	-	-	Y	-	-	Y
IS92c	C	-	-	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
IS92d	C	-	-	-	Y	-	-	-	Y	Y	-	-	-	-	Y	-	-	Y
IS92e	C	-	-	-	Y	-	-	-	Y	Y	-	-	-	-	Y	-	-	Y
IS92f	C	-	-	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
DEC0%	C	Y	-	-	Y	-	Y	-	Y	Y	-	-	-	-	-	-	-	Y
DEC1%	C	Y	-	-	Y	-	Y	-	Y	Y	-	-	-	-	-	-	-	Y
DEC2%	C	Y	-	-	Y	-	Y	-	Y	Y	-	-	-	-	-	-	-	Y
linit	R _T	-	-	-	Y	-	Y	Y	Y	Y	-	-	-	-	-	-	-	Y
lpert	R _T	-	-	-	Y	-	Y	Y	Y	Y	-	-	-	-	Y	-	-	Y
linit	R _O	-	-	-	-	-	-	Y	Y	Y	-	-	-	Y	-	S	-	S
lpert	R _O	-	-	-	-	-	-	Y	Y	Y	-	-	-	Y	-	S	-	S
linit	R _B	-	-	-	-	-	-	Y	-	-	-	-	-	-	-	-	-	-
lpert	R _B	-	-	-	-	-	-	Y	-	-	-	-	-	-	-	-	-	-
S650	E	Y	Y	-	Y	-	Y	Y	Y	Y	Y	-	Y	-	Y	Y	Y	Y
S450	E	Y	Y	-	Y	-	Y	Y	Y	Y	Y	-	Y	-	Y	Y	Y	Y
S350	E	Y	Y	-	Y	-	Y	-	Y	Y	-	-	-	-	Y	Y	-	Y
S550	E	Y	Y	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
S750	E	Y	Y	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
DS550	E	Y	Y	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
DS450	E	Y	Y	-	Y	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
S650	S _{ocean}	Y	Y	-	-	-	Y	Y	Y	Y	Y	?	Y	Y	Y	Y	Y	Y
S450	S _{ocean}	Y	Y	-	-	-	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
S350	S _{ocean}	Y	Y	-	-	-	-	-	Y	Y	Y	-	-	Y	Y	Y	-	Y
S550	S _{ocean}	Y	Y	-	-	-	-	-	Y	Y	Y	-	-	Y	Y	Y	-	Y
S750	S _{ocean}	Y	Y	-	-	-	-	-	Y	Y	Y	-	-	Y	Y	Y	-	Y
DS550	S _{ocean}	Y	Y	-	-	-	-	-	Y	Y	Y	-	-	Y	Y	Y	-	Y
DS450	S _{ocean}	Y	Y	-	-	-	-	-	Y	Y	Y	-	-	Y	Y	Y	-	Y
S650	S _{fert}	Y	-	-	-	Y	-	Y	Y	Y	-	-	-	-	Y	Y	-	Y
S450	S _{fert}	Y	-	-	-	-	-	Y	Y	Y	-	-	-	-	Y	Y	-	Y
S350	S _{fert}	Y	-	-	-	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
S550	S _{fert}	Y	-	-	-	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
S750	S _{fert}	Y	-	-	-	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
DS550	S _{fert}	Y	-	-	-	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y
DS450	S _{fert}	Y	-	-	-	-	-	-	Y	Y	-	-	-	-	Y	Y	-	Y

Table D.1. Calculations contributed. 'Y' denotes calculations contributed, 'S' denotes ocean impulse response function specified directly to define model, '?' denotes calculation flagged as extending the ocean chemistry formalism beyond the range for which it was calibrated. The biotic fluxes S_{fert} are supplied either as net biotic flux or as the fertilisation component only, to be combined with net flux from land-use change.

Model	C(1765)	C(1900)	C(1950)	C(1970)	C(1980)	C(1990)	$\frac{dC}{dt}(1990)$
A ₂		293.70	309.41	326.49	339.91	353.83	1.59
E ₁	281.07	293.53	308.88	325.57	339.34	354.37	1.69
V _A	275.47	290.13	308.36	325.09	338.64	354.17	1.70
V _B	277.95	291.95	308.62	325.15	338.84	354.17	1.67
Z							
Z							

Table D.2 Details of forward initialisations

Model	1850	1900	1950	1960	1970	1980	1990
A1	0.28	0.51	0.82	1.01	1.34	1.81	2.25
A2	0.08	0.32	0.75	1.07	1.46	1.88	2.18
B	0.35	0.63	1.00	1.24	1.65	2.21	2.74
E	0.20	0.40	0.80	1.10	1.50	2.00	2.30
H	0.18	0.36	0.59	0.74	0.99	1.32	1.63
J	0.30	0.54	0.86	1.08	1.45	1.94	2.41
L	0.28	0.51	0.81	1.04	1.44	1.95	2.38
M	0.28	0.48	0.77	0.93	1.19	1.55	1.90
O	0.31	0.57	0.94	1.13	1.45	1.89	2.36
P	0.31	0.56	0.88	1.10	1.46	1.96	2.43
Q	0.26	0.47	0.75	0.91	1.19	1.59	2.01
R						1.42	1.63
T	0.28	0.49	0.78	0.94	1.20	1.59	1.98
W	0.29	0.54	0.89	1.07	1.37	1.81	2.24

Table D.3 Ocean fluxes pre-1990

Model	1850	1900	1950	1960	1970	1980	1990
A1	-0.30	-0.50	-0.68	-0.86	-1.17	-1.64	-2.04
A2	-0.14	-0.36	-0.70	-0.99	-1.35	-1.78	-1.98
T	-0.30	-0.51	-0.76	-0.91	-1.19	-1.69	-2.21
W	-0.30	-0.55	-0.87	-1.00	-1.22	-1.52	-1.95

Table D.4 Fertilisation fluxes pre-1990.

Model	2000	2050	2100	2150	2200	2300
A	7.25	9.17	8.91	6.16	3.03	2.05
B ₁	6.17	8.23	9.59	7.79	5.18	4.51
B ₂	5.98	8.26	9.95	8.12	5.62	5.01
B ₃	6.88	8.99	9.27	7.02	4.35	3.57
E	7.36	10.10	10.13	7.16	3.68	2.15
G	6.10	13.70	6.11	2.34	1.50	
H	7.11	8.57	8.35	5.79	3.16	2.38
J	7.21	9.89	10.10	7.30	3.88	2.31
L	7.33	9.80	9.75	6.80	3.25	1.99
M	6.73	8.68	9.20	5.32	2.88	2.18
O	5.25	7.65	8.31	6.33		
P	5.26	7.20	7.44	5.22	2.65	1.96
Q	4.78	6.81	7.11	4.95	2.39	1.65
T	7.26	9.77	10.44	7.89	4.51	3.04
V _A	8.77	16.89	8.37	1.92	0.37	
W	7.20	10.43	11.19	8.78	5.58	3.74

Table D.5 Industrial emissions S650.

Model	2000	2050	2100	2150	2200	2300
A	6.72	4.86	2.41	1.63	1.36	1.11
B ₁	5.64	4.20	3.61	3.57	3.54	3.40
B ₂	5.47	4.52	4.22	4.15	4.08	3.97
B ₃	6.37	5.25	3.54	3.05	2.81	2.52
E	6.83	5.72	3.18	2.04	1.53	1.11
G	6.10	3.33	1.49	1.49	1.49	
H	6.57	4.89	3.01	2.26	1.89	1.46
J	6.63	4.89	3.01	2.26	1.89	1.46
L	6.78	5.31	2.72	1.76	1.39	1.01
M	6.26	5.35	4.07	1.79	1.55	1.27
O	4.75	4.11	2.76	2.27	1.95	
P	4.76	3.60	1.97	1.53	1.32	1.07
Q	4.33	3.37	1.84	1.39	1.16	0.90
R	5.43	6.71	3.14			
T	6.76	5.32	2.95	1.90	1.53	1.15
V _A	6.89	4.13	3.15	2.86	2.82	
V _B	7.26	6.19	5.78	5.57	5.55	
W	6.66	6.19	4.29	3.31	2.73	2.20
Z						
Z						

Table D.6 Industrial emissions S450.

Model	0	10	25	50	100	200	300	400	500
M-R&H	1.0								
H	1.0	0.782	0.678	0.585	0.474	0.346			
J	1.0	0.697	0.568	0.470	0.376	0.293	0.250	0.223	0.205
L	1.0	0.739	0.629	0.536	0.439	0.345	0.296	0.268	0.253
Q	1.0	0.757	0.619	0.508	0.402	0.310			

Table D.7 Equilibrium response function, ocean only.

Model	0	10	25	50	100	200	300	400	500
H	1.0	0.820	0.731	0.645	0.535	0.399			
J	1.0	0.747	0.658	0.594	0.556	0.526	0.490	0.461	0.439
L	1.0	0.840	0.764	0.709	0.668	0.625	0.575	0.539	0.517
Q	1.0	0.796	0.694	0.623	0.575	0.551			

Table D.8 Perturbation response function, ocean only.

Model	0	10	25	50	100	200	300	400	500
H	1.0	0.517	0.350	0.275	0.230	0.202			
Z	1.0								

Table D.9 Equilibrium response function, biota only.

Model	0	10	25	50	100	200	300	400	500
H	1.0	0.765	0.610	0.496	0.409	0.352			
Z	1.0								

Table D.10 Perturbation response function, biota only.

Assessment of IPCC CO₂ Modelling Study**Appendix E: Figures**

Draft, printed March 31, 1994

Last change 30/3/94

This page is temporary for the purposes of holding production information and will not form part of the final document.

Figure E.1 Calculated emissions for S350 case

Figure E.2 Calculated emissions for S550 case

Figure E.3 Calculated emissions for S750 case

Figure E.4 Calculated emissions for DS450 case

Figure E.5 Calculated emissions for DS550 case

Figure E.6 Concentrations for 1% compound emission reduction

Figure E.7 Concentrations for 2% compound emission reduction

Figure E.8 Concentrations predicted using IS92b

Figure E.9 Concentrations predicted using IS92c

Figure E.10 Concentrations predicted using IS92d

Figure E.11 Concentrations predicted using IS92e

Figure E.12 Concentrations predicted using IS92f

Figure E.13 1995 impulse response relative to S650

Figure E.1 (incomplete)

Figure E.2 (incomplete)

Figure E.3 (incomplete)

Figure E.6

Figure E.7

Figure E.9

Figure E.11

Assessment of IPCC CO₂ Modelling Study

Appendix F: Notation, file a-notat.tex

Draft, printed March 31, 1994

Last change 22/3/94

This page is temporary for the purposes of holding production information and will not form part of the final document.

Appendix F: Notation

C_0	Initial atmospheric CO ₂ concentration.
$C(t)$	Atmospheric CO ₂ concentration at time t , in ppmv.
$D_n(t)$	Net carbon flux from land-use change.
G	Generic notation for response functions.
Q	Generic notation for anthropogenic sources.
Q_{foss}	Fossil carbon source
S	Generic notation for sinks.
S_{ocean}	Net ocean carbon sink
S_{fert}	CO ₂ sink from CO ₂ -fertilisation.
S_{resid}	Residual CO ₂ sink.
t	Time in years

Macro definitions

The following pages are for information during preparation and are not part of the final report. This macros are for use in preparing TeX source. Thses pages are printed from file expand.tex

SYMBOLS : String *COO* expands to $\text{CO}_2, \dots$ String *cotwo* expands to $\text{CO}_2, \dots$
 String *Cziii* expands to $^{13}\text{C}, \dots$ String *cthree* expands to $^{13}\text{C}, \dots$
 String *Cziv* expands to $^{14}\text{C}, \dots$ String *RC* expands to $^{14}\text{C}, \dots$
 String *PY* expands to $\text{y}^{-1}, \dots$ String *GTYC* expands to $\text{Gt C y}^{-1}, \dots$
 String *PM* expands to $\%, \dots$ String *DelRC* expands to $\Delta^{14}\text{C}, \dots$

maths mode expansions

String *Qfoss* expands to $Q_{\text{foss}}, \dots$ String *Soc* expands to $S_{\text{ocean}}, \dots$
 String *Sfert* expands to $S_{\text{fert}}, \dots$ String *Sresid* expands to $S_{\text{resid}}, \dots$
 String *DDT* expands to $\frac{d}{dt}, \dots$

PAGE REFERENCES :

String *onepage* expands to 6, ... String *twopage* expands to 20, ...
 String *threepage* expands to 30, ... String *fourpage* expands to 40, ...
 String *fivepage* expands to 50, ... String *sixpage* expands to 60, ...
 String *sevenpage* expands to 70, ... String *eightpage* expands to 80, ...
 String *ninepage* expands to 90, ... String *tenpage* expands to 100, ...
 String *elevenpage* expands to 110, ... String *twelvepage* expands to 120, ...
 String *thirtpage* expands to 130, ... String *fourtpage* expands to 140, ...
 String *appApage* expands to 300, ... String *appBpage* expands to 400, ...
 String *appCpage* expands to 500, ... String *appDpage* expands to 600, ...
 String *appEpage* expands to 700, ... String *appFpage* expands to 800, ...

Equations String *eqBUDG* expands to 4.1, ...

Files used

B-START TEX	Title pages and contents
B-ISSUE TEX	Sections 1 to 6
B-RES TEX	Sections 7 to 13
B-REFS TEX	References
A-RULES TEX	Appendix A: Instructions
A-DATA TEX	App B: description of data files
A-DESC TEX	App C: Descriptions of models
A-TABLES TEX	App D: Tables
A-FIG TEX	App E: Figures
A-NOTAT TEX	Appendix F: Notation
EXPAND TEX	Details of files, macros
MACRO STY	Macros, inc. macros for cross-references
TEMPHEAD TEX	x
REALHEAD TEX	x
KOALA STY	Style for DAR tech papers, modified for this report

**DATE
FILMED**

10/27/94

END