DOE/ER/40694--1

DE92 018758

NUCLEAR STRUCTURE MODELS: APPLICATIONS AND DEVELOPMENT

Progress Report

November 1, 1991 - June 30, 1992

Paul B. Semmes

Tennessee Technological University

Cookeville, TN 38505

July, 1992

PREPARED FOR THE U. S. DEPARTMENT OF ENERGY

UNDER GRANT NUMBER DE-FG05-92ER40694

TABLE OF CONTENTS

Preface	1
1. Studies of Superdeformed States	2
2. Rotational Structure in Odd-Odd Nuclei	2
2.1. Signature Inversion in Odd-Odd Nuclei: A Fingerprint of Triaxiality?	4
2.1.1. Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction	5
3. Rotational Structure in Odd-A Nuclei	9
3.1 Signatures of γ Deformation in Nuclei and an Application to ^{125}Xe	9
3.2 Nuclear Spins and Moments: Fundamental Structural Information	9
3.3 Electromagnetic Properties of ^{181}Ir : Evidence of β Stretching?	10
References	11
Appendix I: Personnel	12
Appendix II: Publications and Seminars	13
Appendix III: Reprints and Preprints	14

PREFACE

This report summarizes the progress made during the period November 1, 1991 to June 30, 1992 on Grant No. DE-FG05-92ER40694. This is the first annual report made under this grant, and most of the work described here is the product of continuing collaborations with various groups. Therefore, some overlap with earlier work is unavoidable in this report. The report is organized as follows: Superdeformed states are discussed Section 1, including a brief description of results given in a publication and also a view towards future work. Odd-odd nuclei at normal deformation are discussed in Section 2, including a background discussion (Secs. 2 and 2.1) and new work on the problem of signature inversion (Sec. 2.1.1). Major efforts are envisioned in this direction in the future. Finally, odd-mass nuclei at normal deformation are discussed in Section 3, and these applications have benefitted from direct collaboration with experimental colleagues.

The secretarial and accounting duties for this grant have been performed by Gloria Julian of the TTU Physics Department, and her efforts are greatly appreciated. I also wish to thank the personnel at the Department of Mathematical Physics, LTH, Lund, Sweden, and the Niels Bohr Institute in Risø, Denmark for their hospitality and assistance during my visit this year. Finally, extensive discussions with Drs. Ingemar Ragnarsson and Sven Åberg (LTH, Lund, Sweden), Ragnar Bengtsson (UNISOR, Oak Ridge, TN), Ramon Wyss and Witold Nazarewicz (Joint Institute for Heavy Ion Research, Oak Ridge) are gratefully acknowledged.

1. Studies of Superdeformed States

(with I. Ragnarsson and S. Åberg)

A major development was the realization that $\Delta I = 1$ M1 transitions between signature partner bands in superdeformed states in the Hg region should be able to compete favorably with the stretched $\Delta I = 2$ E2 γ -rays that typically dominate the decay paths. Furthermore, at the low transition energies expected at the fairly low spins ($I_0 \approx 10$) observed in this region, the M1 transitions are strongly enhanced by internal conversion. Consequently, the decay within the SD minimum for some bands may be dominated by these "invisible" conversion electrons, and the resulting loss of E2 intensity can mimic unobserved transitions from the SD minimum to the normal deformed states. The possibility of observing such $\Delta I = 1$ transitions between SD states, either directly or indirectly, is exciting because M1 properties are essentially determined by the unpaired nucleon(s) and thus should provide new insights into the single particle structure of SD states. Some implications of this scenario and experimental evidence for it in ^{193}Hg are discussed in publication 1.

Similarly, strong M1 transitions must be expected for some 2-quasiparticle configurations as well. Detailed calculations with the 2-particle + rotor model are planned to investigate which configurations are the best candidates for observable M1 transitions, and whether these calculations will support or clarify configuration assignments previously proposed for this region. So far the configuration assignments have been based mainly on cranking calculations using the Woods-Saxon potential, and so it is desirable to use the Woods-Saxon for the particle-rotor calculations also. Some modifications will be required to the existing 2-qp + rotor computer codes, but for both odd-odd nuclei and negative parity 2-qp states in even-even nuclei these are rather straightforward.

Finally, for both 1-qp and 2-qp states, these calculations can be used to test the reliability of the Berkeley procedure for determining nuclear spins from observed transition energies and expected rotational model behavior by treating the calculated energies as data. Similarly, it should be interesting to compare effective alignments derived from energy differences to expectation values of $R \cdot j$ computed from particle-rotor wave functions.

2. Rotational Structure in Odd-Odd Nuclei

Recently a 2-quasiparticles + rotor model for odd-odd nuclei was developed

(in collaboration with I. Ragnarsson) in which the unpaired proton and neutron are coupled to a rigidly deformed triaxial core [1]. The main features of the model are summarized as follows: The Hamiltonian is written

$$H = \sum_i A_i R_i^2 + \sum_{p,n} (H_{sp} + H_{pair}) + V_{pn} \quad (1)$$

where the first term describes the collective core rotation in terms of the rotational parameters A_i and the core angular momentum R . Inside the parentheses, H_{sp} describes the motion of the single particle in the static deformed field and H_{pair} is a monopole pairing interaction between like nucleons. The modified oscillator potential is used to describe the mean field, and pairing is treated by a standard BCS calculation on the eigenstates of the deformed single particle potential. The resulting adiabatic quasiparticle states are used to construct a basis for diagonalizing the total Hamiltonian as discussed below. The last term in the Hamiltonian is an explicit residual interaction between the odd proton and odd neutron, and the form of V_{pn} has been taken from previous studies of well deformed odd-odd nuclei [2,3]. Both finite range and zero range interactions can be used. The finite range interaction is written

$$V_{pn} = e^{-r^2/r_0^2} (u_0 + u_1 \vec{\sigma}_p \cdot \vec{\sigma}_n + u_2 P_M + u_3 \vec{\sigma}_p \cdot \vec{\sigma}_n P_M + V_T S_{12} + V_{TM} S_{12} P_M) \quad (2)$$

where P_M is the space exchange operator, σ is the Pauli spin matrix and S_{12} is the tensor operator. The different terms are characterized by the strength parameters u_0 , u_1 , u_2 , u_3 , V_T and V_{TM} , and r_0 is the range of the interaction. Similarly, the zero range interaction is written

$$V_{pn} = 4\pi\sqrt{\pi b^3/\sqrt{2}} \delta(\vec{r}_p - \vec{r}_n) (u_0 + u_1 \vec{\sigma}_p \cdot \vec{\sigma}_n) \quad (3)$$

where b is the oscillator length and u_0 and u_1 are strength parameters.

Strong coupling basis states, symmetrized under the D_2 point group, are constructed from intrinsic 2-quasiparticle states. Note that for each pair of adiabatic orbitals $|\nu\rangle$, $|\mu\rangle$ and their conjugates (obtained from a rotation of 180° about the intrinsic x-axis), two distinct intrinsic states ("partner states") can be constructed, i.e.,

$$\Phi_{pn} = \alpha_v^\dagger \beta_\mu^\dagger |\tilde{0}\rangle \quad \text{and} \quad \Phi_{pn} = \alpha_v^\dagger \beta_\mu^\dagger |\tilde{0}\rangle \quad (4)$$

where $\alpha^\dagger$ and $\beta^\dagger$ are the quasiproton and quasineutron creation operators. The BCS vacuum state represents the effective core in this formulation. In the case of axial symmetry, Ω is a good quantum number for each adiabatic orbital, and the two partner states correspond to the familiar K doublets

$$K = \Omega_p + \Omega_n \quad \text{and} \quad K = \Omega_p - \Omega_n \quad (5)$$

Finally, the total Hamiltonian is diagonalized numerically with a sufficient

number of basis states included to insure numerical convergence (typically 5 to 10 proton and neutron Nilsson orbitals close to the Fermi level). From the energy eigenvectors, M1 and E2 electromagnetic matrix elements (diagonal and off-diagonal) can be calculated. Various angular momentum expectation values are also calculated, such as $\langle j_p \cdot R \rangle / |R|$, which represents the projection of the proton angular momentum in the direction of the collective core rotation vector. This quantity is a useful measure of the proton "alignment" in this model, and is meaningful for both triaxial as well as axially symmetric shapes.

A few features of this model are particularly noteworthy:

- The total angular momentum is conserved.
- Normal parity Nilsson orbitals can be selected as easily as unique parity orbitals for both the odd proton and neutron; there is no restriction to single j-shells.
- Triaxial shapes can be treated as well as axially symmetric.
- Both diagonal and off-diagonal matrix elements of the residual interaction between the various 2-qp states are retained, and many 2-qp configurations can be included in the basis.

As a result of these features, this model provides a powerful and flexible framework for studying the properties of rotational states in odd-odd nuclei.

2.1 Signature Inversion in Odd-Odd Nuclei: A Fingerprint of Triaxiality?

(with I. Ragnarsson)

As a consequence of the "turn about" symmetry in rotational nuclei, it is possible to define a signature quantum number α and to classify rotational bands as $\Delta I = 2$ sequences of states with $\alpha = I \bmod 2$. The usefulness of this classification is evident when deviations from strong coupling behavior arise. Such deviations often show a clear signature dependence and simple rules can be derived by considering the perturbations introduced by the Coriolis interaction. For example, in an odd-odd nucleus in which both the unpaired proton and the unpaired neutron occupy unique parity (high-j) orbitals, the even (or odd) spin states are energetically favored compared to the strong coupling $I(I+1)$ energy law if the sum $(j_p + j_n)$ is even (or odd). A similar rule exists for odd-mass nuclei. However, these rules are known to be violated in a number of cases, all involving 2- or 3-quasiparticle states at low rotational frequency. A variety of explanations have been proposed for this signature inversion, usually invoking

either the gamma degree of freedom or proton-neutron residual interactions. Calculations with triaxial shapes or gamma vibrations generally require that the collective rotation occur mainly about the shortest nuclear axis, i.e., "outside the collective sector", as in [4-6], among others. For odd-odd nuclei, the importance of the proton-neutron interaction has been stressed in ref. [7], while somewhat schematic calculations suggest that the inversion can occur for axially symmetric shapes if angular momentum is properly conserved [8,9].

The $\pi h_{11/2} \otimes \nu i_{13/2}$ band in ^{152}Eu is one of the best studied cases in which signature inversion is observed [10,11], and the 2-qp + rotor model has been used to study the necessity of triaxial shapes and/or the influence of the residual interaction. These results have been reported in conference proceedings [12], and a full account is being prepared for publication. Briefly stated, fair agreement with the data could be obtained with no V_{pn} but with a triaxial shape only if the moments of inertia are changed significantly from the usual hydrodynamical estimates so that the core rotation occurs mainly about the shortest nuclear axis. This agrees qualitatively with the cranking results of refs. [5] and [11], but the required moments of inertia are difficult to understand. Excellent agreement with the available data can be obtained for an axially symmetric shape and a zero range residual interaction (Eqn. 3), but only if the u_0 term is included. With an axially symmetric shape but no residual interaction, a signature inversion was obtained in the calculations but at too low spin compared to the data, and furthermore, the signature splitting at higher spin (e.g., $I = 18$) was much too large. If a "standard form" of the residual interaction was included (using either finite range or zero range interactions taken from ref. [2] or [3]) the agreement was even worse. These standard forms omit the u_0 term, which is crucial to our good results; therefore it is imperative to understand whether our V_{pn} is physically reasonable. In most studies of well deformed odd-odd nuclei it is commonly assumed that a single strong coupling basis state dominates the rotational wave function, and that the matrix elements of the residual interaction can be extracted from experimental energies simply from lowest order perturbation theory. The parameters of a residual interaction, e.g. as in Eqns. 2 or 3, are then fitted to the empirically determined matrix elements (Gallagher-Moszkowski splittings and Newby shifts). In ref. [2] it is shown that the u_0 term does not contribute to either the Gallagher-Moszkowski splitting or the Newby shift, and so this term is commonly

omitted in such phenomenological fits. However, in the present calculations, the $\pi h_{11/2} \otimes \nu i_{13/2}$ band contains many strong coupling components, and is not dominated by any one. Thus, there is no reason why a perturbational treatment of V_{pn} should be sufficient and no obvious reason why the u_0 term should be negligible. Additional calculations in other mass regions are certainly needed to confirm and clarify the importance of the u_0 term in the proton-neutron residual interaction.

2.1.1 Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction

(with B. Cederwall, I. Ragnarsson and R. Wyss)

A number of rotational bands were observed in a recent high-spin experiment on the odd-odd nucleus ^{120}Cs , including one band exhibiting a strong signature inversion at low spins. This band was assigned as the $\pi h_{11/2} \otimes \nu h_{11/2}$ structure, and Total Routhian Surface calculations (by R. Wyss) indicate an essentially prolate shape at low spins, with $\beta \approx 0.25$, $\gamma \approx +6^\circ$. However, the TRS calculations also suggest that ^{120}Cs should be much stiffer in the γ plane than ^{152}Eu , and that the signature inversion at low spins cannot be understood in this case within the $\gamma > 0^\circ$ scenario. Calculations were made with the 2-qp + rotor model similar to those described above for ^{152}Eu , and again the signature inversion can be naturally explained with the zero range residual interaction, but only if the u_0 term is included. Calculated and experimental energies within this band are shown as a function of angular momentum I in fig. 1. Note that the smooth curves connecting the two signature sequences (shown by open and filled circles) cross at about $I \approx 16$ in the data (panel "a"). Panel "b" shows the calculated energies for the axially symmetric core shape and no V_{pn} ; clearly the crossing comes too low in spin, and the splitting between the signature sequences is too large for $I > 16$ or so. The agreement is even worse if a "standard form" of the residual interaction is used, i.e., without the u_0 term. The calculated results for a triaxial shape (with $\gamma > 0^\circ$ simulated by interchanging the moments of inertia about the intrinsic x- and y-axes from their hydrodynamical estimates) is shown in "c", and clearly even this large triaxiality cannot reproduce the signature inversion. The calculated energies for an axially symmetric shape and the same zero range V_{pn} as used in ^{152}Eu (including the u_0 term) are shown in "d", and the agreement with the data is now quite good. Finally, the experimental $B(M1; I \rightarrow I-1)/B(E2; I \rightarrow I-2)$ ratios for this band are compared with the calculations in Fig. 2, and these also support the V_{pn} calculations. Overall,

Fig. 1: Level energies as a function of angular momentum I for the $\pi h_{11/2} \otimes \nu h_{11/2}$ band in ^{120}Cs . (Note that the bandhead spin is not firmly established experimentally.)

- a) Experimental values
- b) Calculated level energies with the 2-qp + rotor model for an axially symmetric shape and no residual proton-neutron interaction.
- c) Calculated energies as in b), but with a triaxial shape, $\gamma = +20^\circ$.
- d) Calculated energies as in b), but with the residual interaction included.

Fig. 2: Experimental and calculated $B(M1; I \rightarrow I-1)/B(E2; I \rightarrow I-2)$ ratios as a function of angular momentum I for the $\pi h_{11/2} \otimes \nu h_{11/2}$ band in ^{120}Cs . Experimental values are given by the filled circles (with error bars), and the calculated values are connected with solid lines.

Filled circles: Calculated for an axially symmetric shape and no residual proton-neutron interaction.

Open circles: Calculated energies with the triaxial shape, $\gamma = +20^\circ$.

Diamonds: Calculated with the axially symmetric shape and with the residual interaction included.

these results agree completely with those obtained previously for ^{152}Eu , and again strongly suggest that the u_0 term of the residual interaction cannot generally be neglected in odd-odd nuclei. However, it must be noted that unlike ^{152}Eu , the spins and parities of these states in ^{120}Cs are not well established experimentally.

These results are included in a paper submitted for publication (see Appendix II) that also contains the experimental results.

3. Rotational Structure in Odd-A Nuclei

Three applications of the 1-qp + rotor model that have resulted in publications or written conference contributions are briefly discussed here. A copy or preprint of each publication is also appended to this report.

3.1 Signatures of γ Deformation in Nuclei and an Application to ^{125}Xe

(with D. Lieberz, A. Gelberg, P. von Brentano and I. Ragnarsson)

In this work (publication 2) it is shown that the low energy structure of both the positive and negative parity states in ^{125}Xe is well described by the particle + triaxial rotor model with a "prolatish" triaxial shape specified by the modified oscillator parameters $\epsilon = 0.21$ and $\gamma = 24^\circ$. Furthermore, characteristic features are found in both the positive and negative parity states that determine the degree of triaxiality (γ) rather precisely (these are lifetime of the $I = 7/2+$ isomer and the signature splitting of the $h_{11/2}$ states). The good agreement for both parities with a common shape contradicts earlier claims of shape coexistence [13,14].

3.2 Nuclear Spins and Moments: Fundamental Structural Information

Calculations with the particle + triaxial rotor model using the Woods-Saxon potential have been made in an attempt to predict characteristic features that would signal the onset of strong quadrupole deformation in the region of light rare earth nuclei, with $A \approx 130$. These nuclei are especially interesting because the development of quadrupole shapes has been previously predicted [15,16]. The odd-A calculations build upon those mean field predictions, and all parameters are taken at standard values without any adjustments. Characteristic electromagnetic properties of Nilsson orbitals expected for ^{133}Pm , and $^{133,135}\text{Sm}$ are presented.

This region of nuclei is also one focus of current UNISOR experiments (led by J. L. Wood and J. Brietenbach), and additional calculations are planned to compare with new radioactive decay data for levels in ^{133}Nd .

3.3 Electromagnetic Properties of ^{181}Ir : Evidence of β Stretching?

(with U. Garg and W. Reviol)

Prolate-prolate shape coexistence has been predicted for many neutron deficient Ir isotopes from Total Routhian Surface calculations [17-19], perhaps accompanied by a transition from the less deformed shape ($\beta \approx 0.16$) to the more deformed shape ($\beta \approx 0.25$) as the spin increases. Phenomenological 3-band mixing calculations [20,21] seem to reproduce observed energy levels (alignments) and the $B(M1; I \rightarrow I-1)/B(E2; I \rightarrow I-2)$ ratios for the sequence $^{173,175,177,179,181}\text{Ir}$. However, such mixing calculations requires many parameters, and the M1 and E2 transition rates were estimated from semiclassical formulas rather than calculated from particle-rotor wave functions. That analysis also suggests that the shape transition (β stretching) is present at low spins for all these nuclei, even though the clear irregularities found experimentally in the energy levels of $^{173,175}\text{Ir}$ are not present in the heavier isotopes.

In this contribution, we show that the $BM1/BE2$ ratios in the $[514]9/2^-$ and $[402]5/2^+$ bands of ^{181}Ir are well described (up to $I \approx 29/2$ or so) with a particle-rotor calculation using the Woods-Saxon potential if the shape is fixed to that of the more deformed configuration ($\beta \approx 0.25$). These results contradict the claim of coexistence for ^{181}Ir , and similar results have also been obtained for ^{179}Ir [22]. Articles are being prepared for these results (both ^{179}Ir and ^{181}Ir).

REFERENCES

- [1] I. Ragnarsson and P. B. Semmes, *Hyperfine Interactions* **43** (1988) 425.
- [2] J. P. Boisson, R. Piepenbring and W. Ogle, *Phys. Rep.* **C26** (1976) 99.
- [3] H. Frisk, *Z. Phys.* **A330** (1988) 241.
- [4] I. Hamamoto and B. R. Mottelson, *Phys. Lett.* **127B** (1983) 281; *Phys. Lett.* **132B** (1983) 7.
- [5] R. Bengtsson et al., *Nucl. Phys.* **A415** (1984) 189.
- [6] A. Ikeda and T. Shimano, *Phys. Rev.* **C42** (1990) 149.
- [7] H. Toki, H. L. Yadev and A. Faessler, *Phys. Lett.* **71B** (1977) 1.
- [8] A. J. Kreiner and M. A. J. Mariscotti, *J. Phys.* **G6** (1980) L13.
- [9] I. Hamamoto, *Phys. Lett.* **235B** (1990) 221.
- [10] J. A. Pinston et al., *Nucl. Phys.* **A361** (1981) 464.
- [11] R. Bengtsson and H. Frisk, *Nucl. Phys.* **A437** (1985) 263.
- [12] P. B. Semmes and I. Ragnarsson, in *Conference on High Spin Physics and Gamma-Soft Nuclei*, J. X. Saladin et al. (eds.) (World Scientific, 1991), p. 500; and *International Conference on Future Directions in Nuclear Physics with 4π Detection Systems of the New Generation*, Strasbourg, 1991 (in press)
- [13] A. Luuko, J. Hattula, H. Helppi, O. Knuuttila and F. Donau, *Nucl. Phys.* **A357** (1982) 319.
- [14] H. Helppi, J. Hattula, A. Luuko, M. Jaaskelinen and F. Donau, *Nucl. Phys.* **A357** (1982) 333.
- [15] G. A. Leander and P. Möller, *Phys. Lett.* **110b** (1982) 17.
- [16] B. Kern et al., *Phys. Rev.* **C36** (1987) 1514.
- [17] W. Nazarewicz, M. A. Riley and J. D. Garrett, *Nucl. Phys.* **A512** (1990) 61.
- [18] R. Bengtsson, *Nucl. Phys.* **A520** (1990) 201c.
- [19] R. Wyss, private communication and to be published.
- [20] G. Dracoulis et al., *Phys. Lett.* **257B** (1991) 21.
- [21] G. Dracoulis et al., *Nucl. Phys.* **A534** (1991) 173.
- [22] H.-Q. Jin, Ph. D. Thesis, University of Tennessee, 1991, and to be published.

APPENDIX I

Personnel

FACULTY

Paul B. Semmes

COLLABORATORS

S. Åberg	Lund Institute of Technology (LTH), Lund, Sweden
R. Bengtsson	UNISOR, Oak Ridge TN and LTH, Lund Sweden
B. Cederwall	The Royal Institute of Technology, Stockholm, Sweden
U. Garg	University of Notre Dame
A. Gelberg	University of Köln, Köln, Germany
D. Lieberz	University of Köln, Köln, Germany
I. Ragnarsson	LTH, Lund, Sweden
W. Reviol	University of Notre Dame
P. von Brentano	University of Köln, Köln, Germany
R. Wyss	Joint Institute for Heavy Ion Research, Oak Ridge, TN

APPENDIX II

PUBLISHED ARTICLES

"'Invisible' Decay of Superdeformed States", P. B. Semmes, I. Ragnarsson and S. Åberg, Phys. Rev. Lett. **68** (1992) 460-463.

"Signatures of γ Deformation in Nuclei and an Application to ^{125}Xe ", D. Lieberz, A. Gelberg, P. von Brentano, I. Ragnarsson and P. B. Semmes, Phys. Lett. **282B** (1992) 7-12.

ARTICLES IN PRESS

"Nuclear Spins and Moments: Fundamental Structural Information", P. B. Semmes, Hyperfine Interactions (Contribution to the Second International Conference on On-Line Nuclear Orientation and Related Topics, Oak Ridge, TN, October 16-19, 1991)

"Electromagnetic Properties of ^{181}Ir : Evidence of β -Stretching?", U. Garg, W. Reviol and P. B. Semmes (Contribution to the International Conference on Nuclear Structure at High Angular Momentum, Ottawa, May 18-21, 1992)

ARTICLES SUBMITTED FOR PUBLICATION

"Signature Inversion in ^{120}Cs - Evidence for a Residual p-n Interaction", B. Cederwall, F. Liden, A. Johnson, L. Hildingsson, R. Wyss, B. Fant, S. Juutinen, P. Ahonen, S. Mitarai, J. Mukai, J. Nyberg, I. Ragnarsson and P. B. Semmes, submitted to Nucl. Phys. A

SEMINARS AND LECTURES

"Particle-Core Coupling Models", invited lecture presented at the Workshop on Nuclear Structure Models, Oak Ridge, TN, March 16-25, 1992.

"The Particle - Rotor Model", invited lecture presented at the Hands-On Nuclear Structure Theory Workshop, Niels Bohr Institute Tandem Accelerator Laboratory, Risø, Denmark, June 15-16, 1992.

*2 conf papers
cycled separately;
reprints removed.
cb*

END

**DATE
FILMED**

9 / 14 / 92