THE PHYSICS OF PATTERN FORMATION AT LIQUID INTERFACES

PROGRESS REPORT

for the Period June 1, 1991 to May 31, 1992

DOE/ER/45131--10

James V. Maher

DE92 016971

Department of Physics and Astronomy
University of Pittsburgh
Pittsburgh, PA 15260

June, 1992

Prepared for

The U.S. DEPARTMENT OF ENERGY AGREEMENT NO. DE-FG02-84ER45131

A. Review of Research Accomplishments 6/91-5/92

During the past year we have submitted six papers for publication, three related to the dynamics of macroscopic interfaces, and ultimately all related to solidification, and three related to the internal structure of disorderly materials, with possible applications to the processing of composite materials. Three of the six papers have appeared in print, and their reprints are attached to this report. Two of the remaining papers are scheduled to appear soon in Physical Review A, and the last is still in the refereeing process at Physical Review A. In addition to completing all these projects during the past year, we have begun two new projects, one on pattern formation and one on aggregation within a composite system. The work reported below has been performed by the PI along with his students Patrick Gallagher (who graduated in September, 1991 and is now a postdoc at Boston University), Beth May and Hong Zhao (both of whom will graduate during the next four months), and new students Kathy McCloud and Levent Kurnaz. Where one of the projects required theoretical assistance, we cooperated with and published with two postdocs in the University of Pittsburgh solid state theory group, Chuck Yeung and Jaume Casademunt, neither of whom received any support from this grant.

1. Pattern Formation. We have recently begun an experiment which models the solidification effects of crystalline anisotropy in a Hele-Shaw cell. We have also completed an experiment wherein a constant gradient in the gap of a Hele-Shaw cell has produced far more dramatic effects on pattern formation than were predicted by the mathematical equations which should govern the pattern growth. These two experiments are both fundamental in the sense that we can write down the growth equations with confidence.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

In the second case we can also write down the boundary conditions with confidence and see the close similarities and well-defined differences between our case and directional solidification. In the first case we cannot write down exact boundary conditions, but we can make clear arguments about which experimental effects can arise from the dynamical equation and which must come from the boundary conditions. In a third experiment, we cannot write the equations, but we can exhibit the richness of patterns which can be formed by driving a non-Newtonian fluid, and we can present empirical regularities with the hope that these will lead to useful approximations for understanding the formation of complex patterns.

1a. Regular, Small Scratches as a Model of an Underlying Lattice. We have recently begun measuring the Saffman-Taylor steady-state finger in a Hele-Shaw cell, one plate of which is not smooth but rather is etched with a very regular array of shallow grooves. The grooves are 0.9mm deep and 0.20mm wide, arrayed in one plate in a square pattern of side-length 0.40mm and in another in a rectangular array with sides of lengths 0.40mm and 0.80mm. These scratched plates, used one at a time in conjunction with a smooth plate to form a Hele-Shaw cell with anisotropy allow us to introduce anisotropy into the Hele-Shaw problem in a well controlled way which should be closely analogous to the role of anisotropy from underlying crystal lattices in crystal growth. When we operate our Hele-Shaw cell by driving nitrogen gas to displace paraffin oil, the capillary length is always much larger than the side-lengths of the squares and rectangles, so we can make coarse-graining arguments to allow the use of macroscopic equations for the growth dynamics. S.K. Sarkar and D. Jasnow (Phys. Rev. A39, 5299(1989)) have argued that in the coarse-graining regime the dynamic equation of pattern growth is unaltered by microscopic anisotropy if the underlying lattice has square or hexagonal symmetry (just as the same symmetries give rise to isotropic electrical conductivities). Thus the effects of the plate etching seen with the square-etched plate can confidently be ascribed to the boundary conditions, while extra effects seen only with the rectangular array can be ascribed to alteration of the dynamical equation. Figure 1 shows the width of the Saffman-Taylor finger as a function of the strength of the anisotropy for the square array while Figure 2 shows results for the rectangular array. While small anisotropy gives results indistinguishable from those obtained with cells both of whose plates are smooth, larger values of the anisotropy can be seen to produce wider fingers in the steady state. We have not yet determined whether the same results are obtained when we vary the flow direction relative to the axis of symmetry of the etched plate (the data of the two figures are all for flows along the easy axis of the etched array). However, the preliminary result is that the underlying lattice profoundly affects the selected pattern even for the square array and that the effects are even more pronounced for the

rectangular array, indicating a role for the dynamical equation's change when the symmetry is broken.

1b. Scaling of Radial Patterns in a Viscoelastic Solution. We have driven an aqueous solution of polyethylene oxide out of a circular Hele-Shaw cell by injecting water into the center of the cell. Since the two liquid phases are miscible, there are only dynamical effects to allow patterns to form before diffusion mixes the liquids. The dynamical effects can be expected to be dominated by the known viscoelastic properties of the polymer solution, and this lets us try to relate the pattern formation process for this non-Newtonian case to the scaling rules which are known to apply to the polymer solution. We have four control variables, the rate of injection of water, Q , the gap in the Hele-Shaw cell, b , the molecular weight of the polymer, M_w , and the concentration of the polymer, c . The important results are summarized here and presented in more detail in the preprint appended to this report. Varying the values of the four control parameters results in a dramatic variety of patterns. This can be seen in Figure 2 of the preprint which shows twelve of the approximately sixty combinations we have measured. As happens so often in the study of nonlinear systems, we observe patterns which had previously been thought to be impossible for this problem. For example, several of the patterns show growth in the "fjords" which are normally expected to be shielded from pressure gradients and thus incapable of growing. In addition, several of the patterns show a very sparse structure which has some resemblance to fracture (this is not fracture, however, since the branches appear only by tip-splitting and the branch angles are never greater than approximately 50 degrees). We are able to discern regularity in two length scales, the initial twig width and the peak value of a modal analysis of the overall pattern, by forming a dimensionless number from a ratio of viscoelastic time and hydrodynamic time presented in the preprint (the results of the scaling analysis are presented in Figure 4 of the preprint).

1c. Perturbing Saffman-Taylor Flow with a Small Gap-Gradient. We have completed the first phase of our study of the effects of introducing an extra term into the dynamical equation which governs the growth of Saffman-Taylor patterns by orienting the plates of the Hele-Shaw cell slightly out of parallel. Since the reprint of this work is appended to this report, we will summarize the results succinctly, emphasizing that the equations are very close to those of directional solidification with the added attraction that the three relevant length scales are somewhat easier to calculate and control in this case than in solidification. The most dramatic effect we have seen involves a breakdown of the predictions of linear stability theory for the breakup of an initially flat interface. While the theory works fine for parallel plates, it fails dramatically for very small perturbations, perturbations so

small that their effects should be unobservable for our experiment (see Figure 8 of the reprint). In addition to the interesting effects for the breakup of the flat interface, the single finger which is formed as a quasi steady state is dramatically stabilized if the gap-gradient is negative (cell narrows as the flow continues) and dramatically destabilized if the gradient is positive (see Figures 1 and 2 in the reprint).

2. Microscopic Disorder in Composite Systems. During the past year we have completed the first phase of our work to understand the effects of wetting/adsorption effects inside composite materials. This first experiment concentrated on colloidal suspensions of polystyrene spheres. When the solvent is a mixture whose correlation length for concentration fluctuations is not negligible in comparison with the radius of the spheres, the wetting situation is not unlike that inside a composite material that includes wetted fibres. Our measurements have involved two distinctly different regimes, the temperature range where the solvent is separated into two phases and the range wherein the solvent components are miscible in all proportions. Our solvent was 2, 6 lutidine plus water, which has an inverted coexistence curve with a critical temperature at 33.5° C. Preprints are appended for each case, and we will only summarize the most important results.

2a. Two phase region. In this region, there is a well defined zone, comparable to complete wetting, in which all but one in $\sim 10^4$ of the polystyrene spheres concentrate in the favorable phase of the solvent, leaving the other phase virtually unpopulated. (Which phase is favored depends on the surface charge density of the spheres, but otherwise all phenomena are robust.) Several tens of millikelvin from the solvent critical temperature (exactly how far depends on the surface charge density) the system crosses out of this "complete wetting" regime, and polyballs begin to populate the liquid-liquid interface. As more and more polyballs join the interface, an ordered phase appears at the interface, and this ordered phase can be seen to climb up the side walls of the sample tube. The phase on the side walls can produce Laue rings when it diffracts a laser beam, and under some circumstances, has been seen to produce Laue spots. Our measurements indicate that this phase is correlated over one hundred particle diameters with a mean particle spacing of one particle diameter.

2b. One phase region. In the one phase region on a critical trajectory no peculiar effects are observed in our system. Dynamic light scattering allows us to monitor the size of both the concentration fluctuations and the polyballs as the critical point is approached. There is no evidence of any thick adsorption layer growing on the polyballs and the solvent correlation length grows with a power law exponent close to that expected without the

polyballs (we assume the small difference arises from anomalous viscosity effects since the typical case is to have a polyball not more than ten or so correlation lengths away. The situation changes drastically, however, when we go off the critical trajectory. For all trajectories we have measured on the wetting-unfavorable side of the coexistence curve, we have observed polyball aggregation in the one phase region several hundred millikelvin away from the coexistence curve. This is reminiscent of the aggregations seen by Beysens and Esteve (D. Beysens and coworkers; references 1, 2 and 3 in the appended preprint) in colloids of silica particles. We observe complete consistency between one-phase aggregation and two-phase partitioning; i.e., when we change particle surface charge density to change which phase the particles prefer in the two-phase region, we also change the side of the coexistence curve on which aggregation will appear in the one-phase region. The aggregation region can extend to temperatures well beyond the wetting temperature. We have also slowed the aggregation dramatically by using very low number density of colloidal particles, and we see no evidence of any thick layer of fluid growing on the particles before they aggregate.

B. PLANS FOR THE PERIOD 7/92-6/93

1. Pattern Formation. During the next year, we will work on three particularly favorable cases of clearly definable perturbations to simple Saffman-Taylor pattern formation:

1a. Flow over an Underlying Lattice. Our preliminary work with the etched-array cell reported in section A.1.a has raised several promising questions, all of which we can at least begin to address in the next year. By measuring flows whose growth direction is not parallel to one of the symmetry axes of the underlying lattice, we can accumulate more information on the role of the symmetry in the mobility tensor and its relative importance in comparison with the asymmetric boundary conditions for this flow. In addition, measurement of the detailed shape of the advancing finger tip should let us determine whether the solution nature selects to this problem is the same as that selected when there is no underlying lattice contributing asymmetry to the problem or whether our system has been kicked over into one of the family of other Saffman-Taylor solutions which are known to be unstable when there is no underlying lattice. Finally, we will measure the limits of stability as a function of capillary number and asymmetry. From our preliminary measurements we know that the asymmetry clearly reduces the limits of stability of the growing pattern against macroscopic perturbations. The details of this destabilizing by an underlying lattice could have profound importance for the

understanding of the detailed contributions of kinetic terms to crystal growth instabilities.

1b. Effects of Multiplicative Noise. The results reported above for imposing spatial gradients in a Saffman-Taylor flow indicate that such gradients have a dramatic effect on the growth of patterns. The first-order effect of coupling a perfectly smooth cell to acoustic noise would be to impose time dependent spatial gradients on the cell, gradients which would appear in the growth equations as multiplicative noise since the spatio-temporally varying mobility is operated on by a divergence operator in forming the dynamic equation of growth. We will put an acoustic transducer on the cell during the next year and begin exploring the effect of both pure acoustic waves and random vibrational noise on the growth of patterns. If such effects are as important as the spatial gradient work implies they might be, their analogs may have great importance for limits of control in crystal growth.

1c. Gradient-in-Gap Flow and Directional Solidification. As is discussed in the appended gap-gradient reprint, our gradient-in-gap cell can be made to follow equations which are very close analogs to directional solidification if the flow can be changed to one where the invading fluid is the one which wets the walls of the cell. During the past two years we have made several attempts to measure in this parameter regime. Our success has been limited by the fact that we have not been able to move the contact angle of the wetting preference very far past 90 degrees. Then when the flow velocity exceeds an unfortunately low limit, the dynamic contact angle, which is velocity dependent, drops below 90 degrees and the flow enters the previously studied regime. These failures have produced some fascinating oscillatory pattern trajectories which we want to study eventually, but in the coming year we will emphasize trying to work in the invasion-wetting regime where we can make some direct comparisons with directional solidification.

2. Random Systems and the Microscopic Rearrangement of Composite Materials. The aggregation and segregation of colloidal particles discussed in the two appended preprints raise several issues of interest for basic statistical mechanics and for the processing of composite materials. Since the first system we have studied involved charge-stabilized colloidal particles, we would like to know how much of the rich variety of phenomena the system exhibited came from solvent composition fluctuations and gradients and how much came from counterion distribution effects. During the next year we will attempt to measure aggregation and partitioning effects in a sterically stabilized colloidal dispersion with a binary solvent. Our first attempt will be to use hydrocarbon-stabilized silica particles in near-critical mixtures of cyclohexane and methanol. Such particles are available with

several different hydrocarbon chain-lengths, so if we can see the effect, we should be able to vary the steric-stabilizing parameter in a way which is not analogous to the variations we made in changing the surface charge density of the PLS spheres in our earlier work. The extra control may help us to identify the important physical rules which determine the phenomena.

PUBLICATIONS DURING THE PERIOD 6/1/91-5/31/92

1. Adsorption and Wetting on Moveable Walls, P. D. Gallagher and J. V. Maher, *Physica A* 177, 489 (1991).
2. Capillary Wave Relaxation for a Meniscus between Miscible Liquids, S. E. May and J. V. Maher, *Phys. Rev. Letters* 67, 2013 (1991).
3. Perturbing Hele-Shaw Flow with a Small Gap-Gradient, H. Zhao, J. Casademunt, C. Yeung and J. V. Maher, *Phys. Rev. A* 45, 2455 (1992).
4. Aggregation in Polystyrene-Sphere Suspensions in Near-Critical Binary-Liquid Mixtures, P. D. Gallagher, M. L. Kurnaz and J. V. Maher, submitted to *Phys. Rev. A*.
5. Viscoelastic Effects in Patterns between Miscible Liquids, H. Zhao and J. V. Maher, *Phys. Rev. A*, in press.
6. Partitioning of Polystyrene Latex Spheres in Immiscible Critical Liquid Mixtures, R. D. Gallagher and J. V. Maher, *Phys. Rev. A*, in press.
7. "Universal Features in Transient Patterns," NATO Advanced Research Workshop on "Growth Patterns in Physical Sciences and Biology", Granada, Spain 10/91, (Invited Talk).
8. Polystyrene Latex Spheres in the One Phase Region of a Critical Binary Liquid Mixture, M. L. Kurnaz, P. D. Gallagher and J. V. Maher, *Bull. Am. Phys. Soc.* 37, 393 (1992).
9. The Steady State Regime for Flow in a Hele-Shaw Cell with Anisotropy, K. V. McCloud and J. V. Maher, *Bull. Am. Phys. Soc.* 37, 633 (1992).
10. Polystyrene Latex Spheres in the One-Phase Region of a Critical Binary Liquid Mixture, M. L. Kurnaz, P. D. Gallagher and J. V. Maher, 66th ACS Colloid and Surface Science Symposium, Morgantown, WV, 6/92.

Square Lattice

FIGURE 1

Rectangular Lattice

FIGURE 2

**DATE
FILMED
8/20/92**