AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

March, 1993

Studies in Quantum Field Theory

PI: Daniel G. Caldi

Technical Report

New Phase of QED and the GSI Peaks

I have continued to explore the phenomenon we found in which the expectation value of $\bar{\psi}\psi$, evaluated in certain background classical electromagnetic field configurations, is nonzero. This is a signal of spontaneously broken chiral symmetry, and so, of a phase of QED which is different from the usual perturbative one. Besides being of interest for its own sake as a new aspect of QED and a new way of inducing chiral symmetry breaking in a field theory, this observation is also important for the scenario which Alan Chodos of Yale University and I and our collaborators have proposed as the explanation for the anomalous peaks seen in the low-energy heavy-ion collision experiments at GSI, now being repeated and improved at Argonne and also at GSI. Related experiments are also now being done at SLAC under the direction of A. C. Melissinos of Rochester University. The foundation of our explanation is that the large and rapidly varying background electromagnetic fields generated in the ion collisions, can induce a rearrangement of the QED vacuum so that there is a metastable region where there is a new phase of QED. An important characteristic of this new phase is spontaneous chiral symmetry breaking, so that a new scale is generated, giving the possibility of electron-positron bound states in the range of 1.7 MeV. Motivated by this scenario, as well as for its own sake, we have been searching for direct evidence that background fields, similar, if possible, to those in the experiments, could cause spontaneous chiral symmetry breaking.

In a recent paper, my former graduate student, Saeid Vafaeisefat, who is now a postdoc here at SUNY Buffalo, and I have reported on

positive evidence for this:

D.G. Caldi and S. Vafaeisefat, "Breaking chiral symmetry in QED with external electromagnetic fields," Phys. Lett. B **287** (1992) 185.

This paper gives our results for an external electric field with a gaussian shape in time and one spatial direction. We have continued to explore the parameter space for this configuration, so that we can determine the extent of the region in space and for how long in time, $\langle \bar{\psi}\psi \rangle_{\text{ext}}$ is nonzero. We are currently preparing another paper with the results on the parameter space and also giving details about how the calculations are done.

We have also studied a gaussian configuration varying in three-dimensional space as well as in time, and again we have found that for certain ranges of the widths there is a non-zero value for $\langle \bar{\psi}\psi \rangle_{\text{ext}}$. These results are very encouraging since this 3-D configuration is even closer to the one in the experiments. We are currently writing a paper on this as well.

Along with Alan Chodos and joined by my graduate student, Z. Zhu, Vafaeisefat and I have been looking at some simplified one-dimensional field configurations, which have the advantage of being tractable analytically, in order to understand better the mechanism whereby these varying field configurations may induce chiral-symmetry breaking.

Conformal Field Theories, Quantum Groups, and Integrable Systems

My graduate student Z. Zhu and I are continuing to study the role of quantum groups in making a system integrable and so understandable non-perturbatively. There are a number of systems now known which are integrable in large part because they have a quantum group symmetry. These include systems which do not have conformal symmetry, as well as those which do. The class of systems we have been studying are the affine Toda field theories. We have been especially concerned recently with finding multi-soliton solutions for particular theories and using them to determine the S-matrix. A paper on

our results is in preparation.

Q-Quantum Mechanics

On the study which Chodos and I started of a deformation of quantum mechanics motivated from the type of deformations of commutators one encounters in quantum groups, a graduate student, H. Dietz, and I have been doing numerical calculations as well as exploring further analytical aspects.

**DATE
FILMED**

9 / 2 / 93

END