DOE/ER/40415-19

DOE/ER/40415--19

DE92 016972

MEDIUM ENERGY
NUCLEAR PHYSICS RESEARCH
UNIVERSITY OF MASSACHUSETTS

Progress Report
for the period June 1, 1991 through May 31, 1992

G.A. Peterson, J.F. Dubach, R.S. Hicks, and R.A. Miskimen
Department of Physics and Astronomy
University of Massachusetts
Amherst, MA 01003

June 1992

Prepared for

The U.S. Department of Energy
Under Grant No. DE-FG02-88ER40415.A006

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, expressed or implied, or assumes any legal liability of responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately-owned rights.

TABLE OF CONTENTS

Cover Page and Budget.....	i
Title Page	iv
Table of Contents	vi
I. Introduction	1
II. Experimental Work	7
A. Experiments at the Bates Linear Accelerator Center	7
1. Experiment 87-02: Threshold Electrodisintegration of the Deuteron at High Q^2	7
2. Measurement of the 5th Structure Function in Deuterium and ^{12}C	7
3. Single-Particle Densities of sd-Shell Nuclei	7
4. Experiment 84-28: Transverse Form Factors of ^{117}Sn	9
5. Experiment 82-11: Elastic Magnetic Electron Scattering from ^{13}C at $Q^2 = 1 \text{ GeV}^2/c^2$	10
6. Experiment 89-09: Measurement of the Elastic Magnetic Form Factor of ^3He at High Momentum Transfer	11
7. Experiment 89-15: Coincidence Measurement of the D(e,e'p) Cross-Section at Low Excitation Energy and High Momentum Transfer	12
8. Experiment 87-09: Measurement of the Quadrupole Contribution to the $N \rightarrow \Delta$ Excitation.....	14
B. Experiments at the Stanford Linear Accelerator Center	14
1. Experiment E-140: Measurement of the x-, Q^2 -, and A-Dependence of $R = \sigma_L/\sigma_T$	14
2. PEP Beam-Gas Event Analysis: Physics with the SLAC TPC/ 2γ Detector.....	15
a. Nucleon Resonance Electroproduction in Nuclei	17
b. Target Fragmentation in Deep Inelastic Scattering	20

C.	CEBAF Activities.....	23
1.	Drift Chamber Tests at Brookhaven National Laboratory	23
2.	Experiment PR-89-031: Multi-nucleon Knockout Using the CLAS Detector	24
3.	Electronics Design for the CLAS Region 1 Drift Chamber	24
4.	Color Transparencies in the Electroproduction of Nucleon Resonances	26
5.	Experiment PR-89-015: Study of Coincidence Reactions in the Dip and Delta-Resonance Regions	28
D.	NIKHEF-Lund Activities	29
1.	Trip Reports: Hermes and EEF	29
2.	Photoprotons from ^{10}B	34
III.	Theoretical Work	36
IV.	Publications, etc., from June 1, 1991 through May 31, 1992	38
V.	Nuclear Physics Seminars and Colloquia from June 1, 1991 through May 31, 1992	40
VI.	Comments on Scientific Personnel	42
A.	Postdoctoral Research Associates	42
B.	Faculty	42
C.	Graduate Students	43
VII.	University Related Matters	44.
VIII.	Nuclear Physics Computing Equipment	46
IX.	Comments on the 1993 Budget	47
A.	Faculty Summer Salaries	47.
B.	Postdoctoral Research Associates	48.
C.	Graduate Students	48
D.	^3He Target System Costs	48

E.	Equipment	49
F.	Visiting Russian Scientists	49

I. INTRODUCTION

The University of Massachusetts (UMass) Nuclear Physics Program continues to concentrate upon the use of the electromagnetic interaction in a joint experimental and theoretical approach to the study of nucleon and nuclear properties. During the past year the activities of the group involved data analysis, design and construction of equipment, planning for new experiments, and several short experimental runs. Our work related to the Bates Linear Accelerator Center, as described in part II.A. of this report, largely involved analyzing in fine detail the results of Experiment 87-02, the threshold electrodisintegration of the deuteron at high momentum transfers, and in preparing for Experiment 89-09, the measurement of the elastic magnetic form factor of ^3He . The latter experiment involves the measurement of the very small (10^{-40} cm²/sr) cross sections for elastic scattering at 160° as well as the threshold breakup region of ^3He . We plan to utilize the detector system developed by UMass for Experiment 87-02 on the deuteron breakup, as it has the capacity to measure very small cross sections in the presence of large backgrounds because of its excellent rejection capability. However, a high luminosity target must be developed for the ^3He . This cryogenic high-pressure target is a very complex undertaking for our group which requires state of the art techniques in machining, as well as very complicated engineering design that taxes the resources of a small university group such as ours. For design purposes we have available the expertise of John Mark, the cryogenic target expert at SLAC. This experiment is to be run in the Bates North Hall using the recirculated beam, but not the continuous beam from the South Hall Storage Ring which is under construction at this time. In fact, it is hoped that we will be ready for running when the South Hall Ring is being commissioned in 1993. Up to this time, there has been virtually no support for this experiment from Bates as they are putting all possible efforts into finishing the storage ring. As much work as possible is being done at UMass so that the target assembly, including a new target-changing mechanism, can simply be inserted into the Bates scattering chamber. This does not minimize the effort required at the time of

installation by the Bates staff with regard to the reinstallation of the helium refrigerator unit, new electrical wiring, plumbing, etc. However, UMass has much to do before that stage is reached.

Part II.B. describes our work related to experiments carried out at the Stanford Linear Accelerator Center (SLAC). Our group spent many person-months on site preparing for the E-140 Extension run that was postponed again and again because of beam priorities being given to the Stanford Linear Collider rather than to End Station A physics. Finally the run began in late August when we had teaching commitments as well as a commitment for a (γ, p) run at the Lund University microtron in Sweden. However, our major effort on SLAC-related work had to do with the analysis of so-called beam gas events which had occurred within the PEP storage ring and were detected in the Time Projection Chamber (TPC). Although we were not involved in the data acquisition which had taken place over the past ten years or so during the electron-positron colliding beam work by a collaboration of high-energy physicists, the data were available on summary tapes for our use. In fact the events that we were seeking, collisions of electrons or positrons with the residual gas atoms in the very good vacuum of the storage ring, were considered to be background events by the high-energy physicists. We thought that this work was worth pursuing for two central reasons: (1) The 15-GeV beam energy is considerably higher than is planned for CEBAF, and the data are unique in the entire world at this time, and (2) The experience gained in analyzing these final state multiparticle data acquired with a large detector having a solid angle close to 4π sr should prepare us for data to be received by the CLAS detector at CEBAF when it becomes operational in 1995. The type of physics involved in the TPC analysis, e.g., nucleon resonance excitation, is new to our group, and thus has led us in new directions. A major effort has been put forth in this analysis, as both of our postdoctoral research associates, a graduate student, two Russian students, two senior Russian visitors, and especially Professor Miskimen, have all been involved with this analysis.

Another project that has involved our group in a significant way is the design of the signal and high-voltage translator boards for the innermost drift chamber for the CLAS detector at CEBAF. This is described in section II.C. along with other CEBAF activities. During the past year considerable progress has been made with the translator board project. Since the first beams are expected in very late 1993 from the CEBAF accelerator, and the CLAS detector is not expected to be operational until 1995, and since we have approved experiments which involve the use of the CLAS detector, it is clear that no data will be acquired until at least 1995 on the two approved CEBAF experiments proposed by UMass. Notwithstanding, we are considering proposing another experiment involving color transparency in the electroproduction of nucleon resonances.

In our description of other activities involving collaboration with our colleagues at NIKHEF and Lund, as described in part II.D., we have included a trip report by Ross Hicks who is on sabbatical leave at NIKHEF. He attended meetings of the Hermes Collaboration in Heidelberg, and of the European Electron Facility collaboration at Orsay. These collaborations are planning to carry out several experiments that were similar to those planned by the defunct PEGASYS collaboration at SLAC. This report should be of interest to the D.O.E.

Thus during the past year we have not been involved in many experimental runs, but instead we are undertaking several construction and design projects. This was at least partially the result of there being few accelerators world-wide that were actually producing beams for nuclear physics research. This situation should change in a year or two. Nevertheless, in the meantime we have commitments for several construction projects that must be met. We therefore have dropped out of the SLAC program, at least for the near future, in order to meet our commitments. Also our group has a recollection of the five years devoted to the PEGASYS project at SLAC that came to naught.

Another notable item is described in section VI. The faculty in our group are now all tenured. Also there have been two promotions during the year so that now we have three full

professors and one associate professor. We have seven graduate students active in research activities at this time, and we may acquire one or two more.

In part VIII we describe how our computing facilities have changed in response to economic and scientific needs. In the past year we retired our VAX11/750 and are now using the more powerful VAX cluster at the Center for Computer Services with our own disk drive to be used exclusively by our group. Our old VAX was loaned to the Institute of Theoretical and Experimental Physics (ITEP) in Moscow, Russia, after considerable effort on the part of the principal investigator with U.S. and Russian agencies and airlines.

Our group has been actively sought to join several other collaborations. An example is the Hadronic Electroproduction Collaboration (HELP) at CERN's electron-positron collider LEP. This collaboration is led by Louis Dick who was involved in the defunct PEGASYS collaboration. The spin physics to be pursued and the experimental techniques to be utilized are similar to those outlined in the PEGASYS proposal. Similarly a PEGASYS-like collaboration for the TRISTAN electron-positron collider in Japan may be formed. Both LEP and TRISTAN have higher energy beams than are envisaged for either CEBAF, or for the European Electron Facility (EEF) discussed in part II.D.1., as LEP has 50-GeV beams and TRISTAN has 30-GeV beams. Clearly both facilities are suitable for working in the quark-scaling regime where many questions of great importance remain to be answered. However, the fate of the HELP collaboration is uncertain in the context of other LEP objectives, and similar remarks could be made about the possibility of ventures at TRISTAN.

Another possibility for us is to join Experiment E-781 at Fermilab. This is an approved experiment to study semi-exclusive reactions on nuclear targets with a 600 GeV/c Sigma-minus beam. Searches for exotics such as pentaquarks, hexaquarks, and H-dibaryons could be made, as well as measurements of Compton scattering polarizabilities, radiative transitions, and form factors

of Sigma hyperons. Among the members of the collaboration are our colleagues from Tel-Aviv University, who participated with us in many experiments at Bates and SLAC in past years.

The possibility of carrying out a measurement of pion polarizability has been considered by us during the past year. This is a measurement that bears on tests of the chiral symmetry property of the underlying QCD Lagrangian, as has been pointed out by our high-energy theoretical colleague Barry Holstein. This difficult study may be undertaken in several different types of experiments. Since it is impossible to make a target out of charged pions, the experiments must be performed off-shell, either by radiative pion-nucleus scattering using the Primikoff effect for Compton scattering from a photon in the intense Coulomb field of a heavy nucleus, or by pion photoproduction off of a virtual pion near a proton. The latter appears to be promising at CEBAF energies, and such a measurement is being planned at the LEGS facility at Brookhaven for 300 MeV laser back-scattered photons. A measurement was made at the Lebedev Physical Institute in Moscow several years ago, but a continuous spectrum of bremsstrahlung photons was used. The use of the tagged photon facility at CEBAF would eliminate many of the problems encountered by the Lebedev group. If this experiment were done at CEBAF using the CLAS detector, e.g., at $E_\gamma = 1$ GeV, an additional magnet would be necessary beyond CLAS for measuring the momentum of the far forward-going pions in the $\gamma p \rightarrow n\pi^+\gamma'$ measurement. It is thought that the n and π^+ could be detected, and not the γ' . Clearly this is a second-round CLAS experiment. We have been approached by Lebedev physicists to collaborate in a CEBAF measurement. Obviously a detailed Monte Carlo study must be made before undertaking such a measurement in which many background channels have been opened up at the higher CEBAF energies. We may proceed to seek special funds from the NSF for our Russian colleagues in order to carry out studies preliminary to a proposal. Since Professor Hicks of our group is on sabbatical leave at NIKHEF, and since he has made the most detailed studies of the possibility of pion polarizability

measurements, we will wait until he returns in September to make our decisions about these matters. Also Hicks may be aware of special opportunities at European facilities.

In the meantime we have commitments at Bates and CEBAF that we must honor. However, we realize that delays in the programs at either accelerator could deprive our graduate students of valuable experience at accelerator sites. We therefore are keeping our options open, and a necessary vagueness permeates our plans.

II. Experimental Work

A. Experiments at the Bates Linear Accelerator Center

1. Experiment 87-02: Threshold Electrodisintegration of the Deuteron at High Q^2

Data acquisition for this experiment was completed in the Spring of 1990, and a Physical Review Letter on the experiment was published in November 1991. The two students on this project, K.S. Lee of UMass, and W. Schmitt of M.I.T. are continuing to refine their data analysis and are starting to write their Ph.D. theses. We intend to write one more paper on this experiment, in which we will tie together the two analyses and report more fully on the detector, target, and data analysis aspects of the experiment.

2. Measurement of the 5th Structure Function in Deuterium and ^{12}C :

By detecting the recoil proton in an $(\vec{e}, e'p)$ experiment at an out-of-plane angle and by using a polarized electron beam, the 5th nuclear response function can be measured. The 5th response function is a measure of final state interactions, and vanishes in the absence of final state interactions. The 5th response function in deuterium and carbon in quasi-free kinematics was measured for the first time during an experiment at Bates in the summer of 1991. A. Hotta, X. Jiang, D. Olivieri, R. Miskimen, T. Moreau, and K. Wang participated in the data taking for this experiment in collaboration with groups from the Univ. of Illinois, Arizona State U., and M.I.T. The data are being analyzed by two students from the University of Illinois, the lead institution on this experiment.

3. Single-Particle Densities of sd-Shell Nuclei

For nuclei near the middle of the sd-shell, there exists little evidence for the abrupt filling of the $1d_{5/2}$, $2s_{1/2}$, and $1d_{3/2}$ subshells. Instead, these subshells all have partial occupancies, filling only gradually with increasing A . With its sensitivity to single-particle densities, elastic electron scattering represents a good means of learning about this progressive filling of the subshells, as

well as the radial sizes of the single-particle wave functions. Accordingly, extensive electron scattering measurements have been made at the University of Mainz and at the Bates Electron Accelerator of the elastic Coulomb and magnetic form factors of the $J = 5/2$ nucleus ^{27}Al . Much is also known regarding the Coulomb and magnetic form factors of other nuclei in this region. For example, similar data exists for ^{25}Mg , ^{29}Si and ^{31}P , and the Coulomb form factors have been determined for the $J = 0$ nuclei ^{28}Si , ^{30}S , and ^{32}P , for which magnetic scattering is forbidden.

Nuclear magnetism is largely governed by a relatively small number of nucleons in valence orbits, and hence magnetic scattering has obvious sensitivity to single-particle properties. For example, of the various magnetic multipole components possible in the sd-shell, the highest M5 multipole relies, to a good approximation, only on nucleons in $1d_{5/2}$ orbits. This dependence makes it possible to obtain a precise measure of the radial size of this orbit. On the other hand, because nucleons, predominantly protons, in *all* shell-model orbits contribute to Coulomb elastic scattering, the extraction of single-particle information from Coulomb form factors is less direct. By Fourier-transforming Coulomb monopole (CO) form factors, precise results have been obtained on ground-state charge distributions for nuclei throughout the entire periodic table. In principle, information on single-particle properties can be derived from *differences* between the charge distributions of neighboring nuclei. However, these differences also depend on core polarization and changes in nuclear size parameters, and uncertainties in these properties have impeded many previous interpretations of charge density differences in terms of single-particle densities. On the other hand, these effects appear to be less important for nuclei in the middle of the sd-shell, and because the single-particle orbit parameters are further constrained by the available magnetic scattering data, these nuclei are good candidates for instructive tests of our understanding of configuration mixing in valence shell-model orbits.

Analyses of CO form factors are normally confined to those nuclei having ground state spin $J = 0$ or $J = 1/2$. For $J \geq 1$ nuclei other even Coulomb multipoles enter, beginning with the

quadrupole C2 form factor. Because the cross sections measured in such cases are comprised of an incoherent sum of the various multipole components, the CO form factor is not isolated for transformation. One method of overcoming this difficulty would be to measure the additional form factors which characterize the scattering of polarized electrons from spin-aligned targets. Although facilities for such experiments are currently being developed, the relative complexity of these measurements will likely limit investigations to a relatively small number of cases. For the present, however, the only recourse is to use observed systematics and results from nuclear models to eliminate the contributions of the C2 and higher multipole components. It is this approach that we have taken to extract the charge distribution of ^{27}Al . The reliability of this procedure is supported by comparisons with cross sections measured for *inelastic* transitions where the higher multipoles are more evident.

In this way we have been able to deduce the $^{28}\text{Si} - ^{27}\text{Al}$ charge density difference, in addition to the $^{29}\text{Si} - ^{28}\text{Si}$ and $^{30}\text{Si} - ^{29}\text{Si}$ charge density differences largely based upon data acquired at Bates by UMass. A detailed study is being made to examine the extent to which these results can be interpreted in terms of single-particle densities. Specifically, together with the magnetic scattering data, these results are presently being compared with configuration-mixed calculations made in an expanded shell-model basis.

4. Experiment 84-28: Transverse Form Factors of ^{117}Sn

Several years ago, using the UMass-Bates 180° electron scattering facility, we made measurements of the elastic M1 form factor of the $J^\pi = 1/2^+$ nucleus, ^{117}Sn . Results from particle-transfer reactions^{1,2} show that the dominant configuration in the ^{117}Sn ground state has an unpaired neutron in the $3s_{1/2}$ orbit. Thus, ^{117}Sn might be useful for checking the interpretation of the strongly quenched M1 form factors, such as that³ of ^{205}Tl . *A priori*, it is not clear that similar core polarization (CP) effects would be expected. In ^{205}Tl , the $3s_{1/2}$ valence particle is a proton, and both the proton and neutron shells are almost full. On the other hand, while the ^{117}Sn proton

shell is closed, the valence $3s_{1/2}$ neutron lies midway between the $N = 50$ and $N = 82$ major shell closures. Our analysis of the ^{117}Sn results, therefore, combines $0\hbar\omega$ shell-model evaluations of the configuration mixing in the neutron shell, together with first-order CP calculations based on the same formalism and interaction as applied to ^{205}Tl .

In addition to the elastic M1 form factor of ^{117}Sn , form factors have been measured for transverse excitation of states at 0.159 MeV ($3/2+$), 0.317 MeV ($11/2-$), 0.712 MeV ($7/2+$), 1.01 MeV ($5/2+$, $3/2+$ doublet) and 1.18 MeV ($5/2+$). These transitions appear to have appreciable single-particle character, and therefore, provide additional tests of the suitability of the first-order CP treatment.

It is expected that this work will be completed this summer of 1992. We are awaiting the theoretical calculations of John Dubach in order to complete this long-standing project.

1. E.J. Schneid, A. Prakash, and B.L. Cohen, Phys. Rev. **156**, 1316 (1967).
2. C.R. Bingham and M.L. Halbert, Phys. Rev. C **1**, 294 (1970).
3. C.N. Papanicolas, L.S. Cardman, J.H. Heisenberg, O. Schwentker, T.E. Milliman, F.W. Hersman, R.S. Hicks, G.A. Peterson, J.S. McCarthy, J. Wise, and B. Frois, Phys. Rev. Lett. **58**, 2296 (1987).

5. Experiment 82-11: Elastic Magnetic Electron Scattering from ^{13}C at $Q^2 = 1 \text{ GeV}^2/c^2$

In the process of checking the ELSSY spectrometer detection system built to study the threshold electrodisintegration of the deuteron, and before taking data on a long-liquid deuterium target, data were taken on a compact solid ^{13}C target. The incident beam energy and scattering angle were 540 MeV and 150° corresponding to a three-momentum transfer of $q = 5.08 \text{ fm}^{-1}$ and a four-momentum transfer squared of $Q^2 = 1.01 \text{ GeV}^2/c^2$. *This is the highest Q^2 attained to date for an elastic electron scattering experiment on a nucleus with mass number $A > 4$.* These results were reported in Phys. Rev. C **44**, 1679 (1991).

6. Experiment 89-09: Measurement of the Elastic Magnetic Form Factor of ^3He at High Momentum Transfer

Because calculations involving few-body nuclei are relatively tractable, measurements on these nuclei can provide revealing tests of theoretical notions regarding effects such as meson exchange currents, three-body forces, relativistic dynamics, and possibly the transition at high momentum transfers from the nucleon-meson representation to quark-gluon degrees-of-freedom. In particular, the elastic magnetic form factor of ^3He is of great interest, since this is one of the very few cases in which the meson exchange contributions have been unambiguously identified. Theoretical calculations all lie in reasonable agreement with the existing data. However, beyond $q = 5.5 \text{ fm}^{-1}$, where the data stop, the various predictions begin to diverge. For example, the calculations by Schiavilla *et al.*¹ show a second diffraction minimum at $q = 6.6 \text{ fm}^{-1}$, followed by a maximum near $q = 7.5 \text{ fm}^{-1}$. In contrast, Sauer *et al.*² predict no second diffraction minimum below $q = 8 \text{ fm}^{-1}$, and Hadjimichael *et al.*³ predict a minimum. It is clear, therefore, that the availability of data beyond $q = 5.5 \text{ fm}^{-1}$ would be quite sensitive to the relative importance of the various effects included in the theoretical treatments. The expected cross sections are very small, and require a detector package that would be able to reject comparatively large backgrounds, such as the detector built by us for the deuteron threshold electrodisintegration experiment.

We have been awarded 480 hours of beam time to measure the elastic magnetic form factor of ^3He to $q = 7.4 \text{ fm}^{-1}$. This low counting-rate experiment will use the Bates high-resolution ELSSY spectrometer and the improved detector package mentioned above. The target will consist of a 17 cm long gas cell pressurized to 50 atm and cooled to 20 K. Under these conditions, the effective target thickness will be 0.53 g/cm^2 . Since the breakup threshold of ^3He is at 5.5 MeV, the energy resolution required to isolate the elastic peak can be easily achieved.

Because of the very expensive ^3He target material and exacting design criteria of the target system, we have engaged John Mark of SLAC to work on this project. Through his long

experience on similar projects at End Station A and elsewhere at SLAC, Mr. Mark is an acknowledged leader in the design of high-pressure cryogenic targets. It is estimated that the value assigned to the ^3He inventory needed for this experiment is of the order of \$2 million. We are planning to borrow the ^3He from the Savannah River Laboratory. Every possible means will be used to avoid the loss of any ^3He .

We are making progress towards constructing the main components of the target system. There are two distinct parts of this project: the target and gas recirculator loop inside the scattering chamber, and the gas-handling system and refrigerator on the outside. We will also build the target changing mechanism. Thus the complete target assembly will be built and tested at UMass. We should be able to install it at Bates such that it could be utilized quickly.

1. R. Schiavilla *et al.*, Phys. Rev. C **40**, 2294 (1989).
2. P. Sauer, private communications.
3. E. Hadjimichael *et al.*, Phys. Rev. C **27**, 831 (1983).

7. Experiment 89-15: Coincidence Measurement of the $D(e,e'p)$ Cross Section at Low Excitation Energy and High Momentum Transfer

Much of our research effort during the past few years has been devoted to the measurement of the electrodisintegration of the deuteron near breakup threshold, first at SLAC at coarse resolution, and later at Bates at a higher resolution. The cross section for this reaction is of particular interest because it represents one of the least ambiguous manifestations of the presence of one-pion exchange currents in nuclei. Nevertheless, present interpretations of these inclusive measurements still do not settle questions regarding the relative importance of Δ -isobar contributions, relativistic dynamics, and higher-order exchange currents. In order to resolve these outstanding issues, additional data are required.

In collaboration with Japanese physicists, led by T. Tamae of Tohoku University, and several physicists from Bates, we have an accepted experiment to measure the out-of-plane

coincidence (e,e'p) cross section for the deuteron at low excitation energy and high momentum transfer. This experiment will yield additional structure information in the form of the longitudinal-transverse interference cross section which, according to theoretical treatments, should strongly constrain calculations that consider the effects of meson exchange currents and relativistic dynamics. The enhanced sensitivity to relativistic effects makes this experiment different from other Bates coincidence experiments on the deuteron.

This experiment will require the continuous electron beam from the South Hall storage ring. Electrons scattered out of a special thin-window liquid-deuterium target cell will be detected in the OHIPS spectrometer in coincidence with protons in the MEPS spectrometer. Out-of-plane measurements will be made *without the need of the OOPS spectrometer system* because the kinematic boost near threshold breakup "compresses" the center-of-mass angle in the laboratory frame. The out-of-plane acceptances and angular resolutions of these two spectrometers are adequate for the extractions of the longitudinal-transverse interference cross section with good statistical precision. In addition, the transverse and longitudinal parts of the cross section will also be determined. We do not expect to be able to observe the transverse-transverse interference cross section, since this is predicted to be exceedingly small.

Data-taking for this experiment, approved for 432 hours of beam time, cannot begin before 1993, i.e., before the South Hall ring is completed. In the meantime, the collaboration has many preparations to make. These include the overhaul and reassembly of the T₂₀ target system and the installation of an additional drift chamber in OHIPS for determining ionization and recoil corrections. The responsibility for accomplishing these tasks will be divided between the various collaborating institutions. Professor Hotta of Shizuoka and other Japanese physicists will return to UMass this summer in order to help in planning for the execution of this experiment.

8. Experiment 87-09: Measurement of the Quadrupole Contribution to the $N \rightarrow \Delta$ Excitation

This experiment involves physicists from the University of Illinois (C.N. Papanicolas, spokesperson), Arizona State University, M.I.T., UMass, and the University of Mainz. The participant from UMass is R.A. Miskimen. The purpose of the experiment is to measure the quadrupole contribution of the $N \rightarrow \Delta^+(1232)$ transition by coincident electron scattering. The decay channel to be studied is the $H(e,e'p)\pi^0$ reaction. Protons will be detected simultaneously at four out-of-plane angles, and polarized beams will also be used to enable the detection of the fifth structure function. Components of the out-of-plane proton spectrometers (OOPS) are presently under construction at Illinois, Arizona State, and M.I.T.

B. Experiments at the Stanford Linear Accelerator Center

1. Experiment E-140E: Measurement of the x -, Q^2 -, and A -Dependence of $R = \sigma_L/\sigma_T$, the E-140 Extension

The ratio of the longitudinal to transverse cross sections, $R = \sigma_L/\sigma_T$, is one of the most useful parameters of deep inelastic lepton scattering. Its measurement requires exceedingly high precision. In experiment E-140, which was completed at SLAC in January of 1986, painstaking efforts were made to decrease systematic uncertainties. As a result, data of unprecedented precision were obtained on R for a range of target nuclei in the kinematic region $0.2 \leq x \leq 0.5$ and $1 \leq Q^2 \leq 10$ (GeV/c)². These data, already reported in several publications, ruled out any strong dependence on mass number A , as had been suggested as a possible origin for the EMC effect. Experiment E-140 was also the key to cross-normalization of previous SLAC measurements, and the normalization points of higher Q^2 measurements at other laboratories.

Due to inconsistent linac performance, we did not complete all the objectives of Experiment E-140. As a result, a proposal was made to extend the E-140 measurements. This propos: l

received the approval of the High-Energy Program Advisory Committee (EPAC), and was finally run in late August and all of September of 1991. The Nuclear Physics Injector, funded by the DOE in 1984, was used at normal 2 μ s pulse lengths. The microwave energy doubling system (SLED) with pulse lengths of ~ 0.1 μ s was also used.

In addition to using the SLAC 8 GeV/c spectrometer, as in the previous E-140 measurements, E-140 Extension utilized the 1.6 GeV/c spectrometer, which we upgraded for experiment NE11.

The primary emphasis of the extension of these studies was to determine the dependence of R on the square of the four-momentum transfer Q^2 at values of the Bjorken scaling parameter x , both lower and higher than was done in the initial measurements. In the quark-parton picture, the structure functions for deep inelastic scattering should depend only on x , and not on Q^2 . However, according to perturbative QCD theory, as Q^2 is increased, smaller and smaller parton sizes are probed, with the result that R is predicted to decrease with increasing Q^2 . This Q^2 -dependence of R can be broken down into several parts, one of which derives from the inability to resolve partons into their individual constituents at low Q^2 . Another source of the Q^2 dependence is attributed to the transverse momentum possessed by the struck quark, not only by virtue of its binding inside the nucleus, but also due to the emission of gluons in the scattering process. In addition, there are non-perturbative components of hadronic wave functions which can also give rise to transverse momentum effects, even at very large Q^2 . The analysis of the data is underway at SLAC.

2. PEP Beam-Gas Event Analysis: Physics with the SLAC TPC/ 2γ Detector

At the PEGASYS collaboration meeting of January, 1991, UMass physicists and Dr. Vladimir Gavrilov, from the Institute of Theoretical and Experimental Physics (ITEP) of Moscow, Russia, discussed the possibility of analyzing events in the Time Projection Chamber (TPC) resulting from 15 GeV electrons or positrons interacting with residual gas atoms in the PEP

Storage Ring. It was agreed that Dr. Sergey Shuvalov would come to SLAC in mid-April and stay for about three months in order to look into the feasibility of utilizing old data tapes with such events recorded upon them. UMass physicist Rory Miskimen was to make several trips to SLAC during this period to aid in the preliminary feasibility studies. If there were enough of these "background" events recorded in the e^+e^- colliding beam data summary tapes such that useful information could be extracted, then further analysis would be carried out at UMass, starting in mid-July, 1991. "Tagged events" in which a scattered electron and disintegration products are observed simultaneously outside of the e^+e^- collision region were to be analyzed, and there are even more possibly untagged events.

These plans materialized exactly as outlined. Dr. Sergey Shuvalov spent three months working at UMass, starting in mid-July, 1991. We have involved the two UMass postdoctoral research associates and one graduate student in the analysis. This summer we have Pavel Degtiarenko of ITEP at UMass for three months. Our attitude toward this project has been that it is worth a relatively small amount of effort to extract nuclear physics information from these unique data, already acquired for beam conditions beyond those which will be available at CEBAF, or elsewhere, for years to come.

The Russian physicists have experience with beam-gas event analysis at the ARGUS detector of the DESY laboratory in Hamburg, Germany. UMass physicists think of this analysis as being parallel to future analysis of multiparticle events in the CLAS detector at CEBAF. Thus, we can become acquainted with high-energy multiparticle analysis years before we will encounter similar CLAS data from CEBAF.

The analysis carried out thus far can be broken down into two parts: (a). Nucleon resonance electroproduction in nuclei, and (b). Target fragmentation in deep inelastic electron scattering.

(a). Nucleon resonance electroproduction in nuclei:

The analysis of data for nucleon resonance (N^*) electroproduction in nuclei has focused on two issues: (i) what can be learned from the N^* 's and (ii) what can be learned about medium modifications to the N^* 's. D. Olivieri will work on the analysis of the N^* data for his Ph.D. thesis. In N^* physics the two most topical issues concern the Roper resonance (P_{11}) and its non-observation in electroproduction experiments, and the so-called "missing resonances" which do not couple to the πN system. This is an area of physics that has received intense interest for many years and will form part of the core program for CEBAF. Nevertheless it is possible that because of the TPC's unique particle identification capabilities and acceptance, and because the data bridges the Q^2 gap between the photoproduction measurements and electroproduction experiments at higher Q^2 , that these data can shed some new light on the topics.

In contrast to N^* photoproduction physics, the field of resonance electroproduction in nuclei is wide open with virtually no published coincidence data. One area of interest is in how the N^* -nucleus interaction modifies the mass and width of the delta and higher mass resonances. This information can be used to help constrain models for the N^* -N interaction. Medium modifications are also predicted for p^0 production in the nucleus, with predicted mass shifts of approximately 100 MeV. It is generally believed that the phenomena of multi-nucleon emission is related to resonance absorption. Real photon experiments are involved with this problem, but there is almost no electroproduction data. Finally, Frankfurt, Miller, and Strikman have recently pointed out the importance of looking for color transparency in N^* electroproduction in nuclei as a test of quantum-mechanical models for color transparency. While the Q^2 range of the TPC data is certainly too low to reveal color transparency, these authors have suggested that for $Q^2 \approx 1 \text{ GeV}^2/c^2$, meson exchange currents and final state interactions may be suppressed.

The time-projection-chamber (TPC) is a nearly 4π solid angle colliding beam detector. In addition to tracking, the TPC determines particle identification by a measurement of particle

momentum and dE/dx . The TPC is completely surrounded by electromagnetic calorimeters, which can be used for π^0 identification, and by muon chambers. The " 2γ " spectrometers are positioned in the forward and backward directions along the beam line. Each 2γ spectrometer consists of a magnetic septum, drift chambers for tracking, time-of-flight and Cherenkov counters for particle identification, and NaI shower counters for energy measurement. They cover an angular range out to approximately 10 degrees from the beam axis. For the analysis reported here the 2γ spectrometers were used to detect scattered electrons or positrons and hence tag Q^2 and ν , and the TPC was used to detect hadrons and photons in the final state. The electron and positron beam energies were 14.5 GeV.

The TPC data consist of two types. A small amount of data was taken during a dedicated run on the TPC with injected gases, and a larger amount of data was the beam-gas events extracted from the TPC data tapes acquired over many years of electron-positron collision experiments. During the dedicated run data were taken on deuterium, argon and xenon, for a total of approximately 40,000 events. The trigger for this data was a charged track in the TPC with a good electron tag in the 2γ spectrometer. The beam gas data is a mixture of 30% H_2 and 70% CO by mass. There are approximately 100,000 events with an electron tag, and another 100,000 events without a tag. The trigger requirement for the tagged beam-gas data was two charged tracks in the TPC and a good electron tag.

Semi-exclusive data for the $A(e,e'\pi^-)$ A-1 reaction are shown in Figure 1, where the target is a mixture of carbon and oxygen. In these data the proton and π^- are required to carry all of the energy and momentum for the event within gates that are consistent with Fermi motion and detector resolution. No other charged track or calorimeter hits were allowed. The plot shows the $p\pi^-$ invariant mass distribution. The Δ^0 (1232) peak is clearly visible, as is resonant structure at higher energies. These data clearly demonstrate the feasibility of exclusive resonance electroproduction in nuclei.

FIG. 1

Semi-exclusive data for the $A(e,e'pp)A-2$ reaction are shown in Figure 2. Like the data in the previous figure, the two nucleons are required to carry all of the energy and momentum for the event and no other charged tracks or calorimeter hits were required. The plot shows the photon-nucleon invariant mass distribution for these events. The data show evidence for significant proton-proton emission in the resonance region, particularly near the Δ peak. Two proton emission in the resonance region has been studied in real photon experiments and the reaction is believed to proceed through meson exchange current mediated resonance absorption.

The ratio of pp to $p\pi^-$ yields are plotted in Figure 3 as a function of Q^2 . No acceptance corrections have been made to the data. Also shown in the figure is the result from an absorption calculation for the same ratio. In this calculation it is assumed that $\sigma_{abs} = 40$ mb and the percentage of pp absorption was adjusted to 17% to fit the data at low Q^2 . Corrections have not been made for absorption of the pp or $p\pi^-$ final state in the nuclear medium. The data show a relatively flat response as a function of Q^2 while the absorption calculation shows a rising fraction of pp to $p\pi^-$ yields. An increase of absorption with increasing Q^2 is expected because the resonance propagates further before decay, increasing the probability of absorption. The disagreement between the calculation and data could be interpreted as a decrease of FSI with increasing Q^2 , which is the prediction of Frankfurt, Miller and Strikman. However, because of the preliminary nature of the data, final interpretation must wait until acceptance corrections have been made and final state absorption estimated.

(b). Target Fragmentation in Deep Inelastic Electron Scattering

The largest part of the TPC data sample is in the deep inelastic region where $W > 2$ GeV. Because of the solenoidal nature of the TPC, forward going hadrons from the current fragmentation region go down the beam pipe and miss the detector. Generally the TPC is only able to detect hadrons that come out at wide angles or in the backward direction. These events typically have $x_f < 0$ and are called target fragmentation events.

FIG. 2

FIG. 3

While the physics of target fragmentation is non-perturbative and cannot be used to test QCD in a model-independent way, there is substantial interest in the phenomena. Our analysis will focus on the phenomenology of target fragmentation, and study issues such as the Q^2 and mass number dependence of the process. The TPC data should provide complementary information to streamer chamber results from Fermilab E665, which are at much higher energy loss v . It has also been suggested that the observation of backward going deltas or nucleons may be a signature for exotics such as 6 quark bags or two nucleon correlations. The data will be examined for events of these types.

C. CEBAF Activities

1. Drift Chamber Tests at Brookhaven National Laboratory:

During the summer and fall of 1990 several members of the University of Massachusetts group (S. Churchwell, R.S. Hicks, K.S. Lee, R.A. Miskimen, and K. Wang) participated with other CEBAF users and CEBAF staff in wire chamber studies at Brookhaven National Laboratory. The purpose of these measurements was to study how parallel and non-parallel magnetic field components influence the operation of a hexagonal cell drift chamber, such as will be used in the CLAS detector at CEBAF. Another purpose was to study how admixtures of helium in the chamber gas affect chamber operation. Helium in the chamber gas is useful because it reduces multiple scattering and because magnetic field effects are lessened due to the reduction of drift velocity. Data analysis is now complete and a paper has been submitted to IEEE, where it is presently under review. Based on the results of this study CEBAF is seriously considering the use of helium in the region 2 drift chamber for CLAS, and Bates is planning to use a helium-based gas mixture for the BLAST detector.

2. Experiment PR-89-031: Multi-nucleon Knockout Using the CLAS Detector

This proposal is a collaboration of physicists from UMass, CEBAF, the National Institute of Standards and Technology, and the University of New Hampshire. The spokespersons are R.A. Miskimen, J.W. Lightbody, Jr., and W. Hersman.

The goal of this collaborative proposal is to study the general phenomenology of multi-nucleon emission. To establish the systematics of this process, measurements will be made over a wide range of mass number. The proposed targets are ^3He , ^4He , ^{12}C , and ^{56}Fe . Studies would be made as a function of excitation energy ω , starting from the low excitation side of the quasielastic peak to above the delta resonance. Beyond this general survey of the $(e,e'NN)$ reaction, the proposal focused on four specific topics. These topics are three-body forces, nucleon-nucleon correlations, delta damping through the four nucleon emission process, and evidence for 6-quark bags. Beam time was also requested to study the feasibility of doing σ_L and σ_T separations by the Rosenbluth method.

In 1989 the CEBAF PAC gave conditional approval to this experiment. Full approval was withheld until the spokespersons agreed with the proponents of five other experiments on a common run plan. In June 1991 a workshop was held at CEBAF on the physics that could be studied in a program of $A(e,e'x)$ experiments on nuclear targets in the multi-GeV energy range. Both experimentalists and theorists were invited to the meeting. At the workshop a common run plan for the CLAS electronuclear program was agreed upon, and at the November 1991 PAC meeting experiment PR 89-31 was given final approval. The PAC awarded the electronuclear experiments 33 days of run time in Hall B, with 20% of the time at 0.8 GeV, and 50% at 4.0 GeV.

3. Electronics Design for the CLAS Region 1 Drift Chamber:

The UMass group is responsible for the chamber mounted electronics and signal cables for the CLAS region 1 drift chamber, which is the innermost drift chamber for the CLAS. The

University of Pittsburgh and Carnegie Mellon University are responsible for the construction of the chamber itself. The region 1 drift chamber will contain over 7,000 signal wires.

Our group is designing the signal translator boards and the high voltage (HV) translator boards which must fit in the approximately 1 inch gap between the end-plates of the chamber. The space constraints in the region 1 chamber combined with the mixed high and low voltages that these boards must provide, makes the region 1 electronics one of the most difficult electronics design problems for the CLAS drift chambers. The signal translator boards make the connection with the signal wires and provide signal amplification for driving twisted pair cables to the front-end electronics where the TDC's are located. For a pre-amplifier we are planning to use an amplifier packaged as a single-in-line package (SIP) that has been designed by CEBAF engineers. The signal translator board contains a mix of high voltage and low voltage, which makes high voltage breakdown a great concern. The HV translator boards make the connection with the sense, field, and guard wires and provide the HV biasing. The HV segmentation of the chamber is provided for in the HV translator board.

During the past year we have made substantial progress with the electronics project. We recently moved into a laboratory that will be used specifically for this project. A prototype drift chamber that will be used for testing electronics and cables has been constructed and successfully tested. Our group recently completed the design of its first electronics boards using the CAD system purchased through DOE funds in 1991, and we are waiting for the stuffed boards to be sent to UMass for testing. We are proceeding with the design of the actual region 1 boards and are planning the details of signal and HV segmentation on the circuit board level. The next major milestone as specified in our MOU with CEBAF calls for us to provide the first full size prototype boards to the University of Pittsburgh and C.M.U. groups by the end of the summer.

4. Color Transparency in the Electroproduction of Nucleon Resonances:

In high Q^2 exclusive reactions color transparency predicts that a hadron will have reduced final state interactions (FSI) with the nuclear medium. One measure of FSI suppression is a ratio called the transparency, which is defined as

$$T = \frac{\sigma_A}{A\sigma_N},$$

where σ_A is the nuclear cross section, σ_N is the cross section on the nucleon, and A is the mass number. In general, DWBA calculations (Glauber model calculations) predict transparencies that are significantly less than one and nearly independent of Q^2 . Color transparency predicts that at very high Q^2 the transparency should approach one. In this limit the medium has become "transparent" to the final state hadron, and the FSI have vanished.

In contrast to previous theoretical work in color transparency which was largely semiclassical, theorists are now constructing quantum models for color transparency. In these models the small "object" that is produced in the high Q^2 reaction is a superposition of hadron excited states. The initial state wave function would then be written as

$$|\psi_i\rangle = \sum_h a_h |h\rangle,$$

where a_h are constants and the sum is over hadron excited states. The wave function then evolves with time, and the overlap of the wave function with the hadron ground state or excited state can be calculated. Nucleon resonance (N^*) electroproduction data are urgently needed to test diffractive models of color transparency.

We are preparing a letter-of-intent that will use nucleon resonance electroproduction in nuclei to study color transparency. Calculations suggest that a CEBAF experiment would be in a kinematic range where color transparency would be apparent in oscillating transparency and in the suppression of meson exchange currents. Quantum models of color transparency suggest that weak states (Roper at $Q^2 > 0$ or $\Delta(1232)$ at high Q^2) might increase their strength in the nucleus

relative to other N^* 's, and strong states (S_{11} at high Q^2) might get filtered out. Transparency in quasi-free (e,e'p) might oscillate and be out of phase with transparency in resonance production. The proposed experiment would provide a precision test of quantum models of color transparency that are being developed.

We propose to use the CEBAF CLAS detector with the maximum available beam energy to measure N^* electroproduction in nuclei and the quasi-free (e,e'p) reaction at high Q^2 . $\Delta(1232)$ production and decay will be observed through $p\pi^-$ and $p\pi^0$ final states, and the $S_{11}(1535)$ resonance through the $p\eta$ final state. Direct evidence for final state interactions in the resonance region will be observed through emission of pp and $p\pi^+$ final states. Models for color transparency can be tested by making comparisons of data with different target mass at fixed Q^2 , and by measuring resonance absorption and decay yields as a function of Q^2 . The target required for this measurement are ^2H , ^4He , ^{12}C , and ^{56}Fe . In approximately 400 hours of running at 4.0 GeV incident beam energy, a statistically significant measurement could be made for Q^2 as large as $5 \text{ GeV}^2/c^2$.

No significant modifications of the CLAS detector are required for this experiment. The wide angle calorimeter will be required for electron, π^0 , and η detection. Additional wide angle calorimetry would be highly desirable to increase the acceptance of π^0 and η . A modification to the target holder so that two or more targets could be mounted in the detector would allow the beam to pass through all of them simultaneously. This would be useful in limiting systematic errors, and would permit running this experiment in conjunction with several other experiments. These are the multi-hadron experiments (PR-89-015, -017, -027, -031, -032, and -036) which are approved for 16.5 days of running on ^4He , ^{12}C , and ^{56}Fe at 4 GeV incident energy.

Tentative plans call for submitting a letter of intent to CEBAF to be followed by a full-scale proposal.

5. Experiment PR-89-015: Study of Coincidence Reactions in the Dip and Delta-Resonance Regions

The purpose of this experiment is to do a systematic study of the excitation energy ω and momentum transfer q dependences of the cross sections for coincidence reactions with one or several hadrons in the final state, e.g., (e,N) , $(e,e'NN)$, $(ee'N\pi)$, etc., in the dip and delta resonance regions for several nuclear targets (e.g., ^3He , ^{12}C , ^{40}Ca , and ^{208}Pb). Results of the intermediate energy single arm (e,e') and coincidence $(e,e'p)$ experiments in these regions indicate the importance of the multinucleon processes and the role of $\Delta(1232)$ resonance in the reaction. The goals of the experiment are: (1) to isolate the different multi-nucleon processes and investigate their ω and q dependence; and (2) to study the $\Delta(1232)$ production and its interaction with nucleons. The momentum transfer dependence of the $(e,e'N\pi)$ reaction will be measured to investigate the possible modification of the delta and its magnetic form factor in the nuclear medium. This program will be extended to study reactions with tagged photons.

There is growing evidence to suggest that multi-nucleon processes in inelastic electron scattering may be more important than previously anticipated. Moreover, little is known about the role of Δ -isobars in such interactions. Only by coincidence measurements made by utilizing large-acceptance detectors can one hope to identify and learn about the most important multi-nucleon mechanisms. Together with the extended energy range of the CEBAF accelerator, the particle-identification properties and the almost- 4π acceptance of the CLAS detector will provide unprecedented opportunities to probe these reactions.

This experiment was proposed by H. Baghaei while he was a postdoctoral research associate at UMass. In September of 1990, he moved to the University of Virginia.

D. NIKHEF-Lund Activities

1. Trip Reports: Hermes and EEF

Professor Ross Hicks is spending his sabbatical stay in Amsterdam from January through August of this year. Hicks has attended workshop meetings for two European initiatives during his stay. Neither of these meetings was attended at D.O.E. expense. The first was the January 16-17 meeting of the HERMES collaboration in Heidelberg. The second was February 10-14 workshop at Orsay of European physicists seeking to develop a proposal for the construction of a 15-GeV European Electron Facility (EEF). A report on these meetings follows.

The HERMES collaboration has submitted a proposal to use the HERA ring to measure the spin-dependent structure functions of the proton and neutron with longitudinally-polarized electrons having energies up to 35 GeV. Present knowledge of these spin-dependent structure functions is poor; only g_1^p has been measured, and even these data are relatively imprecise. Nevertheless, the results are exceedingly provocative. One interpretation, which relies upon SU(3) flavor symmetry and the application of two sum rules (Bjorken and Ellis-Jaffe), is that quarks carry only about 12% of the nucleon intrinsic spin. Furthermore, the contribution of s quarks is far from negligible, and this interferes destructively with the spin contributions of u and d quarks. It is the objective of the HERMES collaboration to test this contentious interpretation by making extensive and precise new measurements using internal polarized gas-jet targets of hydrogen, deuterium, and ^3He , in conjunction with a large-aperture, forward magnetic spectrometer. In addition to measurement with polarized beam, HERMES is now considering the possibility of performing experiments with beams of unpolarized electrons. In principle, these could be performed in parasitic mode, without impairing the HERA high-energy program, and hence several months of each year will be available for such work.

About 50 collaboration members were present for the HERMES meeting at Heidelberg. The collaboration is rather broad, as indicated by the following list of institutions.

Experimental Component	Institute
Electromagnetic calorimeter	LAMPF, New Mexico, Cal. Tech, Illinois. Additional possibilities include Frascati, Yerevan, Mainz and NIKHEF.
Transition Radiation Detection	TRIUMF, Univ. of Alberta.
Scintillator hodoscope	Cal. Tech.
Wire chambers	Heidelberg, Zeuthen, Erlangen, St Petersburg, Argonne, MIT, and possibly TRIUMF.
Cerenkov detector	Argonne.
Magnet	St. Petersburg, Sanita.
Silicon strip detectors	Torino, Zeuthen.
Data acquisition system	?
Electron polarimeter	?
Polarized targets	Wisconsin, MIT, Cal. Tech., Argonne, and Heidelberg.

Perhaps the most important issue to emerge from this meeting relates to the qualified approval given to this project by the HERA program advisory committee, which feels that HERMES should be given full approval only after HERA achieves the 80% beam polarization sought in the HERMES proposal. Many of the collaborators, particularly those from the U.S., argued that such conditional approval jeopardized their chances of winning from U.S. funding agencies the support required to develop the facilities for which they were responsible. These groups requested that the collaboration's spokespersons approach the HERA management with the view of obtaining a more affirmative statement of support for HERMES.

Irrespective of this dilemma and the intrinsic merit of HERMES, it was clear that there still remained much to be done if the experimental objectives of HERMES are to be realized. Almost

every aspect of the spectrometer needed attention. For example, flaws had been found in the previous magnet design, and the momentum resolution and background capabilities of the various detector components were still to be resolved. Although much effort is being invested into correcting these deficiencies, it was felt that a judicious expansion of the collaboration would greatly promote chances of success. Other news was more reassuring. In particular, good progress has been made in the development of polarized targets for HERMES. To a large extent, this development was primarily motivated by the need to provide polarized targets for experiments at other laboratories. How these targets would perform in the HERA environment remains something of an open question.

Looking beyond HERMES, European scientists are pressing forward with a new initiative for the construction of EEF, a 15-30 GeV European electron accelerator capable of delivering continuous beam currents in excess of $10\ \mu\text{A}$. It is intended that at 15 GeV the momentum resolution of the beam should be no worse than 3×10^{-4} , and that beams having $>80\%$ polarization be available. The steering committee for this new facility has recently invited the submission of theoretical papers to motivate this project, as well as proposals for specific experiments and detectors. These contributions will be reviewed at Mainz, in early October.

The 50-60 attendees of the Orsay workshop were given a detailed account of the progress that has been made thus far. In particular, the subcommittee responsible for the detector design has decided that the two-linac racetrack CEBAF design will not be appropriate for the higher-energy EEF. Instead, a one-linac, multipass racetrack configuration will likely be adopted. Bonin from Saclay reviewed the progress that has been made on superconducting microwave cavities during recent years. Field gradients of 10 MV/m can be routinely achieved, with surface defects limiting the realization of larger gradients. In principle, 1000 MV/m should be possible, before the onset of the quantum tunnelling that leads to copious electron emission from the cavity walls! Combined, the Saclay and Orsay institutions have about 20 persons working on the development of

microwave cavities, by far the largest group in the world. For EEF, the best unit appears to have 9 cells, and will operate at 1.3 GHz and 1.8 K. Coincidentally, these parameters are identical to those specified by the TESLA collaboration for a new collider facility proposed for construction at DESY. In fact, an agreement has been established to work with DESY to develop the new cavity design.

Compared to the accelerator properties, the ideas regarding the detectors at the new facility are much more preliminary. Nevertheless, conceptual designs were presented for a novel large acceptance detector, as well as more conventional high-resolution magnetic spectrometers. Pierre Bertin from Clermont-Ferrand discussed a possible rationale for the design of the large acceptance detector. At EEF, one would seek to measure not only the hadronization of the quark struck in Deep Inelastic Scattering (DIS), but also the fragmentation functions of diquark target remnants. The Lund String model and Monte-Carlo program can be used to reliably predict the distributions in angle and momentum of the produced hadrons. In particular, it will be useful to anticipate the extent of jet formation with 15 GeV and 30 GeV electron beams.

Saclay's Fournier proposed some design objectives for a large-acceptance spectrometer capable of making such measurements. The angular acceptance should be large, extending from a few degrees to almost 180° . The momenta of charged particles would need to be measured to at least 1%, with an angular precision of better than 1 mr. Precise vertex reconstruction would be required for extended targets. Particle identification should be such that γ -rays, electrons, muons, pions, kaons, and protons can be resolved. The ability to accommodate large luminosities was considered essential, as is the requirement of rejecting backgrounds from such processes as Møller reactions, bremsstrahlung, and secondary photoproduction. Fournier advocated that good background rejection would rely upon a careful design of the detector vacuum chamber, a lack of material in the bending plane of the spectrometer, and the existence of relatively weak trapping fields to confine low-energy background. The task of designing suitable event triggers should not

be underestimated. Fournier presented a novel design for a large acceptance detector, in which the magnetic field derives from a superconducting solenoid, oriented not with its axis along the incoming horizontal beam direction, but vertically. The target is located in an opening at the side of the solenoid, and the momenta of the reaction products are determined by spatial measurements made in banks of wire chambers situated within a 3 T magnetic field. Particle identification is realized by means of dE/dx and time-of-flight measurements, as well as by measurements of shower development in surrounding calorimeters.

Blomqvist presented a preliminary design for a high-resolution 15 GeV/c electron spectrometer of QQDD design. Large-bore, superconducting elements would provide acceptances of 5.4 msr in solid-angle and 10% in momentum. The angular and momentum resolutions would be ≤ 0.5 mr and 0.05%, respectively. Overall, the spectrometer would be 48 m long, about the limit for a movable spectrometer. A spectrometer of comparable specifications for a 30 GeV beam would need to be fixed, and the scattering angle varied by changing the entry angle of the incident electron beam. Synchrotron losses may ultimately limit the performance of such spectrometers. The average energy loss in the spectrometer described by Blomqvist is 8.5 MeV. Monte Carlo calculations are needed to evaluate how the dispersion in this loss impairs the resolution.

The range of theoretical concepts explored at the workshop were primarily concerned with unpolarized electron beams. These topics included hadronization, color transparency, charm production, exclusive virtual Compton scattering, DIS tagged on the observation of slow hadrons emitted at backward angles, and diquark models for strangeness production. The choice of these theoretical subjects underlined the influence of the PEGASYS proposal in motivating the EEF initiative.

The EEF steering committee is chaired by Thomas Walcher (Germany) and includes Jacques Arvieux (France), Enzo de Sanctis (Italy), and Peter de Witt Huberts (Netherlands). The present strategy of the EEF collaboration is to develop arguments that would persuade European

governments of the essential merit of such a project, and to further demonstrate the large number of European scientists who would be active in the endeavor. Indeed, commitments from U.S. and other international scientists are not being actively sought at present; the EEF collaboration wishes to establish, first and foremost, the large number of European scientists dedicated to the project. It is hoped that at least 300 European nuclear scientists would commit substantial research efforts to EEF. These scientists would come mainly from Germany (≈ 150), France (≈ 100), The Netherlands (≈ 50), and Italy (≈ 50). Nevertheless, as one member of the EEF steering committee disclosed, any spontaneous promise of support from U.S. nuclear scientists would be very warmly welcomed. At this early stage, no reliable cost estimates have been made for the project. According to one estimate, it is foreseen that the ultimate request would be in the range \$0.3-0.6B. At first sight, this might appear to be a considerable underestimate, especially since it is of the same order as the cost of CEBAF, a 4 GeV facility. However, it is hoped that the additional expense of a higher-energy EEF beam would be largely offset by the requirement of a smaller beam current. It is envisaged that thick targets can be utilized to maintain good experimental luminosities.

2. Photoprotons from ^{10}B

A short experiment on ^{10}B , requiring only two runs of two weeks each, including detector installation, and beam time both for calibrations as well as measurements, took place in March and August-September of 1991 at the Lund University Microtron in Lund, Sweden. This 100% duty factor accelerator has a tagged photon facility that produced photons of energies 60 and 80 MeV, which were ideal for this photoproton experiment. These measurements will serve as an extension of recent UMass (e,e') and (e,e'p) measurements on ^{10}B previously made at Bates and NIKHEF as a test of the consistency of interpreting these two reactions in terms of a single nucleon wave function. The accurate nuclear structure information obtained from the combined analysis of (e,e') and (e,e'p) data places a constraint on the quasi-free knockout contribution to the $^{10}\text{B}(\gamma,p)$ cross section. Hence, it will be possible to extract the relative importance of two-nucleon processes

(such as those calculated by the Modified Quasi Deuteron model) compared to quasi-free knockout in the (γ, p) reaction.

The class of (γ, p) reactions we seek to study are those in which the knocked-out proton has much more momentum than the incident real photon. In the simplest interpretation, such protons must have been in high-momentum components of the initial state wave function. The cross sections for such processes are, therefore, expected to be sensitive to meson exchange currents and short-range correlations between nucleons. In principle, the sensitivity of the (γ, p) reaction to high-momentum components should complement results from electro-induced reactions. By selecting (γ, p) kinematics to be very close to those of the $(e, e'p)$ experiment, the $(e, e'p)$ data will determine the quasi-free knockout contribution to the (γ, p) reaction. Any differences from the quasi-free knockout description may then point to the need for additional high-momentum components in the nuclear wave function due to N-N correlations.

In parallel with the experimental effort, graduate student Chris Arabadjis is working with Professor Dubach to improve the theoretical description of $(e, e'p)$ reactions.

III. Theoretical Work

Once again the focus of the theoretical work this year has been on weak and electromagnetic interactions in a variety of nuclear processes. Partly in support of this effort, but for its own right as well, we are also working on some aspects of nuclear structure.

One theoretical graduate student, G. Feldman, completed his Ph.D. thesis in January. This thesis discussed weak interactions in the purely hadronic sector and consisted of two parts, the construction of an effective weak parity-violating potential for all nucleon and Δ channels and then application of a similarly-constructed potential for nucleon and Λ channels to the decay of hypernuclei. In the first part of this work he expanded upon a technique based on an $SU(6)_w$ symmetric quark model (originally developed by Desplanques, Donoghue, and Holstein to treat weak interactions in the nucleon-nucleon sector) to construct weak parity-violating nucleon-nucleon-meson, nucleon-delta-meson, and delta-delta-meson vertices. With these vertices, a weak parity-violating potential can be constructed for all nucleon and Δ channels with the exchange of π , ρ , and ω mesons. This work has recently been published in Physical Review C.

We are proceeding to use this weak potential in an analysis of elastic polarized proton-proton scattering, $\vec{p} + p \rightarrow p + \Lambda p$, at intermediate energies where the Δ intermediate states play an important role. While such an analysis would have been the logical second part to Feldman's thesis, this was judged inappropriate since the necessary calculations would be lengthy, technically tedious, and not particularly instructive for a graduate student. Instead, he applied a similar $SU(6)$ -based meson-exchange weak potential in the N - Λ sector (developed earlier by UMass student L. de la Torre) to the simpler case of hypernuclear decay in a Fermi-gas model. In earlier work we had discovered that the addition of K exchange to π exchange in the weak $\Lambda N \rightarrow NN$ potential made little difference to the total hypernuclear decay rate but dramatically modified the predicted ratio of the partial width for decay into np channels to that for decay into pp channels. Feldman explored the K exchange dependence of other observables, particularly

the up-down asymmetry of emitted protons, where he found significant differences between π only and $\pi + K$ exchange models. A manuscript on these results has been drafted and is now being polished for publication.

Graduate student C. Arabadjis continues to make progress on his thesis project, a merging of our large basis shell-model calculations with "standard" calculations of the (e,e'p) reaction. He has recently completed the (p,p') portion of the project (which turned out for a variety of reasons to be surprisingly time-consuming) and is now engaged in putting the various pieces of the project together. We anticipate a completed thesis within a year.

A third graduate student, J. Gallant, has embarked on his thesis work. We have developed a technique which we believe to be novel, for coupling core-polarization techniques with our large-basis shell model codes. He is currently in the process of coding and debugging the programs necessary to implement this technique. Once this is completed, we will be in position to apply this technique to a variety of problems including testing certain assumptions made in previous core polarization calculations, exploring the construction of effective operators for different shell model basis-truncation schemes, and studying a variety of interesting electromagnetic and weak transitions in nuclei.

A review article is being prepared for Physics Reports, in collaboration with T.W. Donnelly, M. Musolf, S. Pollock, E. Beise, and S. Kowalski, on the subject of semi-leptonic tests of the structure of weak neutral currents. This is a topic which has received considerable attention in the last year (particularly in regard to potential CEBAF experiments) and the review article has expanded accordingly. We expect to come to closure on this article by mid-July.

Finally, we have completed an extensive set of shell-model calculations of ^{27}Al , ^{28}Si , ^{29}Si , and ^{30}Si in support of an analysis of electron scattering data being undertaken by Prof. Hicks. The arduous computer task of including meson-exchange current contributions to this analysis will be attempted later this summer.

IV. Publications and Presentations by Members of the University of Massachusetts Medium Energy Physics Group in the Period June 1, 1991 to May 31, 1992

A. Journal Publications:

1. "Elastic Magnetic Electron Scattering from ^{13}C at $Q^2 = 1 \text{ GeV}^2/c^2$," R.A. Miskimen, H. Baghaei, M. Frodyma, R.S. Hicks, K.S. Lee, G.A. Peterson, S.H. Rokni, P.E. Bosted, S.E. Rock, K.A. Dow, W. Turchinets, J.D. Zumbro, B. Frois, J. Martino, S. Platchkov, A. Hotta, and R.L. Huffman, *Phys. Rev. C* **44**, 1679 (1991).
2. "Measurements of Deuteron Threshold Electrodissintegration: A Probe of Short-Range Meson Exchange," K.S. Lee, W. Schmitt, H. Baghaei, D. Beck, P.E. Bosted, S. Churchwell, B. Frois, R. S. Hicks, A. Hotta, J. Martino, R.A. Miskimen, G.A. Peterson, S. Platchkov, M. Spengos, W. Turchinets, K. Wang, C.F. Williamson, T. Yates, and J.D. Zumbro, *Phys. Rev. Lett.* **67**, 2634 (1991).
3. "Electron-Nucleus Interactions," G.A. Peterson, *Nucl. Phys. A* **527**, 367c (1991).

B. Articles Accepted for Publication: None

C. Articles Submitted for Publication:

1. "Systematic Approximations for the Bethe-Salpeter Equation," G.A. Crawford, Submitted to *Phys. Rev. C*.
2. "Effects of Non-Parallel Magnetic Fields on Hexagonal Cell Drift Chambers," M.D. Mestayer, C.L. Tam, K. Wang, H. Baghaei, S. Christo, S.A. Dytman, G.P. Gilfoyle, J.D. Hewitt, F.W. Hersman, R.S. Hicks, R.A. Miskimen, R. Schumacher, and M.F. Vineyard, Submitted to *IEEE Transactions on Nuclear Science*.
3. "Elastic Magnetic Electron Scattering from ^{89}Y and Finite Fermi System Theory," J.E. Wise, F.W. Hersman, J.H. Heisenberg, T.E. Milliman, J.P. Connelly, J.R. Calarco, W. Kim, R.S. Hicks, R.A. Miskimen, G.A. Peterson, R.P. Singhal, S.A. Fayans, A.P. Platonov, and E.E. Saperstein, Submitted to *Phys. Rev. C*.
4. "Measurements of the Electric and Magnetic Form Factors of the Proton from $Q^2 = 1.75$ to 8.83 (GeV/c)^2 ," P.E. Bosted, L. Clogher, A. Lung, L. Stuart, J. Alster, R.G. Arnold, C.C. Chang, F.S. Dietrich, R. Gearhart, J. Gomez, K. Griffioen, R.S. Hicks, C. Hyde-Wright, C. Keppel, S. Kuhn, J. Lichtenstadt, R.A. Miskimen, G.A. Peterson, G. Petratos, S. Rock, S. Rokni, W. Sakumoto, M. Spengos, K. Swartz, Z. Szalata, L.H. Tao, and W. Dodge, Submitted to *Phys. Rev. Lett.*

D. Articles Soon to be Submitted for Publication:

1. "The Elastic M1 Form Factor of ^{117}Sn ," R.S. Hicks, H. Baghaei, R.A. Miskimen, G.A. Peterson, S.H. Rokni, J. Dubach, A. Hotta, and T. Suzuki, To be submitted to *Phys. Rev. C*.

E. Articles in Preparation:

1. "Weak Non-Mesonic Decay of Hypernuclei," L. de la Torre, J.F. Donoghue, J. Dubach, B.R. Holstein, and M. Kimura, To be submitted to *Phys. Rev.*
2. "Probing the Hadronic Neutral Current with Electron and Neutrino Scattering," M.J. Musolf, T.W. Donnelly, J. Dubach, S.J. Pollock, S. Kowalski, and E.J. Beise, Review Article.

F. Published Invited Conference Talks: None

G. Invited Talks to be Published in Conference or Workshop Proceedings:

1. "Measurements of Deuteron Electrodissintegration: A Probe of Short-Range Meson Exchange," Bull. Am. Phys. Soc. **37**, 973 (1992).

H. Abstracts of Contributed Papers:

1. "A Precision Measurement of the Neutron Elastic Electromagnetic Form Factors at High Momentum Transfers," L.M. Stuart, R.G. Arnold, P.E. Bosted, L. Clogher, C. Keppel, A. Lung, S. Rock, M. Spengos, Z.M. Szalata, L.H. Tao, J. Gomez, F. Dietrich, C.C. Chang, G.A. Peterson, R.S. Hicks, R.A. Miskimen, W. Dodge, K. Griffioen, G. Petratos, W. Sakumoto, S. Rokni, R. Gearhart, S. Kuhn, J. Lichtenstadt, J. Alster, C. Hyde-Wright, and K. Swartz, Bull. Am. Phys. Soc. **36**, 2125 (1991).
2. "Precision Measurement of the Neutron Electric and Magnetic Form Factors at High Momentum Transfers," A. Lung, R.G. Arnold, P. Bosted, L. Clogher, C. Keppel, S. Rock, M. Spengos, Z. Szalata, L.H. Tao, J. Gomez, L. Stuart, F.S. Dietrich, G. Chang, G. A. Peterson, R.S. Hicks, R.A. Miskimen, W. Dodge, K. Griffioen, G. Petratos, W. Sakumoto, S. Rokni, R. Gearhart, S. Kuhn, J. Lichtenstadt, J. Alster, C. Hyde-Wright, and K. Swartz, Bull. Am. Phys. Soc. **37**, 884 (1992).
3. "A Design of an out-of-plane Spectrometer Support System for Studying (e,e'x) Reactions," S. Dolfini, L. Bartosek, R. Beck, J. Harrop, R. Laszewski, J. Mandeville, E. Offerman, C. Papanicolas, S. Williamson, A. Bernstein, W. Bertozzi, V. Boeglin, V. Bhushan, D. Dale, S. Gilad, M. Holtrop, D. Jordan, J. Kelsey, S. Kowalski, T. McIlvain, L. Weinstein, K. Dow, G. Dodson, M. Farkhondeh, W. Sapp, D. Tieger, C. Tschalaer, W. Turchinets, J. Zumbro, R. Alarcon, J. Comfort, J. Gorden, D. Martinez, M. Epstein, D. Margaziotis, R.A. Miskimen, D. Mack, and S. Wood, Bull. Am. Phys. Soc. **37**, 885 (1992).
4. "Performance of an out-of-plane Spectrometer for Studying (e,e'x) Reactions," J. Mandeville, R. Beck, S. Dolfini, J. Harrop, R. Laszewski, E. Offerman, C. Papanicolas, S. Williamson, A. Bernstein, W. Bertozzi, V. Bhushan, W. Boeglin, D. Dale, S. Gilad, M. Holtrop, D. Jordan, S. Kowalski, T. McIlvain, L. Weinstein, K. Dow, G. Dodson, M. Farkhondeh, W. Sapp, D. Tieger, C. Tschalaer, W. Turchinets, J. Zumbro, R. Alarcon, J. Comfort, J. Gorgen, D. Martinez, M. Epstein, D. Margaziotis, R.A. Miskimen, S. Wood, and D. Mack, Bull. Am. Phys. Soc. **37**, 885 (1992).
5. "Electroproduction and Absorption of Nucleon Resonances in Nuclei," R.A. Miskimen, J. Button-Shafer, L. Elouadrhiri, D. Olivieri, G.A. Peterson, K. Wang, V. Gavrilov, S. Shuvalov, F. Dietrich, and K. Van Bibber, Bull. Am. Phys. Soc. **37**, 886 (1992).
6. "Target Fragmentation in Deep Inelastic Scattering of 14.5 GeV Electrons from Nuclei," K. Wang, J. Button-Shafer, L. Elouadrhiri, R.A. Miskimen, D. Olivieri, G.A. Peterson, V. Gavrilov, S. Shuvalov, F. Dietrich, and K. Van Bibber, Bull. Am. Phys. Soc. **37**, 886 (1992).

V. An Incomplete List of Nuclear Physics Seminars and Colloquia Given at the University of Massachusetts at Amherst from June 1, 1991 to May 31, 1992

- September 19: Professor Richard Kofler, University of Massachusetts, "High Energy Cosmic Ray Photons"
- September 19: Dr. Sergey Shuvalov, ITEP, Moscow, Russia, "A Proposal for the Measurement of Nuclear Structure Functions at $X_{Bj} > 1$ "
- September 26: Dr. Kebin Wang, University of Massachusetts, "Measuring Pion Polarizability with Tagged Photon Scattering"
- October 2: Professor Rory Miskimen, University of Massachusetts, "Probing the Nuclear Force and the Nucleus at Short Range"
- October 3: Professor John Donoghue, University of Massachusetts, "Testing the Higgs Sector"
- October 10: Dr. Shoba Veeraraghaven, University of Massachusetts, "Cosmological Implications of a 17-keV Neutrino"
- October 10: Dr. Sergey Shuvalov, ITEP-Moscow, Russia, "Deep Inelastic Nuclear Reactions: What New Can We Learn from e-Nucleus Interactions?"
- October 31: Professor Edward Hartouni, University of Massachusetts, "Experimental prospects for the '90's"
- October 31: Professor Howard W. Nicholson, Mount Holyoke College, "Double Beta Decay: Experimental Results"
- November 7: Dr. Latifa Elouadrhiri, University of Massachusetts, "Experimental Study of Pion Electroproduction Near Threshold"
- November 21: Kyong Sei Lee, University of Massachusetts, "A Measurement of the Threshold Electrodisintegration of the Deuteron" (A Ph.D. Dissertation Proposal)
- December 5: Steve Churchwell, University of Massachusetts, "Elastic Magnetic Electron Scattering from ^3He at High Momentum Transfers"
- December 9: Professor Francis Pichanick, University of Massachusetts, "Recent Re-evaluations of the Biological Effects of Ionizing Radiation"
- December 12: Professor Michael Kreisler, University of Massachusetts, "HEP Politics: The Mass of the Neutrino, et al."
- February 3: Professor Moshe Gai, Yale University, "Astrophysics: What's SNU? (Secondary Beams for Nuclear Astrophysics)"
- February 10: Dr. Pat Walsh, Massachusetts Institute of Technology, "Electroproduction of the π^0 Near Threshold"
- February 12: Professor Helmut Vogel, Carnegie Mellon University, "The Beginnings of High Statistics Z^0 Physics"
- February 13: Professor John Donoghue, University of Massachusetts, "Heavy Quark Physics - An Update"

February 18: Dr. Cecile Martin, Orsay, "J/ψ Production with 20 GeV Electrons"

February 20: Professor Monroe Rabin, University of Massachusetts, "A Full Acceptance Detector for SSC?"

February 24: Professor E. Hadjimichael, Fairfield University, "Electromagnetic Form Factors of A=3 Systems"

February 26: Dr. Frank Close, Rutherford Lab, "Cold Fusion: Too Hot to Handle"

March 4: Dr. Roger W. Howell, University of Medicine & Dentistry of New Jersey, "Radionuclides in Medicine: Risks and Benefits"

March 9: Dr. Alan Thompson, Harvard University, "The Quest for the Neutron Electric Form Factor in Polarized ³He"

March 23: Christopher J. Arabadjis, University of Massachusetts, "The (e,e'p) Reaction as a Probe of Nuclear Wave Functions"

March 30: Dr. Pierre Bertin, Clermont-Ferrand, France, "Status of the European Electron Accelerator Project"

April 2: Professor Victor L. Elias, University of Western Ontario, "Quark Condensates"

April 6: Professor Leonid Frankfurt, Michigan State University, "Color Coherence Phenomena"

April 7: Professor Daniel Wyler, University of Zurich, "Exclusive B-Meson Decay"

April 9: Dr. Latifa Elouadrhiri, University of Massachusetts, "Results from the Analysis of Beam-Gas Events from the TPC at SLAC"

April 13: Professor Rory Miskimen and Dr. Kebin Wang, University of Massachusetts, "Pre-presentations for the Washington APS Meeting"

April 16: Professor Arthur Swift, University of Massachusetts, "Technicolor and Moose Diagrams"

April 21: Professor Janos Polonyi, Massachusetts Institute of Technology, "The QCD Vacuum and Localization"

April 23: Antonio Perez, University of Massachusetts, "ε'/ε Measurements at the Φ Factory"

April 27: Professor Nimai Mukhopadhyay, Rensselaer Polytechnic Institute, "Baryon Resonances?"

April 30: Professor Michael Kreisler, University of Massachusetts, "Can High Energy Physics See Final-State Coulomb Interactions?"

May 5: Professor Sangib Mishra, Harvard University, "New Results from the CCFR Fermilab Neutrino Experiment and Future Directions"

May 6: Professor David Rowe, University of Toronto, "Art, Symmetry and Nuclear Physics"

May 21: Dr. Helene Fonvieille, University of Clermont-Ferrand, France, "Status of SLAC Experiment E-142 and the Spin Structure Function of the Neutron"

VI. Comments on Scientific Personnel

A. Postdoctoral Research Associates

Dr. Kebin Wang was a postdoctoral research associate at Stanford University before joining our group in mid-December of 1989. He obtained his Ph.D. degree under the direction of Professor Stanley Hanna at Stanford and continued to do postdoctoral work in Professor Hanna's group. He fills the position vacated by Dr. Sayed Rokni, who is now on the SLAC staff. He is heavily involved in the analysis of the SLAC TPC data.

Dr. Latifa Elouadrhiri joined our group as a postdoctoral research associate on August 20, 1991. She carried out her Ph.D. work at the University of Clermont-Ferrand in France under the direction of Professor Pierre Bertin. Twelve people from his group participated in the now-defunct PEGASYS collaboration at SLAC. Dr. Elouadrhiri's thesis topic involved the difficult measurement at Saclay of electroproduction of positive pions from the proton at threshold. This subject is closely related to experimental tests of chiral symmetry, as suggested by UMass Professor Holstein. We are considering a related experiment for CEBAF, the measurement of the pion's polarizability, perhaps using the CLAS detector. Dr. Elouadrhiri's experience is useful in this endeavor as well as other aspects of our program. She is also heavily involved in the analysis of the SLAC TPC data.

B. Faculty

Professor John Dubach has served as Acting Director of the University Computing Services (UCS) since September of 1989, and will continue in that position through December of 1992. This is a half-time position with funding for the UCS portion coming from the Provost's Office. Because of the hiring freeze mentioned in Part VII, an external candidate could not be hired. Dubach was well-qualified for the position by virtue of his experience using computers to make very large scale theoretical physics calculations. Besides being Acting Director of UCS, and working on his research problems, he also continues to carry a full teaching load, and to act

in an associate editorial capacity for Physical Review Letters. Dubach was promoted to full professor in the spring of 1991.

Professor Ross Hicks spent the spring semester and will spend the summer of 1992 at NIKHEF in Amsterdam. He was promoted this spring to full professor. He will continue to work on the UMass-NIKHEF-Lund related projects outlined in Part II.D, as well as other projects.

Associate Professor Rory Miskimen's tenure case was acted upon in the fall of 1991. Just a few days ago the board of trustees approved his grant of tenure, and his promotion to associate professor. He is playing a major role in the design of electronics for the innermost drift chamber of CEBAF's CLAS detector and in the SLAC TPC beam-gas event analysis with the Soviet physicists, as described in Parts IIC. and II.B., respectively.

C. Graduate Students

Kyong Sei Lee is a Korean working on completing the analysis and writing his thesis on the Bates high-resolution experiment on the threshold-breakup of the deuteron. He will also be involved in the PEP beam-gas event analysis, as described in Part II.B.

Steve Churchwell joined our group at the beginning of the summer of 1989, and worked at SLAC on developing a prototype drift chamber for PEGASYS. He passed the qualifying examination in January of 1990. He is deeply involved in the design of the ^3He target system for the approved Bates experiment on ^3He elastic magnetic scattering.

Geoffrey Feldman was a graduate student in theoretical nuclear physics. He finished his research work on non-leptonic weak interactions in the delta and hypernuclear sectors under Professors Barry Holstein and John Dubach, he wrote his thesis and graduated in January of 1992. He stayed on with our group for the spring semester and at this time has not yet secured a job in physics.

Chris Arabadjis is a graduate student in theoretical nuclear physics. He has passed the qualifying examination, and is working on the theory of the (e,e'p) reaction. We expect him to finish his thesis by January of 1993.

Joseph Gallant is a graduate student in theoretical physics who is working on the problem of core polarization in the shell model. He manages our computer facilities and is supported by University funds until January of 1993, except for summers when he is supported by the grant. Mr. Gallant has passed the qualifying examination.

David Olivieri joined our group in the summer of 1991. He is working on the electronics for the innermost drift chamber of the CLAS detector. He will work on PEP beam-gas event analysis as a thesis topic. Although he has passed the Ph.D. qualifying exam, he still must satisfy a one-semester teaching requirement to be completely qualified.

Xiaodong Jiang, from Beijing, China, joined our group in the summer of 1991. He has passed the qualifying examination and has had experience at Fermilab. He is working on the Bates ^3He experiment and will study the threshold breakup region for which data will be acquired simultaneously with the elastic data.

Thomas Hemmert is a foreign student from Germany who joined our group in January of 1992. He was supported in the spring term by a teaching assistantship and during summer by a research assistantship. He has passed his qualifying examinations and probably will work in nuclear theory.

All of our graduate students have attended the Hampton University Graduate Summer School at CEBAF.

Budget returned.
ds

**DATE
FILMED**
8 / 14 / 92