AIM

Association for Information and Image Management

1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202

Centimeter

Inches

MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

**AN IMPROVEMENT IN THE VALUE OF THE ENERGY
OF THE FIRST EXCITED STATE IN ^{229}Th**

R. G. Helmer and C. W. Reich

Published April 1993

**IDAHO NATIONAL ENGINEERING LABORATORY
EG&G IDAHO, INC.
IDAHO FALLS, IDAHO 83415**

MASTER

Prepared for the
U.S. Department of Energy
Supported through the EG&G Idaho Laboratory
Directed Research & Development Program
DOE Idaho Field Office
Contract DE-AC07-76ID01570.

100

ABSTRACT

It has been known for many years that the first excited state of ^{229}Th lies close to the ground state. Originally this energy was given as < 0.1 keV; later, the authors reported a value of -1 ± 4 eV. Since the existence of a nuclear state at this very low energy is of considerable interest, measurements of its energy, half-life and modes of excitation and deexcitation are important. In an attempt to improve the level energy, we have measured over 175 γ -ray spectra with several combinations of radionuclides. In comparison with our earlier study, we have considered more γ rays in ^{229}Th , used more well-measured reference lines, used more detectors, used detectors with better low-energy resolution, more closely matched peak count rates, and specifically considered certain systematic errors. From this large set of measurements we have deduced a value of 3.5 ± 1.0 eV for the energy of this level.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the assistance of the staff of the Radiation Measurements Laboratory for the use of a counting room and for assistance in optimizing the electronics. They also wish to thank E. W. Killian for modifications of the GINA spectral analysis program, J. D. Baker for preparing most of the sources used, K. D. Watts for assisting with the measurements, and R. C. Greenwood for comments on the manuscript.

CONTENTS

ABSTRACT	iii
ACKNOWLEDGEMENTS	v
1. INTRODUCTION	i
2. METHODOLOGY	4
2.1 Energy Scale and Uncertainties	4
2.2 Calibration Sources	4
2.3 Measurement and Analysis Methods	5
2.4 Detectors	11
3. MEASUREMENT RESULTS AND γ -RAY ENERGIES WITHOUT ^{233}U	12
3.1 Measurement Results for ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am	12
3.2 γ -Ray Energies for ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am	12
4. MEASUREMENT RESULTS AND γ -RAY ENERGIES FOR ^{233}U	17
4.1 Measurement Results	17
4.2 γ -Ray Energies	34
5. ENERGY RELATIONSHIPS IN ^{229}Th	40
6. CONCLUSION	43
7. REFERENCES	44

FIGURES

1. A portion of the low-energy level scheme of ^{229}Th showing the γ rays that are used in this determination of the energy, Δ , of the first excited level	2
2. Portion of a ^{161}Tb and ^{241}Am spectrum from the OR-16 detector showing the γ -ray lines at 25.65 and 26.34 keV. The solid curves show spectral background used in the fit to the 26.34 keV peak and the Gaussian plus background from the fit	10

CONTENTS (CONTINUED)

3.	Portion of the ^{233}U and $^{172}\text{Hf}+^{172}\text{Lu}$ γ -ray spectrum measured with the Si(Li) detector showing the γ rays at 23.93, 25.31, and 29.18 keV	20
4.	Portion of the ^{233}U and ^{241}Am γ -ray spectrum measured with the OR-16 detector showing the γ rays at 25.31, 26.34, and 29.18 keV	21
5.	Portion of the ^{233}U and ^{161}Tb γ -ray spectrum measured with the Si(Li) detector showing the γ rays at 48.91, 53.61, and 54.70 keV	22
6.	Portion of the ^{233}U γ -ray spectrum measured with the OR-16 detector showing the γ rays at 67.94 and 71.81 keV	23
7.	Portion of the ^{233}U and ^{161}Tb γ -ray spectrum measured with from the Si(Li) detector showing the γ rays at 71.81 and 74.56 keV	24
8.	Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with from the OR-16 detector showing the γ rays at 97.13 and 100.10 keV	25
9.	Portion of the ^{233}U γ -ray spectrum measured with the OR-16 detector showing the γ -rays at 117.16 and 118.97 keV	26
10.	Portion of the ^{233}U and ^{169}Yb γ -ray spectrum measured with the PG-16 detector showing the γ rays at 130.52 and 135.33 keV	27
11.	Portion of the ^{233}U γ -ray spectrum measured with the OR-16 detector showing the γ rays at 146.34 and 148.15 keV	28
12.	Portion of the ^{233}U and ^{169}Yb γ -ray spectrum measured with the PG-16 detector showing the γ rays at 177.21, 187.96, and 197.95 keV	29
13.	Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with the OR-12 detector showing the γ rays at 217.15 and 222.10 keV, as well as the fit of a doublet to the former peak	30
14.	Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with the OR-12 detector showing the γ rays at 264.02 and 274.73 keV	31
15.	Gaussian fits to peaks at 59 keV, a singlet from ^{241}Am , and 42 keV, a doublet from ^{233}U	41

CONTENTS (CONTINUED)

TABLES

1. Energies of reference and calibration lines	6
2. Properties of detectors	11
3. Measured spectra that do not include ^{233}U	13
4. Measured γ -ray energy differences between lines of $^{172}\text{Hf}+^{172}\text{Lu}$ and ^{241}Am and lines of ^{161}Tb and ^{241}Am	46
5. Measured γ -ray energy differences between lines of ^{161}Tb and ^{172}Hf and two lines ^{161}Tb	48
6. Measured γ -ray energy differences for two lines from $^{172}\text{Hf}+^{172}\text{Lu}$	50
7. Energies deduced for γ rays from ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am	14
8. Comparison of our results with those previously published in the evaluations of the Table of Radioactive Isotopes (TRI) and the Nuclear Data Sheets (NDS)	15
9. Spectra measured that include ^{233}U	18
10. Measured γ -ray energy differences for ^{241}Am reference lines of 26 and 59 keV	52
11. Measured γ -ray energy difference from the ^{152}Eu reference line of 344 keV	54
12. Measured γ -ray energy differences from the ^{210}Pb reference line of 46 keV	55
13. Measured γ -ray energy differences from the ^{170}Tm reference line of 84 keV	56
14. Measured γ -ray energy differences from the ^{161}Tb reference lines 48 and 74 keV	57
15. Measured γ -ray energy differences from the ^{172}Hf reference lines of 23 and 78 keV	58
16. Measured γ -ray energy differences from the ^{182}Ta reference lines from 31 to 264 keV	60
17. Measured γ -ray energy differences from the ^{169}Yb reference lines of 63 to 307 keV	64

CONTENTS (CONTINUED)

18. Measured γ -ray energy differences between lines from ^{233}U from 25 to 367 keV	68
19. Shifts in calculated peak positions as fitting range is increased	33
20. γ -ray energies for ^{233}U lines	36
21. Difference between crossover and cascade energies	42
22. Energy of ^{229}Th first excited level, Δ	43

AN IMPROVEMENT IN THE VALUE OF THE ENERGY OF THE FIRST EXCITED STATE IN ^{229}Th

1. INTRODUCTION

In γ -ray studies of the level structure of ^{229}Th from the α decay of ^{233}U , Kroger and Reich¹ concluded that the first excited state, with $J^\pi = 3/2^+$, in ^{229}Th occurs within 0.1 keV of the $5/2^+$ ground state. More recently, Reich and Helmer² reported measurements of γ -ray energy differences from which they deduced that this level energy is -1 ± 4 eV. Since it is presumed² that the $3/2^+$ state lies above the $5/2^+$ state, this result also may be stated as indicating that this first excited level lies below 7 eV (at the 2σ level). The relationships of the γ rays involved in these measurements are shown in Figure 1. The energy, Δ , of the first excited level is given by each of the four combinations

$$\begin{aligned}\Delta &= 97-25-71, \\ \Delta &= 97-67-29, \\ \Delta &= 148-118+117-146, \text{ and} \\ \Delta &= 148-76+74-146.\end{aligned}$$

Also of interest are two cascade-crossover combinations

$$\begin{aligned}217 &= 187+29 \text{ and} \\ 164 &= 135+29.\end{aligned}$$

The existence of an excited nuclear state at such a low energy has stimulated considerable interest; see, for example, References 3-8. The basic reason for this interest is that this nuclide may offer the chance to explore new methods of exciting nuclear states. For example, one might be able to excite the electron cloud from an external source (e.g., a laser or a particle) and then have the energy transferred to the nucleus. Such experiments would be of considerable interest in atomic and molecular, as well as nuclear, physics. Since this type of excitation is one of the steps in a basic conceptual design of a γ -ray laser, such measurements would be of interest in that field also.

Because of the interest in this particular level, we undertook a series of measurements to determine more precise energies for some of the γ rays from the decay of ^{233}U in order to reduce the uncertainty in the deduced level energy. These measurements differ from the previous ones² in the following ways:

Figure 1. A portion of the low-energy level scheme of ^{229}Th showing the γ rays that are used in this determination of the energy, Δ , of the first excited level.

- more γ rays in ^{229}Th were considered;
- measurements were made on more detectors, 5 rather than 1;
- more reference lines were used for calibration;
- more measurements were made for each energy difference;
- detectors with better energy resolution were used, especially at the lowest energies;
- the activities of the calibration and ^{233}U sources were carefully matched, so they could be mounted adjacent to each other and thereby reduce the systematic errors due to peak shifts with source position⁹; and
- the peak fitting was done on an interactive system.

In this report, we describe the methods used (sect. 2); the measurement of the energies for some γ rays from ^{161}Tb , $^{172}\text{Hf} + ^{172}\text{Lu}$, and ^{241}Am (sect. 3); the measurement of the energies for some γ rays from ^{233}U (sect. 4); and the application of these results to the level energy in ^{229}Th (sect. 5); and the conclusion (sect. 6). For completeness, and for possible future use, all of the energy differences obtained in this study are reported even though many of them are not applicable to the ^{229}Th level-energy determination.

2. METHODOLOGY

2.1 Energy Scale and Uncertainties

In this report all of the γ -ray energies are on the scale used for the IUPAP-recommended list of energies¹⁰. On this scale the primary γ ray from the decay of ^{198}Au is 411.8044(11) keV. The energy of this line has been determined from very precise measurements of its wavelength¹¹.

There are two uncertainties that apply to this energy value. The first is the combination of all the measurement uncertainties that apply to its wavelength; this uncertainty is 0.37 ppm, or 0.16 eV. The second uncertainty is that from the conversion of the γ -ray wavelength to the electron-Volt scale; this is 2.6 ppm. The total uncertainty is obtained by combining these two contributions in quadrature. If one were to adjust the electron-Volt energy scale to correspond to the most recent values of this wavelength and the fundamental constants, the energy of this 411-keV line would be reduced by 5.8 ppm. Most of the other calibration lines used here would consequently be reduced by a similar amount. This small change would not influence our final result and, therefore, this correction has not been made.

For the analysis in this report, we need consider only the measurement uncertainties for the calibration lines used. That is, we can do all the calculations on what is effectively the wavelength scale. In this way, the measurements relative to different reference lines are properly weighted. In principle, the uncertainty in the conversion between the wavelength and electron-Volt scales is added back in for the final result. But, since the measurement uncertainty in our final result is of the order of 15%, this uncertainty is not significant.

2.2 Calibration Sources

In the search for radionuclides that provide γ -ray lines that would be useful as reference lines, there were two criteria. First, the γ -ray energies should be very well known. Ideally, this means that the uncertainty in the energy is <1 eV and that its measurement was made in the last 15 years or so. Second, the reference lines should be near to, but resolved from, each of the γ rays from ^{233}U of interest.

The energy-calibration sources used were $^{172}\text{Hf}+^{172}\text{Lu}$, ^{161}Tb , ^{241}Am , ^{182}Ta , ^{169}Yb , ^{152}Eu , ^{210}Pb , and ^{170}Tm . The energies of the γ rays used are given in Table 1. Among these lines, those for $^{172}\text{Hf}+^{172}\text{Lu}$ and ^{241}Am were measured¹² specifically for this study. The ^{241}Am lines were remeasured because the

previous values did not meet the above criteria: these values^{17,18} had uncertainties of 1 eV; and the measurements are old (i.e., 1968 and 1970). The values for ^{172}Hf (1.87 y) + ^{172}Lu were measured to allow the use of the 23.9Hf line as an intermediate reference line between the 25.6Tb line and the 25.3U line. (The notation 25Tb means the 25-keV line from the decay of ^{161}Tb . The notation for the radionuclides is: ^{152}Eu , Eu; ^{161}Tb , Tb; ^{169}Yb , Yb; ^{170}Tm , Tm; ^{172}Hf + ^{172}Lu , Hf; ^{182}Ta , Ta; ^{210}Pb , Pb, ^{233}U , U; and ^{241}Am , Am.) The only remaining line that does not meet the above criteria is that for ^{210}Pb , which is from Helmer et al.¹³ and is known only to 1 eV. The radionuclides listed in Table 1 have many other γ rays with well-measured energies, but these lines were either not in an energy range of interest, or there were ^{233}U lines which interfered with their use.

The use of the 25.6Tb line as a reference line is somewhat complicated, since this line is not resolved from the 25.3U line, whose energy we wish to determine. Therefore, we have had to make use of two-step processes to determine this 25.6Tb - 25.3U energy difference. ^{172}Hf has a strong line at 23.9 keV, so we measured the 23.9Hf - 25.3U and 23.9Hf - 25.6Tb differences. Similarly, the ^{241}Am line at 26.3 keV was used to obtain the 25.3U - 26.3Am and 25.6Tb - 26.3Am differences.

The reference-line energies in Table 1 were used in the determination of the energies of 50 γ rays from 25.3 to 367.7 keV in ^{229}Th .

The calibration sources were prepared by depositing a solution containing the radionuclide on a backing of Kaptan film, drying the solution, and covering it with tape and Kaptan film. In most cases, sources were made with a range of activities. The ^{233}U sources were also made with a range of activities. This variety of source strengths was necessary so the ^{233}U and calibration sources could be paired in such a way as to have similar peak count rates for the lines to be compared, and still have the sources be adjacent to each other. In all cases the source holders were in contact, so the sources were separated by distances of ~ 0.3 cm.

2.3 Measurement and Analysis Methods

The typical measurement procedure can be illustrated for the case of comparing the 42U line with the 46Pb line. Particular sources of ^{233}U and ^{210}Pb were chosen so that the count rates in these two peaks were comparable when the sources were placed adjacent to each other. A source-to-detector distance of 10 cm was typically chosen so that the sources were 'far' from the detector compared to the difference in the source position (~ 0.3 cm), but still gave a sufficient count rate. Half of the measured spectra were obtained with the ^{233}U source nearer the detector and the other

Table 1. Energies of reference and calibration lines. Uncertainties include only those from the measurement and not that from the conversion between the wavelength and electron-Volt energy scales (see text).

<u>Parent isotope</u>	<u>Energy (keV)</u>	<u>Parent isotope</u>	<u>Energy (keV)</u>
$^{172}\text{Hf} + ^{172}\text{Lu}^{\text{a}}$	23.9331(2) 78.7426(6)	$^{210}\text{Pb}^{\text{b}}$	46.539(1)
$^{161}\text{Tb}^{\text{c}}$	25.65150(3) 48.91562(6) 74.56711(10)	$^{169}\text{Yb}^{\text{d}}$	63.12077(8) 109.77987(5) 130.52368(4) 177.21402(6) 197.95788(6)
$^{241}\text{Am}^{\text{a}}$	26.3448(2) 59.5412(2)		261.07857(11) 307.73757(9)
$^{182}\text{Ta}^{\text{e}}$	31.7378(7) 84.6808(3) 100.10653(18) 113.6723(2) 152.4308(3) 156.3874(3) 179.3948(3) 198.3530(3) 222.1099(3) 229.3220(6) 264.0755(3)	$^{170}\text{Tm}^{\text{f}}$	84.25510(19)
		$^{152}\text{Eu}^{\text{g}}$	344.2811(17)

^a Energies from Reference 12 and were measured for this study.

^b Energy from Reference 13.

^c Energy from Reference 14.

^d Energies from Reference 15.

^e Energies from Reference 10, except for the 31-keV γ ray, which is computed from the 116.4186(6) - 84.68-8(3) difference.

^f Energy from Reference 10.

^g Energy from Reference 10. See also Reference 16.

half with the ^{210}Pb source nearer the detector; this would cancel out any bias in the energy difference due to differences in the source-detector distances. Since ^{210}Pb has only one γ ray, most of the calibration lines used were from ^{233}U .

A crude estimate was made of the magnitude of the shift in the apparent line energy with source distance for a typical counting arrangement. For the planar Ge detector discussed in Reference 9, we can deduce from the figures therein

- for a 1489-keV γ ray, the change in the apparent energy of a peak as a function of the source-detector distance is ~ 1.2 eV/cm at a distance of 10 cm (Figure 4 of Reference 9);
- the shift at 500 keV is about 0.2 times the shift at 1489 keV (Figure 3 of Reference 9); and
- the shift at energies below 500 keV is approximately proportional to the γ -ray energy.

For a typical measurement made in this study, we have

- a source-detector distance of 10 cm;
- a difference in source distance between sources of 0.3 cm;
- a voltage gradient in the detector of about twice that of the detector in Reference 9; and
- a typical γ -ray energy of 100 keV.

For this case, the estimated peak shift between the sources is the above ~ 1.2 eV/cm, times 2.0 for the larger voltage gradient, times 0.3 for the distance difference of 0.3 cm, times 0.2 to go from 1489 to 500 keV, and times 0.2 to go from 500 to 100 keV. This amounts to 0.03 eV, and thus can be neglected.

In the analysis of these spectra, the energy-channel relationship is taken to be

$$E(x) = a + b \cdot (x - \delta) + c \cdot (x - \delta)^2$$

and the peak widths, FWHM, are

$$W(x) = (\alpha + \beta \cdot x)^{1/2},$$

where x is the channel and E is the energy. The user identifies the γ -ray peaks to be used in these calibrations and provides the γ -ray energies. The parameters a , b , and c are determined by a least-squares fit to the energies and positions of these peaks; and the parameters α and β are from a similar fit to the peak widths and positions. The δ is an estimated zero offset that is computed separately before the parameters a , b , and c are computed. In this step the routine automatically defines the peak fitting region for each peak; that is, the fitting is not interactive.

An interactive routine is then used to fit the peaks of interest in determining γ -ray energy differences. In this peak-fitting routine the user identifies, on a video screen, the channels below and above the peak to be used to determine the linear or step background function, the channel range included in the fit of the peak parameters, and the initial centroid of the peak. The initial peak width is taken from the width-calibration function. The peak parameters are then calculated by a nonlinear least-squares routine.

The peak positions, X_1 and X_2 , from these fits are then used to determine the difference in energy between nearby peaks, namely,

$$\Delta E = E_2 - E_1 = b \cdot (X_2 - X_1) + c \cdot (X_2^2 - X_1^2).$$

The calculation of the uncertainty in this difference includes the variances in b , c , X_1 , and X_2 and the covariance between b and c . This uncertainty is dominated by the two terms involving $\text{var}(X_1)$ and $\text{var}(X_2)$.

For each energy difference where data were taken on different detectors, several weighted averages are computed. These weighted averages are computed with the weights taken as $1/\sigma^2$, where σ is the uncertainty. Averages are computed for each detector separately, and for all of the data together. The reduced- χ^2 value, χ_R^2 , is determined for each average. If the χ_R^2 values are larger than expected, or the averages from different detectors are inconsistent, the data are reviewed to determine if there is an identifiable problem. In some cases a single value, which was an extreme outlier, was removed and the averages recalculated.

The uncertainties given throughout this work, except as influenced by the χ_R^2 values, are statistical. In fitting a set of calibration lines, we use a quadratic relationship, $E_q(X)$, between the energy and the channel number which will differ from the true relationship, $E_t(X)$, by an amount $\delta E(E_y)$. There are several contributions to this difference: (i) the statistical variations in the peak locations; (ii) any errors (not uncertainties, but

actual errors) in the calibration energies; and (iii) the fact that the true relationship may not be adequately represented by any quadratic function (i.e., the true relationship has higher order terms). In addition to these general contributions, for an individual γ -ray peak a fourth factor would be any shift in its position due to other peaks in the spectrum. In our computation of an energy difference, ΔE , between two lines from one spectrum, all of these factors are ignored.

Before discussing our method of addressing these four factors, let us note the precision we are implying in our results. The smallest uncertainty that we have quoted for an average ΔE is 0.1 eV. Since this is from more than 16 measurements, this corresponds to an uncertainty of about 0.4 eV for an individual measurement, if each measured value has the same uncertainty. Many measurements were made with a gain scale corresponding to 20 eV per channel. An uncertainty in a peak position of 0.4 eV then corresponds to 0.02 channels. It is questionable if $\delta E(E_\gamma)$ is this small. However, since we are always determining an energy difference, it is actually only the change in $\delta E(E_\gamma)$ between the two peak positions that needs to be this small.

Our decision to quote only the statistical uncertainty is based on the following. It is assumed that, since the first factor is statistical, in an average over many spectra, it will be averaged out. It is assumed that the second contribution will be identified, or at least partially averaged out, by using different calibration lines whenever possible.

We chose to treat the third factor by making our measurements in the following manner. Each measurement was made on a different gain scale, and measurements were made on different detectors, each with its own ADC and amplifier. Both of these variations in the experimental setup are expected to result in different functions $\delta E(E_\gamma)$. It is assumed that by averaging the results from all of the measurements the average δE will approach zero.

The fourth factor must be dealt with individually for each peak for which it is a concern. A case of particular interest is the 25Tb - 26Am difference. Since these peaks are not completely resolved, as shown in Figure 2, in fitting one of these peaks, the second peak will tend to move the apparent peak location of the first peak closer to that of the second one. Therefore, the measured difference is expected to be less than the actual difference. If this effect is significant, the observed difference should increase as the detector resolution improves. In fact, the values from the Si(Li) and OR-16 detectors agree very well, so this effect is negligible and all of these measurements have been used.

Figure 2. Portion of the ^{161}Tb and ^{241}Am γ -ray spectrum measured with the OR-16 detector showing the γ rays at 25.65 and 26.34 keV. The solid curves show spectral background used in the fit to the 26.34-keV peak and the Gaussian plus background from the fit.

2.4 Detectors

Some properties of the five detectors used in this study are given in Table 2. The detectors include two high-purity Ge coaxial detectors, two Ge planar detectors, and a Si(Li) detector. The resolutions given in the table are not the optimum values for these detectors; rather, they are observed values for actual spectra, where the counting times are typically 72 hours.

Table 2. Properties of detectors.

<u>Detector</u>	<u>Material</u>	<u>Shape</u>	<u>Volume</u> <u>(cm³)</u>	<u>Diam.</u> <u>(mm)</u>	<u>Depth</u> <u>(mm)</u>	<u>FWHM</u> <u>29 keV</u>	<u>(eV) at</u> <u>122 keV</u>
OR-12	Ge	coaxial	114				950
PG-21	Ge	coaxial	63				900
PF-16	Ge	planar		16	10	390	
OR-16	Ge	planar		10	7	270	
Si	Si(Li)	planar		6	4	345	

3. MEASUREMENT RESULTS AND γ -RAY ENERGIES WITHOUT ^{233}U

To simplify the discussion of the results, we have divided the spectral measurements into two groups: those spectra that do not include ^{233}U ; and those that do. The 66 spectra that do not include ^{233}U were measured to obtain improved γ -ray energies for $^{172}\text{Hf}+^{172}\text{Lu}$ and ^{241}Am ; these spectra are listed in Table 3.

In Tables 4-6 the spectra used will be identified by the spectral codes given in Table 3. (To make it easier to follow the text, all tables of the measured energy differences are placed after the references.) In these spectral codes the letters A - E identify the detector, the N indicates that ^{233}U was not present, and the numbers indicate the order in which the spectra were obtained on the particular detector.

3.1 Measurement Results for ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am

Tables 4, 5, and 6 give the individual values for all of the energy differences determined from spectra not containing ^{233}U . Where they are not obvious, the radionuclides present in the source can be deduced from the entry in Table 3 for the specific spectral code. In each of these three tables, the weighted average of the energy difference is given for each detector and then for all of the values together. Any individual value that has been excluded from the averages is identified and the reason is given.

The value adopted for our use is the last value given in each column.

3.2 γ -Ray Energies for ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am

The γ -ray energies deduced from these data are given in Table 7. These values are compared in Table 8 with values from two published evaluations, namely the Table of Radioactive Isotopes¹⁹ and the Nuclear Data Sheets²⁰⁻²². It is clear that the uncertainties quoted here are significantly smaller than the previous ones. It is also interesting that the energy of the 59-keV line from the decay of ^{241}Am has changed by ~ 4 eV. A paper on the results of the analysis of these data has been accepted for publication¹².

There is some statistical objection to including in Table 7 entries which involve the use a line measured in this study as a reference line for a second line. For example, the 78-keV line was measured relative to the 74-keV line from ^{161}Tb . Then the 81-keV line is shown as being measured with respect to both the 74- and 78-keV lines. In spite of this objection, we have included both values in Table 7, and have averaged both values to get the final 81-keV

Table 3. Measured spectra that do not include ^{233}U .

Dates are given in the form month and day, all dates are in 1992.

<u>Detector</u>	<u>Date</u>	<u>Nuclides</u>	<u>Spectral code</u>	<u>Detector</u>	<u>Date</u>	<u>Nuclides</u>	<u>Spectral code</u>
PG-16	0220	Tb & Hf	CN1	Si(Li)	0210	Hf & Am	EN1
	0221	Tb & Hf	CN2		0213	Hf & Am	EN2
	0221	Tb & Hf	CN3		0218	Hf & Am	EN3
	0224	Tb & Hf	CN4		0220	Tb & Am	EN4
	0225	Tb & Hf	CN5		0221	Tb & Am	EN5
	0226	Tb & Hf	CN6		0224	Tb & Am	EN6
	0316	Tb & Hf	CN7		0225	Tb & Am	EN7
	0318	Tb & Hf	CN8		0226	Tb & Am	EN8
	0326	Tb & Hf	CN9		0227	Tb & Hf	EN9
	0330	Tb & Am	CN10		0229	Tb & Hf	EN10
	0401	Tb & Am	CN11		0302	Tb & Hf	EN11
	0409	Tb & Am	CN12		0303	Tb & Hf	EN12
	0415	Tb & Am	CN13		0305	Tb & Hf	EN13
	0415	Tb & Am	CN14		0307	Tb & Hf	EN14
	0416	Tb & Am	CN15		0310	Tb & Hf	EN15
	0417	Tb & Am	CN16		0312	Tb & Hf	EN16
	0427	Hf & Am	CN17		0314	Tb & Hf	EN17
	0429	Hf & Am	CN18		0316	Tb & Am	EN18
	0430	Hf & Am	CN19		0318	Tb & Am	EN19
	0504	Hf & Am	CN20		0319	Tb & Am	EN20
	0506	Hf & Am	CN21		0321	Tb & Am	EN21
	0507	Hf & Am	CN22		0401	Tb & Am	EN22
	0511	Hf & Am	CN23		0409	Tb & Am	EN23
	0514	Hf & Am	CN24		0415	Tb & Am	EN24
	0518	Hf & Am	CN25		0416	Tb & Am	EN25
OR-16	0423	Tb & Am	DN1		0417	Tb & Am	EN26
	0424	Tb & Am	DN2		0521	Hf & Am	EN27
	0428	Tb & Am	DN3		0526	Hf & Am	EN28
	0429	Tb & Am	DN4		0528	Hf & Am	EN29
	0501	Tb & Am	DN5		0601	Hf & Am	EN30
	0504	Tb & Am	DN6		0605	Hf & Am	EN31
	0507	Tb & Am	DN7				
	0511	Tb & Am	DN8				
	0514	Tb & Am	DN9				
	0518	Tb & Am	DN10				

Table 7. Energies deduced for γ rays from ^{161}Tb , $^{172}\text{Hf}+^{172}\text{Lu}$, and ^{241}Am .

Parent isotope	Line (keV)	Reference line	Difference (keV)	<u>γ-ray energy (keV)</u>	
				<u>Individual</u>	<u>Average</u>
^{161}Tb	57.1	48Tb	8.2764(3)	57.1920(3)	
$^{172}\text{Hf}+^{172}\text{Lu}$	23.9	25Tb	1.7184(2)	23.9331(2)	
	78.7	74Tb	4.1755(6)	78.7426(6)	
	81.7	74Ta	7.1841(6)	81.7512(6)	
		78Lu	3.0088(4)	81.7514(7)	81.7513(5)
	90.6	74Tb	16.0787(8)	90.6458(8)	
		81Hf	8.8907(8)	90.6420(9)	90.6440(19)
	112.7	90Lu	22.1340(19)	112.7780(27)	
	114.0	112Lu	1.2827(7)	114.0607(28)	
	122.9	114Hf	8.8549(12)	122.9156(30)	
	125.8	122Hf	2.8966(8)	125.8122(31)	
^{241}Am	26.3	25Tb	0.6933(2)	26.3448(2)	
		23Hf	2.4118(9)	26.3449(9)	26.3448(2)
	59.5	48Tb	10.6257(1) ^a	59.5413(1)	
		74Tb	15.0261(2)	59.5410(2)	59.5412(2)

^a ^{161}Tb has a γ ray at 59.239(13) keV with an intensity 0.012 times that of the 57-keV γ ; this 59.2 γ is not expected to interfere significantly with the 59Am line.

Table 8. Comparison of our results with the evaluated data published in the Table of Radioactive Isotopes¹⁹ (TRI) and the Nuclear Data Sheets (NDS).

<u>Parent isotope</u>	<u>Line (keV)</u>	<u>Present measurement (keV)</u>	<u>Evaluation</u>	
			<u>Ref.</u>	<u>Value</u>
²⁴¹ Am	26.3	26.3448(2)	TRI NDS ²⁰	26.3445(10) 26.345(1)
	59.5	59.5412(2)	TRI NDS ²⁰	59.5364(10) 59.537(1)
¹⁷² Hf	23.9	23.9331(2)	TRI	23.99(5)
	81.7	81.7515(5)	TRI	81.75(5)
	114.0	114.0607(25)	TRI	114.06(10)
	122.9	122.9156(28)	TRI	122.92(10)
	125.8	125.8122(29)	TRI	125.82(5)
¹⁷² Lu	78.7	78.7426(6)	TRI NDS ²¹	78.792(7) 78.738(5)
	90.6	90.6449(13)	TRI NDS ²¹	90.665(17) 90.647(3)
	112.7	112.7780(24)	TRI NDS ²¹	112.757(20) 112.793(5)
¹⁶¹ Tb	57.1	57.1920(3)	TRI NDS ²²	57.212(5) 57.196(2)

energy value. This objection also applies to the values for the 81- and 90-keV lines from $^{172}\text{Hf} + ^{172}\text{Lu}$ and the 26-keV line from ^{241}Am . In the latter case, the 23-keV line from ^{172}Hf is based on the 25-keV line from ^{161}Tb . However, in this case we set out explicitly to provide an alternate path to the 26-keV value because of the small separation, 0.69 keV, between the 25- and 26-keV lines. A proper treatment of the averaging of these values would involve taking into account the correlation between them; this has not been done. But, it should be noted that in all three of these cases the uncertainty for the individual energy value is smaller for the value directly tied to the ^{161}Tb reference line. Therefore, a more complete uncertainty analysis would not change the result significantly.

4. MEASUREMENT RESULTS AND γ -RAY ENERGIES FOR ^{233}U

4.1 Measurement Results

A total of 111 spectra containing ^{233}U were measured for this study; these spectra are listed in Table 9. Tables 10-18 give the individual values for all of the energy differences deduced from these spectra. Table 10 gives the energy differences for the reference lines from ^{241}Am , Table 11 for the reference line from ^{152}Eu , Table 12 from ^{210}Pb , Table 13 from ^{170}Tm , Table 14 from ^{161}Tb , Table 15 from $^{172}\text{Hf}+^{172}\text{Lu}$, Table 16 from ^{182}Ta , Table 17 from ^{169}Yb , and Table 18 for 38 pairs of lines from ^{233}U . (In these tables the lines are from ^{233}U if they are not labeled otherwise.) Although only a few of these energy differences are of interest in the determination of the energy of the first excited state in ^{229}Th , the others are given for completeness and for possible future use. For these tables, the information in Table 9 can be used to identify the nuclides present in any particular spectrum.

For each energy difference in Tables 10-18, the weighted average of the values is given for each detector and for all of the data combined. In many cases a weighted average is also given for the values from the two higher resolution Ge detectors, PG-16 and OR-16. The value adopted for subsequent use is the last value given in the column.

As illustrations of the quality of the spectra obtained, Figures 3-14 show portions of the γ -ray spectra from four of the detectors and for most of the γ rays used in the determination of the energy of the first excited state of ^{229}Th . Some of these examples are from spectra of ^{233}U only, but most of them include calibration sources. These figures show the quality of the data obtained - the typical counting statistics and the resolution obtained for many of the pairs of peaks used.

Figure 5 shows that the peaks at 52 and 53 keV may influence the peak position of the 54-keV peak by influencing the definition of the background below this peak, even though the 53- and 54-keV peaks are almost completely resolved.

From Figure 8 it is clear that the 97-keV peak from ^{233}U has several interfering lines, both at lower and higher energies, under it. These lines, whose influence was not specifically taken into account in the analysis, may introduce a significant systematic error into the determination of the peak position of the 97-keV peak and hence into the measurement of the 97U-100Ta energy difference, especially since the statistical uncertainty in this difference is only 0.3 eV, one of the smallest reported in this study. In order to investigate the possible magnitude of such an effect, four fits were

Table 9. Spectra measured that include ^{233}U .

Dates are given in the form of month, day, and year.

Detec- tor	Calib. Date	Nuclide	Spectral code	Detec- tor	Calib. Date	Nuclide	Spectral code
OR-12	080790	Ta	A1	PG-16 cont.	082390	Am	C11
	080790	Ta	A2		082490	Am	C12
	080890	Ta	A3		082790	none	C13
	080990	Ta	A4		082890	none	C14
	090490	Ta	A5		083190	Eu	C15
	090790	Ta	A6		090490	none	C16
	091090	Ta	A7		091090	Eu	C17
	091390	Ta	A8		091490	Eu	C18
	091990	Yb	A9		091790	Eu, Ta	C19
	092090	Yb	A10		092090	Yb	C20
	092190	Yb	A11		092190	Yb	C21
	092490	Yb	A12		092490	Yb	C22
	100490	Yb, Tm	A13		092890	Yb	C23
	100590	Yb, Tm	A14		102390	Yb, Tm	C24
	100890	Yb, Tm	A15		102690	Yb, Tm	C25
	100990	Yb, Tm	A16		102990	Yb, Tm	C26
	101290	Yb, Tm	A17		110290	Yb, Tm	C27
PG-21	081090	Pb	B1		021392	Hf	C28
	081390	Pb	B2		021892	Hf	C29
	081390	Pb	B3		022792	Tb	C30
	081490	Pb	B4		022992	Tb	C31
	081690	Eu	B5		030292	Tb	C32
	081790	Eu	B6		030392	Tb	C33
PG-16	080890	Ta	C1		030592	Tb	C34
	080990	Ta	C2		030792	Tb	C35
	081090	Ta	C3		031092	Tb	C36
	081390	Ta	C4		031292	Tb	C37
	081390	Ta	C5		031492	Tb	C38
	081590	Ta	C6		040392	Hf	C39
	081690	none	C7		040692	Hf	C40
	081790	none	C8		041092	Hf	C41
	082090	Am	C9		041792	Hf	C42
	082190	Am	C10		042092	Hf	C43
					042392	Hf	C44

Table 9. (Continued)

<u>Detec-</u> <u>tor</u>	<u>Calib.</u> <u>Date</u>	<u>Nuclide</u>	<u>Spectral</u> <u>code</u>	<u>Detec-</u> <u>tor</u>	<u>Calib.</u> <u>Date</u>	<u>Nuclide</u>	<u>Spectral</u> <u>code</u>
OR-16	111390	Ta, Am	D1	Si(Li)	121691	Am	E1
	111490	Ta, Am	D		122091	Am	E2
	111690	Ta, Am	D3		122391	Am	E3
	111990	Ta, Am	D4		010392	Am	E4
	112190	Ta	D5		011092	none	E5
	112690	Ta	D6		011392	none	E6
	112890	Ta	D7		012092	none	E7
	113090	Am	D8		012392	none	E8
	120390	Am	D9		012892	Hf	E9
	120590	Am	D10		020392	Hf	E10
	120790	Am	D11		020792	Hf	E11
	121090	Yb, Tm	D12		022192	Tb	E12
	121290	Yb, Tm	D13		032392	Tb	E13
	121490	Yb, Tm	D14		032592	Tb	E14
	121990	none	D15		032692	Tb	E15
	010291	none	D16		033092	Tb	E16
	010791	none	D17		040292	Tb	E17
	011191	none	D18		040692	Tb	E18
	011691	Pb	D19		041092	Tb	E19
	012191	Pb	D20		041792	Tb	E20
	012391	Pb	D21		042092	Hf	E21
	012591	Pb	D22		060892	Hf	E22
	120391	Am	D23		061192	Hf	E23
	120591	Am	D24		061992	Hf	E24
	120991	Am	D25		062292	Hf	E25
	121291	Am	D26		062592	Hf	E26
	121691	none	D27		062992	Hf	E27
	122091	none	D28		070292	Hf	E28
	010392	none	D29				
	010692	none	D30				
	012792	Hf	D31				
	013192	Hf	D32				
	020392	Hf	D33

Figure 3. Portion of the ^{233}U and $^{172}\text{Hf} + ^{172}\text{Lu}$ γ -ray spectrum measured with the Si(Li) detector showing the γ rays at 23.93, 25.31, and 29.18 keV. The result of a fit to the first peak is shown.

Figure 4. Portion of the ^{233}U and ^{241}Am γ -ray spectrum measured with the OR-16 detector showing the γ rays at 25.31, 26.34, and 29.18 keV, along with a fit to the 26.34-keV peak.

Figure 5. Portion of the ^{233}U and ^{161}Tb γ -ray spectrum measured with the Si(Li) detector showing the γ rays at 48.91, 53.61, and 54.70 keV.

Figure 6. Portion of the ^{233}U γ -ray spectrum measured with the OR-16 detector showing the γ rays at 67.94 and 71.81 keV.

Figure 7. Portion of the ^{233}U and ^{161}Tb γ -ray spectrum measured with the Si(Li) detector showing the γ rays at 71.81 and 74.56 keV.

Figure 8. Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with the OR-16 detector showing the γ rays at 97.13 and 100.10 keV.

Figure 9. Portion of the ^{235}U γ -ray spectrum measured with the OR-16 detector showing the γ rays at 117.16 and 118.97 keV.

Figure 10. Portion of the ^{233}U and ^{169}Yb γ -ray spectrum measured with the PG-16 detector showing the γ rays at 130.52 and 135.33 keV.

Figure 11. Portion of the ^{233}U γ -ray spectrum measured with the OR-16 detector showing the γ rays at 146.34 and 148.15 keV.

Figure 12. Portion of the ^{233}U and ^{169}Yb γ -ray spectrum measured with the PG-16 detector showing the γ rays at 177.21, 187.96, and 197.95 keV.

Figure 13. Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with the OR-12 detector showing the γ rays at 217.15 and 222.10 keV, as well as the fit of a doublet to the former peak.

Figure 14. Portion of the ^{233}U and ^{182}Ta γ -ray spectrum measured with the OR-12 detector showing the γ rays at 264.02 and 274.73 keV.

made to this peak successively adding two channels (one on each side of the peak) to the fitting range, as shown in Table 19. The smallest fitting range, 9 channels, corresponds to the FWHM. In the analyses of the γ -ray peaks in this work, the fits are generally similar to the first two fitting ranges in this table. Similarly, four fits were made to the 100-keV peak from ^{182}Ta in the same spectrum.

Since we fit the peaks with a symmetric Gaussian function, we expect the calculated peak position to shift to lower values as more channels are included on the low-energy side of any singlet peak. If the secondary peak on the low-energy side of the 97-keV line has any significant additional influence on the position of the main peak, we expect the shift of the calculated position of the 97-keV peak to be larger than that of the more nearly singlet peak at 100 keV. The data in Table 19 show that there is no significant difference in the shifts observed between these two peaks as the fitting range is increased. From this, we conclude that any influence on the calculated position of the 97-keV peak due to the presence of the peaks underlying it is less than about 0.5 eV.

In Figure 11, a pair of important peaks is shown; these are at 146 and 148 keV. The 148-keV peak has the difficulty that it is relatively weak so it is hard to get good statistics and there is a small distortion on the low-energy side. In this case, the contribution of this peak to the uncertainty in any ΔE_γ should be dominated by this statistical component. The 146-keV peak has the problem of the presence of the 145-keV line. Only the top portion of the peaks are used in the fit to a singlet, and a Gaussian function without any tailing is used in the fit. If one were to fit the 145-146 pair as a doublet, one would include portions of each peak all the way down to the spectral background; and this would produce an unacceptable bias between the fits to 146- and 148-keV peaks. Therefore, a singlet fit to the 146-keV peak is preferred. The difficulty in fitting the 146-keV peak is important since the energy difference between the 146- and 148-keV lines appears in two of the four energy combinations that are used to deduce Δ .

As discussed above in regard to the 97-keV peak, we have explored the change in the computed positions of the 146- and 152-keV peaks in spectrum of ^{233}U and ^{152}Eu in order to investigate the presence of possible systematic errors in the calculated position of the 146-keV peak due to the presence of the near-lying 145-keV γ peak (which was not specifically included in the analysis). We assume that any influence of the 145-keV peak would cause the shift in the calculated 146-keV peak position to be larger than that for the 152-keV peak, which is a singlet. As shown in Table 19, no such effect is seen. Therefore, we conclude that any such influence is less than about 0.5 eV.

Table 19. Shifts in calculated peak positions as the fitting range is increased.

γ -ray energy (keV)	Peak Channel	Area	χ^2	No. of channels in fit	Change in peak position channels	Change in peak position eV
97	1613.0	16,100	$\leq 1.^a$	9	≈ 0.000	≈ 0.0
			$\leq 1.^a$	11	-0.014	-0.8
			$\leq 1.^a$	13	-0.035	-2.1
			$\leq 1.^a$	15	-0.033	-2.3
100	1667.6	16,200	$\leq 1.^a$	9	≈ 0.000	≈ 0.0
			$\leq 1.^a$	11	-0.013	-0.8
			$\leq 1.^a$	13	-0.022	-1.3
			$\leq 1.^a$	15	-0.033	-2.0
146	2365.9	248,000	$\leq 1.^a$	11	≈ 0.000	≈ 0.0
			$\leq 1.^a$	13	-0.009	-0.6
			2.3	15	-0.049	-3.0
			3.2	17	-0.082	-5.1
152	2464.4	287,000	1.5	12	≈ 0.000	≈ 0.0
			2.2	14	-0.036	-2.2
			2.5	16	-0.055	-3.4
			2.6	18	-0.073	-4.5

^a For values <1.0 , the computer program does not give explicit values.

Since all four values of Δ to be discussed in Section 5 depend on either the 97-keV γ -ray energy or the 146-148 energy difference, our results depend critically on the fits to the lines in Figures 8 and 11.

Figure 13 shows the difficulty of determining the energy of the 217-keV peak, which is part of a multiplet that is less well resolved than the others.

4.2 γ -Ray Energies

From the results in the previous section, the energies for many of the γ rays from ^{233}U have been deduced. These results are given in Table 20.

In considering these results, the reader should understand that these energies have been determined by doing the calculations in a particular order. In some cases, the calculations could have been done in a different order, in which case some of the γ -ray energies would have been slightly different. To illustrate, assume we have the following set of data:

calibration line energies: C_1 and C_2 with $C_1 < C_2$

γ energies to be determined: E_1 and E_2
with $C_1 < E_1 < E_2 < C_2$

measurement results: $\Delta_1(E_1 - C_1)$, $\Delta_2(E_2 - E_1)$, and $\Delta_3(C_2 - E_2)$.

Then one can determine E_1 and E_2 from either of the following two sets of relationships.

$$(1) \quad E_1 = C_1 + \Delta_1 \text{ and}$$

$$E_2 \text{ is the average of } C_1 + \Delta_1 + \Delta_2 \text{ and } C_2 - \Delta_3,$$

or

$$(2) \quad E_2 = C_2 - \Delta_3 \text{ and}$$

$$E_1 \text{ is the average of } C_1 + \Delta_1 \text{ and } C_2 - \Delta_3 - \Delta_2.$$

It would, of course, be possible to remove this bias of choosing a particular order for the calculations by doing a least-squares fit to each of the over-determined sets of data. To do such a fit correctly one would need to develop a covariance matrix for the measured Δ_i values because generally they will be highly correlated. For example, the same spectra might be used to determine Δ_2 and, say, Δ_3 . Therefore, Δ_2 and Δ_3 would be correlated since the peak at E_2 is common to both. Such least-squares fits have not been done, in part, because it is felt that the improvement in the γ -ray energies would not be significant.

In several cases in Table 20, several values are averaged, with $1/\sigma^2$ weighting, to obtain the final value. In most cases this weighting is not strictly correct since the individual values are often correlated, and this correlation has not been taken into account. For example, the average for the 187-keV γ ray involves energy differences with respect to the 177- and 197-keV lines from ^{169}Yb . These two values make use of the same γ -ray spectra and therefore the same 187-keV peaks. Thus, the two energy differences are not independent; and this is not taken into account in the calculation of the average. The effect of neglecting these correlations is that the computed uncertainty is smaller than it would be if the correlations were taken into account.

Table 20. γ -ray energies for ^{233}U lines

^{233}U Line	Reference		Adopted ΔE_γ	γ -ray energy		
	Nuclide	Energy		Value	Average	χ_R^2
25	^{241}Am	26.3448(2)	1.0340(9)	25.3108(9)	25.3106(8)	0.15
	^{172}Hf	23.9331(2)	1.3771(16)	25.3101(16)		
29	^{182}Ta	31.7378(7)	2.5482(10)	29.1896(12)	29.1846(30) ^a	12.9
	^{172}Hf	23.9331(2)	5.2489(9)	29.1820(9)		
	^{241}Am	26.3448(2)	2.8397(4)	29.1845(5)		
42	^{210}Pb	46.539(1)	4.0862(6)	42.453(1)	42.4524(7) ^b	4.27
	^{182}Ta	31.7378(7)	10.7181(9)	42.4559(11)		
	^{161}Tb	48.91562(6)	6.4642(4)	42.4518(4)		
	^{233}U	54.7038(7)	12.2510(4)	42.4528(15)		
53	^{210}Pb	46.539(1)	7.0708(12)	53.610(2)	53.6107(11)	0.20
	^{233}U	54.7038(7)	1.0927(3)	53.6111(14)		
54	^{210}Pb	46.539(1)	8.1640(5)	54.703(1)	54.7038(14) ^a	9.63
	^{241}Am	59.5412(2)	4.8374(2)	54.7036(4)		
	^{169}Yb	63.1208(1)	8.4202(7)	54.7006(7)		
	^{161}Tb	48.91562(6)	5.7890(3)	54.7046(3)		
66	^{241}Am	59.5412(2)	6.5777(6)	66.1189(6)	66.1184(6)	1.69
	^{169}Yb	63.1208(1)	2.9968(8)	66.1176(8)		
67	^{241}Am	59.5412(2)	8.4048(12)	67.9460(12)	67.9460(5)	0.00
	^{169}Yb	63.1208(1)	4.8252(10)	67.9460(10)		
	^{233}U	66.1184(6)	1.8276(4)	67.9460(7)		
70	^{169}Yb	63.1208(1)	7.1614(16)	70.2822(16)	70.2813(13)	0.90
	^{233}U	71.8159(20)	1.5362(6)	70.2797(21)		
71	^{241}Am	59.5412(2)	12.2719(10) ^N	71.8131(10)	71.8159(20) ^a	9.97
	^{169}Yb	63.1208(1)	8.6947(4)	71.8155(4)		
	^{161}Tb	74.56711(10)	2.7492(6)	71.8179(6)		
74	^{182}Ta	84.6808(3)	10.1273(24)	74.5535(24)	74.5390(40) ^a	13.5
	^{170}Tm	84.2551(2)	9.7172(6)	74.5379(6)		
	^{233}U	71.8159(20)	2.7242(3)	74.5401(20)		
	^{233}U	76.3507(27)	1.8099(8)	74.5408(28)		
76	^{182}Ta	84.6808(3)	8.3218(49)	76.359(5)	76.3507(27)	3.07
	^{170}Tm	84.2551(2)	7.9053(16)	76.3498(16)		
83	^{170}Tm	84.2551(2)	1.2423(20)	83.0128(20)		

Table 20. (Continued)

²³³ U Line	Reference		Adopted ΔE_γ	γ -ray energy		
	Nuclide	Energy		Value	Average	χ_R^2
85	¹⁷⁰ Tm	84.2551(2)	1.1673(17)	85.4224(17)		
88	¹⁷⁰ Tm	84.2551(2)	4.2195(14)	88.4746(14)		
89	¹⁷⁰ Tm	84.2551(2)	5.6995(5)	89.9546(5)	89.9568(24) ^a	15.9
	¹⁸² Ta	84.6808(3)	5.2785(10)	89.9593(10)		
	²³³ U	93.3502(3)	3.3924(3)	89.9578(4)		
93	²³³ U	97.1344(3)	3.7842(1)	93.3502(3)		
97	¹⁸² Ta	100.1065(2)	2.9721(2)	97.1344(3)		
117	¹⁸² Ta	113.6723(2)	3.4905(12)	117.1628(12)	117.1628(9)	0.00
	²³³ U	118.9721(15)	1.8092(2)	117.1629(15)		
118	¹⁸² Ta	113.6723(2)	5.2973(12)	118.9696(12)	118.9721(15)	5.99
	²³³ U	120.8194(7)	1.8464(2)	118.9730(7)		
120	¹⁶⁹ Yb	130.5237(0)	9.7036(13)	120.8201(13)	120.8194(7)	0.43
	²³³ U	135.3393(5)	14.5202(6)	120.8191(8)		
123	¹⁶⁹ Yb	130.5237(0)	6.6370(13)	123.8867(13)	123.8860(7)	0.43
	²³³ U	120.8194(7)	3.0663(4)	123.8857(8)		
135	¹⁶⁹ Yb	130.5237(0)	4.8156(5)	135.3393(5)		
139	¹⁶⁹ Yb	130.5237(0)	9.2135(50)	139.7372(50)	139.7278(45)	4.37
	²³³ U	135.3393(5)	4.3863(23)	139.7256(24)		
146	¹⁶⁹ Yb	130.5237(0)	15.8228(25)	146.3465(25)	146.3462(6)	0.06
	¹⁸² Ta	152.4308(3)	6.0848(7)	146.3460(8)		
	¹⁸² Ta	156.3874(3)	10.0410(8)	146.3464(9)		
164	¹⁶⁹ Yb	177.2140(1)	12.6905(10)	164.5235(10)	164.5240(5)	0.27
	¹⁸² Ta	152.4308(3)	12.0932(8)	164.5240(9)		
	¹⁸² Ta	156.3874(3)	8.1372(9)	164.5246(9)		
	¹⁸² Ta	179.3948(3)	14.8711(9)	164.5237(9)		
169	¹⁶⁹ Yb	177.2140(1)	8.2123(54)	169.002(5)		
170	¹⁶⁹ Yb	177.2140(1)	6.4049(24)	170.8091(24)		
174	¹⁶⁹ Yb	177.2140(1)	3.0221(20)	174.1919(20)		

Table 20. (Continued)

²³³ U Line	Reference		Adopted ΔE_γ	γ -ray energy		
	Nuclide	Energy		Value	Average	χ_R^2
187	¹⁶⁹ Yb	177.2140(1)	10.7538(6)	187.9678(6)	187.9669(3)	1.07
	¹⁶⁹ Yb	197.9579(1)	9.9913(5)	187.9666(5)		
	¹⁸² Ta	179.3948(3)	8.5716(7)	187.9664(8)		
	¹⁸² Ta	198.3530(3)	10.3865(8)	187.9665(9)		
193	¹⁶⁹ Yb	197.9579(1)	4.4549(47)	193.503(5)		
208	¹⁶⁹ Yb	197.9579(1)	10.2221(6)	208.1800(6)	208.1795(7)	1.73
	¹⁸² Ta	198.3530(3)	9.8253(10)	208.1783(10)		
	¹⁸² Ta	222.1099(3)	13.9366(54)	208.1733(54)		
217	¹⁸² Ta	222.1099(3)	4.9583(15)	217.1516(15)	217.1519(20)	7.52
	¹⁸² Ta	229.3220(6)	12.1734(10)	217.1486(12)		
	²³³ U	208.1795(7)	8.9754(8)	217.1549(11)		
240	¹⁸² Ta	229.3220(6)	11.0473(30)	240.369(3)	240.3719(17)	1.24
	²³³ U	245.3498(11)	4.9769(14)	240.3729(18)		
245	¹⁸² Ta	229.3220(6)	16.0249(40)	245.347(4)	245.3498(11)	0.52
	²³³ U	248.7242(10)	3.3742(7)	245.3500(12)		
248	¹⁸² Ta	264.0755(3)	15.3513(10)	248.7242(10)		
261	¹⁸² Ta	264.0755(3)	2.1178(39)	261.958(4)		
268	¹⁸² Ta	264.0755(3)	4.5992(21)	268.6747(21)		
274	¹⁸² Ta	264.0755(3)	10.6580(14)	274.7335(14)	274.7347(13)	1.61
	²³³ U	278.1080(9)	3.3718(13)	274.7362(16)		
278	¹⁸² Ta	264.0755(3)	14.0326(11)	278.1081(11)	278.1080(9)	0.02
	²³³ U	291.3561(9)	13.2483(14)	278.1078(17)		
288	²³³ U	278.1080(9)	9.9209(9)	288.0289(13)	288.0292(9)	0.07
	²³³ U	291.3561(9)	3.3267(10)	288.0294(13)		
291	¹⁶⁹ Yb	307.7376(1)	16.3815(9)	291.3561(9)		
293	¹⁶⁹ Yb	307.7376(1)	13.7428(93)	293.995(9)		
302	¹⁶⁹ Yb	307.7376(1)	4.7471(44)	302.990(4)		
317	¹⁶⁹ Yb	307.7376(1)	9.4318(11)	317.1694(11)	317.1689(15)	2.04

Table 20. (Continued)

²³³ U Line	Reference		Adopted ΔE_γ	γ -ray energy		
	Nuclide	Energy		Value	Average	χ_R^2
	¹⁵² Eu	344.2811(19)	27.1168(28)	317.1643(34)		
320	¹⁶⁹ Yb	307.7376(1)	12.8102(11)	320.5478(11)	320.5471(13)	2.32
	¹⁵² Eu	344.2811(19)	23.7410(28)	320.5401(34)		
	²³³ U	317.1689(15)	3.3782(4)	320.5471(16)		
323	¹⁵² Eu	344.2811(19)	20.9013(27)	323.3798(33)	323.3806(14)	0.07
	²³³ U	320.5471(13)	2.8337(9)	323.3808(16)		
328	²³³ U	323.3806(14)	5.3779(44)	328.758(5)		
336	¹⁵² Eu	344.2811(19)	7.6639(19)	336.6172(27)	336.6195(16)	1.06
	²³³ U	323.3806(14)	13.2400(13)	336.6206(19)		
365	¹⁵² Eu	344.2811(19)	21.5394(29)	365.8205(35)		
367	¹⁵² Eu	344.2811(19)	23.5095(95)	367.791(10)	367.795(8)	0.41
	²³³ U	365.8205(35)	1.9809(117)	367.801(12)		

- ^a Calculated uncertainty has been increased by a factor of 2.0 due to the large χ_R^2 value.
- ^b Known doublet (see text), so the measured peak energy may vary with detector resolution.

5. ENERGY RELATIONSHIPS IN ^{229}Th

There are several interesting relationships among the energies of the γ rays in ^{229}Th to be discussed before the energy of first excited state is determined. First, we should comment on the cascade-crossover relationship involving the first three levels in the ground-state rotational band. As shown in Figure 1, this includes the 42- and 54-keV γ -ray cascade and the 97-keV crossover involving the levels at 0, 42, and 97 keV. In our previous article² we did not use this relationship because there was reasonable evidence that the observed 42-keV peak had two components. γ - γ coincidence data¹ imply the existence a weaker 42-keV γ ray between the levels at 71.8 and 29.1 keV, although this γ ray was not directly observed. From the level energies deduced without use of the 42-keV lines, the energy of the weaker γ ray is 200 eV higher than that of the ground-state γ ray. Previous attempts to observe the doublet nature of the 42-keV peak were unsuccessful. However, with improved resolution of the OR-16 detector, we have been able to observe a difference between the shape of the 42-keV peak and those of the neighboring strong peaks. This is illustrated in Figure 15, where 15A shows that the 59-keV peak from ^{241}Am is well fit by a Gaussian function and 15B shows that the 42-keV peak deviates from a Gaussian function in a manner which implies the existence of a second component on the high-energy side of the main peak. Since both of these peaks are in the same spectrum and have similar peak-to-continuum ratios, we are satisfied that the 42-keV peak is a doublet, and that the weaker component has the higher energy, as expected.

The remaining analysis depends critically on deductions related to the relative intensities of certain possible γ rays in the ^{229}Th level scheme. In order to be able to deduce the value of Δ , the energy of the first excited state, the γ rays that populate the levels at 0 or Δ keV must populate only one of them with significant intensity. In Reference 23, this question has been discussed in some detail, and it has been deduced that these lines are sufficiently pure (possible "contaminants" weaker by factors of the order of 10) for the purposes of this analysis. These arguments are not repeated here.

As shown in Figure 1, there are a few relationships among the γ -ray energies reported in Table 20 that can be used to provide checks on the quality or consistency of these results. In the relationships (see Section 1) that we will use to determine the energy of the ^{229}Th first excited state, the first two require that

$$E(71) - E(67) = E(29) - E(25).$$

(a)

(b)

Figure 15. Gaussian fits to peaks at 59 keV, a singlet from ^{241}Am , and 42 keV, a doublet from ^{233}U .

From Table 18a and 18b, these two differences, as directly determined, are 3.8685(4) and 3.8701(4) keV, respectively. Although these two values differ by 0.0016(6) keV, more than the associated uncertainty, this is not unreasonable considering the weakness of the 25- and 67-keV peaks as shown in Figures 3, 4, and 6. (The 29- and 71-keV lines are the ones that, from spin-parity considerations, potentially could be doublets feeding the 0- and Δ -keV levels.)

There are two cascade-crossover combinations that can also be used to test the consistency of the ^{233}U γ -ray energies. These combinations are shown in Figure 1. The corresponding differences between the cascade and crossover energies are shown in Table 21. Both of these values are consistent with zero, and their uncertainties are dominated by that of the 29-keV line, which is 0.0030 keV. If one assumes that these relations are actually zero, one can use them to deduce the energy of this γ ray. The two resulting values are 29.1850(20) and 29.1847(7) keV, and their average is 29.1847(7) keV. Then, from the 25U - 29U difference in Table 18a, one can deduce an energy of 25.3146(8) keV for the 25-keV line.

From the γ -ray energies that we have determined, there are four energy combinations that can be used to obtain the energy of the first-excited level in ^{229}Th . These relationships are shown in Figure 1. The energies from these combinations are shown in Table 22. All four values show a positive energy for this level, in contrast to the result of Reference 2 where two of the three combinations gave negative values. For the first two relations, two values are given - one from the values in Table 20 for $E_\gamma(25)$ and $E_\gamma(29)$ and the other from the values deduced in the previous paragraph for these two lines.

Several average values are given in Table 22. The unweighted averages are given to show the influence of the weight resulting from the smaller uncertainty for the value of 0.0036(6) from the third relation. The next line gives the weighted averages based on the uncertainties given.

Table 21. Difference between crossover and cascade energies.

<u>γ rays (keV)</u>	<u>Difference (keV)</u>
217-187-29	+0.0004(36)
164-135-29	+0.0001(31)

Table 22. Energy of ^{229}Th first excited level, Δ .

γ rays (keV)	Δ (keV)			
	Set #1 ^a		Set #2 ^b	
	Energy	χ^2_{R}	Energy	χ^2_{R}
97-71-25	+0.0069(22)		+0.0029(22)	
97-67-29	+0.0028(31)		+0.0027(9)	
[148-146] - [118-117]	+0.0036(6)			
[148-146] - [76-74]	+0.0029(10)			
Unweighted average	+0.0040(10)		+0.0030(2)	
Weighted average	+0.0036(5)	0.93	+0.0032(4)	0.29

^a The energies for $E_{\gamma}(25)$ and $E_{\gamma}(29)$ are from Table 20.

^b The energies for $E_{\gamma}(25)$ and $E_{\gamma}(29)$ are derived in the text from the computed 217-187 and 164-135 differences for the 29 line and from the measured 29-25 (Table 18a) difference for the 25 line.

6. CONCLUSION

From the results in Table 22, we have adopted an energy of 0.0035(10) keV for the first excited state of ^{229}Th . This value has been taken from the two averages in the last line of the table, with the uncertainty increased by a factor of 2.0 to allow for any unidentified systematic errors.

REFERENCES

1. L. A. Kroger and C. W. Reich, Nucl. Phys. **A259**, 29 (1976); L. A. Kroger, Ph. D. thesis, University of Wyoming, 1971 (unpublished).
2. C. W. Reich and R. G. Helmer, Phys. Rev. Lett. **64**, 271 (1990).
3. V. F. Strizhov and E. V. Tkalya, Sov. Phys. JETP **72**, No. 3, 387 (1991).
4. E. V. Tkalya, Sov. Phys. JETP Lett. **55**, 211 (1992).
5. F. F. Karpeshin, I. M. Band, M. B. Trzhaskowskaya and B. A. Zon, Phys. Lett. **B282**, 267 (1992).
6. E. V. Tkalya, contribution to Proc. III Int. Symposium on Weak and Electromagnetic Interactions, WEIN-92 (Dubna, 1992).
7. E. V. Tkalya, Yad. Fiz. **56**, 2881 (1992).
8. S. Wycech and J. Żylicz, Frontier Topics in Nuclear and Astrophysics (IOP Publishing Ltd., Bristol 1992) p. 365
9. R. G. Helmer, R. J. Gehrke and R. C. Greenwood, Nucl. Instr. and Meth. **123**, 51 (1975).
10. R. G. Helmer, P. H. M. Van Assche, and C. van der Leun, Atomic Data and Nucl. Data Tables **24**, 39 (1979).
11. E. G. Kessler, R. D. Deslattes, A. Henins, and W. C. Sauder, Phys. Rev. Lett. **40**, 171 (1978).
12. R. G. Helmer, accepted for publication in Nucl. Instr. and Meth.
13. R. G. Helmer, A. J. Caffrey, R. J. Gehrke and R. C. Greenwood, Nucl. Instr. and Meth. **188**, 671 (1981).
14. B. Jeckelmann, W. Beer, G. De Chambrier, O. Elsenhans, K. L. Giovanetti, P. F. A. Goudsmit, H. J. Leisi, and A. Rüetschi, Nucl. Instr. and Meth. **A241**, 191 (1985).
15. R. D. Deslattes, E. G. Kessler, W. C. Sauder, and A. Henins, Ann. Phys. **129**, 378 (1980).
16. E. K. Warburton and D. E. Alburger, Nucl. Instr. and Meth. **A253**, 38 (1986).
17. G. C. Nelson and B. G. Saunders, Nucl. Instr. and Meth. **84**, 90 (1970).

18. R. W. Jewell, W. John, R. Massey and B. G. Saunders, Nucl. Instr. and Meth. **62**, 68 (1968).
19. E. Browne and R. B. Firestone, Table of Radioactive Isotopes (John Wiley & Sons, New York, 1986).
20. Y. A. Ellis-Akovali, Nucl. Data Sheets **44**, 407 (1985).
21. W. Gongqing, Nucl. Data Sheets **51**, 577 (1987).
22. R. G. Helmer, Nucl. Data Sheets **59**, 1, (1990).
23. C. W. Reich, Izv. Akad. Nauk SSSR, Ser. Fiz. **55**, 878 (1991).

Table 4. Measured γ -ray energy differences between lines of $^{172}\text{Hf}+^{172}\text{Lu}$ and ^{241}Am and lines of ^{161}Tb and ^{241}Am .

Detector	Spectral code	$^{23}\text{Hf}-^{26}\text{Am}$	$^{25}\text{Tb}-^{26}\text{Am}$	$^{48}\text{Tb}-^{59}\text{Am}$	$^{59}\text{Am}-^{74}\text{Tb}$
PG-16	CN10			10.6254(2)	15.0259(3)
	CN11			10.6256(2)	15.0266(4)
	CN12			10.6256(2)	15.0261(3)
	CN13			10.6260(3)	15.0267(5)
	CN14			10.6256(3)	15.0266(4)
	CN15			10.6258(2)	15.0275(4)
	CN16			10.6259(4)	15.0274(4)
	CN17	2.4103(5)			
	CN19	2.4119(5)			
	CN20	2.4104(6)			
	CN21	2.4111(5)			
	CN23	2.4108(4)			
	CN24	2.4112(5)			
	CN25	2.4101(4)			
OR-16	DN1		0.6927(15)	10.6268(9)	15.0258(15)
	DN2		0.6939(8)	10.6269(7)	15.0253(11)
	DN3		0.6923(14)		15.0236(21)
	DN4		0.6915(11)	10.6266(14)	15.0221(17)
	DN5		0.6927(7)	10.6266(7)	15.0245(9)
	DN6		0.6927(5)	10.6280(6) ^a	15.0245(9)
	DN7		0.6927(7)	10.6290(8) ^a	15.0236(13)
	DN8		0.6928(5)	10.6275(5) ^a	15.0229(7)
	DN9		0.6950(5)	10.6315(6) ^a	15.0237(7)
	DN10		0.6926(6)	10.6300(7) ^a	15.0294(11)
Si(Li)	EN1	2.4126(8)			
	EN2				
	EN3	2.4114(10)			
	EN4		0.6925(7)	10.6274(10)	15.0237(24)
	EN5		0.6935(10)	10.6249(15)	15.0230(35)
	EN6		0.6926(6)	10.6259(10)	15.0276(28)
	EN7		0.6938(6)	10.6269(11)	15.0253(27)
	EN8		0.6935(4)	10.6265(7)	15.0264(15)
	EN18		0.6928(3)	10.6265(6)	15.0268(15)

Table 4. (Continued)

Detec- tor	Spectral code	23Hf-26Am	25Tb-26Am	48Tb-59Am	59Am-74Tb
	EN19		0.6938(4)	10.6266(8)	15.0258(23)
	EN20		0.6939(4)	10.6254(6)	15.0276(17)
	EN21			10.6254(3)	15.0250(6)
	EN22			10.6260(9)	15.0260(23)
	EN23			10.6242(8)	15.0262(15)
	EN24			10.6247(10)	15.0222(17)
	EN25			10.6257(7)	15.0242(11)
	EN26			10.6244(12)	15.0264(16)
	EN27	2.4135(6)			
	EN28	2.4150(8)			
	EN29	2.4141(6)			
	EN30	2.4141(7)			
	EN31	2.4150(6)			

The following are weighted averages and the associated χ_R^2 .

PG-16	2.4108(2) 1.72		10.6256(1) 0.67	15.0266(2) 2.87
OR-16		0.6932(3) 2.25	10.6267(4) 0.04	15.0244(6) 3.35
Si(Li)	2.4139(4) 2.46	0.6933(2) 1.46	10.6257(2) 1.15	15.0253(4) 0.81
All data	2.4118(9) ^b 9.33	0.6933(2) 1.82	10.6257(1) 1.08	15.0261(2) 3.40

^a Not used in average due to grow-in of an interfering peak in the ¹⁶¹Tb source.

^b The weighted average of these 14 differences gives an internal uncertainty of 0.00015 keV and an external uncertainty of 0.00045 keV. The value given here is the latter value increased by an additional arbitrary factor of 2.0 due to the discrepancy between the averages from the individual detectors.

Table 5. Measured γ -ray energy differences between lines from ^{161}Tb and ^{172}Hf and between two lines from ^{161}Tb .

Detec- tor	Spectral code	<u>23Hf-25Tb</u>	<u>48Tb-57Tb</u>	<u>74Tb-78Hf</u>	<u>74Tb-81Hf</u>	<u>74Tb-90Hf</u>
PG-16	CN1	1.7197(6)	8.2820(17)	4.1757(11)	7.1871(16)	16.0805(17)
	CN2	1.7177(7)	8.2749(20)	4.1767(11)	7.1854(15)	16.0792(16)
	CN3	1.7186(3)	8.2830(10) ^a	4.1760(6)	7.1841(7)	16.0799(9)
	CN4	1.7180(4)	8.2758(12)		7.1868(9)	16.0791(10)
	CN5	1.7183(6)		4.1770(10)	7.1844(14)	16.0813(16)
	CN6	1.7176(9)		4.1717(14)	7.1793(17)	16.0713(19)
	CN7	1.7172(4)	8.2692(22)	4.1826(9) ^a	7.1826(9)	16.0755(10)
	CN8	1.7182(6)	8.2787(41)	4.1721(11)	7.1995(13) ^a	16.0688(15) ^a
	CN9	1.7134(9) ^a		4.1722(16)	7.1806(16)	
	CN11		8.2757(7)			
	CN12		8.2764(4)			
	CN13		8.2765(9)			
	CN14		8.2754(9)			
	CN15		8.2774(7)			
	CN16		8.2777(7)			
Si(Li)	EN9	1.7180(3)		4.1755(20)	7.1819(29)	16.0816(45)
	EN10	1.7178(3)		4.1777(27)	7.1868(40)	16.0839(40)
	EN11	1.7187(4)	8.2798(62)	4.1817(42)	7.1909(50)	16.0864(58)
	EN12	1.7188(4)		4.1796(20)	7.1884(35)	16.0801(35)
	EN13	1.7189(3)		4.1808(21)	7.1867(32)	16.0877(37)
	EN14	1.7192(3)		4.1737(23)	7.1788(29)	16.0715(36)
	EN15	1.7189(4)		4.1706(34)	7.1812(40)	16.0837(44)
	EN16	1.7192(4)		4.1721(36)	7.1827(42)	16.0768(46)
	EN17	1.7179(4)		4.1802(38)	7.1835(44)	16.0896(49)
	EN22		8.2757(7)			
	EN23		8.2764(4)			
	EN24		8.2765(9)			
	EN25		8.2754(9)			
	EN26		8.2774(7)			
	EN27		8.2777(7)			

Table 5. (Continued)

The following are weighted averages and the associated χ_R^2 .

PG-16	1.7182(3) 2.22	8.2765(4) 3.02	4.1752(8) 4.13	7.1842(8) 4.12	16.0784(11) 5.12
Si(Li)	1.7186(2) 2.78	8.2758(6) 0.58	4.1771(12) 1.87	7.1839(13) 1.13	16.0817(14) 1.97
All data	1.7184(2) 2.57	8.2764(3) 2.05	4.1755(6) 2.92	7.1841(6) 2.37	16.0787(8) 3.44

^a Value is an outlier, more than 5σ from the average, and was not used in the averages.

Table 6. Measured γ -ray energy differences between pairs of lines from $^{172}\text{Hf}+^{172}\text{Lu}$.

Detec- tor	Spectral code	78Hf-81Hf	81Hf-90Hf	90Hf-112Hf	112Hf-114Hf	114Hf-122Hf	122Hf-125Hf
PG-16	CN1			22.1373(38)	1.2852(46)	8.8619(62)	2.8994(55)
	CN2			22.1334(42)	1.2859(51)	8.8580(60)	2.8986(53)
	CN3			22.1403(19)	1.2830(21)	8.8590(28)	2.9014(25)
	CN4			22.1366(23)	1.2834(26)		
	CN5			22.1450(37)	1.2779(44)		
	CN6			22.1308(34)	1.2927(39)	8.8584(49)	2.8957(43)
	CN7			22.1262(21)	1.2857(24)	8.8535(29)	2.8987(26)
	CN8			22.1123(28) ^a	1.2797(31)	8.8526(43)	2.8938(40)
	CN9				1.2777(36)		2.8898(40)
	CN17	3.0094(10)	8.8924(20)				2.8993(35)
	CN18						
	CN19	3.0078(8)	8.8903(19)	22.1300(65)	1.2825(24)	8.8548(41)	2.8953(24)
	CN20	3.0083(10)	8.8941(22)	22.1304(73)	1.2843(25)	8.8561(49)	2.8969(32)
	CN21	3.0103(9)	8.8878(20)	22.1219(65)	1.2821(24)	8.8497(42)	2.8983(25)
	CN22						
	CN23	3.0090(7)	8.8911(18)	22.1235(63)	1.2819(17)	8.8548(39)	2.8960(22)
	CN24	3.0078(8)	8.8898(19)	22.1256(64)	1.2828(19)	8.8541(40)	2.8945(23)
	CN25	3.0087(7)	8.8896(19)	22.1244(63)	1.2811(17)	8.8503(38)	2.8947(21)
Si(Li)	EN1	3.0103(38)	8.8930(76)				
	EN3	3.0188(63)	8.9153(103)				
	EN27	3.0083(43)	8.8966(63)				
	EN28	3.0068(64)					
	EN29	3.0100(48)	8.8844(65)				
	EN30	3.0003(58)	8.8844(73)				
	EN31	3.0025(45)	8.8928(63)				

Table 6. (Continued)

The following are weighted averages and the associated χ_R^2 .

PG-16	3.0087(3)	8.8906(7)	22.1340(19)	1.2827(7)	8.8549(12)	2.8966(8)
	0.87	1.09	0.98	3.96	1.04	0.71
Si(Li)	3.0102(20)	8.8924(37)				
	1.64	1.62				
All data	3.0088(4)	8.8907(8)				
	1.35	1.19				

^a Value is an outlier, more than 5 σ from the average, and was not used in the averages.

Table 10. Measured γ -ray energy differences for ^{241}Am reference lines of 26 and 59 keV. Lines not otherwise labeled are from ^{233}U .

Detector	Spectral code	25 - 26Am	26Am - 29	54 - 59Am	59Am - 66	59Am - 67	59Am - 71
OR-12	none						
PG-21	none						
PG-16	C9	1.0437(127)	2.8424(27)	4.8372(6)	6.5744(24)		12.2724(13)
	C10	1.0289(136)	2.8444(30)	4.8401(9)	6.5783(29)		12.2705(10)
	C11	1.0316(198)	2.8522(46)	4.8378(9)	6.5786(46)		12.2724(21)
	C12	1.0299(53)	2.8449(16)	4.8403(6)	6.5775(16)		
OR-16	D1	1.0382(139)	2.8421(36)				
	D2	1.0385(45)	2.8383(19)				
	D3	1.0358(95)	2.8389(26)				
	D4	1.0179(88)	2.8380(16)			8.4052(24)	
	D8	1.0274(29)	2.8422(20)	4.8378(5)	6.5787(11)	8.4018(38)	
	D9	1.0292(30)	2.8387(21)	4.8377(5)	6.5764(15)	8.4041(31)	
	D10	1.0298(23)	2.8415(23)	4.8379(5)	6.5769(13)	8.4017(24)	
	D11	1.0316(45)	2.8401(17)	4.8382(4)	6.5784(11)	8.4083(55)	
	D23	1.0325(40)	2.8387(16)	4.8369(6)	6.5799(26)	8.4068(43)	
	D24	1.0324(26)	2.8402(12)	4.8368(5)	6.5762(23)	8.4090(48)	
	D25	1.0332(33)	2.8384(15)	4.8374(6)	6.5738(26)	8.4050(39)	
	D26	1.0269(28)	2.8390(14)	4.8367(5)	6.5816(23)		
	E1	1.0378(19)	2.8387(9)	4.8366(8)		8.4062(138)	
	E2	1.0387(16)	2.8387(10)	4.8383(11)	6.5770(40)	8.4212(93)	
	E3	1.0336(16)	2.8405(9)	4.8351(7)	6.5756(40)	8.4177(90)	
	E4	1.0370(18)	2.8387(9)	4.8368(7)			
Si(Li)							

Table 10. (Continued)

The following are weighted averages and the associated χ^2 values.					
PG-16	1.0316(45) 0.35	2.8448(13) 1.15	4.8388(8) 5.56	6.5770(12) 0.53	12.2719(10) 0.37
PG-16 revised			4.8388(8) ^a 1.39		
OR-16	1.0304(10) 0.95	2.8394(5) 0.54	4.8375(2) 1.40	6.5778(8) 2.67	8.4043(12) 0.52
Si(Li)	1.0367(12) 1.89	2.8392(5) 0.98	4.8364(6) 2.27	6.5763(28) 0.06	8.4170(59) 0.41
All data	1.0340(9) 2.09	2.8397(4) 1.68	4.8376(3) 3.64	6.5777(6) 1.62	8.4048(12) 0.89
All data revised			4.8374(2) ^a 2.10		

^a With all PG-16 uncertainties multiplied by 2.0 due to inconsistency of these values.

Table 11. Measured γ -ray energy differences from the ^{152}Eu reference line of 344 keV.

Detector	Spectral code	317 - 344Eu	320 - 344Eu	323 - 344Eu	336 - 344Eu	344Eu - 365	344Eu - 367
OR-12	none						
PG-21	B5	27.1081(37)	23.7335(38)	20.8981(64)	7.6661(59)		
	B6	27.1177(12)	23.7420(14)	20.8979(29)	7.6648(28)		
PG-16	C15			20.9094(46)	7.6675(57)	21.5344(54)	
	C16				7.6545(72)	21.5432(66)	
	C17				7.6651(61)	21.5437(66)	23.5019(135)
	C18				7.6589(90)	21.5411(68)	23.5168(133)
	C19			20.9042(54)	7.6603(69)	21.5369(72)	
OR-16	none						

The following are weighted averages and the associated χ^2_r values.

PG-21	27.1168(28) 6.09	23.7410(28) 4.41	20.8979(26) 0.00	7.6650(25) 0.04		
PG-16			20.9072(35) 0.54	7.6623(30) 0.61	21.5394(29) 0.45	23.5095(95) 0.62
All data			20.9013(27) 1.67	7.6639(19) 0.49		

Table 12. Measured γ -ray energy differences from the ^{210}Pb reference line of 46 keV.

<u>Detector</u>	<u>Spectral code</u>	<u>42 - 46Pb</u>	<u>46Pb - 53</u>	<u>46Pb - 54</u>
OR-12	none			
PG-21	B1	4.0772(12)		8.1612(16)
	B2	4.0772(19)		8.1657(28)
	B3	4.0776(10)		8.1652(15)
	B4	4.0799(13)		8.1658(16)
PG-16	none			
OR-16	D19	4.0869(12)	7.0706(25)	8.1635(17)
	D20	4.0860(13)	7.0711(23)	8.1647(12)
	D21	4.0856(12)	7.0707(27)	8.1643(16)
	D22	4.0864(11)	7.0708(20)	8.1631(10)
The following are weighted averages and the associated χ^2 values.				
PG-21	4.0780(6)		8.1643(11)	
	0.97		1.75	
OR-16	4.0862(6)	7.0708(12)	8.1638(6)	
	0.21	0.01	0.39	
All data	4.0823(16)		8.1640(5)	
	13.57		0.94	
Adopted	4.0862(6)			

Table 13. Measured γ -ray energy differences from the ^{170}Tm reference line of 84 keV.

Detector	Spectral code	74 - 84Tm	76 - 84Tm	83 - 84Tm	84Tm - 85	84Tm - 88	84Tm - 89
OR-12	A13	9.7057(36)	7.9202(91)				5.6835(23)
	A14	9.7006(27)	7.9097(49)				5.6814(19)
	A15	9.6722(54)	7.9216(153)				5.6886(24)
	A16	9.6773(34)	7.8734(109)				5.6928(16)
	A17	9.6762(35)	7.9010(93)				5.6931(18)
PG-21	none						
PG-16	C24	9.7156(15)	7.9024(25)	1.2358(48)	1.1680(42)		5.7009(11)
	C25	9.7160(16)	7.9038(28)	1.2449(65)	1.1669(999)		5.6991(12)
	C26	9.7182(17)	7.8994(26)	1.2477(51)	1.1661(39)		5.7002(13)
	C27	9.7158(17)	7.9057(27)	1.2445(55)	1.1643(41)		5.6996(14)
OR-16	D12	9.7194(20)	7.9068(38)	1.2472(53)		4.2193(36)	5.6971(18)
	D13	9.7190(23)	7.9070(38)	1.2375(70)	1.1702(44)	4.2172(30)	5.6994(21)
	D14	9.7180(13)	7.9110(21)	1.2394(46)	1.1679(28)	4.2203(17)	5.6989(11)
The following are weighted averages and the associated χ^2 values.							
OR-12		9.6896(67) 18.68	7.9068(65) 3.31				5.6887(25) 8.10
PG-16		9.7163(8) 0.53	7.9027(13) 1.00	1.2427(28) 1.08	1.1661(23) 0.20		5.7000(6) 0.45
OR-16		9.7185(10) 0.20	7.9094(17) 0.72	1.2417(31) 0.84	1.1686(24) 0.19	4.2195(14) 0.41	5.6986(9) 0.46
All data		9.7133(33) 32.11	7.9055(16) 2.54	1.2423(20) 0.83	1.1673(17) 0.28		5.6968(16) 14.08
All data revised ^a		9.7172(6) 0.82	7.9053(16) 2.42				5.6995(5) 0.68

^a With OR-12 data omitted due to its inconsistency with other data.

Table 14. Measured γ -ray energy differences from the ^{161}Tb reference lines 48 and 74 keV.

Detector	Spectral code	42 - 48Tb	48Tb - 54	71 - 74Tb
PG-16	C30		5.7900(7)	2.7492(15)
	C31	6.4643(9)	5.7892(9)	2.7459(18)
	C32	6.4632(10)	5.7885(10)	2.7454(18)
	C33	6.4626(6)	5.7883(5)	2.7489(8)
	C34	6.4618(7)	5.7885(6)	2.7519(11)
	C35	6.4640(7)	5.7895(7)	2.7475(13)
	C36	6.4632(7)	5.7889(6)	2.7519(12)
	C37	6.4594(7) ^a	5.7881(6)	2.7504(11)
Si(Li)	E12	6.4659(9)	5.7905(8)	2.7448(19)
	E13	6.4635(7)	5.7892(8)	2.7535(30)
	E14	6.4657(9)	5.7884(12)	2.7452(42)
	E15	6.4645(10)	5.7927(12)	2.7491(46)
	E16	6.4624(9)	5.7895(11)	2.7486(35)
	E17	6.4634(8)	5.7883(9)	2.7465(24)
	E18	6.4647(11)	5.7922(11)	2.7461(28)
	E19	6.4641(118)	5.7886(120)	2.7520(85)
	E20		5.7867(10)	2.7581(37)

The following are weighted averages and the associated χ^2 values.

PG-16	6.4625(6) 5.12	5.7888(2) 0.98	2.7495(7) 3.08
PG-16 revised	6.4631(4) ^a 1.52		
SiLi)	6.4642(4) 1.81	5.7896(6) 3.16	2.7480(14) 1.91
All data	6.4632(4) 4.18	5.7890(3) 2.23	2.7492(6) 2.42
All data revised	6.4636(3) ^a 2.03		

^a One PG-16 value is an outlier, more than 5σ from the average, and was not used in the averages.

Table 15. Measured γ -ray energy differences from the ^{172}Hf reference lines of 23 and 78 keV.

<u>Detector</u>	<u>Spectral code</u>	<u>$^{23}\text{Hf} - 25$</u>	<u>$^{23}\text{Hf} - 29$</u>	<u>$^{71} - ^{78}\text{Hf}$</u>
PG-16	C28			6.9390(14)
	C29			6.9380(22)
	C39		5.2485(19)	6.9311(15)
	C40		5.2514(21)	6.9260(18)
	C41		5.2480(9)	6.9293(16)
	C42		5.2523(24)	6.9315(13)
	C43		5.2465(26)	6.9329(17)
	C44		5.2518(21)	6.9324(14)
OR-16	D31	1.3796(21)		
	D33	1.3850(16)	5.2531(16)	
Si(Li)	E9	1.3761(42)		6.9428(41)
	E10	1.3787(22)		6.9368(30)
	E11	1.3785(18)		6.9356(27)
	E21	1.3787(23)	5.2539(8)	6.9253(28)
	E22	1.3769(20)	5.2548(9)	6.9367(35)
	E23	1.3730(19)	5.2538(9)	6.9349(24)
	E24	1.3714(27)	5.2549(14)	6.9324(34)
	E25	1.3676(29)	5.2527(8)	6.9299(31)
	E26	1.3688(25)	5.2537(10)	6.9298(28)
	E27		5.2527(11)	6.9286(32)
	E28		5.2538(10)	6.9314(28)

Table 15. (Continued)

The following are weighted averages and the associated χ^2 values.

PG-16	1.3672(100) 1.61	5.2490(8) 1.44
OR-16	1.3830(26) 4.18	
OR-16 revised	1.3830(26) ^a 1.90	
Si(Li)	1.3750(14) 3.51	5.2537(3) 0.67
Si(Li) revised	1.3750(14) ^b 1.59	
All data	1.3771(15) 5.39	5.2528(6) 3.74
All data revised	1.3769(16) 2.58	5.2489(9) 1.49
		6.9326(9) ^c 3.80

^a All OR-16 uncertainties multiplied by 1.50 due to internal inconsistency.

^b All Si(Li) uncertainties have been increased by 1.5 due to internal inconsistency.

^c Value not used; data show two distinct groups with averages of 6.9309(7) and 6.9383(10).

Table 16a. Measured γ ray energy differences from the ^{182}Ta reference lines from 31 to 264 keV.

Detector	Spectral code	29 - 31Ta	31Ta - 42	74 - 84Ta	76 - 84Ta	84Ta - 89	97 - 100Ta	113Ta - 117
OR-12	A1					5.2655(61)	2.9865(35)	3.4140(109)
	A2					5.2617(30)	2.9834(17)	3.4412(41)
	A3					5.2607(27)	2.9818(15)	3.4362(44)
	A4					5.2609(24)	2.9815(17)	3.4412(39)
	A5						2.9826(12)	
PG-21	none							
PG-16	C1					5.2905(88)	2.9750(36)	3.4897(149)
	C2					5.2760(32)	2.9729(14)	3.4936(56)
	C3					5.2748(11)	2.9721(5)	3.4870(17)
	C4					5.2764(19)	2.9719(7)	3.4943(30)
	C5					5.2762(12)	2.9728(7)	3.4925(18)
	C6					5.2770(12)	2.9719(5)	3.4908(19)
OR-16	D1	2.5474(39)	10.7187(35)			5.2810(33)	2.9738(17)	
	D2			10.1308(31)	8.3173(66)	5.2795(14)	2.9711(9)	
	D3	2.5493(17)	10.7170(17)	10.1252(38)	8.3298(103)	5.2828(17)	2.9716(8)	
	D4	2.5502(29)	10.7177(24)	10.1333(56)	8.3249(105)	5.2839(21)	2.9732(10)	
	D5	2.5497(18)	10.7183(20)	10.1309(40)		5.2846(18)	2.9710(9)	
	D6	2.5475(36)	10.7195(33)	10.1229(18)		5.2766(35)	2.9728(17)	
	D7	2.5437(22)	10.7194(22)	10.1402(43)		5.2830(19)	2.9727(12)	
The following are weighted averages and the associated χ^2 values.								
OR-12						5.2613(15)	2.9825(7)	3.4386(33)
						0.19	0.54	2.08
PG-16						5.2761(6)	2.9722(3)	3.4905(12)
						0.93	0.43	1.48
OR-16		2.5482(10)	10.7181(9)	10.1273(24)	8.3218(49)	5.2820(9)	2.9719(4)	
		1.17	0.20	3.70	0.58	1.51	0.93	
All data						5.2769(15)	2.9730(7)	3.4829(62)
						10.81	11.73	48.9
All data revised ^a						5.2785(10)	2.9721(2)	3.4905(12)
						4.31	0.66	1.48

^a All OR-12 values omitted due to poor resolution of these peaks.

Table 16b.

Detector	Spectral code	1131a - 118	146 - 1521a	146 - 1561a	1521a - 164	1561a - 164	164 - 1791a	1791a - 187
OR-12	A1				12.1084(35)	8.1575(60)	14.8600(51)	8.5734(70)
	A2				12.1037(23)		14.8605(27)	8.5720(28)
	A3					8.1524(23)	14.8644(24)	8.5701(24)
	A4				12.1033(18)	8.1587(21)	14.8608(21)	8.5785(25)
	A5						8.5729(10)	10.3851(22)
	A6						8.5692(15)	10.3841(20)
	A7						8.5698(15)	10.3907(22)
	A8						8.5708(13)	10.3886(18)
PG-21	none							
PG-16	C1		6.0769(113)		12.0934(126)	8.1403(127)	14.8655(146)	8.5827(221)
	C2		6.0862(35)	10.0454(44)	12.0978(43)	8.1385(51)	14.8675(57)	8.5632(90)
	C3		6.0875(15)	10.0422(16)	12.0917(16)	8.1370(18)	14.8703(19)	8.5699(32)
	C4		6.0867(18)	10.0436(26)	12.0922(24)	8.1353(30)	14.8739(33)	8.5731(50)
	C5		6.0842(14)	10.0433(18)	12.0957(14)	8.1367(18)	14.8687(20)	8.5769(30)
	C6		6.0859(11)	10.0407(17)	12.0923(14)	8.1374(19)	14.8706(25)	8.5728(35)
OR-16	D1	5.3015(70)	6.0726(63)	10.0310(66)	12.0951(29)	8.1410(37)	14.8711(35)	8.5852(65)
	D2	5.2977(29)	6.0813(24)	10.0354(33)	12.0866(31)	8.1283(36)	14.8796(34)	8.5826(99)
	D3	5.2972(26)	6.0835(25)	10.0418(30)	12.0992(42)	8.1431(44)	14.8721(48)	8.5805(102)
	D4	5.2999(11)	6.0821(31)	10.0382(34)	12.0984(39)	8.1428(41)	14.8679(45)	
	D5	5.2944(29)	6.0819(27)	10.0375(29)	10.0323(58)			
	D6	5.2987(28)	5.2905(61)	5.0774(54)	12.0962(33)	8.1383(38)	14.8730(37)	8.5744(84)
	D7		6.0821(26)	10.0399(32)				
The following are weighted averages and the associated χ^2 values.								
OR-12					12.1041(13) 0.87	8.1559(22) 2.08	14.8618(13) 0.59	8.5712(9) 1.79
PG-16			6.0859(7) 0.69	10.0424(9) 0.49	12.0929(8) 1.18	8.1369(10) 0.11	14.8701(11) 0.43	8.5730(17) 0.80
OR-16	5.2973(12) 0.58		6.0817(11) 0.55	10.0380(13) 0.78	12.0943(23) 2.26	8.1381(28) 2.66	14.8734(20) 1.33	8.5813(42) 0.35
All data			6.0848(7) 1.43	10.0410(8) 1.27	12.0953(13) 5.08	8.1418(24) 10.28	14.8680(14) 3.22	8.5716(7) ^b 1.42
All data revised ^a				12.0932(8) 1.50	8.1372(9) 1.15	14.8711(9) 1.0	8.5742(16) 0.94	10.3869(20) 0.62

^a All OR-16 values omitted due to inconsistency with other data.^b Adopted value.

Table 16c.

Detector	Spectral code	187 - 198Ta	198Ta - 208	208 - 222Ta	217 - 222Ta	217 - 229Ta	229Ta - 240	229Ta - 245
OR-12	A1	10.3735(93)	9.8499(92)		4.9419(76)	12.2055(100)		
	A2	10.3863(33)	9.8262(32)		4.9535(27)	12.1688(33)		
	A3	10.3872(30)	9.8237(30)		4.9488(23)	12.1668(29)		
	A4	10.3807(30)	9.8247(28)		4.9521(25)	12.1686(29)		
	A5	10.3851(22)	9.8247(22)		4.9497(14)	12.1761(17)		
	A6	10.3841(20)	9.8295(20)		4.9482(11)	12.1776(20)	11.0440(35)	16.0127(15)
	A7	10.3907(22)	9.8228(22)		4.9484(14)	12.1740(16)	11.0501(32)	16.0171(13)
	A8	9.8247(18)			4.9469(13)	12.1712(13)		
PG-21	none							
PG-16	C1	10.4114(276)	9.7661(309)		4.9708(205)	12.1765(251)		16.0254(142)
	C2	10.3948(105)	9.8186(123)		4.9647(82)	12.1814(107)		16.0278(37)
	C3	10.3912(39)	9.8191(39)		4.9586(25)	12.1759(27)		16.0309(54)
	C4	10.3910(69)			4.9567(47)	12.1756(51)		16.0293(48)
	C5	10.3851(44)	9.8225(45)		4.9583(30)	12.1714(40)		16.0086(45)
	C6	10.3840(40)	9.8303(39)		4.9561(30)	12.1729(33)		
OR-16	D2	10.3820(71)	9.8259(85)	13.9386(79)				16.0364(78)
	D3	10.3834(102)	9.8163(111)	13.9442(108)				
	D4	10.3755(105)						
	D7	10.3905(91)	9.8326(105)	13.9268(100)	4.9681(87)	12.1855(91)		
The following are weighted averages and the associated χ^2 values.								
OR-12		10.3864(12) 1.80	9.8256(12) 1.91		4.9487(6) 1.21	12.1731(15) 4.28	11.0473(30) 1.65	16.0152(22) 4.91
PG-16		10.3878(22) 0.69	9.8236(33) 1.99		4.9580(15) 0.32	12.1744(17) 0.31		16.0240(44) 3.91
OR-16		10.3831(44) 0.40	9.8253(57) 0.57	13.9366(54) 0.76	4.9681(87) ^c	12.1855(91) ^c		16.0364(78) ^c
All data		10.3865(8) ^b 1.07	9.8253(10) ^b 1.54		4.9500(10) 3.41	12.1734(10) ^b 2.41		16.0169(21) 5.71
All data revised ^a		10.3869(20) 0.62	9.8238(24) 1.31		4.9583(15) 0.49	12.1748(17) 0.50		16.0249(40) 3.60

^a All OR-16 values omitted due to poor resolution and inconsistency with other data.^b Adopted value.^c Only one measurement.

Table 16d.

Detector	Spectral code	<u>248 - 264Ta</u>	<u>261 - 264Ta</u>	<u>264Ta - 268</u>	<u>264Ta - 274</u>	<u>264Ta - 278</u>
OR-12	A2		2.0904(127)		10.6587(78)	14.0368(48)
	A3		2.0923(140)	10.6632(85)	14.0351(49)	14.0314(41)
	A4		2.1108(40)		10.6547(67)	14.0286(27)
	A5		2.1253(40)	4.5966(32)	10.6602(35)	
	A6	15.3531(18)	2.1247(49)	4.6018(48)	10.6573(26)	14.0330(23)
	A7	15.3501(20)	2.1176(49)	4.6008(33)	10.6571(31)	14.0333(17)
	A8	15.3508(15)			10.6579(29)	

The following are weighted averages and the associated χ^2 values.

OR-12	15.3513(10) 0.74	2.1178(39) 3.31	4.5992(21) 0.59	10.6580(14) 0.20	14.0326(11) 0.70
-------	---------------------	--------------------	--------------------	---------------------	---------------------

Table 17a. Measured γ -ray energy differences from the ^{169}Yb reference lines of 63 to 307 keV.

Detector	Spectral code	54 - 63Yb	63Yb - 66	63Yb - 67	63Yb - 70	63Yb - 71	89 - 93Yb	91 - 93Yb
OR-12	A9	-	2.9877(34)	4.8265(99)	7.1827(49)	8.6934(16)	3.4048(12)	
	A10	-	2.9846(35)	4.8334(88)	7.1759(57)	8.6871(17)	3.4029(11)	
	A11	-	2.9856(39)	4.8241(94)		8.6931(18)	3.4087(12)	
	A12	-	2.9908(37)	4.8321(95)	7.1677(55)	8.6898(17)	3.4052(13)	
	A13	-	2.9797(36)	4.8365(93)	7.1811(47)	8.6868(20)	3.3964(11)	
	A14	-	2.9765(24)	4.8402(68)	7.1782(38)	8.6873(15)	3.3974(8)	
	A15	-	2.9965(62)	4.8537(145)	7.1878(99)	8.6883(32)	3.4004(18)	
	A16	-	2.9847(43)	4.8540(118)	7.1723(60)	8.6929(21)	3.4022(12)	
	A17	-	2.9812(35)	4.8546(99)	7.1797(58)	8.6950(18)	3.4014(11)	
PG-21	none							
PG-16	C20	8.4212(4)	2.9938(18)	4.8306(31)	7.1722(45)	8.6957(8)	3.4097(15)	
	C21	8.4220(6)	2.9965(16)	4.8241(36)	7.1694(38)	8.6953(9)	-	
	C22	8.4210(6)	2.9980(21)	4.8294(37)	7.1690(26)	8.6954(12)	3.4116(16)	
	C23	8.4212(5)	2.9937(15)	4.8199(39)	7.1646(36)	8.6958(12)	3.4112(14)	
	C24		2.9953(14)	4.8220(32)	7.1666(23)	8.6931(11)	3.3928(8)	
	C25	8.4179(10)	2.9944(15)	4.8248(29)	7.1652(25)	8.6936(11)	3.3941(8)	
	C26	8.4163(10)	3.0000(17)	4.8264(28)	7.1619(19)	8.6965(11)	3.3940(6)	
	C27	8.4186(9)	2.9961(15)	4.8233(33)	7.1545(19)	8.6925(10)	3.3930(7)	
OR-16	D12	8.4132(15)	3.0009(18)	4.8260(30)	7.1588(23)	8.6946(16)	3.3937(9)	2.3387(56)
	D13	8.4177(14)	2.9972(19)	4.8205(29)	7.1554(26)	8.6926(15)	3.3941(8)	2.3425(29)
	D14	8.4146(11)	2.9996(13)	4.8276(24)	7.1593(15)	8.6946(11)		

The following are weighted averages and the associated χ^2 values.

OR-12		2.9833(19)	4.8401(38)	7.1778(19)	8.6905(11)	3.4049(9)	
		2.54	1.44	0.99	3.19	5.81	
PG-16		2.9958(7)	4.8253(12)	7.1632(20)	8.6948(5)	3.3960(25)	
		1.66	1.11	5.15	1.97	57.9	
OR-16		2.9994(9)	4.8251(21)	7.1584(11)	8.6941(8)	3.3939(6) ^b	2.3417(26)
		1.03	1.85	0.86	0.65	0.11	0.36
All data		2.9948(13)	4.8264(13)	7.1634(17)	8.6937(6)	3.3990(16)	
		8.28	2.29	6.86	4.12	46.7	
All data revised ^a		2.9968(8)	4.8252(10)	7.1614(16)	8.6947(4)	3.3955(19)	
		2.47	1.15	4.88	1.58	44.6	

^a All OR-12 values omitted due to poor resolution of these peaks.

^b Adopted value.

Table 17b.

Detector	Spectral code	93Yb - 94	93Yb - 97	120 - 130Yb	123 - 130Yb	130Yb - 135	130Yb - 139	130Yb - 146
OR-12	A9		3.7612(10)		6.6293(38)	4.8149(17)		
	A10		3.7623(11)		6.6287(32)	4.8157(18)		
	A11		3.7570(12)		6.6348(38)	4.8164(20)		
	A12		3.7630(11)		6.6293(38)	4.8170(21)		
	A13		3.7709(12)		6.6325(38)	4.8077(32)		
	A14		3.7704(11)		6.6334(25)	4.8149(17)		
	A15		3.7662(14)		6.6273(55)	4.8148(35)		
	A16		3.7636(13)		6.6244(42)	4.8148(22)		
	A17		3.7650(12)		6.6277(36)	4.8138(20)		
PG-21	none							
PG-16	C20		3.7670(12)		6.6345(27)	4.8174(18)		
	C21			9.7031(13)	6.6320(34)	4.8188(21)	9.2116(119)	15.8298(17)
	C22		3.7640(14)	9.7014(13)	6.6343(26)	4.8182(21)		15.8222(19)
	C23		3.7652(16)	9.7104(32)	6.6367(28)	4.8166(18)	9.2198(90)	15.8165(32)
	C24		3.7815(7)		6.6452(39)	4.8120(33)		15.8176(34)
	C25		3.7822(6)	9.7058(29)	6.6390(38)	4.8132(34)	9.2220(107)	15.8176(29)
	C26		3.7811(5)	9.7116(37)	6.6370(33)	4.8170(30)	9.2011(94)	15.8146(37)
	C27		3.7822(6)		6.6412(41)	4.8098(38)		
OR-16	D12	1.3014(31)	3.7830(6)	9.7068(69)	6.6427(73)	4.8133(70)		
	D13	1.2980(29)	3.7826(6)		6.6463(80)	4.8091(77)		
	D14	1.3019(23)						

on

The following are weighted averages and the associated χ^2 values.

OR-12	3.7643(15) 14.35				6.6303(12) 0.81	4.8149(7) 0.89		
PG-16	3.7798(22) 61.5			9.7035(15) 3.19	6.6366(13) 1.39	4.8166(9) 1.15	9.2135(50) 0.96	15.8228(25) 6.28
OR-16	1.3006(16) 0.59		3.7828(4) ^b 0.22	9.7068(69)	6.6443(54) 0.11	4.8114(52) 0.16		
All data			3.7756(22) 108.4	9.7036(13) 2.60	6.6339(11) 1.93	4.8156(5) 1.01		
All data revised ^a			3.7807(16) 50.7		6.6370(13) 1.31			

^a All OR-12 values have been omitted due to inconsistency with other data.^b Adopted value.

Table 17c.

Detector	Spectral code	164 - 177 b	168 - 177 b	170 - 177 b	174 - 177 b	177 b - 187	187 - 197 b	193 - 197 b
OR-12	A9	12.6843(15)			10.7535(19)	9.9924(18)	9.9939(18)	
	A10	12.6841(18)			3.0253(74)	10.7512(18)	9.9904(17)	
	A11	12.6821(13)		6.4025(116)	3.0130(77)	10.7546(17)	9.9928(19)	
	A12	12.6810(17)		6.4067(152)	3.0073(74)	10.7524(21)	9.9879(21)	
	A13	12.6845(22)	8.2227(161)	6.3964(91)	3.0225(66)	10.7548(23)	9.9901(12)	
	A14	12.6823(15)	8.2243(104)	6.4095(57)	3.0158(47)	10.7556(14)	9.9914(31)	
	A15			6.4078(168)	3.0153(130)	10.7491(33)	9.9897(21)	
PG-21	A16	12.6833(20)	8.2201(231)	6.4087(113)	3.0364(83)	10.7544(22)	9.9923(17)	
	A17	12.6813(16)	8.2604(201)	6.3978(100)	3.0270(71)	10.7530(19)		
	none							
PG-16	C20	12.6878(17)	8.2607(227)	6.4037(123)	3.0207(112)	10.7550(30)	9.9884(31)	
	C21	12.6885(18)	8.2076(214)	6.4043(117)	3.0213(97)	10.7524(29)	9.9897(31)	
	C22	12.6893(23)	8.2238(356)	6.3972(198)	3.0075(95)	10.7539(29)	9.9901(28)	
	C23	12.6881(19)	8.2120(283)	6.3898(142)	3.0177(89)	10.7536(33)	9.9883(33)	
	C24	12.6936(29)	8.2109(147)	6.4151(93)	3.0293(81)	10.7613(185)	9.9943(33)	
	C25	12.6952(31)	8.2111(151)	6.3981(99)	3.0319(77)	10.7537(35)	9.9945(32)	4.4547(121)
	C26	12.6961(31)	8.2102(148)	6.4128(80)	3.0217(73)	10.7514(35)	9.9953(36)	4.4479(92)
	C27	12.6949(30)	8.2138(133)	6.4083(83)	3.0256(65)	10.7577(34)	9.9966(33)	
	D12	12.6984(74)	8.2218(208)	6.4001(138)	3.0382(114)	10.7553(78)	9.9882(65)	4.4521(107)
	D13	12.6903(67)		6.3804(130)	3.0115(114)	10.7617(71)	9.9868(61)	4.4612(76)
OR-16	D14	12.6957(41)	8.1969(123)	6.4082(77)	3.0339(71)	10.7569(44)	9.9974(38)	

66

The following are weighted averages and the associated χ^2 values.

OR-12	12.6827(6) 0.65	8.2286(76) 0.98	6.4049(36) 0.37	3.0198(30) 1.46	10.7537(6) 0.86	9.9912(6) 0.98	
PG-16	12.6902(11) 2.04	8.2154(63) 0.64	6.4065(37) 0.56	3.0231(29) 0.75	10.7540(12) 0.33	9.9919(12) 1.07	4.4504(73) 0.20
OR-16	12.6950(32) 0.37	8.2034(109) 1.06	6.4008(78) 1.69	3.0299(70) 1.73	10.7577(34) 0.22	9.9875(44) 0.01	4.4581(62) 0.48
All data	12.6855(10) 4.78	8.2178(44) 0.37	6.4049(24) 0.60	3.0221(20) 1.23	10.7538(6) 0.58	9.9913(5) 0.87	4.4549(47) 0.44
All data revised ^a	12.6905(10) 1.72	8.2123(54) 0.72					

Table 17d.

Detector	Spectral Code	197Yb - 208	291 - 307Yb	294 - 307Yb	302 - 307Yb	307Yb - 317	307Yb - 320
OR-12	A9	10.2229(16)	16.3817(25)		4.7400(132)	9.4338(19)	12.8125(28)
	A10	10.2204(18)	16.3814(20)		4.7342(144)	9.4329(24)	12.8083(23)
	A11	10.2227(17)	16.3816(33)		4.7767(141)	9.4318(19)	12.8125(36)
	A12	10.2205(17)	16.3784(34)		4.7534(147)	9.4294(33)	12.8144(37)
	A13	10.2199(19)	16.3781(23)		4.7324(117)	9.4341(35)	12.8144(23)
	A14	10.2218(12)	16.3860(40)		4.7486(73)	9.4374(23)	12.8107(45)
	A15	10.2207(29)	16.3822(26)		4.7444(283)	9.4277(39)	12.8039(29)
	A16	10.2188(20)	16.3815(26)		4.7594(157)	9.4240(25)	12.8081(27)
	A17	10.2201(18)			4.7634(134)	9.4303(24)	
	none						
	C20	10.2258(30)	16.3853(46)	13.7365(186)	4.7402(265)	9.4283(47)	12.8072(52)
	C21	10.2237(28)	16.3890(51)	13.7425(179)	4.7293(259)	9.4259(54)	12.8048(69)
	C22	10.2224(33)	16.3805(50)	13.7422(182)	4.7061(318)	9.4397(48)	12.8138(62)
	C23	10.2204(31)	16.3836(52)	13.7512(198)	4.7016(284)	9.4323(43)	12.8085(63)
	C24	10.2340(33)					
	C25	10.2260(34)					
	C26	10.2282(35)					
	C27	10.2270(34)					
OR-16	D-12	10.2238(66)					
	D-13	10.2308(66)					
	D-14	10.2220(38)					
The following are weighted averages and the associated χ^2_R values.							
OR-12		10.2211(6) 0.60	16.3811 0.59		4.7494(45) 1.14	9.4318(13) 2.43	12.8104(13) 1.67
PG-16		10.2257(14) 1.63	16.3846(25) 0.49	13.7428(93) 0.10	4.7212(139) 0.44	9.4319(29) 1.49	12.8086(30) 0.36
OR-16		10.2241(29) 0.67					
All data		10.2221(6) 1.64	16.3815(9) 0.67		4.7471(44) 1.18	9.4318(11) 1.99	12.8102(11) 1.19

Table 18a. Measured x-ray energy differences between lines from ^{233}U from 25 to 367 keV.

Detector	Spectral code	25 - 29	29 - 42	42 - 54	53 - 54	54 - 66	66 - 67	66 - 70	66 - 71
OR-12	A9						1.8388(104) ^k	4.1950(59) ^l	
	A10						1.8487(94) ^k	4.1913(66)	
	A11						1.8384(101) ^k		5.7075(42)
	A12						1.8613(101) ^k	4.1769(65) ^l	5.6991(39)
	A13						1.8568(98) ^k	4.2013(55) ^l	5.7070(35)
	A14						1.8637(70) ^k	4.2017(42) ^l	5.7108(24)
	A15						1.8572(156) ^k	4.1913(114) ^l	5.6919(66)
	A16						1.8693(125) ^k	4.1877(72) ^l	5.7082(44)
	A17						1.8734(104) ^k	4.1985(67) ^l	5.7138(38)
PG-21	B1	3.8903(135)			1.0995(50)		1.8690(279) ^k	4.1961(106) ^l	5.6934(68)
	B2	3.9098(249)			1.0906(82)		1.8709(130) ^k	4.1696(94) ^l	5.7017(124)
	B3				1.1172(53)		1.8611(153) ^k		5.7037(67)
	B4				1.1031(48)		1.8662(89) ^k	4.1908(44) ^l	5.7150(34)
	B5						1.8577(68) ^k	4.1899(22) ^l	5.7084(15)
	B6								
PG-16	C1	3.8624(135) ^a			1.0839(67)				
	C2	3.8719(65) ^a		12.2522(17)	1.0922(26)				
	C3	3.8640(21) ^a		12.2522(17)	1.0901(9)				
	C4	3.8410(48) ^a		12.2512(18)	1.0924(15)				
	C5	3.8727(25) ^a	13.3042(25) ^c	12.2544(18)	1.0922(9)				
	C6	3.8494(35) ^a	13.2985(21) ^c	12.2547(14)	1.0912(10)				
	C7	3.8364(34) ^a	13.2927(27) ^c	12.2462(14)	1.0917(8)		1.8176(40)		
	C8	3.8676(23) ^a			1.0925(8)	11.4167(22)	1.8273(43)		
	C9		13.2696(17)	12.2480(12)	1.0935(16)		1.8358(51)		5.6922(34)
	C10				1.0821(21)		1.8192(73)		
	C11				1.0944(32)		1.8234(98)		
	C12				1.0889(13)		1.8295(35)		
	C13				1.0878(30)	11.4108(27)	1.8142(42)		
	C14				1.0917(8)		1.8199(43)		
	C15	3.8693(32) ^a			1.0913(9)		1.8255(16)		
	C16	3.8740(22)			1.0902(15)	11.4183(22)	1.8265(23)		
	C17	3.8764(40)			1.0900(21)	11.4181(18)	1.8306(22)		
	C18	3.8748(37)			1.0869(42)		1.8274(20)		
	C19	3.8713(42)						4.1784(48)	5.7019(19)
	C20	3.8733(34)	13.2765(16)	12.2520(12)	1.0862(45)		1.8368(36)	4.1729(41)	5.6988(17)
	C21	3.8662(57)	13.2762(13)	12.2518(11)	1.0891(31)		1.8277(38)	4.1711(33)	5.6974(23)
	C22	3.8606(65)	13.2765(13)	12.2535(10)	1.0882(30)		1.8315(42)	4.1709(39)	5.7021(19)
	C23	3.8794(63)	13.2755(14)	12.2533(11)	1.0913(24)		1.8262(41)		
	C24	3.8806(60)	13.2773(15)	12.2533(11)					
	C25	3.8777(46)	13.2667(10)		1.0914(8)		1.8304(29)	4.1708(26)	5.6993(13)
	C26	3.8700(49)	13.2691(9)	12.2485(7)	1.0907(6)		1.8265(30)	4.1619(22)	5.6965(16)
	C27	3.8827(42)	13.2696(15)	12.2494(11)	1.0901(6)		1.8272(35)	4.1584(22)	5.6964(14)
	C28	3.8674(40)	13.2673(9)	12.2474(7)					
	C29	3.8785(54)	13.2695(22)						
	C30	3.8657(73)	13.2701(23)						
	C31		13.2754(17)						
	C32		13.2713(17)				1.8176(67)		
	C33		13.2721(12)				1.8250(35)		
	C34		13.2672(14)				1.8217(55)		
	C35		13.2705(14)				1.8282(48)		
	C36		13.2677(14)				1.8273(50)		
	C37		13.2605(18)				1.8307(48)		

Table 18a. (Continued)

Detector	Spectral code	25 - 29	29 - 42	42 - 54	53 - 54	54 - 66	66 - 67	66 - 70	66 - 71
OR-16	D1		13.2661(29)	12.2475(24)	1.0917(71)				
	D2		13.2647(16)	12.2464(15)	1.0901(18)				
	D3	3.8739(95)	13.2663(16)	12.2478(13)	1.0901(26)				
	D4		13.2679(26)	12.2454(17)	1.0860(37)				
	D5	3.8730(76)	13.2680(20)	12.2450(18)	1.0849(24)				
	D6	3.8813(153)	13.2669(27)	12.2451(22)	1.0733(50)				
	D7	3.8831(66)	13.2631(18)	12.2482(15)	1.0880(29)		1.8265(26)	4.1575(20)	
	D8		13.2631(12)	12.2515(11)	1.0938(6)		1.8254(40)	4.1628(28)	
	D9	3.8678(23)	13.2654(12)	12.2509(10)	1.0936(10)		1.8272(33)	4.1604(22)	5.6972(15)
	D10	3.8723(21)	13.2650(13)	12.2519(12)	1.0908(8)		1.8233(26)	4.1576(20)	5.6960(12)
	D11	3.8699(17)	13.2636(13)	12.2518(11)	1.0905(5)		1.8231(29)	4.1579(22)	5.6937(14)
	D12	3.8717(20)	13.2601(11)	12.2499(11)	1.0930(9)		1.8233(29)	4.1583(26)	5.6954(15)
	D13	3.8689(29)	13.2631(11)	12.2505(10)	1.0930(6)		1.8280(23)	4.1598(14)	5.6950(9)
	D14	3.8692(19)	13.2612(11)	12.2498(11)	1.0929(5)		1.8280(18)	4.1604(12)	5.6967(9)
	D15	3.8702(15)	13.2649(14)	12.2519(12)	1.0942(6)		1.8267(11)	4.1590(8)	5.6960(6)
	D16	3.8683(9)	13.2621(12)	12.2513(11)	1.0941(5)		1.8271(12)	4.1593(9)	5.6959(6)
	D17	3.8685(9)	13.2629(12)	12.2511(11)	1.0941(6)		1.8298(13)	4.1595(8)	5.6966(6)
	D18	3.8673(10)	13.2629(12)	12.2509(11)	1.0940(4)				
	D19	3.8694(75)	13.2619(29)	12.2504(21)	1.0929(28)				
	D20	3.8694(79)	13.2611(23)		1.0936(24)				
	D21	3.8561(82)	13.2650(27)		1.0936(30)				
	D22	3.8551(60)	13.2620(22)		1.0923(20)		1.8196(110)	4.1542(98)	5.6946(61)
	D23	3.8712(40)			1.0948(17)		1.8284(60)	4.1717(83)	5.6948(61)
	D24	3.8726(25)	13.2639(15)	12.2548(11)	1.0953(12)		1.8306(48)	4.1523(70)	5.6911(48)
	D25	3.8717(32)	13.2656(18)	12.2529(13)	1.0944(14)		1.8351(54)		
	D26	3.8659(27)	13.2634(15)	12.2549(11)	1.0947(14)		1.8234(45)		
	D27	3.8692(12)	13.2635(24)	12.2544(16)	1.0943(4)	11.4136(11)	1.8285(18)		
	D28	3.8706(10)	13.2632(21)	12.2544(14)	1.0946(3)	11.4139(9)	1.8283(9)		
	D29	3.8699(12)	13.2633(21)	12.2546(15)	1.0940(3)	11.4144(10)	1.8286(17)		
	D30	3.8680(11)		12.2546(14)	1.0934(4)	11.4142(10)	1.8306(16)		
	D31	3.8681(13)							
	D33	3.8706(16)	13.2654(15)						
Si(Li)	E1	3.8764(19)	13.2639(12)	12.2507(10)	1.0917(18)		1.8443(98)		
	E2	3.8774(18)	13.2596(18)	12.2548(12)	1.0998(18)		1.8421(98)		
	E3	3.8740(17)			1.0987(17)				
	E4	3.8757(18)			1.0957(18)				
	E5	3.8702(15)	13.2648(24)	12.2485(14)	1.0937(9)	11.4142(28)	1.8262(72)		
	E6	3.8709(16)	13.2646(26)	12.2492(16)	1.0947(10)	11.4148(29)	1.8237(65)		
	E7	3.8744(14)	13.2660(20)	12.2505(13)	1.0947(9)	11.4093(25)	1.8436(65)		
	E8	3.8725(12)	13.2669(20)	12.2508(13)	1.0941(10)	11.4119(25)	1.8397(52)		
	E9	3.8786(34)	13.2730(23)						
	E10	3.8756(17)	13.2669(23)						
	E12		13.2775(13) ^e		1.0995(9)	11.4124(25)	1.8357(67)		
	E22	3.8779(20)							
	E23	3.8809(19)							
	E24	3.8835(27)							
	E25	3.8852(29)							
	E26	3.8849(26)							
	E27	3.8813(26)							
	E28	3.8859(23)							

The following are weighted averages and the associated χ^2 values.

a. n = 14, p = 14 values omitted including six run with ^{182}Ta , and two p-21 values omitted.

^a Nine pg-16 values omitted, including six runs

c. All Si(L1) uncertainties multiplied by 2.0.
Four PG-16 values omitted.

Four PG-16 values omitted.
All PG-16 uncertainties multiplied by 2.0.

e: One Si(Li) value omitted.

f All PG-16 uncertainties multiplied by 2.0.

9 All OR-16 uncertainties multiplied by 1.5.

h; All OR-16 uncertainties multiplied by 1.5.

All Si(Li) uncertainties multiplied by 1.5.

j All four PG-21 values omitted.

K All fourteen OR-12 and PG-21 values omitted
L All fourteen OR-12 and PG-21 values omitted
M All fourteen OR-12 and PG-21 values omitted
N All fourteen OR-12 and PG-21 values omitted
O All fourteen OR-12 and PG-21 values omitted
P All fourteen OR-12 and PG-21 values omitted
Q All fourteen OR-12 and PG-21 values omitted
R All fourteen OR-12 and PG-21 values omitted
S All fourteen OR-12 and PG-21 values omitted
T All fourteen OR-12 and PG-21 values omitted
U All fourteen OR-12 and PG-21 values omitted
V All fourteen OR-12 and PG-21 values omitted
W All fourteen OR-12 and PG-21 values omitted
X All fourteen OR-12 and PG-21 values omitted
Y All fourteen OR-12 and PG-21 values omitted
Z All fourteen OR-12 and PG-21 values omitted

l All twelve OR-12 and PG-21 values omitted.
m All PG-14 uncertainties multiplied by 1.5.

iii) All PG-16 uncertainties multiplied by 1.5.

Table 18b.

Detector	Spectral code	67 - 70	67 - 71	70 - 71	71 - 74	74 - 76	89 - 93	93 - 97	117 - 118
OR-12	A1								
	A2								
	A3								
	A4								
	A5								
	A9	2.3562(110) ^a	3.8669(100) ^c	1.5107(51) ^d	2.7456(30) ^g	1.7794(78)	3.3905(42)	3.7669(36) ^h	1.8746(107) ⁱ
	A10	2.3426(104) ^a	3.8537(89) ^c		2.7543(28) ^g	1.7739(68)	3.3934(20)	3.7713(18) ^h	1.8084(33)
	A11		3.8690(95) ^c		2.7648(38) ^g	1.8029(84)	3.3945(17)	3.7748(16) ^h	1.8065(30)
	A12	2.3156(109) ^a		1.5222(56) ^d	2.7806(34) ^g	1.7804(105)	3.3970(14)	3.7721(17) ^h	1.8101(32)
	A13	2.3445(102) ^a	3.8502(93) ^c	1.5057(46) ^d	2.7539(31) ^g	1.7856(93)			1.8143(11)
	A14	2.3380(77) ^a	3.8471(68) ^c	1.5091(38) ^d	2.7617(23) ^g	1.7909(50)			
	A15	2.3341(174) ^a	3.8346(147) ^c	1.5005(101) ^d	2.7847(57) ^g	1.7507(160)			
	A16	2.3183(132) ^a	3.8389(119) ^c	1.5205(61) ^d	2.7651(35) ^g	1.8039(113)			
	A17	2.3251(114) ^a	3.8405(100) ^c	1.5153(60) ^d	2.7681(34) ^g	1.7753(96)			
PG-21	B1			1.4991(93) ^d					1.8039(39)
	B2			1.4973(175) ^d					1.8094(71)
	B3			1.5299(77) ^d					1.8154(32)
	B4			1.5372(89) ^d					1.8090(34)
	B5			1.5242(34) ^d					
	B6	2.3321(66) ^a	3.8520(64) ^c	1.5185(18) ^d					
PG-16	C1								
	C2								
	C3								
	C4								
	C5								
	C6								
	C7	2.3530(49)	3.8779(37)	1.5249(34)	2.7224(18)	1.8106(39)	3.3925(72)	3.7794(50)	1.8166(142)
	C8	2.3404(46)	3.8689(41)	1.5286(24)	2.7264(15)	1.8126(33)	3.3948(28)	3.7818(19)	1.8082(49)
	C9	2.3302(60)		1.5319(42)	2.7228(23)	1.8092(54)	3.3927(10)	3.7854(8)	1.8112(17)
	C10	2.3442(83)	3.8730(70)	1.5287(51)	2.7295(31)	1.7974(67)	3.3936(16)	3.7827(11)	1.8073(23)
	C11	2.3441(101)	3.8703(89)	1.5262(53)	2.7232(53)	1.8173(92)	3.3934(10)	3.7843(7)	1.8106(16)
	C12	2.3399(44)	3.8686(33)	1.5287(30)	2.7250(15)	1.8062(41)	3.3925(11)	3.7842(8)	1.8081(18)
	C13	2.3649(107)	3.8788(54)	1.5139(105)	2.7225(37)	1.8104(29)	3.3917(10)	3.7840(7)	1.8110(15)
	C14	2.3571(61)	3.8791(39)	1.5220(48)	2.7275(19)	1.8042(38)	3.3934(12)	3.7821(8)	1.8090(15)
	C15	2.3442(29)	3.8706(14)	1.5264(26)	2.7255(8)	1.8077(14)	3.3901(13)	3.7833(11)	1.8073(22)
	C16	2.3459(38)	3.8698(22)	1.5238(32)	2.7277(11)	1.8061(24)	3.3905(22)	3.7847(12)	1.8064(28)
	C17	2.3414(40)	3.8679(18)	1.5265(36)	2.7262(11)	1.8100(19)	3.3920(13)	3.7830(13)	1.8148(40)
	C18	2.3428(32)	3.8715(18)	1.5288(27)	2.7247(8)	1.8122(20)	3.3929(20)	3.7838(16)	1.8098(18)
	C19			1.5221(28)	2.7271(12)	1.8102(28)	3.3933(9)	3.7834(7)	1.8068(16)
	C20	2.3416(55)	3.8651(32)	1.5235(45)	2.7264(16)	1.8068(33)	3.3893(12)	3.7824(10)	1.8107(20)
	C21	2.3452(52)	3.8712(36)	1.5259(38)	2.7293(17)	1.8068(47)	3.3923(14)	3.7858(9)	
	C22	2.3396(45)	3.8659(38)	1.5263(28)	2.7271(17)	1.8037(33)	3.3920(11)	3.7835(9)	
	C23	2.3447(53)	3.8759(41)	1.5312(38)	2.7267(18)	1.8140(45)	3.3920(13)	3.7830(13)	
	C24	2.3446(37)	3.8710(31)	1.5264(23)	2.7247(12)	1.8132(25)	3.3919(15)	3.7847(11)	
	C25	2.3404(36)	3.8688(28)	1.5285(24)	2.7246(13)	1.8123(28)	3.3916(11)	3.7839(8)	
	C26	2.3355(32)	3.8700(27)	1.5346(18)	2.7214(12)	1.8188(26)	3.3932(9)	3.7838(7)	
	C27	2.3312(37)	3.8692(33)	1.5380(18)	2.7254(12)				
	C28								1.8072(15)
	C29			1.5298(32)					1.8064(20)
	C30	2.3350(54)	3.8648(46)						1.8110(20)
	C31		3.8631(63)						1.8102(21)

Table 18b. (Continued)

Detector	Spectral code	67 - 70	67 - 71	70 - 71	71 - 74	74 - 76	89 - 93	93 - 97	117 - 118
	C32	2.3462(72)	3.8807(62)	1.5346(42)					1.8088(23)
	C33	2.3339(39)	3.8698(33)	1.5359(23)					1.8103(13)
	C34	2.3338(59)	3.8706(52)	1.5368(30)					1.8068(19)
	C35	2.3352(53)	3.8691(46)	1.5339(31)					1.8084(23)
	C36	2.3340(55)	3.8652(46)						1.8079(19)
	C37	2.3410(54)	3.8696(45)	1.5285(33)					1.8102(18)
	C39								1.8085(25)
	C40								1.8084(32)
	C41								1.8102(20)
	C42								1.8154(20)
	C43								1.8124(25)
	C44								1.8094(26)
OR-16	D1			1.5339(73)	2.7310(33)	1.8159(74)	3.3897(43)	3.7802(35)	
	D2				2.7405(41)	1.7954(110)	3.3899(17)	3.7857(15)	
	D3				2.7314(59)	1.8084(118)	3.3836(19)	3.7855(13)	
	D4				2.7353(44)	1.7950(96)	3.3848(24)	3.7837(17)	
	D5				2.7374(108)		3.3833(21)	3.7855(15)	
	D6				2.7257(46)		3.3893(40)	3.7847(28)	
	D7				2.7233(11)	1.8132(114)	3.3858(20)	3.7825(16)	
	D8	2.3310(30)	3.8695(25)	1.5289(75)	2.7233(11)	1.8088(23)	3.3925(7)	3.7843(6)	1.8108(12)
	D9	2.3373(44)	3.8731(38)	1.5385(18)	2.7207(16)	1.8106(39)	3.3936(9)	3.7841(7)	1.8100(15)
	D10	2.3332(36)	3.8700(32)	1.5358(25)	2.7250(15)	1.8082(30)	3.3923(8)	3.7844(6)	1.8078(16)
	D11	2.3342(29)	3.8727(24)	1.5385(17)	2.7225(10)	1.8142(23)			1.8082(11)
	D12	2.3328(32)	3.8686(27)	1.5358(20)	2.7228(13)	1.8125(36)	3.3924(6)	3.7848(5)	1.8090(15)
	D13	2.3349(34)	3.8721(27)	1.5372(23)	2.7223(14)	1.8120(35)			1.8097(16)
	D14	2.3318(24)	3.8670(22)	1.5352(12)	2.7233(8)	1.8070(19)			1.8077(9)
	D15	2.3324(19)	3.8688(17)	1.5363(10)	2.7236(7)	1.8119(16)	3.3930(5)	3.7847(4)	1.8092(8)
	D16	2.3324(12)	3.8693(10)	1.5369(7)	2.7222(6)	1.8114(12)	3.3930(6)	3.7841(4)	1.8084(6)
	D17	2.3322(13)	3.8688(11)	1.5366(8)	2.7233(6)	1.8118(12)	3.3927(6)	3.7846(4)	1.8086(7)
	D18	2.3297(14)	3.8669(13)	1.5372(8)	2.7240(6)	1.8106(14)	3.3925(5)	3.7843(3)	1.8100(5)
	D19			1.5403(88)	2.7207(57)		3.3931(30)	3.7854(25)	1.8165(64)
	D20			1.5232(73)	2.7244(55)		3.3932(30)	3.7836(23)	1.8121(60)
	D21			1.5319(85)	2.7194(57)		3.3958(37)	3.7833(24)	1.8075(61)
	D22	2.3327(116)		1.5387(60)	2.7229(44)		3.3871(25)	3.7877(22)	1.8088(52)
	D23	2.3278(69)	3.8653(57)	1.5375(45)					1.8105(43)
	D24	2.3273(51)	3.8646(44)	1.5373(29)					1.8065(24)
	D25	2.3320(59)	3.8666(50)	1.5446(37)					1.8097(29)
	D26	2.3275(49)	3.8719(41)	1.5444(32)					1.8095(27)
	D27	2.3246(19)	3.8665(17)	1.5419(11)					1.8072(9)
	D28	2.3292(11)	3.8668(9)	1.5376(7)					1.8088(6)
	D29	2.3240(19) ^b	3.8661(17)	1.5421(11)					1.8083(9)
	D30	2.3211(17)	3.8646(15)	1.5435(10)					1.8091(8)
	D31								1.8089(13)
	D33								1.8092(15)

Table 18b. (Continued)

	Si(Li)	E1	3.8681(140)	1.5298(73)	1.8142(89)
		E2	2.3282(115)	1.5327(79)	1.8191(70)
		E3	2.3253(118)	1.5453(41)	1.8154(65)
		E5	2.3242(77)	1.5343(51)	1.8083(59)
		E6	2.3332(76)	1.5343(38)	1.8071(41)
		E7	2.3222(65)	1.5353(40)	1.8097(64)
		E8	2.3191(60)	1.5695(45)	1.8111(53)
		E10			1.8104(58)
		E11			
		E12	2.3296(73)	1.5371(42)	
The following are weighted averages and the associated χ^2 values.					
OR-12		2.3357(47)	3.8514(40)	1.5119(25)	1.8132(30)
		1.54	1.42	1.56	10.5
OR-12 revised					1.8127(16)
					2.91
PG-21		2.3321(66)	3.8520(64)	1.5199(26)	1.8101(26)
				2.87	1.79
PG-16		2.3408(12)	3.8702(7)	1.5301(9)	1.8094(4)
		1.78	1.07	2.78	1.07
PG-16 revised					
OR-16		2.3296(9)	3.8679(4)	1.5381(5)	1.8089(2)
		3.45	1.27	3.67	0.79
OR-16 revised		2.3303(7) ^b		1.5377(4) ^e	
		2.13		2.46	
Si(Li)		2.3251(29)	3.8612(24)	1.5411(45)	1.8107(20)
		0.47	0.93	7.49	0.45
Si(Li) revised				1.5367(18) ^f	
				1.05	
All data		2.3320(8)	3.8683(5)	1.5359(7)	1.8092(2)
		4.17	1.92	9.33	1.71
All data reduced set		2.3326(8) ^{a,b}	3.8685(4) ^c	1.5362(6) ^{d,e,f}	1.8092(2) ⁱ
		3.92	1.45	4.86	0.96

a All eight OR-12 values and the one PG-21 value omitted.
b One OR-16 value omitted.
c All eight OR-12 values and the one PG-21 values omitted
d All seven OR-12 values and all six PG-21 values omitted
e One OR-16 value omitted.
f One Si(Li) value omitted.
g All nine OR-12 values omitted.
h All four OR-12 values omitted.
i One OR-12 value omitted.

Table 18c.

Detector	Spectral code	118 - 120	120 - 123	120 - 135	123 - 125	135 - 139	135 - 146	145 - 146	146 - 148
OR-12	A1	1.8437(111)							
	A2	1.8456(19)							
	A3	1.8481(17)							
	A4	1.8438(19)							
	A5	1.8467(10)							
	A16					4.3983(171)			
	A17					4.4005(165)			
PG-21	A18					4.4031(137)			
	B1	1.8462(36)							
	B2	1.8382(67)							
PG-16	B3	1.8478(30)							
	B4	1.8418(32)							
	C1	1.8435(119)	3.0833(257)				11.0050(48)	0.9863(75)	1.8318(211)
	C2	1.8521(47)	3.0729(97)				11.0008(16)	0.9993(30)	1.8148(55)
	C3	1.8491(15)	3.0635(38)				11.0050(25)	0.9973(35)	1.8081(104)
	C4	1.8472(22)	3.0634(50)				11.0025(20)	0.9958(31)	1.8120(73)
	C5	1.8458(16)	3.0664(32)				11.0057(18)	0.9990(31)	1.8236(61)
	C6	1.8448(15)	3.0681(31)				11.0041(23)	0.9977(27)	1.8082(55)
	C7	1.8447(15)	3.0650(30)				11.0025(19)	0.9977(27)	1.8221(52)
	C8	1.8474(15)	3.0644(35)			4.4042(122)	11.0025(19)	1.0009(36)	1.8119(77)
	C9	1.8499(22)	3.0671(43)			4.3907(216)	11.0050(29)	1.0079(46)	1.8259(116)
	C10	1.8449(32)	3.0671(56)				11.0020(40)	0.9954(43)	1.8166(124)
	C11	1.8485(39)	3.0586(82)				11.0082(46)	1.0090(54)	1.8169(63)
	C12	1.8463(16)	3.0658(26)		1.4757(217)		11.0027(22)	1.0025(32)	1.8190(44)
	C13	1.8480(22)	3.0632(34)			4.3868(194)	11.0050(25)	1.0057(31)	1.8247(54)
	C14	1.8455(20)	3.0668(35)				11.0028(27)	0.9929(58)	1.8110(23)
	C15					4.3877(128)	11.0040(21)	1.0008(33)	1.8141(31)
	C16							1.0005(37)	1.8121(30)
	C17							1.0039(39)	1.8150(35)
	C18							1.0016(35)	1.8234(31)
	C19							1.0021(30)	1.8154(64)
	C20						11.0044(14)	1.0037(38)	1.8160(47)
	C21						11.0073(20)	1.0030(38)	1.8144(50)
	C22				1.5335(217)		11.0017(24)	1.0035(42)	1.8119(52)
	C23							1.0013(46)	1.8155(45)
	C24						11.0045(13)	1.0031(21)	1.8155(45)
	C25						11.0044(17)	1.0003(21)	1.8206(36)
	C26						11.0006(15)	1.0006(24)	1.8159(35)
	C27					4.4050(105)	11.0048(14)	0.9991(19)	1.8200(34)
	C28	1.8451(14)				4.3913(88)	10.9977(36)	0.9988(24)	1.8111(36)
	C29	1.8436(20)		14.5181(39)				0.9971(38)	1.8081(54)
	C30	1.8447(22)						1.0047(36)	1.8883(61) ^b
	C31	1.8439(20)							
	C32	1.8470(25)						0.9985(24)	1.8145(45)
	C33	1.8442(13)						0.9962(31)	1.8077(55)
	C34	1.8468(18)						1.0063(33)	1.7928(60)
	C35	1.8466(21)						0.9988(33)	1.8003(55)
	C36	1.8457(20)						1.0065(29)	1.8040(60)
	C37	1.8494(17)							
	C39	1.8459(28)							
	C40	1.8513(29)							

Table 18c. (Continued)

Detector	Spectral code	118 - 120	120 - 123	120 - 135	123 - 125	135 - 139	135 - 146	145 - 146	146 - 148
OR-16	C41	1.8427(21)							
	C42	1.8457(21)							
	C43	1.8442(26)							
	C44	1.8479(26)							
	D1	1.8452(88)	3.0537(86)	14.5186(41)			11.0100(37)	1.0103(53)	1.8197(144)
	D2	1.8491(35)	3.0658(133)				11.0068(66)	1.0024(71)	
	D3	1.8493(36)	3.0435(115)				11.0018(60)	1.0081(82)	
	D4	1.8503(45)	3.0509(109)				11.0050(106)	1.0175(73)	
	D5	1.8501(44)					11.0021(55)	1.0441(157)	
	D6	1.8473(86)					11.0021(16)	1.0042(66)	
	D7	1.8471(39)	3.0439(108)				11.0048(20)	1.0024(20)	
	D8	1.8460(12)	3.0676(23)				11.0046(20)	1.0064(28)	
	D9	1.8477(16)	3.0637(30)				11.0055(15)	1.0072(23)	
	D10	1.8456(17)	3.0682(31)	14.5207(22)	1.5369(181)	4.3615(163)	11.0040(20)	1.0072(23)	
	D11	1.8473(12)	3.0709(22)	14.5184(17)	1.5404(157)	4.4002(148)	11.0025(20)	1.0057(19)	
	D12	1.8481(15)	3.0640(28)		1.5384(149)	4.3677(103)	11.0035(13)	1.0065(25)	
	D13	1.8456(17)	3.0672(29)			4.3772(125)	11.0041(11)	1.0053(28)	
	D14	1.8464(10)	3.0685(18)			4.3872(134)	11.0035(13)	1.0076(16)	
	D15	1.8453(7)	3.0682(14)	14.5214(13)	1.5276(72)	4.3955(96)	11.0030(8)	1.0057(15)	
	D16	1.8470(6)	3.0642(11)	14.5234(10)	1.5511(61)	4.3858(62)	11.0037(9)	1.0032(13)	
	D17	1.8457(6)	3.0659(15)	14.5190(8)	1.5514(71)	4.3858(55)	11.0025(9)	1.0054(13)	
	D18	1.8458(5)	3.0639(11)	14.5194(9)	1.5598(55)	4.3840(56)	11.0025(9)	1.0066(13)	
	D19	1.8315(63)	3.0684(14)			4.3804(52)	11.0018(93)	1.0018(93)	
	D20	1.8501(63)	3.0520(13) ^a				1.0111(88)	1.0111(88)	
	D21	1.8409(65)					1.0138(96)	1.0138(96)	
	D22	1.8484(48)	3.0579(88)				1.0162(80)	1.0162(80)	
	D23	1.8485(35)							
	D24	1.8458(25)							
	D25	1.8457(27)							
	D26	1.8535(28)							
	D27	1.8478(10)							
	D28	1.8465(7)							
	D29	1.8469(9)							
	D30	1.8484(9)							
	D31	1.8464(13)							
	D33	1.8445(16)							
Si(Li)	E3	1.8419(85)							
	E5	1.8465(60)							
	E6	1.8554(68)							
	E7	1.8468(58)							
	E8	1.8504(43)							
	E10	1.8481(58)							
	E11	1.8498(57)							
	E12	1.8434(61)							
				14.5177(29)			11.0022(36)	1.0074(13)	1.8131(16)
							10.9990(27)	1.0067(11)	1.8150(24)
								1.0061(14)	1.8098(26)
								1.0081(14)	1.8047(39)

Table 18c. (Continued)

The following are weighted averages and the associated χ^2 values.

OR-12	1.8464(7) 0.80					4.4010(90) 0.02			
PG-21	1.8448(18) 1.00								
PG-16	1.8461(4) 1.02	3.0669(7) 0.60	14.5181(39)	1.5200(39) 7.37	4.3897(45) 1.76	11.0037(4) 0.95	1.0007(6) 1.13	1.8157(19) 6.08	1.8145(10) ^b 1.77
PG-16 revised									
OR-16	1.8465(2) 1.11	3.0644(12) 7.59	14.5202(7) 2.31	1.5484(46) 2.35	4.3835(24) 0.89	11.0034(4) 0.71	1.0060(4) 1.15	1.8119(8) 1.87	
OR-16 revised		3.0661(7) ^a 1.98							
Si(Li)	1.8483(21) 0.40								
All data	1.8464(2) 0.96	3.0651(8) 4.14	14.5202(6) 2.06	1.5472(52) 3.16	4.3863(23) 1.35	11.0035(3) 0.83	1.0042(5) 2.22	1.8133(10) 4.91	1.8128(6) ^b 1.91
All data reduced		3.0663(4) ^a 1.24							

^a One OR-16 value omitted.^b One PG-16 value omitted.

Table 18d.

Detector	Spectral code	146 - 164	164 - 165	208 - 217	216 - 217	240 - 245	245 - 248
OR-12	A6					4.9745(26)	3.3723(13)
	A8					4.9671(31)	3.3745(14)
	A9					4.9874(50)	3.3705(27)
	A10					4.9968(49)	3.3693(27)
	A11					4.9793(58)	3.3679(24)
	A12					4.9720(50)	3.3726(27)
	A13					4.9811(45)	3.3742(22)
	A14					4.9737(29)	3.3762(15)
	A15					4.9733(86)	3.3744(44)
	A16					4.9817(60)	3.3686(35)
	A17					4.9777(50)	3.3753(26)
PG-21	none						
PG-16	C1		0.9787(281)	8.9908(294)	1.0628(295)		3.3439(215)
	C2		1.0212(89)	8.9773(124)	1.0647(76)		3.3803(51)
	C3			8.9767(34)	1.0719(163)		
	C4			8.9695(57)	1.0708(91)		3.3653(74)
	C5		1.0234(74)	8.9754(37)	1.0646(92)		3.3691(75)
	C6		1.0231(79)	8.9771(36)	1.0658(84)		
	C7		1.0250(66)		1.0534(97)		
	C8		1.0354(75)	8.9814(42)			
	C9		1.0174(91)				
	C10		1.0422(121)				
	C12		1.0315(60)				3.3812(48)
	C13		1.0325(63)				3.3783(58)
	C14		1.0099(83)	8.9751(48)	1.0740(63)		
	C15		1.0262(65)	8.9762(20)	1.0587(84)		
	C16		1.0300(46)	8.9784(25)	1.0677(35)		
	C17		1.0214(41)	8.9764(24)	1.0622(60)		
	C18		1.0311(57)	8.9788(25)	1.0662(63)		
	C19			8.9785(23)	1.0741(48)		
	C20		1.0266(85)	8.9776(35)	1.0746(52)		
	C21	18.1784(19)	1.0271(91)	8.9733(29)	1.0729(71)	4.9861(83)	3.3633(47)
	C22		1.0426(96)	8.9739(37)	1.0564(65)	4.9703(78)	3.3681(44)
	C23		1.0322(98)	8.9810(38)	1.0659(76)	4.9921(87)	3.3751(44)
	C24	18.1815(13)	0.9926(49) ^a		1.0619(96)	4.9774(83)	3.3853(45)
	C25	18.1801(12)	1.0170(48)			4.9824(82)	3.3839(35)
	C26	18.1803(17)	1.0004(49)			4.9809(73)	3.3773(37)
	C27	18.1822(12)	1.0130(48)			4.9877(72)	3.3738(37)
						4.9876(69)	3.3796(33)

Table 18d. (Continued)

Detector	Spectral code	146 - 164	164 - 165	208 - 217	216 - 217	240 - 245	245 - 248
OR-16	D2	18.1750(32)	1.0206(61)	8.9586(130)	1.0829(58)		3.3995(152)
	D7		1.0297(45)	8.9721(36)	1.0708(68)		3.3777(65)
	D8	18.1785(20)	1.0225(65)	8.9763(40)	1.0787(72)	4.9857(105)	3.3792(72)
	D9	18.1741(18)	1.0181(58)	8.9700(39)	1.0815(54)	4.9723(87)	3.3809(49)
	D10	18.1774(17)	1.0275(53)	8.9741(33)	1.0851(65)	4.9808(112)	3.3727(61)
	D11		1.0316(58)	8.9779(35)	1.0845(74)	4.9849(104)	3.3664(65)
	D12	18.1763(16)	1.0215(57)	8.9677(44)	1.0770(41)	4.9815(65)	3.3789(42)
	D13	18.1785(10)	1.0251(34)	8.9709(23)	1.0748(29)	4.9787(58)	3.3737(32)
	D14		1.0224(28)		1.0781(29)	4.9775(42)	3.3743(23)
	D15		1.0235(20)	8.9649(16) ^b	1.0782(27)	4.9659(44)	3.3754(30)
	D16		1.0283(24)	8.9630(15) ^b		4.9724(44)	3.3781(24)
	D17		1.0273(23)				
	D18		1.0337(210)				
	D22						

The following are weighted averages and the associated χ^2 values.

OR-12						4.9764(23)	3.3730(8)
						3.57	1.53
PG-16	18.1806(6)	1.0205(27)	8.9770(8)	1.0674(16)	4.9831(27)	0.76	3.3766(18)
	1.08	3.97	0.52	0.78			2.20
PG-16 revised		1.0228(22) ^a					
		2.49					
OR-16	18.1773(7)	1.0254(10)	8.9676(16)	1.0780(13)	4.9750(21)	1.09	3.3760(11)
	1.17	0.74	3.78	0.61			0.85
OR-16 revised			8.9675(16) ^c				
			1.72				
All data	18.1791(7)	1.0237(14)	8.9726(12)	1.0737(13)	4.9769(14)		3.3742(7)
	2.32	3.00	4.27	1.67	2.07		1.79
All data revised		1.0246(11) ^a	8.9754(8) ^{b,c}				
		1.87	1.48				

^a One PG-16 value omitted.

^b One OR-16 value omitted.

^c All OR-16 uncertainties multiplied by 1.5.

Table 18e.

Detector	Spectral code	274 - 278	278 - 288	278 - 291	288 - 291	291 - 293	291 - 295	291 - 317	317 - 320
OR-12	A1								3.3768(29)
	A2								3.3754(29)
	A3								3.3745(27)
	A4								3.3762(12)
	A5	3.3684(35)	9.9166(22)		3.3277(17)				3.3811(13)
	A6	3.3728(38)	9.9256(24)		3.3217(18)				3.3800(11)
	A7	3.3759(31)	9.9216(22)		3.3254(18)				3.3769(24)
	A8	3.3740(32)			3.3254(34)				3.3796(21)
	A9	3.3717(51)			3.3296(33)				3.3765(19)
	A10	3.3826(56)	9.9177(40)		3.3283(33)				3.3831(22)
	A11	3.3754(49)	9.9212(40)		3.3281(38)				3.3803(20)
	A12	3.3703(66)	9.9186(46)						3.3771(13)
	A13	3.3750(52)	9.9213(40)	13.2467(31)	3.3283(23)				3.3830(35)
	A14	3.3702(35)	9.9222(28)	13.2485(30)	3.3178(57)				3.3799(22)
	A15	3.3686(106)	9.9266(71)	13.2505(20)	3.3305(41)				3.3777(18)
	A16	3.3723(55)	9.9186(49)		3.3263(38)				3.3799(94)
	A17	3.3720(56)	9.9181(45)						3.3930(174)
PG-21	B1								3.3672(62)
	B2								3.3797(65)
	B3								3.3747(27)
	B4								3.3757(12)
	B5								3.3821(53)
	B6								3.3851(59)
PG-16	C13	3.3884(105)	9.9192(112)	13.2637(103)	3.3445(81)	2.6489(541)		25.8471(179)	3.3769(31)
	C14	3.3666(133)		13.2372(79)	3.2983(171)	2.6492(553)		25.8337(68)	3.3794(49)
	C15	3.3589(56)	9.9198(123)	13.2355(49)	3.3295(57)	2.6558(563)	4.5994(564)		3.3765(34)
	C16		9.9198(63)	13.2493(41)	3.3335(63)	2.7034(554)	4.6108(502)	25.8134(31)	3.3793(27)
	C17	3.3611(81)	9.9180(73)	13.2515(49)					3.3798(34)
	C18	3.3723(77)	9.9121(61)	13.2473(40)	3.3298(59)	2.6488(185)		25.8123(34)	3.3790(50)
	C19	3.3485(81)	9.9194(64)	13.2491(46)	3.3204(69)	2.6465(178)			3.3789(67)
	C20	3.3791(100)			3.3370(83)	2.6382(181)			3.3742(56)
	C21	3.3624(88)			3.3444(86)	2.6324(199)			3.3762(61)
	C22	3.3844(105)			3.3438(81)				
	C23	3.3509(129)	9.9139(91)						

The following are weighted averages and the associated χ^2 values.

OR-12	3.3731(12) 0.59	9.9214(11) 1.22	13.2492(15) 0.56	3.3261(9) 1.31				3.3787(6) 1.63
PG-21								3.3755(11) 0.71
PG-16	3.3653(39) 2.01	9.9171(29) 0.20	13.2469(26) 1.84	3.3329(32) 1.74	2.6441(88) 0.25	4.6150(313) 0.11	25.8155(42) 3.87	3.3785(12) 0.32
All data	3.3718(13) 1.45	9.9209(9) 0.90	13.2483(14) 1.45	3.3267(10) 1.79				3.3782(4) 1.19

Table 18f.

Detector	Spectral code	320 - 323	323 - 328	323 - 336	328 - 336	365 - 367
OR-12	A1					
	A2	2.8262(55)				
	A3	2.8263(51)				
	A4	2.8404(53)				
	A5					
	A6	2.8356(21)				
	A7	2.8298(22)				
	A8	2.8292(19)				
	A9	2.8362(40)				
	A10	2.8324(42)				
	A11	2.8386(36)				
	A12	2.8325(39)				
	A13	2.8352(38)				
	A14	2.8356(23)				
	A15	2.8283(68)				
	A16	2.8340(41)				
	A17					
PG-21	B1	2.8428(165)				
	B2	2.8010(358)				
	B3	2.8380(119)				
	B4	2.8205(120)				
	B5	2.8354(61)				
	B6	2.8441(30)				
PG-16	C13	2.8221(89)				
	C14	2.8315(107)				
	C15	2.8386(44)				
	C16	2.8277(71)				
	C17	2.8334(73)				
	C18	2.8257(71)				
	C19	2.8397(55)				
	C20	2.8244(95)				
	C21	2.8391(91)				
	C22	2.8412(83)				
	C23	2.8285(99)				
The following are weighted averages and the associated χ^2 values.						
OR-12		2.8329(10)	5.3790(45)	13.2395(14)	7.8530(26)	
		1.39	0.57	1.27	1.16	
PG-21		2.8410(28)		13.2330(39)		
		1.23				
PG-16		2.8338(22)	5.3619(169)	13.2469(30)	7.8788(170)	1.9809(117)
		0.86	0.35	0.83	0.09	2.19
All data		2.8337(9)	5.3779(44)	13.2400(13)	7.8536(24)	
		1.41	0.55	1.35	0.98	

**DATE
FILMED**

9/8/93

END