PROGRESS REPORT
ON
RESEARCH IN NUCLEAR PHYSICS

June 1, 1991 - June 30, 1992

R. L. Kozub and M. M. Hindi

DEPARTMENT OF PHYSICS

TENNESSEE TECHNOLOGICAL UNIVERSITY

COOKEVILLE, TENNESSEE 38505

June, 1992

**PREPARED FOR THE U. S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-87ER40314**

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREFACE

The progress on Grant No. DE-FG05-87ER40314 from June 1, 1991 to June 30, 1992, is summarized in this report. The main emphases of the program are studies of the structure of neutron-rich nuclei and rare electron capture processes.

Most of our activities for the past year were centered on the rare electron capture component of our program, which is based on the campus of Tennessee Technological University (TTU). We also continued with feasibility studies pertaining to the search for bound clusters of neutrons. The Fragment Mass Analyzer (FMA) at the ATLAS, together with Compton-suppressed Ge detectors and a Bragg detector in the FMA focal plane, is necessary for our proposed work on neutron-rich nuclei. This combination is just now becoming available for use.

Five undergraduate students, all majoring in physics, are currently working on the project. They are rookies Robert Boyd and Alex Altgilbers and veterans Danny Bardayan, Yuriko Taketani, and Jim Parker. All are working with enthusiasm and making positive contributions.

The secretarial and accounting duties for this project have again been performed flawlessly by Gloria Julian. We appreciate her efficient, accurate service and abundant patience.

We would like to thank the many scientists and support personnel at other institutions and laboratories who have assisted us with this work. In particular, MMH is deeply appreciative of the hospitality and support shown him during his visits to Lawrence Berkeley Laboratory and Montana State University.

R. L. Kozub

M. M. Hindi

CONTENTS

Preface	1
Electron Capture Decay of ^{179}Ta	2
Search for 17-keV Neutrinos in the Internal Bremsstrahlung Spectrum of ^{125}I	8
β^+ Decay and Cosmic-ray Half-life of ^{91}Nb	12
Publications and Seminars	15
TTU Personnel	16
Appendix:	
Yrast Decays in ^{43}K (preprint)	
Comment on 'Double <i>K</i> -shell Ionization in the Electron Capture Decay of ^{55}Fe ' (reprint)	
Reply to 'Coherent Radiation, Not a Heavy Neutrino?' (reprint)	
Double <i>K</i> -Shell Vacancy Production in the Electron Capture Decay of ^{125}I (reprint)	

Electron Capture Decay of ^{179}Ta

M.M. Hindi, J.G. Parker*, R.L. Kozub, B. Sur,[†] K.R. Czerwinski,[†] E.B. Norman,[†]
K.T. Lesko,[†] and R.-M. Larimer[†]

We continue to study rare (secondary) processes in the electron capture decays of nuclei with low decay energies. Last year we reported¹ the production of a ^{179}Ta source at the Lawrence Berkeley Laboratory. After chemical purification the source was sent to Tennessee Tech for counting. We conducted singles and coincidence experiments to determine the EC Q -value, and to measure the probability of producing K and L holes during electron capture, and to determine the intensity of internal bremsstrahlung radiation associated with capture from the various shells. We report here preliminary results on the determination of the first two quantities.

1. Measurement of the Q value

The Q value for decays with low transition energy can be deduced from L -to- K capture ratios. To measure that ratio we recorded the spectrum of Hf x rays produced following the EC decay of ^{179}Ta . The x rays were detected in a planar Ge detector with a 16-mm diameter, 10-mm depth and 0.128-mm-thick Be window. The source was kept at a distance of $\simeq 7$ cm from the detector to minimize the effect of summing of K x rays with their associated L x rays. The resulting spectrum of Hf L x rays is shown in Fig. 1. We extracted the yield of L x rays by fitting this spectrum with Gaussian peaks with fixed centroids on top of a quadratic background. The yield $N_{L_{1i}}$ of a radiative transition i that fills a hole in the L_1 shell is related to the number N_0 of EC decays by

$$N_{L_{1i}} = N_0 (P_K n_{KL_{1i}} + P_{L_1} n_{L_1 L_{1i}}) \epsilon_{L_{1i}}, \quad (1)$$

where P_K is the K capture fraction, $n_{KL_{1i}}$ is the number of L_{1i} x rays per K hole, P_{L_1} is the L_1 capture fraction, $n_{L_1 L_{1i}}$ is the number of L_{1i} x rays per L_1 hole, and $\epsilon_{L_{1i}}$ is the efficiency for detecting L_{1i} x rays. Similarly, the yield $N_{L_{2i}}$ of a radiative transition i that fills a hole in the L_2 shell is related to the number N_0 of EC decays by

$$N_{L_{2i}} = N_0 (P_K n_{KL_{2i}} + P_{L_1} n_{L_1 L_{2i}} + P_{L_2} n_{L_2 L_{2i}}) \epsilon_{L_{2i}}. \quad (2)$$

For the allowed decay of ^{179}Ta the capture fractions are related to the Q value by

$$P_K = C \beta_K^2 B_K (Q - E_K)^2 \quad (3)$$

*TTU student

[†]Nuclear Science Division, Lawrence Berkeley Laboratory.

¹R.L. Kozub and M.M. Hindi, *Progress Report on Research in Nuclear Physics, 1990-1991*, (DOE/ER/40314-5), 11 (1991).

Figure 1: Measured spectrum of Hf L x rays following the EC decay of ^{179}Ta

$$P_{L_1} = C \beta_{L_1}^2 B_{L_1} (Q - E_{L_1})^2 \quad (4)$$

$$P_{L_2} = C \beta_{L_2}^2 B_{L_2} (Q - E_{L_2})^2, \quad (5)$$

where C contains the nuclear matrix elements, β_x is the Coulomb amplitude of the bound-state electron radial wavefunction in orbital x , B_x is the associated electron exchange and overlap correction, and E_x is the electron binding energy in the daughter (Hf) atom.

Because the probability of a $K \rightarrow L_1$ hole transfer is small (i.e., the term n_{KL_1} in eq. (1) is small), N_{L_1} is dominated by the P_{L_1} term, while N_{L_2} in eq. (2) is dominated by the P_K term. Therefore the ratio N_{L_1}/N_{L_2} is a sensitive measure of P_{L_1}/P_K , which in turn is a sensitive measure of the Q value (for Q values that are not much larger than E_K , which is the case for ^{179}Ta).

In this preliminary analysis we deduced the Q value from the ratio of $L\gamma_3$ to $L\gamma_1$ x rays. While the yield of $L\gamma$ x rays is smaller than the yield of $L\beta$ x rays (see Fig. 1), the $L\gamma$ group is less complicated than the $L\beta$ group and the separation between L_1 and L_2 x rays is largest in that group. From the measured ratio $N_{L\gamma_3}/N_{L\gamma_1} = 0.693 \pm 0.011$ we deduce $Q = 117.6 \pm 1.3$ keV. In obtaining this result we used the tabulations of Prowse and Firestone² for the x-ray intensities and the Coulomb amplitudes. The error on the Q

² Table of Radioactive Isotopes, by E. Browne and R.B. Firestone, V.S. Shirley, ed., (Wiley, New York, 1986).

value reflects only the statistical error on the x-ray yields. Because the tabulated x-ray intensities were derived from relative emission probabilities and from radiative atomic yields, there is probably a large correlation between the derived values and it is not yet clear to us how to propagate the error due to the tabulated uncertainties in the x-ray intensities.

Our deduced value of $P_L/P_K = 0.687 \pm 0.017$ is in fair agreement with Jopson *et al.*'s³ value of 0.63 ± 0.06 , but not with Bisi *et al.*'s⁴ value of 1.4 ± 0.4 .

Our deduced Q value seems to indicate that decay to the first excited state of ^{179}Hf at 122.7 keV is energetically forbidden; however, we need a better assessment of the uncertainty in the Q value before a firm conclusion can be made.

2. Probability of producing K and L holes in the EC decay of ^{179}Ta

In addition to the singles measurement described above, we conducted a coincidence measurement to determine the probability of K -shell ionization accompanying L capture and L -shell ionization accompanying K capture. Although theoretical estimates indicate a small, but observable, probability ($\sim 10^{-5}$) for this process, there has been, to our knowledge, no experimental measurement of this probability.

For this measurement the source was sandwiched between the planar detector described above and an intrinsic coaxial Ge detector with a 44-mm diameter, 49-mm depth and a 0.5-mm-thick Be window. A 24 mg/cm²-thick Aluminum foil was inserted between the source and the coaxial detector to eliminate the detection of L x rays in this detector. The energy and timing signals from each detector were recorded event by event on magnetic tape over a period of several months. To measure the probability of ionization of both the K and L shells we look for coincidences between $K\beta$ and L x rays. As Fig. 2 shows, a $K\beta$ x ray is associated with the transfer of a K hole to an M or higher shell, while an L x ray is associated with an L hole; thus coincidences between $K\beta$ and L x rays would not occur unless there was simultaneous ionization of both shells. For ^{179}Ta there is no interference from $K\beta$ - L coincidences that arise in the filling of a double K -shell vacancy, since the Q_{EC} value of ^{179}Ta is less than twice the K -shell binding energy. Figure 3 shows L x-ray spectra recorded with the planar detector (a) in singles, (b) in true time coincidence with $K\beta$ x rays (recorded in the coaxial detector), and (c) in accidental coincidence with $K\beta$ x rays. Figure 4 shows K x-ray spectra recorded with the coaxial detector (a) in singles, (b) in true time coincidence with $L\beta$ x rays, and (c) in accidental coincidence with $L\beta$ x rays. Fig. 4(b) shows a substantial continuum of photons above the $K\alpha$ peaks in true coincidence with $L\beta$ x rays. These coincidences arise mostly from internal bremsstrahlung (IB) associated with capture from the L_2 and L_3 shells followed

³R.C. Jopson, H. Mark, C.S. Swift, and J.H. Zenger, Phys. Rev. **124**, 157 (1961).

⁴A. Bisi, L. Zappa, and E. Zimmer, Nuovo Cimento **4**, 307 (1956).

Figure 2: X-ray transitions used in determining probability of simultaneous creation of L and K holes.

by L_2 and L_3 x rays, and, to a much lesser extent, IB associated with L_1 capture followed by L_1 x rays.

The yield $N(L\gamma_3 \otimes K\beta)$ of $L\gamma_3$ x rays in coincidence with $K\beta$ x rays was extracted from a fit to the $L\gamma$ group shown in Fig. 3(b). This yield is related to the probability P_{KL_1} of double hole creation during electron capture by

$$N(L\gamma_3 \otimes K\beta) = N_0 P_{KL_1} n_{L_1 L\gamma_3} \epsilon_{L\gamma_3} n_{KK\beta} \epsilon_{K\beta}. \quad (6)$$

By dividing the above equation by the one relating the yield $N(L\gamma_1 \otimes K\alpha_2)$ of $L\gamma_1 \otimes K\alpha_2$ coincidences to the probability P_K of K capture,

$$N(L\gamma_1 \otimes K\alpha_2) = N_0 P_K n_{L_2 L\gamma_1} \epsilon_{L\gamma_1} n_{KK\alpha_2} \epsilon_{K\alpha_2}, \quad (7)$$

we get

$$\frac{P_{KL_1}}{P_K} = \frac{N(L\gamma_3 \otimes K\beta)}{N(L\gamma_1 \otimes K\alpha_2)} \frac{n_{L_2 L\gamma_1} n_{KK\alpha_2} \epsilon_{L\gamma_1} \epsilon_{K\alpha_2}}{n_{L_1 L\gamma_3} n_{KK\beta} \epsilon_{L\gamma_3} \epsilon_{K\beta}}. \quad (8)$$

In this preliminary analysis we have assumed $\epsilon_{L\gamma_1} \simeq \epsilon_{L\gamma_3}$ and $\epsilon_{K\alpha_2} \simeq \epsilon_{K\beta}$. From the measured values $N(L\gamma_3 \otimes K\beta) = 5 \pm 13$ and $N(L\gamma_1 \otimes K\alpha_2) = 9.01 \times 10^6$ we get $P_{KL_1}/P_K = (1.0 \pm 2.5) \times 10^{-5}$. Using our measured value of $P_{L_1}/P_K = 0.605$, we determine a probability per L_1 capture of producing a K and an L_1 hole during the EC

Figure 3: (a) Sample singles spectrum showing the L x-ray region recorded in the planar Ge detector. (b) Spectrum of L x rays in true coincidence with a gate set on the $K\beta$ region in the coaxial detector. (c) Same as (b) except for the time gate, which was set on accidental coincidences.

decay of ^{179}Ta of $P_{KL_1}/P_{L_1} = (1.7 \pm 4.1) \times 10^{-5}$. This limit is of the same order of magnitude as the theoretical result of Mukoyama, Kitahara and Shimizu⁵ (1.1×10^{-5}), but obviously not sensitive enough to confirm it.

Another limit on P_{KL} was deduced from the yield of $K\beta$ x rays in coincidence with $L\beta$ x rays. This yield was extracted by fitting the $K\beta$ region of the spectrum gated by $L\beta$ (Fig. 4(b)) with Gaussians superimposed on a quadratic background. Since all three L subshells contribute to these coincidences, the quantities P_{KL_i} cannot be deduced separately from this yield; rather the following combination of the P_{KL_i} can be extracted:

$$\frac{\sum_i P_{KL_i} \sum_j n_{L_i L \beta_j}}{P_K} = \frac{N(K\beta \otimes L\beta)}{N(K\alpha_2 \otimes L\beta)} \frac{n_{KK\alpha_2}}{n_{KK\beta}} \frac{\sum_j n_{L_2 L \beta_j} \epsilon_{K\alpha_2}}{\epsilon_{K\beta}}, \quad (9)$$

where we have assumed the efficiency for all $L\beta$ x rays to be the same. From our extracted yields we get $0.146P_{KL_1} + 0.209P_{KL_2} + 0.038P_{KL_3} = (2.2 \pm 8.6) \times 10^{-6}$ per L_1 capture. Mukoyama *et al.* find $P_{KL_1} = 1.1 \times 10^{-5}$, $P_{KL_2} = 0.375 \times 10^{-5}$, and $P_{KL_3} = 0.165 \times 10^{-5}$, which leads to a linear combination of 2.45×10^{-6} . Again, although our result is of the

⁵T. Mukoyama, T. Kitahara, and S. Shimizu, Phys. Rev. C 9, 2307 (1974).

Figure 4: (a) Sample singles spectrum showing the K x-ray region recorded in the coaxial Ge detector. (b) Spectrum of K x rays in true coincidence with a gate set on the $L\beta$ region in the planar detector. (c) Same as (b) except for the time gate, which was set on accidental coincidences.

same order of magnitude as the theoretical estimate of Mukoyama, it is not sensitive enough to test it quantitatively.

We plan to reanalyze the coincidence peaks using our two-dimensional fitting routines, after we adapt those to our new DECStation 5000/200 workstation. This should improve our results slightly; however, the dominant source of statistical uncertainty arises from the accidentals subtraction. Our time resolution (50 ns FWHM) is poor because of the very low photon energies we are detecting. It is unlikely that we can improve substantially upon this result with our present system.

Search for 17-keV Neutrinos in the Internal Bremsstrahlung Spectrum of ^{125}I

M.M. Hindi, R.L. Kozub, and S.J. Robinson

There have been claims of observation of a massive (≈ 17 keV) neutrino in the beta spectra of ^3H , ^{14}C , ^{35}S , ^{63}Ni , and in the internal bremsstrahlung (IB) spectra of ^{55}Fe and ^{71}Ge .¹ These claims are contradicted by several experiments, mostly performed on ^{35}S . The issue is certainly far from being settled. If neutrino mixing does occur, then it should be a universal phenomenon and should occur in electron capture decay, as well as in beta decay. We have begun a study of the IB spectrum of ^{125}I . A search for neutrino mixing in the IB spectrum of ^{125}I has been conducted before,² but that experiment was designed to test for a mixing fraction of about 2–4%; it does not reject the current claims of $\approx 1\%$ mixing fraction with a high degree of statistical confidence. Because of the low Q value (186.1 keV) and the exclusive decay to the first excited state of ^{125}Te at 35.5 keV, whose decay radiation is easily attenuated relative to the higher energy IB radiation, ^{125}I presents a favorable case for study. Furthermore, high intensity, relatively contaminant free, commercial sources of ^{125}I are readily available. Thus we decided to measure the IB spectrum of ^{125}I , with the aim of obtaining enough statistics to test the hypothesis of a 1.0% admixture of a 17 keV neutrino at a high confidence level.

A 200 mCi source was purchased from Amersham Corporation. Radiation from the source was detected in an intrinsic planar Ge detector with a 16-mm diameter, 10-mm depth and a 0.128-mm-thick Be window. Four copper sheets, each 0.5-mm thick, and backed by several Al foils with a total thickness of 0.5 mg/cm², were placed between the source and the detector. In addition to reducing the count rate of the Te x rays and the 35.5-keV γ ray in the detector, the copper served to reduce the summing probability of IB with x rays to negligible levels. (Since ^{125}I decays 100% of the time to the first excited state of ^{125}Te , the IB is always accompanied by internal conversion x rays or 35.5-keV γ rays from the decay of that level; thus, true summing of x rays or 35.5-keV γ rays with the IB is possible, leading to a distortion of the measured IB spectrum.) The transmission of the 35.5-keV γ rays through the copper sheets was measured to be 6×10^{-6} and that of the Te x rays to be 2×10^{-8} . These transmission probabilities then result in a total summing probability of less than 1×10^{-7} , which is totally negligible.

The detector and source were surrounded by a graded shield made of 0.1-cm-thick Al plates, 0.95-cm-thick Cu plates, and 20.3-cm-thick lead bricks. The signal from the preamplifier was fed into two amplifiers (ORTEC models 570 and 572) with 1- μs shaping times, but with different gains. One of the amplifiers was equipped with pileup rejection circuitry. The amplifier signals were digitized in two channels of a 13 bit (8196 channels)

¹See, for example, I. Žilimen *et al.*, Phys. Rev. Lett. **67**, 560 (1991), and references therein.

²M.J.G. Borge *et al.*, Phys. Scr. **34**, 591 (1986).

CAMAC analog-to-digital converter (ADC, ORTEC model AD413), with dispersions of 33 eV and 39 eV per channel. This duplication of signals should serve as a check that any observed distortion in the spectrum is not an artifact of the amplifiers or the ADC.

Four separate spectra were collected (pileup-accepted and pileup-rejected spectra for each of the two amplifiers) using a VAX 11/730 computer. The spectra were saved to disk once every eight hours. The ^{125}I was counted for a total of 31.5 d during a counting period that lasted 40 d. The count rate at the beginning of the counting period was 1300 s^{-1} . Background spectra were collected for a total counting time of 10.3 d.

The efficiency and response of the detector were determined from calibrated sources of ^{106}Cd and ^{57}Co ; for energy calibrations ^{133}Ba was used in addition to these sources. The energy resolution of the system at 122 keV was 0.560 keV. To test the stability of the system, several calibration spectra were collected before, during, and after the measurements. The standard deviation of the centroids of peaks in the calibration spectra and of the 35.5-keV γ ray peak was less than 20 eV. Hence, all the ^{125}I spectra and background spectra were summed together, without applying any shifts. The resulting spectra (compressed to 0.156 keV/channel) are shown in Fig. 1. There are 7.4×10^6 counts in the last 17 keV of the IB spectrum, and $\approx 1.1 \times 10^6$ counts/keV 17 keV below the end point.

Figure 1: Spectra of the ^{125}I source and of background collected for 31.5 d and 10.3 d, respectively, in a planar Ge detector. γ -ray energies are given in keV.

The dynamic range of the preamp-amp combination on the planar detector allowed a maximum photon energy of ≈ 320 keV to be observed. To check for contaminant lines at higher energies we counted the source with a 91 cm³ coaxial Ge detector for 10.7 d and the background for 6.5 d. The resulting background-subtracted spectrum is shown in Fig. 2. We found lines due to ¹²⁶I, which Amersham had indicated was present in the source at a level $< 5 \times 10^{-5}$. We confirmed that the lines are due to ¹²⁶I by following the decay curve of their activity and finding a half-life of 12.2 ± 2.1 d, which agrees with the ¹²⁶I half-life of 13.0 d. We also found a longer lived line at 662 keV, which we ascribe to ¹³⁷Cs. Since the source was made in a reactor by neutron capture on enriched ¹²⁴Xe gas ($^{124}\text{Xe} + n \rightarrow ^{125}\text{Xe}(17\text{h}) \xrightarrow{\text{EC}} ^{125}\text{I}$), the ¹³⁷Cs could have been produced by neutron capture on residual amounts of ¹³⁶Xe in the gas ($^{136}\text{Xe} + n \rightarrow ^{137}\text{Xe}(3.8\text{m}) \xrightarrow{\beta^-} ^{137}\text{Cs}$). We have searched for other contaminants, especially those that can be produced from neutron interactions with the other stable isotopes of Xe; we found no evidence for any other contaminant lines.

Figure 2: Background-subtracted ¹²⁵I spectrum measured with a 91 cm³ coaxial Ge detector. γ -ray energies are given in keV. The spectrum was shifted upwards by 200 counts for plotting purposes.

We are in the process of analyzing the data to search for the presence of a “kink” that can be ascribed to 17-keV neutrinos. We have also started another set of runs with the same source. The count rate at the start of these new runs was 650 s⁻¹, so that the pileup

contribution should be about half of its value in the previous set of runs. Also, the ^{126}I should have decayed away to completely negligible levels. The contribution of ^{137}Cs can be subtracted away by measuring the response of the planar detector to a ^{137}Cs source.

β^+ Decay and Cosmic-ray Half-life of ^{91}Nb

M.M. Hindi, B. Sur,[†] K.L. Wedding,[‡] K.R. Czerwinski,[†]
E.B. Norman,[†] K.T. Lesko,[†] and R.-M. Larimer[†]

Last year¹ we reported a measurement of the β^+ decay branch of ^{91}Nb (decay scheme shown in Fig. 1), to determine its suitability as a cosmic ray chronometer. This branch determines the lifetime of ^{91}Nb as a cosmic ray, since then it would be stripped of its

Figure 1: Decay schemes of ^{91}Nb and $^{91\text{m}}\text{Nb}$. Level energies are in keV.

electrons and the faster electron capture decay would not be possible. The branching ratio was determined from two sets of experiments, one to measure the K -capture decay rate (by detecting K x rays), and the other to measure the β^+ decay rate (by detecting 511-511 keV coincidences). Because the 61-day $^{91\text{m}}\text{Nb}$ isomer contributes to both types of radiation (Fig. 1), it was necessary to repeat the experiments over a period of several months to isolate the contribution of the shorter-lived isomer to the measured rates. Since the isomer gave the dominant contribution to the initial rates, we decided to perform

[†]Lawrence Berkeley Laboratory

[‡]Carlton College

¹R.L. Kozub and M.M. Hindi, *Progress Report on Research in Nuclear Physics, 1990-1991*, (DOE/ER/40314-5), 6 (1991).

another set of measurements 8 months later (4 isomer half-lives) at which point the contribution of isomer was expected to be small.

We performed a 511-511 keV coincidence measurement at LBL in July, 1991, to determine the β^+ decay rate of the source, and another set of coincidence and singles measurements at TTU during August and September, 1991, to determine the K x ray and β^+ rates. The results for ^{91}Nb and $^{91\text{m}}\text{Nb}$ are summarized in Tables 1 and 2, respectively. While the β^+ branching ratio extracted from the 1991 runs is consistent with that extracted from the 1990 runs, the absolute decay rates are $\sim 30\%$ lower ($\sim$ two standard deviations lower) in the later runs. The absolute rates for $^{91\text{m}}\text{Nb}$ in the 1991 runs are only $\sim 10\%$ lower than the rates we measured in 1990 (assuming a half-life of 60.87 ± 0.22 d for the isomer; if we adopt our absolute rates over the two sets of runs we get a half-life of 58.8 ± 1.1 d, in marginal agreement with the accepted value).

The disagreement between the 1990 and 1991 results ($\sim 2\sigma$) is not severe enough to permit a firm conclusion as to whether the decrease in rate attributed to ^{91}Nb is real or not. To insure that the β^+ activity is not due to contaminants, we have searched for γ rays from all known long-lived β^+ emitters. We found evidence for ^{22}Na (0.70 ± 0.07 pCi on August 19, 1991) and ^{152}Eu (2.1 ± 0.2 pCi on August 19, 1991). (The contribution of these has been taken into account in the values reported in Table 1.) The limit on the contribution of all other β^+ emitters is 5%.

We intend to perform another set of measurements on the same source this summer, to determine if the decrease in rate is real.

Table 1: Decay rates of the ^{91}Nb .

Date of measurement	Total rate (Bq)	β^+ rate (Bq)
November 16, 1990	270 ± 30	$(2.8 \pm 0.7) \times 10^{-2}$
July 24, 1991	-	$(1.4 \pm 0.5) \times 10^{-2}$
September 14, 1991	200 ± 11	$(1.8 \pm 0.4) \times 10^{-2}$

Table 2: Decay rates of the $^{91\text{m}}\text{Nb}$ isomer.

Date of measurement	K -conversion rate (Bq)	1205-keV rate (Bq)
November 16, 1990	1992 ± 60	(59 ± 2)
September 14, 1991	56.5 ± 3.7	(1.73 ± 0.04)

PUBLICATIONS AND SEMINARS

(June 1, 1991 - June 30, 1992)

Journal Articles:

Comment on 'Double K-shell Ionization in the Electron Capture Decay of ^{55}Fe '

M. M. Hindi, R. L. Kozub, H. J. Nagy, and G. Schupp, Phys. Rev. C **44**, 2237-2238 (1991).

Reply to 'Coherent Radiation, Not a Heavy Neutrino?'

B. Sur, E. B. Norman, F. E. Wietfeldt, K. T. Lesko, M.M. Hindi, R.-M. Larimer, P. N. Luke, W. L. Hansen, and E. E. Haller, Phys. Rev. Letters **68**, 1435 (1992).

Double K-Shell Vacancy Production in the Electron Capture Decay of ^{125}I

M. M. Hindi and R. L. Kozub, Phys. Rev. C **45**, 1070-1077 (1992).

Submitted Paper:

Yrast Decays in ^{43}K

R. L. Kozub, C. R. Bybee, M. M. Hindi, J. F. Shriner, Jr., R. Holzmann, R.V.F. Janssens, T.-L. Khoo, W. C. Ma, M. W. Drigert, U. Garg, and J. J. Kolata, Phys. Rev. C (submitted, 1992).

Invited Seminars:

Evidence for the Emission of a 17-keV Neutrino in Nuclear β Decay

M. M. Hindi, presented at Los Alamos National Laboratory (June 13, 1991), Roanoke College (October 9, 1991), and the University of Kentucky (October 11, 1991).

Double K-Shell Ionization in the Electron Capture Decays of ^{139}Ce and ^{125}I

M. M. Hindi, Sigma Xi Research Award Lecture, Tennessee Technological University (November 7, 1991).

TTU PERSONNEL

<u>Principal Investigators</u>	<u>Sources of Support</u>
R. L. Kozub, Professor	TTU, DOE
M. M. Hindi, Associate Professor	TTU, DOE
<u>Student Assistants</u>	
A. Altgilbers	DOE
D. Bardayan	DOE
R. Boyd	DOE
J. Parker	DOE
Y. Taketani	DOE
<u>Secretary/Account Clerk</u>	
G. J. Julian	TTU
<u>Other Nuclear Physics Faculty</u>	
S. Ayik, Associate Professor	TTU, DOE
K. Kumar, University Professor	TTU
D. P. Murdock, Assistant Professor	TTU, IU
S. J. Robinson, Assistant Professor	TTU, ORAU
P. B. Semmes, Assistant Professor	TTU
J. F. Shriner, Jr., Associate Professor	TTU, DOE
J. C. Wells, Jr., Professor	TTU, ORAU

DOE = Department of Energy
IU = Indiana University
ORAU = Oak Ridge Associated Universities
TTU = Tennessee Technological University

*preprints + reprints
removed, ok*

**DATE
FILMED**

8 / 14 / 92