2
BNL- 49182
AGS/AD/93-8
INFORMAL REPORT

RECEIVED
JUL 27 1993

OSTI

**CONSTRUCTION Of The COURANT-SNYDER INVARIANTS For The
NON-LINEAR EQUATIONS OF MOTION And CRITERION For The
LONG-TERM STABILITY Of The BEAM In A STORAGE RING**

V. Garczynski

April 26, 1993

Alternating Gradient Synchrotron
BROOKHAVEN NATIONAL LABORATORY
Upton, New York 11973

Under Contract No. DE-AC02-76CH00016 with the
United States Department of Energy

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Contents

Abstract	iv
1. Introduction	1
2. The Model	2
3. The Courant-Snyder Invariants as the Lyapunov Functions and the Long-Term Stability Criterion	3
4. Generalization of the Courant-Snyder Invariants to the Non-Linear Case and the Stability Criterion	7
5. Formal Solution of the Perturbed Equations of Motion	9
6. Formal Solution of the Equations for the Generalized CST Parameters	10
Acknowledgments	11
References	11

Abstract

The Courant-Snyder invariants become Lyapunov functions when the β -functions admit non-zero lower, and finite upper bounds. The long-term stability of motion then follows. This alternative criterion for the long-term stability of motion can be generalized to the non-linear case. A single particle subjected to an arbitrary static magnetic field is considered in some detail, as an example.

1. Introduction

The long-term stability of the beam in a storage ring has long been an unsolved problem.^{1,6,15-22} We would like to present here a new approach to this problem by extending classic Courant-Snyder invariants to the full, nonlinear equations of motion case. The point is that the invariants are, at the same time, Lyapunov functions⁷ when the relevant β -functions admit non-zero lower, and finite upper-bounds. The long-term stability of motion then follows. This criterion of stability can be easily transferred to the nonlinear case, in contradistinction to the usual criterion formulated in terms of the trace of a transfer matrix. Hence we replace the long-term stability problem for nonlinear equations of motion with the problem of existence of the non-zero lower bound, and the finite upper bound of the generalized β -functions. The latter problem can be approached, both analytically or numerically using electronic computers. In the last event one gains an insight into asymptotic regions of the survival plots which lie far beyond the reach of the conventional particle tracking techniques.

2. The Model

We start with the simplest case of interest of a single, on momentum, charged particle motion subjected to the Lorentz force produced by a static magnetic field. Possible effects of acceleration, linear coupling, beam intensity, power supplies ripples, residual gas scattering and other complications will not be treated here. We hope that this is excusable for the first paper on the subject. In the right (x, y, s) -particle accelerators coordinate system, the equations of motion can be written as follows⁹

$$\begin{aligned} \text{a.} \quad x' &= a + \varphi_x, \\ \text{b.} \quad a' &= -k_x x + \varphi_a, \\ \text{c.} \quad y' &= b + \varphi_y, \\ \text{d.} \quad b' &= -k_y y + \varphi_b. \end{aligned} \tag{2.1}$$

where we have used the usual notations

$$k_x = h^2 + hb_1, \tag{2.2}$$

and

$$k_y = -hb_1, \tag{2.3}$$

and

$$h = \frac{1}{\rho} = \frac{eB_0}{cp_0}, \tag{2.4}$$

are periodic functions of the position s in a ring, while the coordinates a, b

$$a = \frac{p_x}{p_0}, \quad b = \frac{p_y}{p_0}, \tag{2.5}$$

where p_0 is a kinetic momentum of the reference particle and p_x, p_y are its transversal components. In short, the equations of motion can be written as

$$z' = A(s)z + \varphi(z, s), \tag{2.6}$$

where

$$z = \begin{bmatrix} x \\ a \\ y \\ b \end{bmatrix} = \begin{bmatrix} X \\ Y \end{bmatrix}, \quad (2.7)$$

and, in absence of the linear coupling, the A -matrix is quasidiagonal

$$A = \begin{bmatrix} A_x & 0 \\ 0 & A_y \end{bmatrix}, \quad (2.8)$$

where

$$A_{x,y} = \begin{bmatrix} 0 & 1 \\ -k_{x,y} & 0 \end{bmatrix}. \quad (2.9)$$

The functions $\varphi(z, s)$ represent nonlinearities introduced by a static magnetic field, and by the relativistic kinematics factors, as well. The condition (2.4) ensures that the reference curve, $z = 0$, is a solution of the equations of motion

$$\varphi(0, s) = 0. \quad (2.10)$$

We are interested in the stability of the reference trajectory, (trivial, $z = 0$ solution), that is we ask whether nearby curves stay that way for sufficiently long time. More exactly, we will ask, first, the simpler question whether curves nearby the reference curve stay that way for all times. This problem is classic and over hundred years old by now. The first method, due to Poincaré,⁸ of solving the stability problem is to consider only the infinitesimally close trajectories which amounts to neglecting the higher order than linear terms in the equations of motion. The resulting linear equations, with periodic coefficients, (Floquet case),^{11,12}

$$z' = A(s)z, \quad (2.11)$$

are basic for the Courant-Snyder theory. The second method, introduced by Lyapunov, connects the stability issue with the existence of a function $v(z, s)$ satisfying the requirements¹⁴

$$1^\circ \quad v(z, s) \leq w(z), \text{ for all } s \geq s_0, \quad (2.12)$$

where

$$2^\circ \quad w(0) = 0, \quad (2.13)$$

and

$$3^\circ \quad w(z) > 0 \text{ when } z \neq 0, \quad (2.14)$$

and

$$4^\circ \quad \frac{dv}{ds} = \frac{\partial v}{\partial s} + \sum_{k=1}^4 \frac{\partial v}{\partial z_k} [A(s)z + \varphi(z, s)]_k \leq 0. \quad (2.15)$$

The very existence of such the function, called - Lyapunov function, entails the stability of the reference trajectory. Moreover, the stability is in the large, which means that the nearby curves may stay on some finite distance from the reference curve. We note, in

passing, that only knowledge of the equations of motion itself is sufficient to settle the question of stability of the zero solution, without actually solving the equations.

Following our strategy, we will briefly rederive essential elements of the Courant-Snyder theory and formulate an alternative, to the usual, condition for long-term stability of motion.

3. The Courant-Snyder Invariants as the Lyapunov Functions and the Long-Term Stability Criterion

The linearized equations of motion are of the Hamiltonian form

$$\begin{aligned} x' &= a, \\ a' &= -k(s)x, \quad k(s) = k(s+C). \end{aligned} \quad (3.1)$$

Similar equations hold for the vertical betatron motion. Since the motions are decoupled, it will be sufficient to consider only one of them, say – horizontal motion, in order to illustrate our point. General solution of the above equations is given by the transfer matrix

$$X(s) = M(s, s_0) X_0, \quad (3.2)$$

which is symplectic, and satisfies the conditions

$$\tilde{M} \sigma M = \sigma, \quad \sigma = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad (3.3)$$

$$M(s'', s) M(s, s') = M(s'', s'), \quad \text{for any } s, \quad (3.4)$$

$$M(s'', s') = M^{-1}(s', s''), \quad (3.5)$$

$$M(s'' + C, s' + C) = M(s'', s'), \quad C\text{-circumference.} \quad (3.6)$$

The matrix M admits convenient parameterization^{1,10} which is particularly simple for the single turn transfer matrix

$$M(s) = M(s+C, s) = \cos \mu \mathbf{1} + \sin \mu J(s), \quad (3.7)$$

where

$$J = \begin{bmatrix} \alpha & \beta \\ -\gamma & -\alpha \end{bmatrix}, \quad (3.8)$$

and where the Courant-Snyder-Twiss, (CST), parameters α, β, γ are periodic functions while the tune $-\mu$ is position independent. The parameters are constrained by the symplecticity requirement

$$\gamma = \frac{1}{\beta} (1 + \alpha^2). \quad (3.9)$$

It follows from the relations (3.4) - (3.6) that the single turn transfer matrices are connected as follows

$$M(s'') = M(s'', s') M(s') M^{-1}(s'', s'). \quad (3.10)$$

We may infer from this the following differential equations for the parameters

$$\beta' = -2\alpha, \quad (3.11)$$

$$\alpha' = k\beta - \gamma, \quad (3.12)$$

$$\gamma' = 2k\alpha. \quad (3.13)$$

The constraint (3.9) is compatible with these equations, and, in fact, the last equation follows from the previous ones, for any function k . The function β can always be assumed to be non-negative. Moreover, because of the equation

$$2\beta\beta'' - (\beta')^2 + 4k\beta^2 = 4, \quad (3.14)$$

which follows from the above equations, the assumption that, $\beta(s_0) = 0$, leads to an apparent contradiction. Hence, the β -function is strictly positive, periodic function which is continuous, together with its first derivative.

Knowing a solution of the linearized equations motion (3.1) one can construct the Courant-Snyder invariant¹

$$\epsilon = -(\sin \mu)^{-1} \tilde{X} \sigma M X = -\tilde{X} \sigma J X, \quad (3.15)$$

$$= \tilde{X} \begin{bmatrix} \gamma & \alpha \\ \alpha & \beta \end{bmatrix} X, \quad (3.16)$$

$$= \gamma x^2 + 2\alpha x a' + \beta a'^2, \quad (3.17)$$

$$= \frac{1}{\beta} [x^2 + (\alpha x + \beta a')^2] \geq 0. \quad (3.18)$$

Bringing the real symmetric matrix of this quadratic form to the diagonal form, by an orthogonal transformation,

$$R X = \xi, \quad (3.19)$$

one finds for the invariant the expression

$$\epsilon = \lambda_+ \xi_1^2 + \lambda_- \xi_2^2, \quad (3.20)$$

where the eigenvalues $\lambda_{\pm}$ satisfy the equation

$$\begin{vmatrix} \gamma - \lambda & \alpha \\ \alpha & \beta - \lambda \end{vmatrix} = \lambda^2 - \lambda(\beta + \gamma) + 1 = 0. \quad (3.21)$$

Hence, we get

$$\lambda_{\pm} = \frac{1}{2}(\beta + \gamma) \pm \left\{ \left[\frac{1}{2}(\beta + \gamma) \right]^2 - 1 \right\}^{1/2}. \quad (3.22)$$

Both roots are real because of the inequality

$$\frac{1}{2}(\beta + \gamma) \geq 1, \quad (3.23)$$

which follows from the well know inequality

$$\beta + \frac{1}{\beta} \geq 2, \quad (3.24)$$

valid for any non-negative real number β . Denoting

$$\frac{1}{2}(\beta + \gamma) = ch\chi = 1 + \kappa, \quad (3.25)$$

we get the estimates

$$\lambda_+ = ch\chi + (ch^2\chi - 1)^{1/2} = e^\chi \leq 2ch\chi, \quad (3.26)$$

and

$$\lambda_- = ch\chi - (ch^2\chi - 1)^{1/2} = e^{-\chi} \geq \frac{1}{2ch\chi}. \quad (3.27)$$

Also we get the inequalities

$$\lambda_+ = 1 + \kappa + (\kappa^2 + 2\kappa)^{1/2} \geq 1, \quad (3.28)$$

and

$$\lambda_- = 1 + \kappa - (\kappa^2 + 2\kappa)^{1/2} \leq 1. \quad (3.29)$$

As the result, the inequalities hold

$$0 \leq \lambda_- \leq 1 \leq \lambda_+ \leq \infty. \quad (3.30)$$

It follows from this and from the orthogonality of the matrix R , that the following inequalities are satisfied

$$\lambda_- (x^2 + a^2) \leq \epsilon \leq \lambda_+ (x^2 + a^2). \quad (3.31)$$

It is thus clear that the Courant-Snyder invariant will become a Lyapunov function when the eigenvalue λ_+ admits an upper bound, independent of s . It is obvious from the formula (3.22) that λ_+ attains its maximum value when the $\beta + \gamma$ does. This, in turn, implies an extreme of the β -function because of the relation

$$(\beta + \gamma)' = \beta' \left\{ 1 - \frac{1}{\beta^2} \left[1 + \left(\frac{\beta'}{2} \right)^2 - \frac{\beta\beta''}{2} \right] \right\}. \quad (3.32)$$

The expression in the curly bracket vanishes only when $k = 1$ as it follows from the equation (3.14) for the β -function. Assuming that $k \neq 1$, which can happen rather accidentally, we conclude that maximum of $ch\chi$ is attained when β -function has its extremum, i.e. when β' vanishes. Thus we obtain

$$\sup(ch\chi) = \sup \frac{1}{2} \left(\beta + \frac{1}{\beta} \right) \equiv \frac{1}{2}m, \quad (3.33)$$

where m is some fixed positive number, or infinity, which is greater of the two numbers

$$m = \max \left\{ \hat{\beta} + \frac{1}{\hat{\beta}}, \check{\beta} + \frac{1}{\check{\beta}} \right\}, \quad (3.34)$$

and where we have denoted, as usual

$$\hat{\beta} = \sup(\beta), \quad \check{\beta} = \inf(\beta). \quad (3.35)$$

Assuming that the number m is finite

$$0 < m < \infty, \quad (3.36)$$

which in turns implies that

$$\check{\beta} > 0 \text{ and } \hat{\beta} < \infty, \quad (3.37)$$

we get from (3.26) and (3.27) the inequalities

$$\lambda_+ \leq m, \quad (3.38)$$

and

$$\lambda_- \geq \frac{1}{m}. \quad (3.39)$$

As a result we obtain the estimates

$$\frac{1}{m} (x^2 + a^2) \leq \epsilon \leq m (x^2 + a^2). \quad (3.40)$$

Hence, the Courant-Snyder invariant becomes the Lyapunov function and the long-term stability of motion follows. In fact, the first inequality implies

$$|x(s)| \leq (\epsilon m)^{1/2} \leq A, \quad (2A - \text{aperture}), \quad (3.41)$$

when the emittance is restricted to

$$\epsilon \leq \frac{A^2}{m}. \quad (3.42)$$

This, in fact, makes it unnecessary to go any deeper into Lyapunov's theory itself, as the stability region became completely known. Thus the existence of bounds $\check{\beta}$ and $\hat{\beta}$, implies the long-term stability of motion, in the large. Hence, the essential limitation of Poincaré's approach, which secures the stability of only orbits which are infinitesimally close to the reference trajectory, is remedied. In the case when

$$m = \hat{\beta} + \frac{1}{\check{\beta}}, \quad (3.43)$$

one gets the well known condition on the emittance

$$\epsilon \leq \frac{A^2}{\hat{\beta} + \frac{1}{\check{\beta}}} \leq \frac{A^2}{\check{\beta}}. \quad (3.44)$$

If, however,

$$m = \check{\beta} + \frac{1}{\hat{\beta}}, \quad (3.45)$$

we obtain instead the restriction

$$\epsilon \leq \frac{A^2}{\tilde{\beta} + \frac{1}{\tilde{\beta}}} \leq \tilde{\beta} A^2. \quad (3.46)$$

To summarize, the long-term stability of motion is guaranteed when the β -function admits finite lower and upper bounds

$$0 < \tilde{\beta} \leq \hat{\beta} < \infty. \quad (3.47)$$

This, in turn, imposes restriction on the focusing function $k_x(s)$ which appears in the equations of motion (3.1), and in the equation for the β -function (3.14), as well.

Let us recall that the usual condition for stability, which requires that the tune μ to be a real number

$$\text{tr} M(s) = 2 \cos \mu \leq 2, \quad (3.48)$$

and thus entails that the characteristic multipliers, $e^{\pm i\mu s}$, became benign phase factors. However the phase factors do not play any role in the nonlinear case, and the above condition become meaningless when the nonlinearities are present.

We will assume in the sequel that the stability conditions (3.47) are satisfied on the level of the linearized equations of motion (3.1). We will then perturb the system, by including the nonlinear terms and we will ask for the stability conditions of the perturbed motion. Before doing so we will need to generalize the Courant-Snyder invariants to the nonlinear case.

4. Generalization of the Courant-Snyder Invariants to the Non-Linear Case and the Stability Criterion

Let us define the "quadratic" form

$$E = \tilde{X} \begin{bmatrix} \Gamma_x & A_x \\ A_x & B_x \end{bmatrix} X, \quad (4.1)$$

$$= \Gamma x^2 + 2Axa + Ba^2, \quad (4.2)$$

and similar for the vertical motion, where the new CST parameters B, A, Γ are assumed to be periodic functions of s satisfying the conditions similar to those from the linearized case

$$\Gamma B - A^2 = 1, \quad (4.3)$$

and

$$\begin{aligned} \frac{dB}{ds} &= -2A, \\ \frac{dA}{ds} &= KB - \Gamma, \\ \frac{d\Gamma}{ds} &= 2KA, \end{aligned} \quad (4.4)$$

where the function K is determined by the requirement that E is a constant of motion of the full equation (2.6)

$$\frac{dE}{ds} = 0. \quad (4.5)$$

As the result we get for the quantity K the expression

$$K = k + \Delta k, \quad (4.6)$$

where

$$\begin{aligned} \Delta k &= -\frac{1}{Ax^2 + Bxa} [(Aa + \Gamma x) \varphi_x + (Ax + Ba) \varphi_a], \\ &= \Delta k [\Omega(s), z(s), s]. \end{aligned} \quad (4.7)$$

The B -function has the properties similar to those of the original β -function. It is positive and satisfy the differential equation which follow from the equations (4.4)

$$2BB'' - (B')^2 + 4KB^2 = 4. \quad (4.8)$$

The difference is that now B, A, Γ depend on $z(s)$, and the last equation becomes highly nonlinear since the new coefficients enter the quantity K , also. Nevertheless, the generalized Courant-Snyder invariant is positive definite due to the constraint (4.3). The same sort of considerations, *mutatis mutandis*, as in the linear case apply here and we may write the estimates

$$\frac{1}{M} (x^2 + a^2) \leq E \leq M (x^2 + a^2), \quad (4.9)$$

where, as before [cf (3.40)],

$$M = \sup \left(B + \frac{1}{B} \right) = \max \left\{ \hat{B} + \frac{1}{\hat{B}}, \check{B} + \frac{1}{\check{B}} \right\}, \quad (4.10)$$

where again as before, $\hat{B}$ is maximum of B , and $\check{B}$ is its minimum value. Assuming that

$$M < \infty, \quad (4.11)$$

which holds when the bounds of B -function satisfy the restriction

$$0 < \check{B} \leq \hat{B} < \infty, \quad (4.12)$$

the long-term stability of motion, in the large, then follows. Assuming that

$$E \leq \frac{A^2}{M}, \quad (4.13)$$

we get the stability condition, for all times

$$|x(s)| \leq A, \quad s \geq s_0. \quad (4.14)$$

Similar considerations apply, obviously, to the vertical motion.

5. Formal Solution of the Perturbed Equations of Motion

We will apply the method of variation of constants when dealing with the equations (2.6) and (4.4). One starts with the linearized equations of motion (2.11) and finds a closed orbit solution first

$$z_{co}(s) = T(s, s_0) z_0 = z_{co}(s + C). \quad (5.1)$$

Then, one is seeking a solution of the full equation (2.6) in the form

$$z(s) = T(s, s_0) z_0(s), \quad (5.2)$$

where now the function $z_0(s)$ solves the resultant equation

$$z_0'(s) = T^{-1}(s, s_0) \varphi[T(s, s_0) z_0(s), s]. \quad (5.3)$$

Integrating and iterating one gets the result

$$z_0(s) = z_0 + \int_{s_0}^s T^{-1}(s', s_0) \varphi[T(s', s_0) z_0(s'), s'] ds' = \lim_{n \rightarrow \infty} z_0^{(n)}(s), \quad (5.4)$$

where we denoted

$$\begin{aligned} z_0^{(0)}(s) &= z_0, \\ z_0^{(1)}(s) &= z_0 + \int_{s_0}^s T^{-1}(s', s_0) \varphi[z_{co}(s'), s'] ds, \\ &\dots\dots\dots \end{aligned} \quad (5.5)$$

$$z_0^{(n)}(s) = z_0 + \int_{s_0}^s T^{-1}(s', s_0) \varphi[T(s', s_0) z_0^{(n-1)}(s'), s] ds', n = 1, 2, \dots$$

Applying the transfer matrix to $z_0(s)$ we thus obtain the solution for $z(s)$

$$z(s) = \lim_{n \rightarrow \infty} z^{(n)}(s), \quad (5.6)$$

where

$$\begin{aligned} z^{(0)}(s) &= z_{co}(s), \\ z^{(1)}(s) &= z_{co}(s) + \int_{s_0}^s T(s, s') \varphi[z_{co}(s'), s'] ds', \text{ etc.} \end{aligned} \quad (5.7)$$

As the result we may write

$$z(s) = z_{co}(s) + \Delta z[z_{co}^{(s)}, s], \quad (5.8)$$

where $z_{co}^{(s)}$ denotes a functional dependence of the Δz on the $z_{co}(s')$, for $s_0 \leq s' \leq s$.

6. Formal Solution of the Equations for the Generalized CST Parameters

Denoting

$$\omega = \begin{bmatrix} \beta \\ \alpha \\ \gamma \end{bmatrix}, \quad \Omega = \begin{bmatrix} B \\ A \\ \Gamma \end{bmatrix}, \quad \psi = \begin{bmatrix} \psi_B \\ \psi_A \\ \psi_\Gamma \end{bmatrix}, \quad (6.1)$$

and

$$\mathcal{A} = \begin{bmatrix} 0 & -2 & 0 \\ k & 0 & -1 \\ 0 & 2k & 0 \end{bmatrix}, \quad \text{periodic matrix}, \quad (6.2)$$

where

$$\begin{aligned} \psi_B &= 0, \\ \psi_A &= \Delta k B, \\ \psi_\Gamma &= 2\Delta k A, \end{aligned} \quad (6.3)$$

we may write the equations (4.4) in the form

$$\Omega' = \mathcal{A}(s) \Omega + \psi[\Omega, z(s), s]. \quad (6.4)$$

The same method of solving it as before yields

$$\Omega(s) = \lim_{n \rightarrow \infty} \Omega^{(n)}(s), \quad (6.5)$$

where

$$\begin{aligned} \Omega^{(0)}(s) &= \omega(s), \\ \Omega^{(1)}(s) &= \omega(s) + \int_{s_0}^s \mathcal{M}(s, s') \psi[\omega(s'), z(s'), s'] ds', \text{ etc.} \end{aligned} \quad (6.6)$$

Here, $\omega(s)$ is the periodic solution of the linearized equation

$$\omega' = \mathcal{A}(s) \omega, \quad (6.7)$$

and

$$\omega(s) = \mathcal{M}(s, s_0) \omega_0 = \omega(s + C). \quad (6.8)$$

Similar considerations apply to the vertical motion and yield the analogous result

$$\Omega(s) = \omega(s) + \Delta \omega[\omega(s), z(s), s]. \quad (6.9)$$

Knowing the B -function one may approach the question of existence of the finite lower, $\tilde{B}$, and upper, $\hat{B}$, bounds. This in turn imposes restrictions of the nonlinearities under which the long-term stability of the beam holds.

We postpone applications of the scheme, to some concrete accelerating and steering structures, to future publications.

Acknowledgments

I would like to thank Ernest Courant and Bill Weng for their interest in the work and comments. Many thanks to Dr. G. Swiatek of the Mathematics Department of SUNY at Stony Brook for several illuminating discussions on Lyapunov's method. I would like to thank Melvin Month for his interest, encouragement and patience, and for several informative and critical discussions on the long-term stability problem.

References

1. E.D. Courant and H.S. Snyder, *Ann. Phys.* 3 (1958) 1.
2. A.N. Kolmogorov, *Dokl. Acad. Nauk USSR*, 98(4), 527 (1954).
3. V.I. Arnold, *Mathematical methods in classical mechanics*, Moscow, 1974.
4. J. Moser, *Stable and Random Motion in Dynamical Systems* (Princeton University Press, Princeton, New Jersey, 1973).
5. E.D. Courant, R.D. Ruth and W.T. Weng, *Stability in Dynamical Systems*, SLAC- PUB-3415 (1984).
6. J. Wisdom, *Astron. J.* 87, 577 (1982).
7. A.M. Lyapunov, "Probleme général de la stabilité du mouvement." *Ann. of Math. Studies* 17, Princeton University Press, 1947, p. 259. [Original Russian, dated 1892. This is a reproduction of the French translation dated 1907.]
8. H. Poincaré, *Methods nouvelle de la mécanique céleste*, Vol. 1, Gauthier-Villars, Paris, 1892.
9. J. Bengtsson, "Non-linear transverse dynamics for storage rings with applications to the low-energy antiproton ring (LEAR) at CERN", CERN 88-05, 1 August 1988.
10. R.Q. Twiss and N.H. Frank, *Rev. Sci. Instr.* 20, 1 (1949).
11. G. Floquet, *Ann. Ec. Normale* 12, 47-88, 1883.
12. N. Minorsky, *Nonlinear oscillations*, Robert E. Krieger Publishing Company, Malabar, Florida, 1987.
13. A. Lichtenberg, M. Lieberman, *Regular and Stochastic Motion*, Springer-Verlag, New York, Inc. 1983.
14. Y. Yoshizawa, *Stability Theory and the Existence of Periodic Solutions and Almost Periodic Solutions*, Springer-Verlag, New York Inc. 1975.
15. A. Ando, *Proc. 12th Int. Conf. on High-Energy Accelerators* (Fermilab, 1983).
16. F. Willeke, in "Accelerator Physics Issues for Superconducting Super Collider", M. Tigner, editor, U. Michigan Report UM HE 84-1 (Dec. 1983).
17. T. Collins, "Distortion Functions", FNAL Tech Note Fermilab-84/114 (Oct. 23, 1984).
18. A.J. Dragt, "Nonlinear Lattice Functions," in *Proc. 1984 Summer Study on the Design and Utilization of the Superconducting Super Collider*, R. Donaldson and J.G. Morfin, editors (Snowmass, 1984).
19. E. Forest, "Lie Algebraic Methods for Charged Particle Beams and Light Optics", Ph.D. thesis, U. Maryland (unpublished, 1984), and "Normal Form Algorithm on Nonlinear Symplectic Maps," Superconducting Super Collider Central Design Group Report SSC- Report 29.
20. L. Michelotti, *Particle Accelerators* 16, 233-252 (1985) and 19, 205-210 (1986).

21. D.R. Douglas, "Dynamic Aperture Calculations for Circular Accelerators and Storage Rings", in AIP Conference Proceedings 153, Physics of Particle Accelerators, M. Month and M. Dienes, editors, Vol. 1, p. 392-473, New York 1987.
22. N. Merminga and K-Y. Ng, "Hamiltonian Approach to Distortion Functions," FN-493, Fermilab (August, 1988).

END

DATE
FILMED
9 / 10 / 93