UNITED STATES DEPARTMENT OF THE INTERIOR GEOLOGICAL SURVEY



AIRBORNE RADIOACTIVITY SURVEY OF PART OF SARATOGA NW QUADRANGLE, CARBON COUNTY, WYOMING

FEBRUARY 1954

PREPARED IN COOPERATION WITH UNITED STATES ATOMIC ENERGY COMMISSION

106° 45'

TRACE ELEMENTS MEMORANDUM **REPORT 746**

RELEASED TO OPEN FILE:

JUNE 25, 19 HALLANA

201115 65 AP26

41° 30 Walter Krd 11 20

AIRBORNE RADIOACTIVITY SURVEY OF PART OF SARATOGA NW QUADRANGLE, CARBON COUNTY, WYOMING

by J. R. Henderson

The accompanying map shows the results of an airborne radioactivity survey in 133 square miles of Saratoga NW quad rangle, Wyoming. This area is part of a larger survey made in southern Carbon and Sweetwater Counties by the U. S. Geological Survey, November 9-24, 1953. The work was undertaken as part of a cooperative program with the U. S. Atomic Energy Commission.

The survey was made with scintillation detection equipment mounted in a C-47 aircraft and consisted of parallel east-west flight lines spaced at quarter mile intervals, flown approximately 500 feet above the ground. Aerial photographs were used for pilot guidance, and the flight path of the aircraft was recorded by a gyrostabilized, continuousstrip-film camera. The distance of the aircraft from the ground was measured with a continuously recording radio altimeter.

The width of the zone on the ground from which the anomalous radiation is measured at the nominal 500 foot flight altitude varies with the areal extent and the intensity of radioactivity the width of the zone may be as much as 1,400 feet. Thus, quarter mile spacing of the flight lines could be adequate to detect anomalies from strong sources of radioactivity; however, small areas of considerable radioactivity midway between the flight lines may not be noted.

The approximate locations of twenty-two anomalies are shown on the accompanying map. The plotted position of these anomalies may be in error by as much as a quarter mile owing to errors in available base maps or to the existence of areas on the base maps up to several square miles in which it is impossible to find and plot recognizable landmarks.

The present technique of airborne radioactivity measurement does not permit distinguishing between activity due to thorium and that due to uranium. An anomaly, therefore, may represent radioactivity due entirely to one or to a combination of these elements. The radioactivity anomalies shown on the accompanying map cannot be interpreted in terms of either the radioactive content or the extent of the source materials. Any particular anomaly may represent slightly greaterthan-average radioactivity over an area of a few thousand square feet or high radioactivity over an area of a few hundred square feet.

These radioactivity anomalies indicate localities of more-than-average radioactivity and, therefore, suggest areas in which uranium or thorium deposits are more likely to occur.



This map has been receased without editorial and technical review for conformity with U-S- Geological Survey standards and nomenclature.

metadc1393117

Der man

.

.

And the second second

RELEASED TO OPEN FILE: JUNE 25. 1954

.

