1 of 1

BNL-49409

Exact Physical Model for Magnets in Storage Rings*

D.Maletić, A.G.Ruggiero

Brookhaven National Laboratory, Upton, NY 11793

1 Motivation

It is common practice to perform computer simulation to determine the stability of motion of charged particles circulating in storage rings over long periods of time. For this purpose several computer codes have been written.¹ In the past, great effort was spent to insure that the representation of the motion used during simulation was symplectic;² that is, for conservative systems of particles, like protons, antiprotons and heavy ions, the equations of motion not only are to be derived from a Hamiltonian, but they also have to satisfy a variety of momentum and energy conservation laws. If these should not become fully satisfied, the results of such computer trackings, especially long ones, could be invalid. Simulated motion could show instability where it should not occur, and vice versa.

A particle encounters several types of elements during its motion in a storage ring. Some of them have linear properties, like drifts and pure quadrupoles. In these cases, the approximate status of motion of the particle at the end of the particular element can be easily given as an integrable function of the status of motion at the beginning of that element. Nevertheless, even in these linear or quasi-linear elements the exact equations of motion have kinematic terms that are difficult to integrate.

Other elements encountered by the particle during its motion in a storage ring are nonlinear, like sextupoles, intentionally used to correct chromatic effects, and the nonlinear imperfections of magnets, especially superconducting magnets. The motion in these nonlinear elements cannot be generally integrated. Therefore one has to use some numerical technique to perform integration by computer.

One method is to lump each nonlinear element into a thin lens of zero length. Motion through such an element is then represented by a kick. The

*Work performed under the auspices of the U.S. Department of Energy

MASTER

position coordinates of the particle remain unchanged, while velocity receives a sudden change that depends on the particle position and on the properties and strength of the nonlinear lens. The numerical codes that make use of this method are called *kick codes* or *exact codes*. Since the motion they represent with a kick is truly symplectic, their approximation of reality is physically correct. TEAPOT³ is one of these codes which also represents integrable elements, like dipoles and quadrupoles, as a series of thin lenses and drifts. Since the motion in a drift can be calculated exactly, it seems that the representation of any accelerator element as a sequence of drifts and thin lenses, linear and nonlinear, is indeed very powerful.

It should be understood that in order to predict the stability of the motion over very long periods of time it is mandatory that all kinematic terms be properly included in the model. Neglecting some of them may invalidate the results of very time-consuming exercises on the computer.

Adopting the thin lens representation requires discussion of one more issue. Forces acting on a particle are to be derived exactly and consistently from Maxwell's equations. Generally, ordinary kick codes do not enforce a complete description of the fields, and this is to be corrected. Not only do magnets exhibit edge effects due to their discontinuity in space, but the thin lens model of either linear or nonlinear elements is a three-dimensional field representation that requires careful estimation of the field components. Usually only the longitudinal component of the vector potential is retained for the field estimate. Since this has discontinuous behaviour, transverse components are also present, especially at the edges of the elements; these are usually neglected. If the transverse components of the field are neglected, the divergence of the vector potential does not vanish and the divergence and the curl of the magnetic field differ from zero, though the corresponding equations of motion remain symplectic. This, again, may invalidate results of long computer simulations. Thus, we consider it very important that the field representation be exact and complete and that it satisfy Maxwell's equations.

In this report we try to make estimates of both kinematic and field effects on the stability of a particle motion, by employing a truly Maxwellian representation of the magnetic field in exact equations of motion. For this purpose we adopt a simple FODO cell model, which repeats periodically to infinity. This model includes only quadrupoles and drifts, leaving out the bending magnets to avoid the problem of the trajectory curvature. We think this model is a physically consistent approximation of a storage ring. We derive several models with different levels of approximation and compare them by evaluating the importance of these effects. The relevance to long-term stability is being investigated in the meantime by comparing the different models with extensive computer simulations. The results will be shown in a subsequent report.

2 The Test Model

To determine the importance and the magnitude of kinematic effects, and of the effects of the field components required for complete Maxwellian representation of the field, we adopt the following test model. It consists of an infinitely long sequence of FODO cells, all identical, as shown in Figure 1.

Figure 1: The test model as an infinite sequence of identical FODO cells

The FODO cell has the following structure:

$$QF/2 \quad L \quad QD/2 \quad QD/2 \quad L \quad QF/2 \quad (1)$$

which begins from the middle of the horizontally focusing quadrupole (QF) and ends in the middle of the following focusing quadrupole. The vertically focusing quadrupole (QD) is located halfway. The quadrupoles have the same length l and are separated by a drift of length L . Quadrupole magnets are described by the field gradient. The gradient is $+G$ in QF and $-G$ in QD. The only other parameter required for calculating the motion of a charged particle of charge q and momentum p is the particle magnetic rigidity

$$B\rho = pc/q \quad (2)$$

where c is the speed of light.

Our test model is made only of drifts and quadrupoles. For simplicity we did not include dipole magnets to avoid the issue of the curvature they introduce.

The reference line is the straight line that coincides with the symmetry axis of the quadrupole. We assume that quadrupoles are perfectly aligned so that their axes coincide. The quadrupole fields vanish on this axis.

This model is basically an approximation of a storage ring where the reference orbit is a straight line. We introduce a longitudinal coordinate s along the reference line and a pair of rectangular transverse coordinates, x

for horizontal and y for vertical, which measure the distance of a particle trajectory from the reference line; that is, on the reference line $x = y = 0$.

3 Equations of Motion

To describe the motion of a particle it is convenient to consider the plane perpendicular to the main axis of motion (the reference orbit) which includes the particle position at time t . It is on this plane that the distances x and y from the reference axis are evaluated at time t . To the same instant t one can associate the longitudinal coordinate s , which is given by the interception of the $x-y$ plane with the reference orbit. Since there is a unique correspondence between s and t , it is possible and actually more convenient to take the longitudinal coordinate s as the independent variable, rather than the time t . The displacements are then taken as functions of s , that is, $x = x(s)$ and $y = y(s)$.

In the following we use a prime to denote the derivative with respect to s , and we denote the components of the magnetic field as B_x , B_y and B_s .

It can be proven⁴ that the exact equations of motion for the FODO test model are

$$x'' = -\frac{q}{cp}\sqrt{1+x'^2+y'^2}[B_y(1+x'^2) - B_x x' y' - B_s y'] \quad (3)$$

$$y'' = \frac{q}{cp}\sqrt{1+x'^2+y'^2}[B_x(1+y'^2) - B_y x' y' - B_s x'] \quad (4)$$

It can also be proven⁴ that the following relation holds:

$$x' x'' + y' y'' = \frac{q}{p}(1+x'^2+y'^2)^{3/2}(y' B_x - x' B_y) \quad (5)$$

which may be found useful in a variety of situations.

4 A Symplectic Integration

The general integration of the system consisting of Eqs. (3) and (4) is very difficult. It may not be possible to prove their integrability in general. Two features that contribute to the difficulty of the problem are:

1. There are kinematic terms represented by the x' and y' factors which are usually neglected.
2. There are field contributions, represented by the components B_x , B_y and B_s . Usually the longitudinal component is neglected as well as the edge effects.

We need to find a symplectic integration method that allows us to solve the equations of motion, satisfying all energy and momentum conservation laws. Conventional methods, like Newton, Lagrange, Runge-Kutta, etc., may not be valid for this purpose. We find the “TEAPOT-like” method of replacing each element with a sequence of thin lenses and drifts, as shown in Figure 2, a valid one.

Figure 2: Replacing a quadrupole with a sequence of thin lenses and drifts

The length l of the quadrupole magnet is divided into N steps, each consisting of a drift of length $\lambda = l/N$ with a thin lens quadrupole in the middle. Each of these N thin lenses is a quadrupole magnet with zero length. In the limit $\lambda \rightarrow 0$ or $N \rightarrow \infty$, one can recover the original model of the full-length quadrupole.

In a drift $B_x = B_y = B_s = 0$, so that the equations of motion are simply

$$x'' = y'' = 0 \quad (6)$$

with solutions represented by straight-line trajectories.

The integration through the thin lens quadrupoles is shown in Figure 3. The trajectory of the particle consists of straight lines before and after the thin lens. The trajectory is deflected by the thin lens, leaving the position of the particle unchanged and altering the velocity by an amount that depends on x , y , x' and y' as well as on the field gradient in the magnet.

Figure 3: The trajectory of a particle going through a thin lens

5 Exact Field Expression

We wish to calculate the exact expression for the magnetic field distribution of a finite-length n -pole magnet, including the distribution at the edges. For this purpose we introduce two functions: the dimensionless function $f(s)$ which defines the longitudinal shape of the magnet and

$$g_n(x, y) = (a_n + ib_n)(x + iy)^n \quad (7)$$

which describes the longitudinal field potential inside the n -pole magnet. It is easily verified that

$$\nabla^2 g_n(x, y) = 0. \quad (8)$$

Let $\mathbf{B}$ and $\mathbf{A}$ be respectively the magnetic field and the magnetic vector potential. They satisfy

$$\nabla \cdot \mathbf{A} = 0, \quad (9)$$

$$\nabla^2 \mathbf{A} = 0, \quad (10)$$

and

$$\mathbf{B} = \nabla \times \mathbf{A}. \quad (11)$$

Since we are using a rectangular coordinate system (x, y, s) Eq. (10) is equivalent to

$$\nabla^2 A_x = \nabla^2 A_y = \nabla^2 A_s = 0. \quad (12)$$

It can be proven by direct substitution that the following expressions solve Eqs. (9) and (10) for any chosen functions $f(s)$ and $g_n(x, y)$:

$$A_s(x, y, s) = g_n \sum_{m=0}^{\infty} C_m (x^2 + y^2)^m f^{(2m)}, \quad (13)$$

$$A_x(x, y, s) = -g_n(x + iy) \sum_{m=0}^{\infty} \frac{C_m}{2(m + n + 1)} (x^2 + y^2)^m f^{(2m+1)}, \quad (14)$$

$$A_y(x, y, s) = -g_n(y - ix) \sum_{m=0}^{\infty} \frac{C_m}{2(m + n + 1)} (x^2 + y^2)^m f^{(2m+1)}, \quad (15)$$

where the coefficients C_m are related to each other according to

$$C_{m-1} = -4m(m+n)C_m , \quad (16)$$

$$C_0 = 1 , \quad (17)$$

and

$$f^{(m)} = \frac{d^m f(s)}{ds^m} . \quad (18)$$

Inserting Eqs. (13) to (15) into Eq. (11) yields

$$B_x = ig_n \sum_{m=0}^{\infty} C_m [(2m+n)x - iny](x^2 + y^2)^{m-1} f^{(2m)} , \quad (19)$$

$$B_y = ig_n \sum_{m=0}^{\infty} C_m [(2m+n)y + inx](x^2 + y^2)^{m-1} f^{(2m)} , \quad (20)$$

$$B_s = ig_n \sum_{m=0}^{\infty} C_m (x^2 + y^2)^m f^{(2m+1)} . \quad (21)$$

As a special case let us consider a quadrupole magnet in normal orientation, that is, $n = 2$, $b_2 = 0$ and $a_2 = -G/2$. After taking the imaginary part to obtain the real field representation, we have

$$B_x = Gy f^{(0)} + G \sum_{m=1}^{\infty} C_m (x^2 + y^2)^{m-1} [(2m+1)x^2 y + y^3] f^{(2m)} , \quad (22)$$

$$B_y = Gx f^{(0)} + G \sum_{m=1}^{\infty} C_m (x^2 + y^2)^{m-1} [(2m+1)xy^2 + x^3] f^{(2m)} , \quad (23)$$

$$B_s = G \sum_{m=0}^{\infty} C_m xy (x^2 + y^2)^m f^{(2m+1)} . \quad (24)$$

If we retain only the linear terms in x and y , we get

$$B_x = Gy f(s) , \quad (25)$$

$$B_y = Gx f(s) , \quad (26)$$

$$B_s = 0 . \quad (27)$$

6 Linear Model

Let us neglect all the kinematic terms in Eqs. (3) and (4) and retain only the linear terms in the field expansion, that is, Eqs. (25) to (27), where $f(s)$ is a step function equal to 1 in the quadrupole and zero everywhere else.

The resulting equations of motion in a quadrupole are

$$x'' = -Kx , \quad (28)$$

$$y'' = Ky , \quad (29)$$

where, for convenience,

$$K = \frac{q}{cp} G = \frac{G}{B\rho} . \quad (30)$$

In the drifts, between quadrupoles, the equations of motion are

$$x'' = y'' = 0 . \quad (31)$$

These equations are linear. They can be easily solved and their solution is usually represented by the matrix notation which involves the lattice functions⁵ β_H and β_V as well as the phase advance functions⁵ Ψ_H and Ψ_V .

As an example we take the following values, which approximate regular cells in the arcs of RHIC⁶:

$$B\rho = 840 \text{ Tm} , \quad (32)$$

$$G = 81 \text{ T/m} , \quad (33)$$

$$L = 14 \text{ m} , \quad (34)$$

$$l = 1 \text{ m} . \quad (35)$$

The lattice functions for this system, obtained with the SYNCH code,⁷ are plotted in Figure 4. The quadrupole gradient has been chosen to make the phase advances across one FODO cell equal to 90° in both planes.

7 The TEAPOT Model

Let us now examine the approximation of a quadrupole magnet by a sequence of thin lens magnets and drifts, as shown in Figure 2. We still adopt the linear approximation, as in the TEAPOT code.³ The only difference from the equations of motion of the linear model is that the function $f(s)$ is a delta function at each thin lens, positioned at $s = s_i$.

Figure 4:
The amplitude
lattice functions
for the FODO cell

$$f(s) = \lambda \delta(s - s_i) \quad (36)$$

where $\lambda = l/N$ is the length of a step approximated by one thin lens and a drift.

The position coordinates x and y of a particle crossing the thin lens are unchanged, that is

$$\Delta x = \Delta y = 0. \quad (37)$$

At the same time the angles x' and y' receive a change

$$\Delta x' = -K \lambda x, \quad (38)$$

$$\Delta y' = K \lambda y. \quad (39)$$

This model can be easily integrated and investigated. The results are given for comparison in Table 1. The maximum and minimum values of the amplitude functions are reported as well as the phase advance across the FODO cell, $\Delta\Psi/2\pi$. The results for the linear model in Table 1 were derived with the SYNCH code.⁷ The results of the approximation of the whole quadrupole by one thin lens were computed manually, since for this case the following relations hold:

$$\Lambda = \frac{L + l}{2}, \quad (40)$$

$$\sin\left(\frac{\Delta\Psi}{2}\right) = \frac{\Lambda|K|}{2}, \quad (41)$$

Table 1: Comparison of different linear models

MODEL	β_{max}	β_{min}	$\Delta\Psi/2\pi$
Linear Model	50.80254	8.86074	0.24992
Thin Lens	51.75140	8.31240	0.25734
TEAPOT (2)	50.71639	8.77329	0.25176
TEAPOT (4)	50.78029	8.83885	0.25038
TEAPOT (10)	50.79895	8.85724	0.24999
TEAPOT (20)	50.80164	8.85987	0.24994
TEAPOT (100)	50.80231	8.86005	0.24992
TEAPOT (200)	50.80250	8.86071	0.24992
Simpson-Bode (2)	51.15743	8.73205	0.25115
Simpson-Bode (3)	50.78102	8.83882	0.25038
Simpson-Bode (4)	50.75739	8.84859	0.25027

$$\beta_{max} = \frac{2}{l|K|} \sqrt{\frac{2 + \Lambda l|K|}{2 - \Lambda l|K|}}, \quad (42)$$

$$\beta_{min} = \frac{2}{l|K|} \sqrt{\frac{2 - \Lambda l|K|}{2 + \Lambda l|K|}}. \quad (43)$$

The other entries in Table 1 were obtained with a TEAPOT type of code, for several different numbers of sequences into which each quadrupole is subdivided. The last three entries correspond to the Simpson-Bode⁸ method of integration, where magnets are subdivided into thin lenses each carrying a different weighted strength. Several different numbers of subdividing steps (shown in parentheses) were used.

It can be seen that the single thin lens model provides results only a few percent from those of the exact linear model. Subdividing the quadrupole into more thin lenses gives a closer approximation. Nevertheless, even if we divide the magnet into 200 thin lenses the results are off by an error of 10^{-5} for the amplitude function.

It has been assumed that a more effective way of integrating the motion through a magnet is the Simpson-Bode method. This method is commonly used in tracking particle motion through nonlinear elements. Its effectiveness cannot be easily proven except in a linear case, where the results can be compared to the exact solution. The results in Table 1 show that the Simpson-Bode method has better convergence than the method using TEAPOT procedure.

Let (x, x') and (y, y') be the particle position and angle in the middle of

any of the two quadrupoles, QF and QD. Then the following quantities are invariant⁵

$$\epsilon_H = \frac{x^2}{\beta_H} + x'^2 \beta_H, \quad (44)$$

$$\epsilon_V = \frac{y^2}{\beta_V} + y'^2 \beta_V. \quad (45)$$

The values of the amplitude functions β_H and β_V vary somewhat from model to model as shown in Table 1. In the following we take the linear model as the reference one.

8 The Kinematic Model

The kinematic model is defined by the set of exact equations (3) and (4), where kinematic terms are retained to any order, but the field expansion is truncated beyond the linear terms, Eqs. (25) to (27). This model is truly symplectic.

Investigation of this model is relevant to determining the magnitude and the importance of the effects of the kinematic terms. This can be done by comparison with the linear model.

We continue to approximate a quadrupole with a sequence of thin lenses and drifts. Indeed this is the only method we know for numerical symplectic integration.

The equations of the motion through a thin quadrupole now become

$$x'' = -K\lambda\sqrt{1+x'^2+y'^2}(x+xx'^2-yy'y')\delta(s), \quad (46)$$

$$y'' = K\lambda\sqrt{1+x'^2+y'^2}(y+yy'^2-xx'y')\delta(s) \quad (47)$$

and can be integrated exactly, yielding kicks

$$\Delta x' = -K\lambda\sqrt{1+x'^2+y'^2}(x+xx'^2-yy'y'), \quad (48)$$

$$\Delta y' = K\lambda\sqrt{1+x'^2+y'^2}(y+yy'^2-xx'y') \quad (49)$$

where x , x' , y and y' on the right correspond to the instant just before entering the lens, as proven in the Appendix. At the same time x and y remain unchanged.

We have not been able to derive invariants similar to those given for the linear model. Nevertheless, it can be seen that the kinematic terms introduce coupling between horizontal and vertical motion. The coupling disappears only if the motion is purely horizontal ($y = y' = 0$) or purely vertical ($x = x' = 0$). In these cases the equations of the motion reduce to either

$$x'' = -K\lambda x(1 + x'^2)^{3/2}\delta(s) , \quad (50)$$

$$y'' = 0 , \quad (51)$$

or

$$x'' = 0 , \quad (52)$$

$$y'' = K\lambda y(1 + y'^2)^{3/2}\delta(s) . \quad (53)$$

Let us take, for instance, the purely horizontal motion. In this case we can decouple our second-order differential equation of motion, Eq. (50), into two first-order differential equations:

$$x' = \frac{p_x}{\sqrt{1 - p_x^2}} , \quad (54)$$

$$p'_x = -K\lambda x\delta(s) . \quad (55)$$

which can be derived from the Hamiltonian

$$H = 1 - \frac{K\lambda}{2}x^2\delta(s) - \sqrt{1 - p_x^2} . \quad (56)$$

9 The Maxwellian Field Model

The Maxwellian field model is obtained by neglecting all the kinematic terms in Eqs. (3) and (4) but retaining the exact field series representation, Eqs. (22) to (24). Only B_x and B_y field components enter the equations:

$$x'' = -\frac{q}{cp}B_y , \quad (57)$$

$$y'' = \frac{q}{cp}B_x . \quad (58)$$

Inserting Eqs. (22) to (24) we obtain

$$x'' = -K\lambda\{x\delta(s) + \sum_{m=1}^{\infty} C_m(x^2 + y^2)^{m-1}[(2m+1)xy^2 + x^3]\delta^{(2m)}(s)\} , \quad (59)$$

$$y'' = K\lambda\{y\delta(s) + \sum_{m=1}^{\infty} C_m(x^2 + y^2)^{m-1}[(2m+1)x^2y + y^3]\delta^{(2m)}(s)\} , \quad (60)$$

where again a thin lens representation has been assumed.

The integration of Eqs. (59) and (60) is cumbersome, but it can be exactly derived. If we retain only the linear, $m = 1$ and $m = 2$, terms we obtain

$$\begin{aligned} \Delta x' = & -K\lambda\left(x - \frac{1}{2}xy'^2 - yy'x' - \frac{1}{2}xx'^2 + \right. \\ & \left. \frac{5}{16}xy'^4 + \frac{5}{4}yy'^3x' + \frac{5}{8}xy'^2x'^2 - \frac{1}{4}yy'x'^3 - \frac{3}{16}xx'^4\right) , \end{aligned} \quad (61)$$

$$\begin{aligned} \Delta y' = & K\lambda\left(y - \frac{1}{2}yx'^2 - xx'y' - \frac{1}{2}yy'^2 + \right. \\ & \left. \frac{5}{16}yx'^4 + \frac{5}{4}xx'^3y' + \frac{5}{8}yx'^2y'^2 - \frac{1}{4}xx'y'^3 - \frac{3}{16}yy'^4\right) . \end{aligned} \quad (62)$$

The particle coordinates x , x' , y and y' on the right again have values describing a particle just before it enters the thin lens.

The factors in x' and y' which appear in these equations do not originate from the kinematic terms. They are introduced by the integration across the derivatives of the delta function.

Note that higher-order terms in the field expansion cause higher-order terms in the equations of motion. Therefore, raising the truncation in the expansion of the field beyond $m = 2$ will not change the terms in the equations of motion up to the fifth order but will only generate terms higher than fifth order.

We like to point out that this model is also symplectic, whatever the order of truncation is in the expression of the magnetic field.

10 Comparison and Conclusion

Let us introduce the general model, where the equations of motion retain the exact representation of the kinematic terms, and include the Maxwellian field series expansion to a sufficiently high order. In particular we retain only the field expansion up to and including the $m = 2$ terms of Eqs. (22) to (24).

Integration through a thin lens then leads to the following changes of the particle velocity:

$$\begin{aligned} \Delta x' = & -K\lambda\sqrt{1+x'^2+y'^2} \left(x + \frac{1}{2}xx'^2 - 3yy'x' - \frac{3}{2}xy'^2 + \right. \\ & \left. \frac{13}{4}yy'^3x' + \frac{25}{8}xy'^2x'^2 + \frac{3}{4}yy'x'^3 + \frac{13}{16}xy'^4 - \frac{3}{16}xx'^4 \right), \quad (63) \end{aligned}$$

$$\begin{aligned} \Delta y' = & K\lambda\sqrt{1+x'^2+y'^2} \left(y + \frac{1}{2}yy'^2 - 3xx'y' - \frac{3}{2}yx'^2 + \right. \\ & \left. \frac{13}{4}xx'^3y' + \frac{25}{8}yx'^2y'^2 + \frac{3}{4}xx'y'^3 + \frac{13}{16}yx'^4 - \frac{3}{16}yy'^4 \right). \quad (64) \end{aligned}$$

One can see that the square-root term is not expanded but retained exactly to preserve the nature of the kinematic terms. This model also preserves the symplectic properties. The terms within parentheses are proper to the field series representation and truncation.

We have thus developed four models given by the sets of equations (38)–(39), (48)–(49), (61)–(62) and (63)–(64). Each of these models represents different approximations, and they can be easily tested against each other to determine the importance of either neglecting or including the kinematic as well as the field expansion terms for the stability of particle motion over long periods of time.

It is easy to estimate the correction to the usual linear approximation for each of the models. Retaining only the lowest-order part without cross terms, we have

$$\Delta x'_1 = -K\lambda x \left(1 + \frac{3}{2}x'^2 \right), \quad (65)$$

$$\Delta x'_2 = -K\lambda x \left(1 - \frac{1}{2}x'^2 \right), \quad (66)$$

$$\Delta x'_3 = -K\lambda x (1 + x'^2), \quad (67)$$

corresponding respectively to the kinematic, the Maxwellian and the general model. These equations are not to be used in real simulation but only to estimate the magnitude of the correction introduced, as they are not symplectic.

According to the linear model

$$\Delta x' = -K\lambda x. \quad (68)$$

Using parameters typical of the arc FODO cell in RHIC⁶ each model gives a correction of magnitude

$$\frac{\Delta x'_i - \Delta x'}{\Delta x'} \approx 10^{-8} - 10^{-9} . \quad (69)$$

These correction factors are not important in determining the motion of only a few revolutions and for ring design. They may be very important and cannot be ignored in determining the stability of motion of a particle for a long period of time.

Appendix

We are interested in the integration of the following equation:

$$x'' = F[x(s), x'(s)]\delta(s) \quad (A.1)$$

where F is an analytical function in a sufficiently large interval around $s = 0$.

Let us replace the delta function with a step function $H(s, \epsilon)$, which is zero everywhere except in the interval from $s = -\epsilon$ to $s = +\epsilon$, where it is equal to one,

$$x'' = F(x, x') \frac{H(s, \epsilon)}{2\epsilon} . \quad (A.2)$$

The exact solution of the original equation is obtained by taking the limit $\epsilon \rightarrow 0$.

In particular

$$\Delta x' = \lim_{\epsilon \rightarrow 0} \frac{1}{2\epsilon} \int_{-\epsilon}^{\epsilon} F(s) ds . \quad (A.3)$$

This integral can be estimated by subdividing the 2ϵ interval into N substeps of length $h = 2\epsilon/N$, so that

$$\Delta x' = \lim_{\epsilon \rightarrow 0} \left\{ \frac{1}{2\epsilon} \lim_{N \rightarrow \infty} h \left[\sum_{n=1}^N F(-\epsilon + nh) \right] \right\} . \quad (A.4)$$

Using the analytical characteristics of the $F(s)$ in the interval of integration, we can expand:

$$F(-\epsilon + nh) = F(-\epsilon) + F'(-\epsilon)nh + \frac{F''(-\epsilon)}{2}(nh)^2 + \dots . \quad (A.5)$$

Substituting this expansion into Eq. (A4) yields

$$\Delta x' = \lim_{\epsilon \rightarrow 0} \lim_{N \rightarrow \infty} \frac{1}{N} [N F(-\epsilon) + h F'(-\epsilon) \sum_{n=1}^N n + \dots + \frac{F^{(m)}(-\epsilon)}{m!} h^m \sum_{n=1}^N n^m + \dots]. \quad (\text{A.6})$$

As $\sum_{n=1}^N n^m$ is a polynomial in N of order $m+1$, in the limit $N \rightarrow \infty$ we derive

$$\Delta x' = \lim_{\epsilon \rightarrow 0} [F(-\epsilon) + 2\epsilon F'(-\epsilon) + \dots + \frac{(2\epsilon)^m}{m+1} \frac{F^{(m)}(-\epsilon)}{m!} + \dots]. \quad (\text{A.7})$$

In the limit $\epsilon \rightarrow 0$ finally

$$\Delta x' = F(x, x') \quad (\text{A.8})$$

where F is evaluated just at the entrance of the integration interval.

The generalization to a system of two second-order differential equations is straightforward.

References

1. F.C. Iselin, Computer Programs for Accelerators, CAS Proceedings, CERN, Geneva (1987).
2. H. Wiedemann, PEP Note 220 (1976).
3. R. Talman, L. Schachinger, SSC-52 (1985).
4. K.L. Brown, SLAC-75, Stanford (1972).
5. E.D. Courant, H.S. Snyder, Theory of the Alternating Gradient Synchrotron, *Ann. Phys.* **3**, 1-48 (1958).
6. Conceptual Design of RHIC, BNL 52195 (1989).
7. E.D. Courant, A.A. Garren, A.S. Kenney, M.J. Syphens, A Users Guide to SYNCH, BNL (1985).
8. Handbook of Mathematical Functions, Eds. M. Abramowitz and A. Stegun, NBS, Washington (1970).

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**DATE
FILMED**

11 / 5 / 93

END

[REDACTED]

1

1

[REDACTED]