PROGRESS REPORT

1992

Rangeland - Plant Responses to Elevated CO₂

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Progress Report 1992

Rangeland - Plant Response to Elevated CO₂

Submitted By:

Dr. Clenton E. Owensby
Dr. Patrick I. Coyne
Dr. Jay M. Ham
Dr. William Parton
Dr. Charles Rice
Ms. Lisa M. Auen
Mr. Neal Adam

Department of Agronomy
Kansas State University
Manhattan, KS 66502-5501
913-532-7232

Program of the:

U.S. Department of Energy
Atmospheric and Climate Research Division
Office of Health and Environmental Research
Office of Energy Research
Washington, D.C. 20545

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

TABLE OF CONTENTS

I. Introduction	3
II. Biomass Production in a Tallgrass Prairie Ecosystem Exposed to Elevated CO ₂	5
III. Nitrogen and Phosphorus Dynamics of a Tallgrass Prairie Ecosystem Exposed to Elevated Carbon Dioxide	35
IV. Forage Quality for Ruminants on a Tallgrass Prairie Ecosystem Exposed to Elevated Carbon Dioxide	61
V. Effects of Elevated Atmospheric CO ₂ and Soil Nitrogen on Canopy Leaf Area Distribution in a Tallgrass Prairie	77
VI. Research Initiated in 1992	96
A. Above- and Belowground Biomass Sampling	98
B. Soil Microbial Biomass and Respiration under Elevated CO ₂	99
C. Physiology of <i>Andropogon gerardii</i> under Elevated CO ₂	100
D. Simulation of C and N Dynamics under Elevated CO ₂ in Tallgrass Prairie	105
E. Fumigation Chamber Effects	110
F. ¹³ C/ ¹² C ratios in Above-and Belowground Biomass under Elevated CO ₂	120
VII. Literature Cited	121

Preprints removed

Introduction

This project report incorporates manuscripts prepared in 1992 and submitted for publication on research completed in 1989-1991 and a description of the work begun in 1992. Several broad conclusions can be drawn from the work that was accomplished during the first 3-year phase of the study.

- ♦ In prairie ecosystems dominated by C_4 grasses, it is likely that elevated atmospheric CO_2 will increase ecosystem level productivity, with a greater increase in belowground productivity. The increased productivity will primarily result from increased water use efficiency due to the anti-transpirant action of CO_2 .
- ♦ Fumigation chambers are directly confounded with elevated CO_2 effects, in that both reduce evapotranspiration. The reduced evapotranspiration of the fumigation chamber is primarily through reduced wind speeds and reduced radiation. In very dry years, fumigation chamber effects are negligible, but in years with normal precipitation, chamber effects and elevated CO_2 effects are essentially equal with respect to reduced evapotranspiration effects.
- ♦ Increased production under elevated CO_2 results in reduced nitrogen concentration in the herbage and increased fiber concentrations. Consequently, digestibility of the herbage is reduced, and microbial degradation of surface litter and soil organic matter is slowed. On the negative side, ruminant productivity will likely be reduced substantially, but increased carbon storage in the soil may buffer against future rise in atmospheric CO_2 .
- ♦ Tallgrass prairie will not likely change greatly in botanical composition, since the C_4 dominants responded to elevated CO_2 more than the C_3 subdominants.

Based on the above conclusions, the research initiated in 1992 is aimed at determining the partitioning of carbon in the various pools in the carbon cycle. The aim is to estimate the net carbon flux of the tallgrass prairie ecosystem and to apply those data to parameterize models which will simulate long-term impacts of elevated CO₂ on ecosystem processes.. Further, we have initiated research to determine the effects of elevated CO₂ on the physiology of the principal C₄ dominants. Since practicality requires the use of fumigation chambers in elevated CO₂ research, we have begun work to characterize the effect of fumigation chambers of carbon, heat, and water fluxes in tallgrass prairie.

Research Initiated in 1992

Carbon input into the ecosystem is affected primarily by environment and photosynthetic pathway (C_3 vs C_4). Differential input of carbon between CO_2 -enriched and ambient CO_2 regimes is largely a function of photosynthetic pathway and water relations, although nutrient limitations may, in some ecosystems, be important as well. Our initial studies indicate that increased water-use-efficiency with elevated CO_2 may have a greater impact on carbon acquisition by plants than differences in photosynthetic pathway. Nitrogen addition to the system did not cause us to alter those conclusions. We plan to better characterize the physical environment, with particular emphasis on those factors affecting water use and gas exchange. We plan to dedicate four chambers and two unchambered plots to that task. In order to estimate the net carbon input into the ecosystem, we will do whole chamber gas exchange on the large chambers. Use of these large chambers will reduce the variability associated with the heterogeneous nature of natural grassland communities. By harvest, we will be able to determine the amount of carbon remaining in aboveground biomass. Using estimates of aboveground respiration and aboveground biomass, we will generate an estimate of the amount of carbon translocated below ground. To further validate the differential response of the plant community to CO_2 concentration, we will measure single leaf photosynthetic rates and leaf water potentials of the dominant C_3 and C_4 plants in the natural ecosystem. We will further examine responses of those species to elevated CO_2 under variable water and light regimes in the laboratory. We will examine short- and long-term acclimation potential in leaf-level gas exchange in *A. gerardii* to elevated CO_2 . Short-term responses will be measured by exposing tillers from ambient atmospheres to CO_2 concentrations up to 1000 ppm and measuring photosynthesis (A) and stomatal conductance (g). Long-term acclimation will be assessed using tillers in the field that have been exposed to elevated CO_2 over three growing seasons. Additionally, the relationship between A and intercellular CO_2 concentration (C_i) will be

determined and non-stomatal limitations estimated. Photosynthetic capacity ($A_{\max}$) of leaves, defined as the maximum photosynthetic rate that can be achieved at saturating light levels with no stomatal limitation, will be compared between plants from enriched and ambient chambers. We will also measure standard photosystem II fluorescence kinetics which will enable us to evaluate, in greater detail, other non-stomatal limitations to CO_2 uptake in *A. gerardii* grown in ambient and enriched chambers. We will estimate relative differences in root biomass by using root ingrowth bags in each treatment. Preliminary indications are that root growth under elevated CO_2 is almost twice that with an ambient CO_2 environment.

With the above measures we anticipate that we will be able to characterize and quantify the carbon input into the ecosystem under ambient and elevated CO_2 .

Loss of carbon to the atmosphere from the ecosystem will likely be affected by changes in the C:N ratio and fiber content of surface litter. Our previous research has indicated that plant material collected from plots with elevated CO_2 had a higher C:N ratio and higher fiber than plots with ambient CO_2 . We will continue to measure the decomposition rate of litter harvested from all treatments. Those data will enable us to characterize the residence time for carbon in surface litter. We will determine the amount and chemical composition of surface litter throughout the course of the study

Loss of carbon through root respiration is affected by the belowground environment, with temperature, water relations, nutrient status, and amount of carbon allocated to the belowground biomass affecting carbon losses. With increased root biomass under elevated CO_2 , total root respiration will undoubtedly be higher. We will estimate belowground CO_2 losses by measuring soil respiration and microbial biomass. Measuring CO_2 evolution from the soil surface aids in describing energy flow patterns i.e. nutrient mineralization and decomposition rates. We will also characterize the chemical composition of root material and use those data to estimate decomposition rates. The effect of elevated CO_2 on long-term ecosystem source-sink relationships will

be estimated using the Century Model which has been used extensively in modeling fire effects in the tallgrass prairie ecosystem. Data collected previously and during the proposed study will be used to parameterize the Century Model to deal with elevated CO₂. The Century Model is particularly adept at estimating changes in soil organic matter levels over a wide range of environments. Validation by data collected here will be extremely useful in extending its ability to look at potential responses of other ecosystems to elevated CO₂.

Specific Research Plan

Biomass Accumulation

Aboveground Biomass. Due to the relatively small area available for sampling, we will determine peak biomass in August and collect smaller samples for growth dynamics analysis to determine differences in growth patterns among treatments. Aboveground peak biomass will be estimated by clipping 4, 0.5 m² subplots in each experimental plot in early August. Biomass will be separated into leaf and stem categories for *A. gerardii*, *P. pratensis*, forbs, warm-season grasses, and cool-season grasses. Leaf blade area and sheath area of each group will be determined and the samples dried to moisture-free as leaf blade, leaf sheath, stem tissue and weighed. Nitrogen, phosphorus, and fiber analyses will be run on the dried samples. Twenty tillers each for *A. gerardii* and *P. pratensis* will be harvested biweekly for growth dynamics analysis. The growth dynamics samples will be treated similarly to the biomass samples. The aboveground biomass sampling will be used in the estimation of respiration and as a part of whole-chamber water flux measurements.

Belowground Biomass.

Roots. Relative differences in belowground biomass among treatments will be assessed by buried nylon bags to a 15 cm depth in each plot. Eight 5-cm dia bags, filled with fine sand, will be buried in each plot in late March each season and

recovered in early November. Roots which have grown into the bags will be extracted, dried to moisture-free, and ground through a 1 mil screen. A C:N ratio and lignin content will be determined.

Microbial biomass. Microbial biomass carbon and nitrogen will be determined by the fumigation-incubation technique as proposed by Jenkinson and Powlson (1976). Following a preincubation period of 5 days at 25 °C, soil samples from each plot to a depth of 15 cm will be subjected to chloroform fumigation and CO₂ evolution between unfumigated and fumigated samples during a 10-day incubation at 25 °C. Nitrogen will be determined by KCl extraction and colorimetric determination of NH₄-N and NO₃-N.

In 1991, we evaluated the effects of doubled atmospheric concentrations of CO₂ on soil microbial biomass C and N, and soil respiration in tallgrass prairie under field conditions. Microbial biomass C (MBC) and N (MBN) tended to increase under high levels of CO₂, but in most cases these differences were not significant. Addition of N increased the differences of MBC and MBN between high and normal levels of CO₂. High levels of CO₂ increased soil respiration in N fertilized prairie.

Carbon Fixation

Responses in CO₂ flux in tallgrass prairie to CO₂ enrichment will be examined at two complementary scales: through whole canopy/ecosystem measurements of CO₂ exchange and through more intensive leaf-level photosynthetic studies in *Andropogon gerardii*, the dominant C₄ grass in the region. In addition to providing a more mechanistic understanding of grassland responses to elevated CO₂, these basic ecosystem and physiological measurements will be incorporated into modeling efforts that can extend our predictive capabilities beyond the temporal scope of this study. A. K. Knapp will take primary responsibility for these CO₂ exchange studies.

Whole Chamber Measurements.

Net Carbon Exchange. CO₂ enrichment effects on ecosystem carbon exchange and canopy carbon flux (photosynthesis minus rhizosphere respiration which will be estimated separately) will be determined by using experimental chambers as open "cuvettes". Net canopy carbon flux will be determined by differential CO₂ levels between inlet air and air leaving the chamber, monitored continuously through sequential sampling of the chambers during the 24-hr day. Photosynthetically active radiation, air temperature and vapor pressure deficit will be measured concurrently so that environmental relationships with canopy photosynthesis and ecosystem carbon exchange (aboveground and belowground carbon flux) can be established. The chamber and canopy boundary layer resistances measured as a part of whole-chamber water flux measurements will be incorporated into calculations after adjustments are made for differences in diffusivity of CO₂ and H₂O under chamber conditions. Night ecosystem respiration in enriched and ambient chambers will be estimated using data collected by the continuous differential monitoring of CO₂ concentrations in the chambered plots. Leaf/stem level estimates of respired CO₂ (see below) coupled with whole chamber biomass estimates can be used to calculate the contribution of aboveground vegetation to ecosystem respiration. By subtraction, rhizosphere respiration can be calculated under chambered conditions. Again by monitoring temperature (soil and air) and soil moisture, environmental relationships with respiration can be established.

Leaf level gas exchange in A. gerardii. Results from the initial phase of this research program indicate that enriched CO₂ increased aboveground productivity in *A. gerardii*. Direct effects of CO₂ on photosynthetic capacity have usually been small or not detectable in C₄ species (Long and Woolhouse 1978; Kramer 1981; Carlson and Bazzaz 1982; Rogers et al. 1983; Riechers 1983; Ziska et al. 1990; Mooney et al. 1991). A small amount of preliminary data collected in 1989 also failed to show any increase in photosynthesis in *A. gerardii* at the leaf level. Nonetheless, one of our goals

is a more comprehensive investigation of photosynthetic responses in *A. gerardii* to enriched CO₂. Earlier analyses by Coyne indicated that the relative growth rates of *A. gerardii* remained constant longer into the season in enriched chambers than in ambient. Calculated Unit Leaf Rate (an estimate of photosynthetic activity of leaves) also remained higher in tillers in enriched chambers. These results suggest that important interactive effects involving CO₂ uptake and water status, light responses and potential stomatal/biochemical limitations need to be quantified.

Field studies. In the field, net photosynthesis (A) will be measured in upper, middle and lower (20 cm each) portions of the canopy at saturating light levels. Diurnal measurements of A (and stomatal conductance, g) will be made in enriched and ambient chambers as well as in adjacent, unchambered populations. Measurements will be made at ca. 2 hr intervals every two weeks during the growing season. Leaf level measurements of A and g (4 tillers per chamber) will be made with a LI-COR 6200 (field portable) gas exchange system. Concurrently, plant xylem pressure potential will be measured with a Scholander-type pressure chamber on 5-7 tillers per chamber. Routine measures of plant tissue N concentration will allow us to calculate both nitrogen-use-efficiency and water-use-efficiency for plants in enriched and ambient chambers. Dark respiration rates in leaves and stems stratified according to canopy position will also be made at night, and during the day with shrouded chambers, using the LICOR gas exchange system.

Laboratory Experiments in Gas Exchange Physiology. Several aspects of gas exchange responses to enriched CO₂ will be examined in *A. gerardii*. Tillers will be grown in enriched and ambient chambers and transported to a nearby laboratory. We have found that removal (and potting) of 30 cm deep soil cores with intact *A. gerardii* tillers allows us to measure gas exchange responses in the lab identical to those measured in undisturbed field plants (Knapp unpubl. data). Major objectives of carbon fixation research are listed below (*italics*) along with our proposed methods.

We propose to *examine short- and long-term acclimation potential in leaf-level gas exchange in *A. gerardii* to elevated CO₂*. Short-term responses will be measured by exposing tillers from ambient atmospheres to CO₂ concentrations up to 1000 ppm and measuring photosynthesis (A) and stomatal conductance (g). Additionally, the relationship between A and intercellular CO₂ concentration (C_i) will be determined and non-stomatal limitations estimated (Jones 1985). Long-term acclimation will be assessed by comparing similar measurements made on tillers grown in enriched chambers to those grown under ambient conditions. A null-balance gas exchange system (PAC-SYS 9900, Data Design Group) with a cuvette capable of controlling temperature, humidity and CO₂ partial pressure at specified levels will be used. Thus, interactions between response to CO₂, temperature and vapor pressure deficit can be evaluated.

We propose to *determine the photosynthetic capacity (A_{max}) of leaves of *A. gerardii* grown under ambient and enriched CO₂ conditions*. Leaf level gas exchange measurements of CO₂ uptake are a product of both stomatal and non-stomatal limitations to photosynthesis. Thus, differences in A measured in the field or lab between enriched and ambient plants may be partially attributable to differences in g between treatments. Stomatal conductance may be influenced by water status, direct effects of CO₂ on stomatal aperture, or feedback from mesophyll demands for CO₂ through alterations in C_i (Jarvis and Mansfield 1981; Schulze and Hall 1982; Morison 1985). Photosynthetic capacity (A_{max}) of leaves, defined as the maximum photosynthetic rate that can be achieved at saturating light levels with no stomatal limitation, will be compared between plants from enriched and ambient chambers. A_{max} will be determined with an O₂-probe photosynthetic system (Hansatech Inc.) that measures O₂ evolution from leaf discs at 3-4% CO₂ concentrations. The high CO₂ concentration used in this system overcomes diffusional limitations due to stomatal aperture and, thus, provides an estimate of maximum photosynthetic capacity. Several studies have shown that starch may accumulate in leaves at high CO₂ (Kramer 1981; Azcon-Bieto 1983; Czarnowski and Starzecki 1989; but see Riechers 1983) and end-product

(non-stomatal) inhibition of A may result. We will document diurnal patterns (dawn-to-dusk) in $A_{\max}$ at 2-hr intervals to assess such limitations in enriched and ambient tillers. This photosynthetic system also is capable of measuring standard photosystem II fluorescence kinetics (Delieu and Walker 1983) which will enable us to evaluate, in greater detail, other non-stomatal limitations to CO_2 uptake in *A. gerardii* grown in ambient and enriched chambers.

We propose to *evaluate interactive effects of light level and water status on gas exchange in A. gerardii from enriched and ambient chambers*. Greater biomass and canopy development in enriched chambers may potentially lead to greater light limitations to photosynthesis and greater respiration costs. However, several studies have reported that quantum yields and A at low light levels are increased by elevated CO_2 (Kramer 1981; Sionit et al. 1982; Starzecki and Czarnowski 1989). Moreover, light compensation points may be decreased under high CO_2 (Kramer 1981). We will measure these parameters with the null-balance gas exchange system on whole leaves and on leaf discs with the O_2 -probe system (excluding stomatal effects). Neutral density filters will be used to vary incident sunlight from compensation points to full sunlight. We anticipate that the increase in production in *A. gerardii* under elevated CO_2 may be due in part, not to higher A at high light levels, but to higher A at the low light levels found within the plant canopy.

Previous work from this project and research by others (see review by Dahlman et al. 1985) suggest that the water status of plants (and soil moisture) at elevated CO_2 is enhanced because stomata close partially in response to high CO_2 (Morison 1985). Thus, improved plant water relations may also explain, in part, the increase in production of *A. gerardii* in enriched chambers. Field measurements described above will document patterns in plant water relations during the growing season, but a laboratory assessment of the relationship between leaf water potential and gas exchange in ambient and enriched plants is necessary to precisely identify interactions. The effect that plant water stress has on CO_2 uptake will be evaluated in several ways. Transplanted (potted) tillers will be kept in their respective chambers and allowed to

dry-down. As plant water potential decreases, leaf level gas exchange will be measured with the null-balance system and $A_{\max}$ determined with the O_2 -probe (fluorescence kinetics will also be assessed). Because water availability fluctuates frequently in extant tallgrass prairie ecosystems (Knapp 1984), and it may vary more in a "greenhouse world" (Karl et al. 1991), we will also document differences in the ability of *A. gerardii* tillers (enriched and ambient) to recover following alleviation of water stress. In these studies, droughted plants will be re-watered and the time course of recovery in A , g and $A_{\max}$ will be measured at several hour and daily intervals. We anticipate that plants grown under enriched CO_2 may be able to recover more rapidly following drought stress than ambient plants. Finally, light response curves (as described above) will be developed for enriched and ambient tillers under different levels of water stress to examine the 3-way interaction among CO_2 , light and water.

Gas Exchange Responses to Variable Sunlight. Plants in open environments experience substantial variability in sunlight due to broken cloud cover and within canopy shading (Knapp and Smith 1987). Increases in grassland canopy development under conditions of enriched CO_2 and predicted increases in cloud cover in a greenhouse world suggest that these grasses will experience even greater sunlight variability in the future. Recent studies indicate that significant differences in CO_2 uptake, transpiration and water use efficiency may result from different (species specific) patterns in the kinetics of stomatal and photosynthetic responses during periods of fluctuating sunlight (see review by Knapp and Smith 1990). For example, stomatal conductance in *A. gerardii* responds rapidly to variations in sunlight under conditions of ambient CO_2 concentrations (Knapp, unpubl. data). This response pattern conserves water, but also results in reduced CO_2 uptake. Other studies have shown that plant water status may influence the response kinetics of stomata such that plants under no water stress maintain high levels of stomatal conductance during sun and shade periods (Knapp and Smith 1990). This type of response pattern maximizes CO_2 uptake, but at reduced water-use-efficiency. In enriched CO_2 chambers, where the water relations of *A. gerardii* are improved, we can predict that stomatal response

Current modeling capabilities allow for the simulation of management practices such as fire, grazing, and cultivation on the ecosystem dynamics of natural and managed grasslands. Our recent modeling efforts in the tallgrass prairie have focused on the impact of fire on soil organic matter dynamics and plant productivity, especially related to allocation patterns of C and N. We have found that the tallgrass system responds to fire by improving growing conditions for dominant tallgrass species enhancing biomass production and nitrogen use efficiency. The belowground system was significantly altered with fire so that greater root production and lower nitrogen root concentration resulted in a nitrogen-conserving environment. We found that appropriate modeling of the belowground system was fundamental to understanding the impact of changing management and environmental conditions on nutrient cycling and production in the tallgrass ecosystem (Ojima *et al.* 1987, 1990).

A comparison of observed and simulated annual aboveground plant production from 1930 to 1968 for a tallgrass prairie site was made and results indicate good correlation with interannual variability. The results showed that the model correctly predicted the short- and long-term effects of annual burning on ecosystem properties. Our results suggest that most of the influential direct effect of fire was the increase in nitrogen use efficiency (NUE) produced by burning. This response was the major factor which influenced the long-term sustained productivity of annually burned tallgrass prairie. Sensitivity of ecosystem processes and fluxes of C and N to changes in climate and CO₂ levels have also been evaluated with CENTURY. Results from simulating the effect of a CO₂ doubling scenario suggest that plant production in the Kansas tallgrass ecosystem may decrease because of changes in nutrient availability due to changes in decomposition rates and due to changes in precipitation (Schimel *et al.* 1990, Kittel *et al.*, in prep.). The effect of elevated CO₂ levels on water-use-efficiency (WUE) and NUE were evaluated with CENTURY (Kittel *et al.* in prep), and results indicate that in the Flint Hills tallgrass ecosystem plant production is expected to increase. However, despite this apparent increase in plant production, soil C declines due to increased decomposition. The direct effect of CO₂ enrichment, as represented by changes in WUE and NUE, indicate that these impacts tend to mollify the decline in soil C. These

simulations support the contention that predictions of climate and CO₂ effects on production and carbon storage must consider the interactions with decomposition and nutrient cycling, as well as the ecophysiological effects. These studies highlight the value of ecosystem models to investigate the effects of elevated CO₂ and climate change on total system C balance.

The plant production submodel simulates the dynamics of C, N, and P in the live and dead aboveground plant material and live and dead roots. Maximum plant production is controlled by soil temperature and available water. Production is further modified by the current amount of aboveground plant material (e.g., mulch effect) and reduced as a function of available soil nutrients. The allocation of C to shoots and roots change as a function of the climate, grazing rate, and fire.

Both fire and grazing effects on plant production are included (Ojima *et al.* 1990, Holland *et al.*, submitted). The major impact of fire is to increase the root-to-shoot ratio, lower the N content of the live shoots and roots, remove vegetation, and return nutrients during the years when fire occurs. Grazing removes vegetation, returns nutrients to the soil, alters root-to-shoot ratio, and increases the N content of live shoots and roots. These impacts of fire and grazing on plant production are analogous to certain effects of elevated CO₂ fertilization on nutrient use efficiency and allocation patterns of C and N in the tallgrass ecosystem.

In the SOM submodel, plant residues are decomposed by microbes, and the resulting microbial products become the substrates for SOM formation. We divide SOM into three fractions:

1. an active soil fraction consisting of live microbes and microbial products (1 to 4 y turnover time),
2. a protected fraction that is more resistant to decomposition as result of physical or chemical protection (20 to 40 y turnover time), and

patterns to natural variations in sunlight would shift towards less rapid kinetics so that CO₂ uptake is increased. However, the effect that high CO₂ concentration may have on the kinetics of stomatal responses to fluctuations in sunlight has not been investigated. As noted earlier, increased CO₂ concentrations generally reduce stomatal conductance (Morison 1985), but we need to investigate the effect that CO₂ enrichment has on non-steady-state response kinetics during periods of variable sunlight. Because periods of variable sunlight are more common than periods of constant sunlight, especially for leaves within canopies, evaluating the effect that CO₂ enrichment has on gas exchange may provide key insights into overall system responses to enriched CO₂. Methods for these studies are similar to those used to study steady-state responses except that incident light is varied frequently (every 5-10 min) while gas exchange is monitored (Knapp and Smith 1990).

CO₂ Source/Sink Relationships

Century Model

In order to determine the effect of elevated CO₂ on ecosystem dynamics in the tallgrass ecosystem a fuller understanding of the coupling between biogeochemical and ecophysiological processes which control the fate of C being sequestered or respired from the system must be attained. One way to investigate the complex interaction of these processes is the utilization of simulation models. A tallgrass prairie model (CENTURY) which incorporates these processes has been developed to simulate long-term trends in productivity and the biogeochemistry of C, N and P in the plant-soil system of natural grasslands and agroecosystems (Parton *et al.* 1987, Schimel *et al.* 1990, Burke *et al.* 1990, Ojima *et al.* 1990). Carbon sequestering in plant and soil components of the system and soil organic matter (SOM) decomposition and plant respiration are well represented in this model and provide a starting point to further develop our understanding of how elevated CO₂ concentrations will affect the grassland ecosystem.

3. a fraction that is physically protected or chemically resistant and has a long turnover time (800 to 1200 y).

The flow diagrams for C, N and P submodels of CENTURY show that the organic C, N and P flows have a similar structure with the N and P following the carbon flows. The N and P submodels also include the flows of mineral N and P compounds. The C:P ratio of the SOM fractions float as a function of the labile P level (lower values with higher labile P levels). The driving variables for the model include soil texture, monthly precipitation, average monthly maximum and minimum air temperature and the plant lignin content. A detailed description of the CENTURY model is presented by Parton *et al.* (1987, 1988).

The CENTURY model has been modified to include the direct effects of burning on the tallgrass ecosystem. The changes were primarily based on an analysis of the observed fire effects at Konza prairie and the Aldous plots located in the Flint Hills near Kansas State University. The modifications include:

1. calculating biological N_2 fixation rates as a function of the ratio of mineral N to labile P (high N_2 fixation with low ratios) based on Eisele *et al.* (1989).
2. developing a plant production model where maximum plant production was a function of monthly precipitation and soil temperature and the standing dead biomass,
3. allowing the C:N ratio of the active SOM to float from 6 to 14 as a function of the mineral N level (low values for high mineral N levels),
4. increasing the nitrogen use efficiency for burned plants (higher C:N ratios for burned plant material), and
5. incorporating the combustion losses of aboveground C, N and P into the model.

We have refined CENTURY in order to understand the impact of elevated CO₂ on the tallgrass ecosystem. The model needs to be made more sensitive to direct ecophysiological response relative to WUE on plant production and CO₂ utilization as indicated from earlier observation by Knapp (1984, 1985), Schimel *et al.* (1990), and Owensby *et al.* (1991). Observations collected over the past two years indicate that significant enhancement of C₄ plant production counter to *a priori* predictions that C₄ production would be little affected. These plant responses are critical to understanding the interaction between nutrient availability and plant production to elevated CO₂. The CO₂ fertilization effect on plant production may not be expressed if water or nutrients are limited. Changes in NUE and changes in plant residue quality (e.g., N or lignin content) will also modify the extent to which plant production will respond to CO₂ fertilization. Previous modeling experiments and field observations suggest a strong interaction between processes influencing carbon allocation and nutrient cycling. Inclusion of interactions between CO₂ concentration and NUE and WUE on whole system C storage will be evaluated with CENTURY. In addition, the interaction of climate changes (*i.e.* alterations in precipitation pattern and temperature) will be assessed. The observations by Owensby *et al.* (1991) will provide the opportunity to make these refinements in our current model of plant production relative to altered CO₂ and to make an assessment of the whole ecosystem response relative to these altered levels.

After incorporation of these refinements to our current model, we will be able to provide insight into the potential for sequestering of atmospheric C within this highly productive system, especially in the belowground compartment. In addition, we will be able to provide conditional constraints on the behavior of the system relative to the interactive nature of altered moisture and temperature regime in conjunction with increased CO₂ levels.

In 1992, preliminary results indicate that the simulated increase in plant production and litter quality modified the decomposition cycle. Soil C increased, N mineralization decreased, and litter decay rates decreased in the CO₂-enriched chambers. Changes

in climate related to wet and dry years modify the soil C dynamics due changes in decomposition.

Chamber effect on water relations and net carbon assimilation

Increasing concentrations of atmospheric carbon dioxide may influence plant productivity and survivability in the future. Extensive field research will be required to quantify the potential impact of increased CO₂ gas concentrations on managed and natural ecosystems. Results from this crucial research will be used to make management and policy decisions that will impact global productivity in the 21st century. Unfortunately, existing methodology for studying plant responses to the gaseous environment is often inadequate, inaccurate, or in the worst case, misleading. The techniques used to study the effect of elevated CO₂ often disrupt the plant environment, and climatological impacts of the methodology have a greater influence on ecosystem response than the trace gas under investigation. Thus, basic research is needed to quantify the climatic impact of enrichment methodology so that the true effect of elevated CO₂ can be evaluated. Here, the investigators propose to evaluate the microclimate influence of open top CO₂ enrichment chambers on a tallgrass prairie ecosystem, and separate the 'chamber effect' from the impact of CO₂ enrichment. Novel theory, experimentation, and simulation will be combined to formulate a new approach for quantifying the chamber effect in CO₂ enrichment studies.

Background

Experimentally determining plant response to gaseous CO₂ requires that air around the plant or plant leaf be modified to contain CO₂ concentrations above ambient levels. Typically this has been accomplished by enclosing the leaf, plant, or plant community in a chamber where aerial gaseous conditions can be controlled. Leaf chambers, phytotrons, greenhouses, tracking chambers, closed chambers, and open-top field

chambers have all been used to study trace gas effects on plants. Early work was predominated by leaf chamber, green house, and phytotron studies since gas concentrations are easily controlled and monitored. Various different types of cuvette and chambers have been developed for this work (e.g. Carlson and Bazzaz, 1980; Valle *et al.*, 1985). While laboratory studies are useful for discovering basic physiological responses to trace gases, the unnatural, sterile, environment limits the utility of the results. Research must be conducted under field conditions so the plants experience temporal environmental variations, and can interact with other organisms (Mooney *et al.*, 1991). Research of this nature requires that the chamber or gas delivery system be constructed around the field site of interest. Small, closed chambers (2 m²) have been used to measure plant response to CO₂ and measure ecosystem gas exchange (Chauduri *et al.*, 1987; Drake *et al.*, 1989). Prudhomme *et al.* (1984) constructed a closed, tracking- chamber that was capable of mimicking conditions outside the chamber by using feedback from external sensors. Unfortunately, almost all closed chambers reduce radiation, increase air temperature, and have low wind speeds. Reduced air flow can create large boundary layer resistances that limit plant uptake of the pollutant (Black, 1982), or nullify physiological impacts of the trace gas (Shaer and Van Bavel, 1987; Kimball, 1986). In an attempt to overcome the radiation and aerodynamic limitations of small, closed chambers, researchers developed larger, open-top chambers that allow more natural conditions at the field surface, while giving up the ability to make mass balance calculations. Open-top chambers (OTC) have been widely used in both CO₂ and ozone studies (Mandl *et al.*, 1973; Heagle and Philbeck, 1978; Rogers *et al.*, 1983). Chambers of this type were selected for ozone research by the EPA-sponsored National Crop Loss Assessment Network (NCLAN), and are also used by the Department of Energy's CO₂ evaluation program. Open-top chambers typically consist of plastic-covered, open-ended cylinders 3 m in diameter and approximately 3 m in height. The opening in the top of the chamber is often reduced to inhibit the effect of vertical eddy movement on gas concentrations near the canopy. A electric fan is used to aspirate the chamber, exchanging the air every 20 to 30 s. Although widely used,

several studies have shown that the OTC can increase temperatures 1 to 2 C and reduce radiation by 10 to 15% (Heagle *et al.*, 1979; Olszyk *et al.*, 1980) when compared to nearby companion plots. Other researchers have shown that modified wind speed profiles within OTC alter pollutant concentrations in the canopy (Colls and Baker, 1988). Other aerodynamic studies by Unsworth *et al.* (1984a, 1984b) found that canopy boundary layer resistance to O₃ within the chamber was similar to conditions in nearby companion plots (Wesely *et al.*, 1982). This comparison, however, was limited to one location over a short time period, and insufficient data are given by either Unsworth *et al.* (1984) or Wesely *et al.* (1982) to generalize their results.

While altered microclimates within OTC have been observed, the effect of these alterations on crop growth and yield has been difficult to quantify. Typically researchers run a chamber without adding trace gases and then compare plants inside the control chamber with plants in unchambered companion plots. Research has shown that the chamber effect varies between years, and sometimes alters morphology but not yield (Heagle *et al.*, 1979; Owensby *et al.*, 1991). Heggestad *et al.* (1980) reported that the chamber effect increased growth of certain plant organs and reduced the biomass of others. Owensby *et al.* (1991) found that the chambers increased crop growth during normal rainfall years, but had little effect during low rainfall years. Inconsistent reports of the chamber effect on crop growth and yield lead Drake *et al.* (1985) to conclude that climatological impact of OTC on plant growth was negligible. The inability to quantify an observation, however, does not justify its dismissal. More research is needed to understand the interaction between the impact of the chamber and the influence of the trace gas under investigation.

The apparent inconsistent effect of the chamber in OTC studies is very similar to data reported for crops grown near wind breaks and other shelters. Early microclimate research on sheltered crops showed that differences between sheltered and unsheltered plants varied drastically from year-to-year, and that crop yield, morphology, and physiology could be differentially but inconsistently affected by the shelters (Grace, 1977). Similarity between sheltered and OTC experiments suggests that the same

biophysical processes are involved. In a review of sheltered crop experiments, Grace (1977) suggested that the year- to-year variability of a shelter on yield is due to complex relationships between plant water status, soil water, and climate. During high rainfall years, lower windspeeds (higher boundary layer resistance) near the sheltered crop reduce evapotranspiration and gives the sheltered crop a more favorable water balance and higher yield. During dry years, however, increased stomatal resistance controls water loss from the crop and the aerodynamic influence of the shelter has little effect. This hypothesis is consistent with data reported by Owensby *et al.* (1991), who found that the effect of OTCs on plant growth was more noticeable in normal rainfall years, but not evident during low rainfall periods. It is probable that variability in the chamber effect observed in OTC studies is due to year-to-year variability in climate at individual research sites, and inherent differences in climate-ecosystem interactions between sites. For example, normally high wind speeds present on a tallgrass prairie will be reduced by a chamber enrichment system (Owensby *et al.* 1991), while chamber enrichment may actually increase normal wind speeds at other sites (Unsworth *et al.*, 1984). Thus, quantifying the chamber microclimate effect in CO₂ enrichment studies requires a complete understanding of the biophysical processes involved rather than empirical approximation by comparison of chambered and unchambered plots.

Problems With CO₂ Enrichment: A Simulation Study of Open-Top Chambers.

The Department of Energy (DOE), in collaboration with the Department of Agronomy at Kansas State Univ., established a CO₂ enrichment site near Manhattan, KS to study the effect of CO₂ on rangeland plants. Traditional OTC methodology (Rogers *et al.*, 1983) was used in the study. One of the proposed investigators, Dr. J. Ham, has been collaborating with scientists on the DOE project for the past year, and has developed an energy and water balance model of the OTC system. The model solves the surface energy balance by finding the equilibrium canopy temperature that satisfies the governing energy budget equation for the plant canopy. An iterative Newton-Raphson procedure is used to find the solution temperature (Bristow, 1987). Aboveground

radiation and aerodynamic transfer are modeled using a uniform, one-dimensional approach (e.g. Van Bavel and Lascano, 1987; Ham and Heilman, 1991). Aerodynamic and surface resistances to heat and mass transfer are described with bulk parameters. Stomatal action is a function of radiation, CO_2 , temperature, and leaf water potential, and is based on data from experiments with Eastern gammagrass (Coyne and Bradford, 1985). Absorption and redistribution of water in the root zone is modeled using a matric flux potential approach similar to that described by Campbell (1985). The belowground portion of the model predicts soil water content and water potential with depth, by simulating concurrent liquid and vapor flow. Leaf water potential is also simulated by solving the liquid water flow equation, assuming root, xylem, and soil resistances. While the model needs refinement, preliminary results can provide insight into the environmental physics of chamber microclimate.

The energy and water balance simulations of chambered and unchambered plant canopies clearly show that OTC reduce water losses in the chamber, and that these savings eventually result in a more favorable leaf water potential for chambered plants. Preliminary findings indicate that decreased wind speed in the chambers is the primary factor governing this response. Data also suggest that the important antitranspirant action of CO_2 is moderated or masked inside the chambers. Improved data on plant response to light, soil water, temperature, and CO_2 , is needed to refine the simulation for predicting specific ecosystem responses to chamber enrichment. Additionally, improved microclimate theory and data for OTC methodology will be required. As demonstrated here, a good working model of the OTC system can be a valuable tool when analyzing data from CO_2 enrichment studies.

Rationale and Objectives

A review of the literature, recent data of Owensby *et al.* (1991), and the simulation study suggest that several climatological aspects of open top chamber CO_2 enrichment need qualification. The overall objective of the proposed research will be to evaluate the environmental physics of open-top CO_2 enrichment chambers, with emphasis on

separating the biophysical impact of CO₂ enrichment from the climatological influence of the chamber. Specific objectives of the study are:

- ♦ quantify aerodynamic transport of heat, water vapor, and CO₂ in enriched-chambers, ambient-chambers, and unchambered companion plots,
- ♦ separate the climatological effect of the chamber from the antitranspirant action of CO₂ by direct, simultaneous measurements of plant water use in enriched-chambers, ambient- chambers, and unchambered companion plots,
- ♦ document soil water dynamics inside and outside the chambers over the growing season to see if differences in soil water potential develop as predicted by simulation, and
- ♦ develop and refine a numerical energy and water balance model of open-top enrichment chambers.

The experimental portion of the study will be conducted in four open-top chambers specifically designated for microclimate research. This approach will prevent meteorological measurements from disrupting other aspects of the study. CO₂ concentrations in two of the microclimate chambers will be enriched to twice ambient levels while the remaining two chambers will be left at ambient concentrations.

Chamber Aerodynamics, Radiation, Temperature, Vapor Density, and CO₂.

An important first step in the analysis of the OTC system will be the comparison of basic meteorological parameters inside the chamber with those in the unchambered companion plots. While some preliminary information on chamber environment is already present (Owensby et al., 1991), a more detailed analysis will be required to meet new project objectives. Characterization of the chamber environment will be conducted during the first year of the project, and will typically consist of short intensive

measurement periods using temporary sensors and apparatus rather than the installation of a continuous measurement/data- acquisition system.

Air Speed and Turbulent Intensity: Heat balance anemometers of the Simmons design (Kanemasu and Tanner, 1968) will be used measure air velocities. Vertical and horizontal gradients in wind speed will be determined by positioning sensors at multiple locations in the chambers and in the companion plots. The magnitude, frequency and turbulent intensity (coefficient of variation) of mean wind speed will be sampled using a high resolution datalogger (CR7X, Campbell Scientific). Turbulent intensity and speed frequency will be used to analyze aerodynamic conditions in the chamber since the lack of a defined equilibrium sublayer (logarithmic wind profile) precludes the use of traditional flux profile theory (K-Theory) inside the chamber (Corrsin, 1974). The research team has already constructed and calibrated over 30 anemometers that can be used for this purpose.

Radiation Regime: The radiation environment in the chambers will be evaluated by measuring all forms of radiation that influence either photosynthesis or the surface energy balance. The measurements and corresponding instruments used for the analysis will include: global irradiance (model 8-48, Eppley Laboratories), photon flux density (model LI 190, Li-Cor, Inc.), longwave radiation (model PIR, Eppley Laboratories), and net radiation (model Q6, Radiation Energy Balance Systems). The spatial variation in radiation within the plot will be determined by mounting radiometers on actuating platforms that traverse the surface of the plot (e.g. Ham et al., 1991). The photometric properties of the plastic used to cover the chambers will also be measured with a spectral radiometer.

Temperature and Heat Flux: Air and soil temperatures will also be measured at multiple locations within the chamber and in the companion plots using thermocouples. Soil heat flux will also be measured using the combination method (Kimball and Jackson, 1979).

Water Vapor and CO₂ Concentrations: Vertical and horizontal gradients in CO₂ and H₂O will be measured within the chamber and the companion plots. A dedicated gas sampling system connected to a dual H₂O/CO₂ gas analyzer (model LI 6262, Li-Cor Inc.) will be used to measure gas concentrations at various locations in the chamber. Quantifying the spatial and temporal variation of CO₂ in the chamber will be crucial in determining the feasibility of estimating net carbon exchange in the chamber by measuring inflow and outflow gas concentrations.

***Quantifying Ecosystem Evapotranspiration, Carbon Exchange, and
Resistance to Mass Transport by Chamber Scale
Porometry/Gas-Analysis***

The effect of elevated CO₂ on ecosystem water and carbon exchange will be quantified by measuring whole-chamber evapotranspiration and net carbon flux. This will be accomplished by closing the chambers for brief periods, and measuring the time rate change in water vapor and CO₂ concentration inside the system. The four microclimate chambers will first be modified so that a temporary barrier to air flow can be installed approximately 1.5 m above the canopy. The fans and associated duct work will also be modified such that air can be recirculated inside the closed area without contamination from outside air. A portable sensor platform will be positioned near the center of the plot before the chamber is closed. The sensor platform will consist of a gas collection manifold, infrared transducer, air temperature sensor, and pyranometer. A portable instrument/control system, located outside the chamber, will be used to measure dynamic changes in CO₂, water vapor, and temperature after the chamber is closed. The instrument/control system will consist of a dual CO₂/H₂O analyzer (LI 6262, Li-Cor Inc.), a communication-based data acquisition system (uMega-1050, Omega Engineering), a laptop microcomputer, gas pumps, and power supply. The computer-based data acquisition system will allow for real-time processing of canopy gas exchange and transpiration in the field. Evapotranspiration (ET) and carbon exchange rate (CER) will be estimated from the time rate change in CO₂ and H₂O after the chamber is closed. The measurement system described here will be

capable of estimating the time rate change in both CO_2 and water vapor every 1 s using on-board numerical algorithms that are part of the data acquisition system (Labtech Notebook, Omega Engineering). Thus, by making almost instantaneous measurements of CER and ET, quasi steady-state conditions can be identified that are representative of chamber gas exchange.

During the draw down cycle inside the closed chamber, the system can also be operated as a chamber-scale, ventilated diffusion porometer. Given the time rate change in water vapor, and the surface temperature of the plant canopy (from the IRT) the canopy resistance to vapor transport (r_v) can be estimated. It will be possible to characterize the aerodynamic resistance of the system by first wetting the surface of the canopy with a sprinkler, then closing the chamber and measuring the time rate change in water vapor. Since stomatal resistance will not be influencing transport during this period, the resistance term will represent aerodynamic resistance only. Since air flow and canopy size are very similar in all chambers it can be assumed that the aerodynamic measured in one chamber is representative of all chambers. Once the aerodynamic chamber resistance is known, it can be subtracted from the total resistance computed during a regular "dry-canopy" measuring cycle to determine canopy surface resistance to vapor transport (canopy surface resistance = total resistance (r_v) - chamber aerodynamic resistance). It will also be possible to back calculate ET using the resistance measurements, as typically performed in porometry. Water use efficiency can be computed using the acquired data. If CO_2 is indeed affecting the plant stomates, large differences in surface resistance and ET should be evident between the enriched and ambient plots. Canopy surface resistance, ET, and CER will be measured using the closed chamber method near midday in the four microclimate chambers during the 1993 and 1994 growing season. Canopy resistance to vapor transport, and its response to changes in leaf area, and CO_2 , has never been quantified in a CO_2 enrichment study, although this information is crucial when attempting to develop a comprehensive model of ecosystem carbon exchange.

Model Development

The energy and water balance model of the open top chambers and prairie ecosystem described earlier, will continue to be modified and refined. Base line data on rangeland plant response to environmental stimuli will be used to make the model more representative of the ecosystem under investigation. This data can also be used to develop photosynthesis module for the model. The microclimate research will provide important inputs on canopy aerodynamic and radiative properties that are needed to simulate mass transport in the aerial environment. Hopefully the model can be used by all the proposed researchers to develop more accurate measurement techniques, and to develop an test new theory on ecosystem carbon balance. Programming modules developed for the rangeland ecosystem could easily be transported to other large scale models for further exploration of global carbon relations.

Plant Growth, Photosynthesis, and Ecosystem Dynamics Models

The ecosystem-level experiment proposed here will provide information needed to parameterize the ecosystem level models being developed at Duke University under the direction of Jim Reynolds for critical assessment of future impacts of elevated carbon dioxide on tallgrass prairie ecosystems. While we do not plan under the scope of this project to utilize these models directly, we plan an active interaction with the SDSU group in generating model output to predict future ecosystem-level changes in tallgrass prairie. Further, the photosynthesis work we propose will provide input into the plant growth model which can provide predictive insight into carbon fixation, carbon allocation, and carbon losses from the ecosystem. Since the modelling effort at Duke centers around compartmented models (comprised of modules), we anticipate that the modelling efforts described earlier in this proposal, i.e. Century and the energy and water balance models, will be integrated into the models currently being developed for tallgrass prairie at Duke. We anticipate that as model needs arise, we will initiate experiments, when possible, in cooperation with the Duke modeling group to obtain the necessary data.

$^{13}\text{C}/^{12}\text{C}$ Ratio of Above- and Belowground Biomass

Differential $^{13}\text{C}/^{12}\text{C}$ ratios between the gas fumigating the CO_2 -enriched chambers and the current atmospheric CO_2 may be used to estimate carbon input into the soil complex. We are currently determining the $^{13}\text{C}/^{12}\text{C}$ ratios of above- and belowground biomass using a ratioing mass spectrometer in cooperation with Dr. Steve Leavitt at the University of Arizona.

For carbon isotope analysis of soils, HCl was first used to remove CaCO_3 , plant fragments were removed by initially floating each soil sample in a concentrated NaCl solution, and then remaining fragments were picked out under 20X magnification. Using isotope mass balance calculations, the $^{13}\text{C}/^{12}\text{C}$ ratio differences in organic carbon between enriched and ambient plots can be ascertained.

Literature Cited

- Azcon-Bieto, J. 1983. Inhibition of photosynthesis by carbohydrates in wheat leaves. *Plant Physiology* 73: 681-686.
- Black, V.J. 1982. Effects of sulfur dioxide on physiological processes in plants. In M.H. Unsworth & N.P. Ormrod (eds.) *Effects of gaseous air pollution in agriculture and horticulture*, pp. 67-92, Butterworths, London.
- Bristow, K.L. 1987. On solving the energy balance equation for surface temperature. *Agric. For. Meteorol.* 39:49-54.
- Burke, I. C., D. S. Schimel, C. M. Yonker, W. J. Parton, L. A. Joyce, and W.K. Lauenroth. 1989. Regional modeling of grassland biogeochemistry using GIS. *Landscape Ecology* 4:45-54.
- Campbell, G.S. 1985. *Soil physics with BASIC: Transport models for soil-plant systems*. Elsevier, New York.
- Carlson, R.W., and F.A. Bazzaz. 1980. Elevated CO₂ concentrations differentially increase growth, photosynthesis, and water use efficiency of plants.. p. 608-623. In J.J. Singh and A. Deepak (eds.) *Environmental and climatic impact of coal utilization*. Academic Press, New York.
- Carlson, R.W. and F.A. Bazzaz. 1982. Photosynthetic and growth responses to fumigation with sulfur dioxide at elevated carbon dioxide for carbon 3 and carbon 4 plants. *Oecologia* 54: 50-54.
- Chaudhuri, U.N., R.B. Burnett, M.B. Kirkham, and E.T. Kanemasu. 1987. Effect of elevated levels of CO₂ on winter wheat under two moisture regimes. *Progress Rep. of Res. No. 040. Response of vegetation to carbon dioxide*. U.S. Dep.

Energy, Carbon Dioxide Res. Div., Office of Energy Res., Washington, DC. 70 p.

Colls, J.J., and C.K. Baker. 1988. The methodology of open-field fumigation. p. 361-371. In P. Mathy (ed.) Air pollution and ecosystems. Proc. of an international symposium, Grenoble, France. D. Reidel Pub. Co., Dordrecht, Holland.

Corrsin, S. 1974. Limitations of gradient transport models in random walks and turbulence. *Advances in Geophysics*. 18A:25-60.

Coyne, P.I., and J.A. Bradford. 1985. Comparison of leaf gas exchange and water use efficiency in two eastern gammagrass accessions. *Crop Sci.* 65-75.

Czarnowski, M. and W. Starzecki. 1989. The relationship between photosynthesis inhibition and starch accumulation in tomato leaves. *Bulletin of the Polish Academy of Sciences (Biological Sciences)* 37: 217-222.

Delieu, T.J. and D.A. Walker. 1983. Simultaneous measurement of oxygen evolution and chlorophyll fluorescence from leaf pieces. *Plant Physiology* 73: 534- 541.

Drake, B.G., H.H. Rogers, and L.H. Allen, Jr. 1985. Methods of exposing plants to elevated carbon dioxide. p. 11-31. In B.R. Strain and J.D. Cure (eds.) *Direct effects of increasing carbon dioxide on vegetation*. U.S. Dep. of Energy Report, DOE/ER- 0238.

Drake, B.G., P.W. Leadley, W.J. Arp, D. Massiry, and P. Curtis. 1989. An open top chamber for controlling CO₂ concentration and measuring net ecosystem gas exchange. *Funct. Ecol.* 3:363-371.

Eisele, K., D. S. Schimel, L. A. Kapustka, and W. J. Parton. Effects of available P and N:P ratios on non-symbiotic nitrogen fixation in prairie soils. *Oecologia* 79:471-474.

Grace, J. 1977. Plant response to wind. Academic Press, New York, NY.

Hall, D.O. 1989. Carbon flows in the biosphere: present and future. *J. Geol. Soc.* 146:175-181.

Ham, J.M., and J.L. Heilman. 1991. Aerodynamic and surface resistances affecting energy transport in a sparse crop. *Agric. Forest Meteorol.* 53:267-284.

Heagle, A.S., and R.B. Philbeck. 1978. Exposure techniques. In W.W. Heck, S.V. Krupa, & S.M. Linzon (eds.) *Methodology for the assessment of air pollution effects on vegetation*, Ch. 6, pp. 1-19, Air Pollution Contr. Assoc., Pittsburg.

Heagle, A.S., R.B. Philbeck, & M.B. Letchworth. 1979. Injury and yield responses of spinach cultivars to chronic doses of ozone in open-top field chambers. *Environ. Qual.* 8:368-373.

Heggestad, H.E., A.S. Heagle, J.H. Bennett, and E.J. Kock. 1980. The effects of photochemical oxidants on the yield of snap beans. *Atmos. Environ.* 14:317-326.

Hobbs, N.T., D.S. Schimel, C.E. Owensby, and D.S. Ojima. 1991. Fire and grazing in the tallgrass prairie: Contingent effects on nitrogen budgets. *Ecology* 72: (in press).

- Holland, E. A., W.J. Parton, J. K. Detling, and D. L. Coppock. Physiological responses of plant populations to herbivory and their consequences for nutrient flows. *American Naturalist* (submitted).
- Jarvis, P.G. and T.A. Mansfield. 1981. *Stomatal Physiology*. Cambridge University Press, Cambridge.
- Jones, H.G. 1985. Partitioning stomatal and non-stomatal limitations to photosynthesis. *Plant, Cell and Environment* 8: 95-104.
- Kanemasu, E.T., and C.B. Tanner. 1969. A heat transport anemometer. *Bioscience* 18:327-329.
- Kimball, B.A., and R.D. Jackson. 1979. Soil heat flux. p. 211-229. In B.J. Barfield and J.F. Gerber (ed) *Modification of the aerial environment of plants*. ASAE Monograph No. 2, American Society of Agricultural Engineers.
- Kimball, B.A. 1986. Carbon dioxide stimulation of growth and yield under environment restraints. In H.Z. Enock and B.A. Kimball (eds.) *CO₂ Enrichment of greenhouse crops*. CRC Press, Boca Raton, FL.
- Kittel, T.G.F., D. S. Schimel, and W. J. Parton. Sensitivity of the North American Central Grassland ecosystems to climate change. (in prep).
- Knapp, A.K. 1984. Water relations and growth of three grasses during wet and drought years in a tallgrass prairie. *Oecologia* 65:35-43.
- Knapp, A.K. 1984. Post-burn differences in solar radiation, leaf temperature and water stress influencing production in a lowland Tallgrass prairie. *American J. of Botany* 71:220-227.

- Knapp, A.K. 1985. Early season production and microclimate associated with topography in a C₄ dominated grassland. *Acta Oecologia* 6:337-346.
- Knapp, A.K. and W.K. Smith. 1987. Stomatal and photosynthetic responses during sun/shade transitions in subalpine plants: influence on water use efficiency. *Oecologia* 74: 62-67.
- Knapp, A.K. and W.K. Smith. 1990. Stomatal and photosynthetic responses to variable sunlight. *Physiologia Plantarum* 78: 160-165.
- Karl, T.R., R.R. Heim, Jr., and R.G. Quayle. 1991. The greenhouse effect in central North America: If not now, when? *Science* 251: 1058-1061.
- Kramer, P. 1981. Carbon dioxide concentration, photosynthesis and dry matter production. *Bioscience* 31: 29-33.
- Long, S.P. and H.W. Woolhouse. 1978. The responses of net photosynthesis to vapour pressure deficit and CO₂ concentration in *Spartina townsendii* (*sensu lato*), a C₄ species from a cool temperate climate. *Journal of Experimental Botany* 29: 29: 567-577.
- Mandl, R.H., L.H. Weinstein, D.C. McCune, & M. Keveny. 1973. A cylindrical, open-top chamber for the exposure of plants to air pollutants in the field. *J. Environ. Qual.* 2:371-376.
- Morison, J.I.L. 1985. Sensitivity of stomata and water use efficiency to high CO₂. *Plant, Cell and Environment* 8: 467-474.

- Ojima, D. S. 1987. The short-term and long-term effect of burning on tallgrass ecosystem properties and dynamics. Ph.D. Dissertation, Colorado State University, Fort Collins.
- Ojima, D.S., Parton, W.J., Schimel, D.S., and C.E. Owensby. 1990. Simulated impacts of annual burning on prairie ecosystems. In: Fire in North American Tallgrass Prairies. S.L. Collins and L.L. Wallace (eds.). University of Oklahoma Press Norman and London, 175p.
- Olszyk, D.M., T.W. Tibbitts, and W.M. Hertsberg. 1980. Environment in open-top field chambers utilized for air pollution studies. J. of Environ. Qual. 9:610-615.
- Owensby, C.E., P.I. Coyne, L.M. Auen, and N. Sionit. 1991. Rangeland - plant response to elevated CO₂: Large-chamber system. U.S. Dep. of Energy Greenbook Report. No. 059.
- Parton, W. J., D. S. Schimel, C. V. Cole, and D. S. Ojima. 1987. Analysis of factors controlling soil organic levels of grasslands in the Great Plains. Soil Sci. Soc. Am. J. 51:1173-1179.
- Parton, W. J., J. W. B. Stewart and C. V. Cole. 1988. Dynamics of C, N, P and S in grassland soils: A model. Biogeochemistry 5:109-131.
- Prudhomme, T.I., W.C. Oechel, S.J. Hastings, and W.T. Lawrence. 1984. Net ecosystem gas exchange at ambient and elevated CO₂ concentrations in the Tussock Tundra at Toolik, Alaska: An evaluation of methods and initial results. pp. 155-162. In J.H. McBeath (ed.) Potential effects of carbon dioxide induced climatic changes in Alaska. Proc. of a conference (Miscellaneous Pub. 8-31). School of Agriculture and Land Resources Management, Univ. of Alaska, Fairbanks, AK.

- Rogers, H.H., N. Sionit, J.D. Cure, J.M. Smith, and G.E. Bingham. 1983. Influence of elevated carbon dioxide on water relations of soybeans. *Plant Physiol.* 74:233-238.
- Rogers, H.H., J. Thomas and G. Bingham. 1983. Response of agronomic and forest species to elevated CO₂. *Science* 220: 428-429.
- Sakuratani, T. 1981. A heat balance method for measuring water flux in the stem of intact plants. *J. Agric. Meteorol.* 37:9-17.
- Shaer, Y.A. and C.H.M. Van Bavel. 1987. Relative role of stomatal and aerodynamic resistances in transpiration of a tomato crop in a CO₂-enriched greenhouse. *Agric. and Forest Meteorol.* 41:77-85.
- Schimel, D.S., W.J. Parton, T.G.F. Kittel, D.S. Ojima, and C.V. Cole. 1990. Grassland biogeochemistry: Links to atmospheric processes. *Climate Change* 17:13-25.
- Schimel, D.S., T.G.F. Kittel, A.K. Knapp, T.R. Seastedt, W.J. Parton, and V.B. Brown. 1991. Physiological interactions along resource gradients in a tallgrass prairie. *Ecology* 72: in press.
- Schlentner, R.E. and K. Van Cleve. 1985. Relationships between CO₂ evolution from soil, substrate temperature, and substrate moisture in four mature forest types in interior Alaska. *Can. J. For. Res.* 15:97-106.
- Schulze, E.D. and A.E. Hall. 1982. Stomatal responses, water loss and CO₂ assimilation rates of plants in contrasting environments. Pages 181-230 In: *Physiological Plant Ecology II. Encyclopedia of Plant Physiology Vol. 12B*, O.L. Lange, P.S. Nobel, C.B. Osmond and H. Ziegler, eds.

- Sionit, N., H. Hellmers, and B.R. Strain. 1982. Interaction of atmospheric CO₂ enrichment and irradiance on plant growth. *Agronomy Journal* 74: 721-725.
- Starzecki, W and M. Czarnowski. 1989. Increase in photosynthetic rate in tomato leaves induced by compensation of low intensity irradiation by increased CO₂ concentration. *Acta Physiologiae Plantarum* 11: 215-221.
- Tanner, C.B., and Y. Shen. 1990. Water vapor transport through a flail chopped corn residue. *Soil Sci. Soc. Am. J.* 54:945-951.
- Unsworth, M.H., A.S. Heagle, and W.W. Heck. 1984a,b. Gas exchange in open-top field chambers I. Measurement and analysis of atmospheric resistances to gas exchange. II. Resistances to ozone uptake by soybeans. *Atmos. Environ.* 18(2): 373-385.
- Valle, R., J.W. Mishoe, W.J. Campbell, J.W. Jones, L.H. Allen, Jr. 1985. Photosynthetic responses of 'Bragg' soybean leaves adapted to different CO₂ environments. *Crop Sci.* 25:333-339.
- Van Bavel, C.H.M. and R.J. Lascano. 1987. ENWATBAL: A numerical method to compute the water loss from a crop by transpiration and evaporation. Soil and Crop Science Department, Texas Agr. Expt. Sta., Texas A&M University, College Station, TX 77843.
- Wesely, M.L., J.A. Egtmon, D.H. Stedmon, and E.D. Yalvao. 1982. Eddy correlation measurement of NO₂ flux to vegetation and comparison to O₃ flux. *Atmos. Environ.* 16:815-820.
- Wiant, H.V. 1967. Influence of wind on the concentration of CO₂ at the forest floor. *J. For.* 65:563.

Ziska, L.H., B.G. Drake and S. Chamberlain. 1990. Long-term photosynthetic response in single leaves of a C₃ and C₄ salt marsh species grown at elevated atmospheric CO₂ *in situ*. *Oecologia* 83: 469-472.

**DATE
FILMED**

12/18/92