DOE/ER/40145--7

DE93 001798

RESEARCH IN THEORETICAL NUCLEAR PHYSICS

Progress Report

November 1, 1989 - September 30, 1992

Takeshi Udagawa
Theoretical Nuclear Physics Group
Department of Physics
University of Texas
Austin, Texas, 78712

September 1992

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-84ER40145

MASTER
ds

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

TABLE OF CONTENTS

i	Introduction	1
2	Overview	2
	2.1 Low Energy Nuclear Reactions Induced by Light and Heavy Ions	2
	2.2 Medium Energy Physics	18
	2.3 Nuclear Structure	26
3	Title Pages of Published Papers	43
4	Title pages of Submitted Papers	71
5	Personal of Group	76

1. INTRODUCTION

This report describes the accomplishments in basic research in nuclear physics carried out by the theoretical nuclear physics group in the Department of Physics at The University of Texas at Austin, during the period of November 1, 1989 to September 30, 1992 under grant DE-FG05-84ER40145 between the University of Texas at Austin and the United States Department of Energy.

The major part of this report is contained within sections 2, 3, and 4. Section 2 is a presentation of our research achievements and their significance. Both work completed and in progress are covered. In section 3, title pages of papers published during the period are reproduced, while in section 4, first pages of papers submitted but not yet in print are reproduced. These papers are numbered as 3-1 through 3-26 and 4-1 through 4-4, according to the order which they appear in these two sections.

Section 2 gives an overview of our work, its interrelations, and its relation to work done by other groups. As seen, a variety of subjects are included in our work, and thus the description is presented in three separate parts; (2.1) through (2.3). Each part includes brief introductory remarks, followed by the description of work done on specific topics. The main topic in each part can be found in the list of contents given at the beginning of this report. In the presentations in section 2, frequent references are made to papers in sections 3 and 4, and also to many other papers.

2. OVERVIEW

2.1 Low Energy Nuclear Reactions Induced by Light and Heavy Ions

One of the most important goals of nuclear reaction theories is to establish an approach for analyzing experimental data from different types of nuclear reactions, i.e., complete fusion (CF), incomplete fusion (IF) and direct reactions (DR), in a systematic way based on a single theoretical framework. In an attempt to achieve this goal, we have developed a few techniques for calculating cross sections for CF and various types of IF reactions within the framework of DR theory. As is well known, the DR theory itself can be formulated on the basis of the unified theory of nuclear reactions. Within this theory, the fusion part of the reaction process is incorporated in the imaginary part of the optical potential (though such a description can account for the energy averaged behavior only). This is the key ingredient in our approach.

The technique developed for the IF reaction calculation is the breakup-fusion (BF) approach. In this approach, we assume that the reaction proceeds as breakup of the incident particle a into, say, $b + x$, followed by absorption of x by the target A , b being the emitted particle. In other words, the IF reaction is viewed as a partial fusion (or more precisely, reaction) taking place in three (or more) body channels that are reached by breakup (or by some other DR, knockout, for instance). We then treat this partial fusion using the imaginary part of the optical potential W_x for the $x + A$ system. Since the first breakup reaction is itself a typical DR, it can be treated by the usual DR theory. It is thus possible to calculate the whole BF process within the framework of the DR theory.

The technique developed for calculating the fusion cross section, which is called the DR approach to fusion (DRAF), also utilizes the imaginary part of the optical potential. More specifically, fusion is treated as absorption by the inner part of the imaginary potential W . We call this inner portion of W the fusion potential W_F . The remainder (outer part) of W may then be ascribed to the IF-plus-DR processes, which we call the DR part. We then calculate the total fusion and DR cross sections, σ_F and σ_D respectively, as $\sigma_i = (2/\hbar v) \langle \chi^{(+)} | W_i | \chi^{(+)} \rangle$ ($i=F$ or D), where $\chi^{(+)}$ is the optical model wave function.

The major part of the work described in the following subsections concerns the application, testing, and improvement of these theoretical methods.

2.1.1 Calculations of Complete Fusion, Incomplete Fusion and Direct Reaction Cross Sections within the Framework of Direct Reaction Theories

Using the two techniques discussed in the introductory remarks, i.e., the BF approach¹ and DRAF², together with the multistep direct reaction (MSDR) method,³ developed also by our group, it is possible to calculate the total non-compound nuclear reaction cross section σ_D by two different methods. One approach is to use DRAF, while the other is to calculate cross sections of all possible IF and DR reactions using the BF and DR methods, and then sum the individual cross sections. Let us write the σ_D 's obtained by these two methods as σ_D^{DRAF} and σ_D^{IF+DR} , respectively. An important issue here is whether these two cross sections mutually agree or not. Furthermore, we may question whether the regions from whence the dominant contributions to these cross sections originate are the same in the two approaches.

In collaboration with Dr.Mastroleo at Instituto de Estudos Avancados, Sao Jose dos Campos, Brazil and Dr.Mustafa at Lawrence Livermore National Laboratory, we recently tried to answer the above question,⁴ taking as an example the $d+^{93}\text{Nb}$ system. Fig. 1 shows the results of the calculated σ_D^{DRAF} and σ_D^{IF+DR} for $E_d=10\sim 25$ MeV. Use was made of a fusion potential radius of $R_F=7.2$ fm (or $r_F=1.58$ fm) in obtaining σ_D^{DRAF} . With this choice for the R_F -value, good agreement between the two theoretical predictions was obtained. We then examined the radial regions where the elastic breakup (EB) and BF reactions dominantly occur. The results showed that the centers of the radial region where the EB and BF reactions take place are 10 fm and 8.8 fm respectively. The resulting centers are well outside the fusion potential radius $R_F=7.2$ fm, used for the DRAF calculations. Thus both approaches reach a consistent conclusion about the regions where the IF and DR predominantly occur in the incident channel. This shows that the two approaches are mutually consistent.

-
1. T. Udagawa and T. Tamura, Phys. Rev. C **24**, 1348 (1981).
T. Udagawa and T. Tamura, Phys. Rev. C **33**, 494 (1986).
T. Udagawa, X.-H. Li and T. Tamura, Phys. Rev. C **37**, 429 (1988).
 2. T. Udagawa, B. T. Kim and T. Tamura, Phys. Rev. C **32**, 124 (1984).
 3. T. Tamura, T. Udagawa, and H. Lenske, Phys. Rev. C **26**, 379 (1982).

4. R. C. Mastroleo, T. Udagawa, and M. Mustafa, *Phys. Rev. C* **42**, 683 (1990). (See also 3-1.)
 R. C. Mastroleo, T. Udagawa, and M. Mustafa, *Proc. of Intern. Symp. on Nuclear Collective Motion and Nuclear Reaction Dynamics*, held at Riken, Japan, eds. by K.-I. Kubo, M. Ishihara, M. Ichimura, and S. Yamaji (World Scientific, 1991), p.270. (See also 3-2.)

2.1.2 Dynamical Calculations of Linear Momentum Transfer in α -Induced Reactions

One remarkable result of the calculations reported in the previous subsection is that the sum of the calculated total incomplete fusion (IF) and direct reaction (DR) cross sections, σ_{IF} and σ_{DR} , respectively, amounts to about 50% of the total reaction cross section, σ_R . Since $\sigma_{IF} + \sigma_{DR}$ well approximates the total non-compound nuclear cross section, we reach the conclusion that the complete fusion (CF) cross section, σ_{CF} , evaluated as $\sigma_{CF} = \sigma_R - \sigma_{IF} - \sigma_{DR}$ is about 50% of σ_R . This is a sizable reduction of σ_{CF} from σ_R . Consider that σ_{CF} is a crucial ingredient in the usual statistical model calculation of emission cross sections of various particles from the decay of the compound nucleus,¹ where σ_R is customarily used in place of σ_{CF} . It is expected then that if use is properly made of σ_{CF} , the calculated emission cross sections will significantly change. In this respect, it is important to test the validity of the above result for σ_{CF} .

Unfortunately, there are presently no experimental data available for σ_{CF} . Therefore, it is not possible to make a direct test of the calculated σ_{CF} against the data. It is, however, possible to make an indirect test via comparison of the calculated linear momentum transfer per nucleon, p/a , with experiment, since p/a is very sensitive to σ_{CF} . Note that we can calculate p/a in a consistent manner by using the σ_{CF} calculated as $\sigma_{CF} = \sigma_R - \sigma_{IF} - \sigma_{DR}$ and also various DR and IF cross sections evaluated in the course of obtaining the total σ_{IF} and σ_{DR} .

Such a test was indeed recently attempted,² taking the $\alpha + {}^{59}\text{Co}$ system³ with an incident energy $E_a = 80$ MeV, as an example. The calculated p/a -value was found to be 163 MeV/c, which agreed well with the experimental value of 160 MeV/c. We note that the dominant contribution (69%) of the calculated p/a comes from CF. The good agreement of the calculated p/a with the experimental data thus strongly supports the validity of the calculated

σ_{CF} (=967 mb, which is 58% of σ_R). We also note that the various DR and IF cross sections used in obtaining the above p/a -value have also been tested against the data taken for the $\alpha+^{58}\text{Ni}$ system.⁴

Using the calculated value of σ_{CF} =967 mb, one can go one step further to estimate the radius, R_F , of the fusion potential, W_F , introduced in our direct reaction approach to fusion (DRAF).⁵ (See also the discussion given in the introductory remark and previous subsection.) R_F can be determined by fitting the calculated σ_{CF} within DRAF to the above σ_{CF} =967 mb. The R_F -value thus deduced is $R_F = 6.57$ fm, which corresponds to a reduced radius parameter of $r_F = 1.69$ fm. The r_F -value thus obtained is not very far from the value of 1.58 fm determined previously for the $d+^{93}\text{Nb}$ system with $E_d = 10\text{-}25$ MeV,⁶ as discussed in the previous subsection 2.1.1. Note that these two values are both larger than the typical nuclear radius parameter, say, $r_0 = 1.2$ fm, indicating that fusion begins at a fairly large distance. This conclusion is in very good agreement with what was concluded for heavy-ion fusion.⁶

As already argued above, the fact that σ_{CF} is significantly smaller than σ_R has an important influence on the calculated cross sections of particles emitted from the highly excited compound nucleus. This is most likely the case since we along with other research groups have demonstrated that some of the difficulties encountered can be removed by the use of σ_{CF} or equivalently the transmission coefficient evaluated from DRAF.^{1,5} The work described in this and previous subsections, together with that discussed in the subsequent sections, was summarized⁷ recently at the International Conference held in honour of the Late Professor Taro Tamura, at RIKEN, Japan, December 18-20, 1989.

-
1. N. G. Nicolis *et al.*, Phys. Rev. C **45**, 2393 (1992).
 2. R. C. Mastroleo, T. Udagawa, and M. G. Mustafa, Phys. Lett. B **245**, 329 (1990).
(See also 3-3.)
 3. J. Jastrzebski, P. P. Singh, T. Mróz, S. E. Vigdor, M. Fatyga, and H. J. Karwowski, Phys. Rev. C **34**, 60 (1986).
 4. J. R. Wu *et al.*, Phys. Rev. C **20**, 1284 (1979).
 5. T. Udagawa, B. T. Kim, and T. Tamura, Phys. Rev. C **32**, 124 (1981).
 6. M. G. Mustafa, T. Tamura, and T. Udagawa, Phys. Rev. C **35**, 2067 (1989).

7. T. Udagawa, *Proc. of Intern. Symp. on Nuclear Collective Motion and Nuclear Reaction Dynamics*, held at Riken, Japan, eds. by K.-I. Kubo, M. Ishihara, M. Ichimura, and S. Yamaji (World Scientific, 1991), p.181. (See also 3-4.)

2.1.3 Breakup-Fusion Analyses of the $^{40}\text{Ca}(^6\text{Li},d)^{44}\text{Ti}$ Reactions and α -Cluster Structure in ^{44}Ti

The breakup-fusion (BF) approach employed in the studies reported in the previous subsections was originally developed for the description of the incomplete-fusion reaction, i.e., a stripping-type transfer reaction to the unbound region. The approach was, however, shown^{1,2} to be applicable to calculate cross section for stripping reactions to the bound region also. In fact, the approach provides a unified description of stripping reactions to both bound and unbound regions. The method was then applied to analyze single nucleon transfer reaction data, demonstrating that it is even capable of extracting spectroscopic information for high-lying single particle states.

Recently, we have extended the analysis^{3,4} to data of the α -transfer $^{40}\text{Ca}(^6\text{Li},d)^{44}\text{Ti}$ reactions at incident energies of 28, 50, and 60 MeV.⁴⁻⁶ The aim is to extract spectroscopic information, particularly information about the α -cluster structure in ^{44}Ti , which is of current interest.⁷⁻¹² Since many of the α -cluster states appear in the unbound region, use of the BF method is most appropriate for the purpose. From the analysis of the data, transferred angular momenta $\ell_{\alpha,i}$ and spectroscopic factors S_i have been extracted for the individual final states up to $E_{ex} \approx 10$ MeV.

The relative S -factors deduced from the data with the three different incident energies agreed well with each other. The most remarkable result is an identification of a 0^+ state at $E_{ex} = 9.32$ MeV, (identified only in the analysis of the 60 MeV data). We have assigned the state to be the 0^+ member of the $N=14$ band. From this assignment we were further able to estimate the absolute magnitude of the S -factors for the lowest 0^+-12^+ states of the $N=12$ band. The values are $S=0.45, 0.38, 0.19, 0.30, 0.21, 0.19$ and 0.32 , respectively, which agree fairly well with the theoretical values of $S=0.36, 0.41, 0.40, 0.29, 0.35, 0.31$ and 0.23 obtained by Arima and Tomoda.¹³ We were also able to confirm for the first time in the

(${}^6\text{Li}, d$) reaction that the 8.04 MeV state is the 12^+ member of the $N=12$ band. The S -factors have been obtained for a fairly large number of $J^\pi=0^+, 2^+, 4^+$ and 3^- states. The results, together with observation of the $N=14$ 0^+ state at $E_{ex}=9.32$ MeV, strongly suggest that the cluster strength of $N=12$ and 13 is widely spread into many states.

The conclusions drawn in the study described above are still not sufficiently solid. In order to test them further, additional studies are needed, particularly in the determination of the absolute magnitude of the S -factors. One possible way to accomplish this is to extend the analysis to the continuum region. There it is expected that the α -cluster picture of the ${}^6\text{Li} \rightarrow \alpha + d$ process becomes more valid, and therefore we may obtain more reliable information on the overall normalization constant from the analysis of such data. Unfortunately, such data are not available at present. The extension of the present measurement to highly excited continuum states is thus highly desirable.

-
1. T. Udagawa, Y. J. Lee, and T. Tamura, Phys. Lett. B **196**, 291 (1987).
 2. T. Udagawa, Y. J. Lee, and T. Tamura, Phys. Rev. C **39**, 1 (1989).
 3. Chong-Yeal Kim and T. Udagawa, Phys. Rev. C **46**, 532 (1992) (See also 3-5.)
 4. P. Guazzoni, M. Jaskola, L. Zetta, Chong-Yeal Kim, T. Udagawa, and G. Bohlen, to be published. (See also 4-1.)
 5. H. W. Fulbright, C. L. Bennet, R. A. Lindgren, R. G. Markham, S. C. McGuire, G. C. Morrison, U. Strohmusch, and J. Töke, Nucl. Phys. **A284**, 329 (1977).
 6. T. Yamaya, S. Oh-ami, M. Fujiwara, T. Itahashi, K. Katori, M. Tosaki, S. Saito, S. Hatori, and S. Ohkubo, Phys. Rev. C **42**, 1935 (1990).
 7. F. Michel, G. Reidemeister, and S. Ohkubo, Phys. Rev. Lett. **57**, 1215 (1986).
 8. F. Michel, G. Reidemeister, and S. Ohkubo, Phys. Rev. C **38**, 2377 (1988).
 9. A. C. Merchant, K. F. Pál, and P. E. Hodgson, J. Phys. G. **15**, 605 (1989).
 10. A. Arima, *Proc. Topical Conference on Physics of Medium Light Nuclei*, ed. by P. Blasi and R. Ricci (Editrice Compositori, Bologna, 1979) p.19.
 11. T. Wada, and H. Horiuchi, Phys. Rev. C **38**, 2063 (1988).
 12. S. Ohkubo, *Proc. of 6th International Conference on Nuclear Reaction Mechanisms*, ed. by E. Gadioli (Universita Degli Studio Di Milano, 1991) p.396.
 13. T. Tomoda and A. Arima, Nucl. Phys. **A303**, 217 (1978).

2.1.4 Breakup of ${}^7\text{Li}$ into α - t Continuum at Relative Energy of $0 \sim 2$ MeV

This is a continued effort in collaboration with the experimental group at Texas A&M University. The subject is the breakup of ${}^7\text{Li}$ into α and t , which has been studied by several authors^{1,2}, because of the interest in the value of the so-called astrophysical S -factor for very small relative energies ε ($=10\sim 200$ keV) in the α - t continuum. In principle, the S -factor could be determined by measuring the cross section of the direct γ -capture process $\alpha(t,\gamma){}^7\text{Li}$.^{3,4} Since the cross section is extremely small at small ε , however, it has been difficult to measure the cross section in this way.

Baur, *et al.*, proposed to use Coulomb breakup data to determine the S -factor indirectly.² Using a newly developed technique, Utsunomiya, *et al.*⁵⁻⁷, at Texas A&M performed measurements of the Coulomb breakup of ${}^7\text{Li}$. There 42 and 63 MeV beams of ${}^7\text{Li}$ were bombarded on various targets such as ${}^{208}\text{Pb}$, ${}^{144}\text{Sm}$, ${}^{120}\text{Sn}$, ${}^{58}\text{Ni}$, and ${}^{27}\text{Al}$. The breakup cross section were determined as functions of the relative energy ε and the angle of the $\alpha - t$ system.

We have analyzed the data for the yields to the continuum at $\varepsilon = 0 \sim 1.0$ MeV, attempting to extract the astrophysical S -factors as a function of ε . In the most recent analysis,⁷ we introduced an idea that the nuclear-Coulomb interference effects do not significantly change the ε -dependence of the Coulomb excitation cross section. We tested the validity of this idea by making a DWBA calculation, confirming that it is indeed valid. The energy dependence of the resultant S -factors then agreed very well with those deduced from the (γ,p) reaction data of Ref. 4.

-
1. A. C. Shotter *et al.*, Phys. Rev. Lett. **53**, 1539 (1981)
 2. G. Baur, C. A. Bertulani and H. Rebel, Nucl. Phys. **A458**, 188 (1986).
 3. G. M. Griffiths *et al.*, Can. J. Phys. **39**, 1397 (1961).
 4. U. Schroder *et al.*, Phys. Lett. **B192**, 55 (1987).
 5. H. Utsunomiya, R. P. Schmitt, Y.-W. Lui, D. R. Haenni, H. Dejbakhsh, L. Cooke, P. Heimberg, A. Ray, T. Udagawa and T. Tamura, Phys. Lett. **B211**, 24 (1988).
 6. H. Utsunomiya, Y.-W. Lui, L. Cooke, H. Dejbakhsh, D. R. Haenni, P. Heimberg, A. Ray, B. K. Srivastava, R. P. Schmitt, and T. Udagawa, Nucl. Phys. **A511**, 379 (1990). (See also 3-6.)
 7. H. Utsunomiya, Y.-W. Lui, D. R. Haenni, H. Dejbakhsh, L. Cooke, B. K. Srivastava, W. Turmel, D. O'Kelly, R. P. Schmitt, D. Shapira, J. Gomez del Campo, A. Ray, and T. Udagawa, Phys. Rev. Lett. **65**, 847 (1990). (See also 3-7.)

2.1.5 Dispersive Optical Model Potentials for Subbarrier Heavy Ion Fusion and Scattering

Analyses of the subbarrier fusion cross section σ_F based on the direct reaction approach to fusion (DRAF)¹ have been one of the major subjects of our heavy ion reaction studies during the past several years. Very recently, we have initiated a new type of analysis, i.e., the simultaneous χ^2 -analysis²⁻⁴ of elastic scattering (σ_{EL}), fusion (σ_F), and total direct reaction (DR) cross section (σ_D) data. The optical potential used in such an analysis consists of a volume-type real potential, $V_R(r)$, a volume-type imaginary fusion potential, $W_F(r)$, and a surface-derivative-type imaginary DR potential, $W_D(r)$. The parameters of these potentials are then fixed from the χ^2 -analysis of σ_{EL} , σ_F , and σ_D . In DRAF, σ_i ($i=D$ and F) is calculated within the optical model as¹ $\sigma_i = (2/\hbar v) \langle \chi^{(+)} | W_i(r) | \chi^{(+)} \rangle$, where v is the incident velocity and $\chi^{(+)}$ is the elastic distorted wave function.

Such analyses have so far only been made for the two systems, $^{16}\text{O} + ^{208}\text{Pb}$ and $^{32}\text{S} + ^{58,64}\text{Ni}$. The results obtained are, however, very encouraging, revealing many interesting features in V_R , W_F and W_D , which in turn provide valuable information on the dynamics of the nuclear reactions involved. For instance, the radius, r_F , of the fusion potential W_F turned out to be quite large ($r_F=1.40$ fm), larger than the typical radius (1.30 fm) for the Coulomb-barrier top, indicating that fusion starts to occur at a large distance. Extracted values for W_F at the strong absorption radius R_A , $W_F(R_A)$, showed very clearly the so-called threshold anomaly⁵, i.e., $W_F(R_A)$ showed a sudden decrease when the incident energy E went below the Coulomb barrier height. Further, W_D showed a very interesting system-dependence, reflecting the dynamical features of the systems considered. Moreover, the real and imaginary potentials satisfied very well the required dispersion relation^{6,7} at $r = R_A$. (Very recently, it came to our attention that some of the older data used in the analyses of Ref. 3 do not agree with new data taken very recently at different laboratories.^{8,9} This means that we need to repeat our analysis. We believe, however, that the qualitative features of W_D and W_F determined in Ref. 3 will remain unchanged in the new W_D and W_F .)

Recently, we went one step further to obtain a fully dispersive optical potential, i.e., an optical potential that satisfies the dispersion relation over all space.⁴ This was done for the $^{16}\text{O} + ^{208}\text{Pb}$ system. The optical potential used consisted of the Coulomb potential, $U_C(r)$,

the energy-independent bare (Hartree-Fock or folding) potential, $V_0(r)$, and the real and imaginary polarization potentials, $V(r)$ and $W(r)$, respectively, that describe the coupled-channel (CC) effects. The polarization potentials are further assumed to have volume-type fusion and surface-derivative-type DR parts; $V(r) = V_F(r) + V_D(r)$ and $W(r) = W_F(r) + W_D(r)$.

The optical potential assumed above involves eleven parameters, excluding the Coulomb radius parameter in U_C , which we chose as $r_c = 1.25$ fm. It was impossible to determine all eleven parameters uniquely from a χ^2 -analysis. We thus fixed *a priori* some of the parameters; the parameters of the bare potential $V_0(r)$ were fixed as $V_0=60.4$ MeV, $r_0=1.176$ fm, and $a_0=0.658$ fm, while all the geometrical parameters in the polarization potential $V(r) + iW(r)$ were fixed at $r_F=1.40$ fm, $a_F=0.25$ fm, $r_D=1.50$ fm, and $a_D=0.45$ fm. These values are chosen as the best possible values, taking into consideration the results of earlier studies.^{3,10} Note, however, that these values are still not unique, and this non-uniqueness might introduce a certain ambiguity in the analyses made in Ref. 4.

Since the geometrical parameters in $V_i(r)$ and $W_i(r)$ ($i = F$ and D) are assumed to be the same, the dispersion relations hold for the strength parameters V_i and W_i . The relation reads^{6,7}

$$V_i(E) = V_i(E_s) + \frac{E - E_s}{\pi} P \int_0^\infty dE' \frac{W_i(E')}{(E' - E_s)(E' - E)}, \quad (1)$$

where P stands for the principal value and $V_i(E_s)$ is the value of the potential at a reference energy $E = E_s$. Using the above relation, it is possible to evaluate values of V_F and V_D , once the values of $W_F(E)$ and $W_D(E)$ are known. This means that we now have only two parameters to be fixed from the analysis.

Fig. 2 shows values of $W_F(E)$ and $W_D(E)$ determined from the χ^2 -fit. These $W_F(E)$ and $W_D(E)$ values can be well represented by the following functions of E (in units of MeV):

$$W_F(E) = \begin{cases} 0 & \text{for } E \leq 77.5 \\ 0.52(E - 77.5) & \text{for } 77.5 < E \leq 84 \\ 3.4 & \text{for } 84 \leq E. \end{cases} \quad (2)$$

$$W_D(E) = \begin{cases} 0 & \text{for } E \leq 70 \\ 0.185(E - 70) & \text{for } 70 < E \leq 90 \\ 0.37 & \text{for } 90 \leq E \end{cases} \quad (3)$$

The solid lines shown in Fig. 2 are the W_F and W_D values predicted by the above Eqs. (2) and (3). As seen, the lines fit the empirical values (shown by the filled circles) rather well. In obtaining V_F and V_D from Eq. (1) use was thus made of the above functional forms for $W_F(E)$ and $W_D(E)$, together with $V_F(E_s)=1.8$ MeV at $E_s=84$ MeV and $V_D(E_s)=0.5$ MeV at $E_s=90$ MeV.

In Fig. 2, it is seen that the slope (0.185) of W_D is much smaller than that of W_F (0.52). This means that the E-dependence of W_D is much weaker than that of W_F . This clearly reflects the observed fact that the energy-dependence of the experimental σ_D is much weaker than that of σ_F . This, together with the fact that the W_F dominates the total imaginary potential, $W (=W_F + W_D)$, at least in the higher energy part of the Coulomb-barrier energy region, shows that the rapid energy change in W originates from fusion. Namely, the main cause of the threshold anomaly comes from the coupling of the elastic channel with fusion channels.

In Figs. 3 and 4, we present the final fit of the calculated $R = \sigma_{EL}/\sigma_{RUTH}$ (where σ_{RUTH} is the Rutherford cross section), σ_F and σ_D to the experimental data. As seen, the fit is good. The dotted line shown for one case in Fig. 3 is the R -value obtained by setting $V_D=0$. As seen, when V_D , which is one of the new elements in the analyses, is set to zero, the fit of the calculated R to experiment becomes worse, showing that V_D plays an important role. Note that the deviation of the dotted line from the full line is seen only in the angle region between, say, θ_1 and θ_2 , where θ_1 is the angle at which R starts to decrease while θ_2 is the angle where the value of R becomes $R \sim 10^{-2}$. This region corresponds to the radial region, where V_D has an appreciable value.

The work described above was presented¹¹ at the international workshop on subbarrier fusion held at Daresbury, June 5-7, 1990.

1. T. Udagawa, B. T. Kim and T. Tamura, Phys. Rev. C **32**, 124 (1984).
2. S. W. Hong, T. Udagawa and T. Tamura, Nucl. Phys. **A491**, 492 (1989).
3. T. Udagawa, T. Tamura and B. T. Kim, Phys. Rev. C **39**, 1840 (1989).
4. B. T. Kim, M. Naito, and T. Udagawa, Phys. Lett. B **237**, 19 (1990). (See also 3-8.)
5. J. S. Lilley, B. R. Fulton, M. A. Nagarajan, I. J. Thompson, and D. W. Banes, Phys. Lett. B **151** 181 (1985).
6. C. Mahaux, H. Ngo and G. R. Satchler, Nucl. Phys. **A449** (1986) 354; Nucl. Phys. **A456**, 134 (1986).
7. M. A. Nagarajan, C. Mahaux and G. R. Satchler, Phys. Rev. Lett. **54**, 1136 (1985).
8. R. J. Tighe *et al.*, Phys. Rev. C **42**, 1530 (1990)
9. Y. Sugiyama, private communication.
10. G. R. Satchler, M. A. Nagarajan, J. S. Lilley and I. J. Thompson, Ann. of Phys. **178**, 110 (1987).
11. T. Udagawa and B. T. Kim, *Inst. Phys. Conf. Ser. No 110*, (IOP Publishing Ltd, 1991), 103. (See also 3-9.)

2.1.6 Systematics of Sub-Barrier Fusion Cross Sections and the Threshold Anomaly

One of the consequences found in the study reported in the previous subsection 2.1.5 is that the main cause of the threshold anomaly¹ is fusion, i.e., the coupling of the elastic channel to fusion channels. We have found additional evidence of this, which was reported recently.² The evidence is that the threshold energy E_F of the subbarrier fusion cross section, σ_F , agrees very well with the threshold energy E_0 of the imaginary potential W . The threshold anomaly is of course seen in W , which is extracted from an analysis of elastic scattering data. As noted previously, W at the strong absorption radius R_A can be very well represented in the sub-barrier region by a linear function of E as $W(R_A) = \alpha(E - E_0)$. The zero intercept E_0 then defines the threshold energy of W . Stelson *et al.*³ observed the quantity S defined as $S = (E\sigma_F)^{1/2}$ can also be very well represented by a linear function of E as $S = \beta(E - E_F)$ in the sub-barrier region. E_F then defines the threshold energy E_F of the sub-barrier fusion.

We have tried¹ to collect information on the (E_F, E_0) -value. We have so far been able to obtained such values for three systems, $^{16}\text{O} + ^{60}\text{Ni}$, $^{16}\text{O} + ^{63}\text{Cu}$ and $^{16}\text{O} + ^{208}\text{Pb}$. The values obtained are: $(E_F, E_0) = (29.2, 29.3)$, $(30.2, 31.0)$ and $(71.3, 69.0)$ MeV, respectively, for the above three system. As seen, the E_F - and E_0 -values agree well with each other.

E_F is a quantity characterizing the threshold of the fusion reaction. The above agreement of E_0 with E_F thus indicates that the strong energy dependence seen in $W(E, R_A)$ in the threshold energy region (threshold anomaly) indeed comes from fusion.

-
1. J. S. Lilley, B. R. Fulton, M. A. Nagarajan, I. J. Thompson, and D. W. Banes, Phys. Lett. B **151**, 181 (1985).
B. R. Fulton, D. W. Banes, J. S. Lilley, M. A. Nagarajan, I. J. Thompson, Phys. Lett. B **162**, 55 (1985).
 2. T. Udagawa, M. Naito, and B. T. Kim, Phys. Rev. C **45**, 876 (1992). (See also 3-10.)
 3. P. H. Stelson, H. J. Kim, M. Beckerman, D. Shapira, and R. L. Robinson, Phys. Rev. C **41**, 1584 (1990).
P. H. Stelson, Phys. Lett. B **205**, 190 (1988).

2.1.7 Dynamical Fusion and Direct Reaction Potentials For Heavy-Ion Scattering and Fusion at Sub-Coulomb-Barrier Energies

In subsection 2.1.5, we have reported results of our simultaneous χ^2 -analyses¹⁻³ of elastic scattering, fusion and total direct reaction (DR) cross section data taken at sub- and near-Coulomb-barrier energies. One important result obtained from the analyses is that the fusion potential $W_F(r)$, assumed to have a Woods-Saxon shape should have a large radius parameter $r_F \approx 1.4$ fm, placing the resulting radius R_F beyond the peak in the Coulomb barrier. The value is much larger than that usually assumed in the barrier penetration model⁴ or coupled-channels (CC) calculations^{5,6} for fusion. Satchler, *et al.*,⁷ then proposed that the large r_F values have their origin in the known importance of couplings to DR channels. In a CC approach, the so-called “bare” imaginary fusion potential $W_{0F}(r)$ may still be confined inside the Coulomb barrier and have a small radius, $r_{0F}=1.0$ fm. If one projects, however, this CC problem onto the elastic channel, there appears an additional term, ΔW_F , in the resultant fusion potential. This is a part of the dynamical polarization potential (DPP), ΔU , which appears when such a projection is done. The proposal made by Satchler, *et al.*,⁷ is that r_F becomes large because of ΔW_F in ΔU .

A study⁸ was recently made to test whether $\Delta W_F(r)$ indeed makes the r_F -value of the resultant fusion potential ($W_{0F}+\Delta W_F$) much larger than r_{0F} . For this purpose, we calculated

ΔU , particularly the fusion and DR potentials, ΔW_F and ΔW_D , respectively, where ΔW_D is the DR part in the total imaginary potential in ΔU . The actual numerical calculations were carried out for the $^{16}\text{O}+^{208}\text{Pb}$ system at $E_{lab}=80$ MeV, taking into account couplings with the 2.61 MeV 3^- , 3.20 MeV 5^- , and 4.07 MeV 2^+ states in ^{208}Pb , and to the 6.13 MeV 3^- state in ^{16}O using the collective form factor.

The resultant ΔW_F and ΔW_D are found to be highly nonlocal and ℓ -dependent. Because of this, it was difficult to reach a conclusion on Satchler's hypothesis. To see the area where fusion and DR take place more clearly, we thus examined the following absorption densities $A_i(r)$ ($i=F$ and D),

$$A_i(r) = \int_0^\infty \Delta W_i(\ell; r, r') \text{Re}[\rho_\ell(r, r')] dr'. \quad (\rho_\ell(r, r') \equiv \chi_\ell^*(r) \chi_\ell(r')) \quad (1)$$

The above expression can be derived from the divergence of the current density. Figs. 5a and 5b show the calculated $A_F(r)$ and $A_D(r)$ as functions of r for the $\ell=10$ case. The DR absorption density $A_D(r)$ is appreciable in the region of $r > 12$ fm. It has an oscillatory long range tail. The period of oscillation reflects that of the distorted wave function $\chi_\ell(r)$, and the long range tail comes from Coulomb excitation. The nuclear excitation contribution is confined in the peak region of $12 \text{ fm} < r < 14 \text{ fm}$. The center of the peak is located at $r \approx 13$ fm. This region is simply the grazing region where DR are supposed to occur.

The fusion absorption density $A_F(r)$ also has an oscillatory structure. The period of oscillation reflects that of the nonlocal density $\text{Re}[\rho(r, r')]$ in Eq. (1). The envelope of oscillation, peaked at $r \approx 9.7$ fm, is roughly proportional to $v_{\ell\ell'}^2(r) W_{0F}(r) |\chi_\ell(r)|^2$, where $v_{\ell\ell'}^2(r)$ is the channel coupling potential, the limit of $A_F(r)$ obtained when the Green's function is assumed to be a δ -function (adiabatic assumption). The region where $A_F(r)$ takes an appreciable value is about $8 \text{ fm} < r < 12 \text{ fm}$, where $v_{\ell\ell'}(r)$ and $W_{0F}(r)$ strongly overlap. (Note that the peak of $v_{\ell\ell'}(r)$ appears at $R_0=9.8$ fm of $V_0(r)$, while $W_{0F}(r)$ has $R_{0F}=8.4$ fm.) Particularly, the region where $A_F(r)$ takes a large positive value (absorption region) is limited to $r \leq 10.2$ fm. This absorptive region is somewhat shifted outward as compared with the "bare" fusion region defined by $W_{0F}|\chi_\ell|^2$, but still stays inside the barrier (the top of the Coulomb barrier ≈ 11.4 fm). It is much inside the fusion region predicted in our model³ with $r_F=1.4$ fm ($R_F=11.8$ fm). Note that, next to this absorptive region, there appears a

creative region ($10 < r < 12$ fm) where $A_F(r)$ becomes negative. The existence of such a creative region may effectively reduce the overall effects from the channel-couplings on fusion. Therefore, the result does not support the idea proposed in Ref. 7.

In Ref. 8, we have also studied in some details the question whether the calculated ℓ -dependent, non-local potentials can be localized in a reasonable manner. The answer was negative; it is improbable that such a highly nonlocal DPP, like the one obtained in the study of Ref. 8, can be localized in a meaningful manner.⁹ As remarked, the DPP is also ℓ -dependent. In Ref. 9 a very elaborate effort was made to obtain localized and ℓ -averaged DPP's. Unfortunately, the potentials obtained are unable to reproduce consistently the calculated CC elastic scattering cross sections.

In contrast to this, phenomenologically we have always been successful in finding a local, ℓ -independent optical potential that can reproduce the experimental elastic scattering, fusion and total DR cross section data. Apparently, it has not been possible so far to derive such a local, ℓ -independent phenomenological potential directly from CC calculations. One possible origin of this failure may be ascribed to the insufficient number of channels included in the CC calculations and the use of a too simplified form of the coupling scheme (neglecting the coupling between inelastic channels, etc.). One might generally expect¹⁰ that if one includes more channels and uses a more realistic coupling scheme, the complicated nonlocality may be washed away. The Green's functions in the different inelastic channels have different oscillations in r -space and the resulting potential becomes more local because of the random-phase nature of the contributions from a large number of terms. The ℓ -dependence may also be averaged out. It would be interesting and important if such calculations could be done. The work described here was presented at the international conference on nuclear dynamics held at Nikko, Japan.¹¹

1. T. Udagawa, B. T. Kim and T. Tamura, Phys. Rev. C **32**, 124 (1985). (See also 3-8.)
2. T. Udagawa, T. Tamura and B. T. Kim, Phys. Rev. C **39**, 1840 (1989).
3. B. T. Kim, M. Naito, and T. Udagawa, Phys. Lett. B **237**, 19 (1990).
4. J. R. Birkelund and J. R. Huizenga, Ann. Rev. Nucl. Part. Sci. **33**, 265 (1983).
5. S. G. Steadman and M. J. Rhodes-Brown, Ann. Rev. Nucl. Part. Sci. **36**, 649 (1986).
S. Landowne and S. C. Pieper, Phys. Rev. C **29**, 1352 (1984).
6. G. R. Satchler, *et al.*, Ann. Phys. **178**, 110 (1987)
7. G. R. Satchler, *et al.*, Phys. Rev. C **41**, 1869 (1990).
8. B. T. Kim and T. Udagawa, Phys. Lett. B **273**, 37 (1991). (See also 3-11.)
9. I. J. Thompson, *et al.*, Nucl. Phys. A **505**, 84 (1989).
10. L. Carlton, Y. Hahn, and G. Cattapan, Phys. Rev. C **43**, 2441 (1991).
11. B. T. Kim and T. Udagawa, Proc. of Inter. Conf. on *Towards a Unified Picture of Nuclear Dynamics*, ed. by Abe *et al.*, (AIP, New York 1991), p.133. (See also 3-12.)

2.1.8 A Rapid Method for Nonlocal Optical Model Calculations

Theoretical nuclear optical potentials are known to be nonlocal in general.¹ The nonlocality arises from the exchange of particles between the projectile and target (knockon exchange) and from coupling to non-elastic channels. The contribution from the latter channel-coupling (CC) is often referred to as a dynamic polarization potential, and is given by the familiar Feshbach formula.¹

The nonlocality of the polarization potential U_p often becomes very large.²⁻⁷ An example is provided for the $^{16}\text{O} + ^{208}\text{Pb}$ system at sub- and near-Coulomb barrier energies⁸ as discussed in the previous subsection. As remarked there, the nonlocality of the potential calculated in Ref. 8 is so large that it is difficult to localize it in a reasonable manner. In such cases, it is necessary to do a nonlocal calculation.

In a recent article⁹ we proposed a new, rapid, and reliable method for solving the nonlocal optical model Schrödinger equation: $(E - H_\ell)|\chi_\ell\rangle = U'|\chi_\ell\rangle$, where χ_ℓ is the partial wave function that we wish to evaluate, H_ℓ is the partial wave Hamiltonian, which may include a local optical potential U_0 , and $U' \equiv U - U_0$, U being the nonlocal potential. The basic idea of the approach is to modify the above Schrödinger equation into the following integral equation for $|\lambda_\ell\rangle$ defined as $|\lambda_\ell\rangle = U'|\chi_\ell\rangle$;

$$|\lambda_\ell\rangle = |\rho_\ell\rangle + U'G_\ell|\lambda_\ell\rangle \quad (|\rho_\ell\rangle \equiv U'|\chi_\ell^{(0)}\rangle), \quad (1)$$

where $G_\ell = (E - H_\ell + i\epsilon)^{-1}$, while $|\chi_\ell^{(0)}\rangle$ is the solution of the local optical model Schrödinger equation defined by H_ℓ .

We solve the above Eq. (1) using the Lanczos method,¹¹ in which $|\lambda_\ell\rangle$ is expanded in terms of an orthonormal set of basic vectors $|D_i\rangle$. We generate such basic vectors iteratively starting with the doorway state vector $|\rho\rangle$; $|D_0\rangle = (1/N_0)|\rho\rangle$, where N_0 is a normalization constant. The validity of this approach depends on the number N of the basic vectors needed in the expansion in order to obtain enough accuracy. We examined this point for two example cases; one being the nonlocal problem considered by Perey and Buck¹² and the other was the problem encountered when a coupled-channels problem is reduced to a one dimensional problem. For the later problem, we took the $^{16}\text{O}+^{208}\text{Pb}$ system at the incident energy of $E_{lab}=80$ MeV, and considered the simple case of coupling to the 3^- state in ^{208}Pb at an excitation energy of 2.6 MeV. The first example is a rather simple nonlocal problem, while the second case is a highly nonlocal problem. The N -values required for obtaining a good accuracy (0.1%) for the first and second example cases were 3 and 8 for, respectively. The N -value for the second is larger than that for the first example, which is due to the complex nature of the nonlocality of the potential considered in the later case. Nevertheless, the N -values are rather small for both cases, indicating the method is valid for both. A computer program has been developed for numerical calculations.¹³

-
1. H. Feshbach, Ann. of Phys. (N. Y.) **5** 357 (1958), *ibid*, **19**, 287 (1962).
 2. C. L. Rao, M. Reeves, and G. R. Satchler, Nucl. Phys. **A207**, 182 (1973).
 3. P. W. Coulter and G. R. Satchler, Nucl. Phys. **A293**, 269 (1977).
 4. N. Vinh Mau and A. Bouyssy, Nucl. Phys. **A257**, 189 (1976).
 5. F. Osterfeld, J. Wambach, and V. A. Madsen, Phys. Rev. C **23**, 179 (1981).
 6. R. S. Mackintosh and S.G. Cooper, Nucl. Phys. **A494**, 123 (1989).
 7. I. J. Thompson, M. A. Nagarajan, J. S. Lilley, and M. J. Smithson, Nucl. Phys. **A505**, 84 (1989).
 8. B. T. Kim and T. Udagawa, Phys. Lett. B **273**, 37 (1991) (See also 3-11.)
 9. B. T. Kim and T. Udagawa, Phys. Rev. C **42**, 1147 (1990). (See also 3-13.)
 10. T. Udagawa and B. T. Kim, Phys. Rev. C **40**, 2271 (1989).
 11. R. R. Whitehead, A. Watt, B. J. Cole, and I. Morrison, Adv. Nucl. Phys. **9**, 123 (1977).
 12. F. Perey and B. Buck, Nucl. Phys. **32**, 353 (1962).
 13. B. T. Kim, M. C. Kyum, S. W. Hong, M. H. Park, and T. Udagawa, to be published in Com. Phys. Comm. (See also 4-2.)

2.2 Medium Energy Physics

The work reported here was done in collaboration with Osterfeld at KFA in Jülich, Germany and Hong at Sung Kyung Kwan University, Seoul. The work concerns nuclear medium effects on Δ excited in nuclei by means of the (p,n)-, (d,2p)- and (^3He ,t)-charge-exchange reactions and electron scattering at intermediate energies. We shall report on our calculations of the inclusive and exclusive cross sections, and spin observables taking into account the nuclear mean field and two-body delta-hole interactions.

2.2.1 Inclusive Spectra of Δ -Excitations in Nuclei

The shift of the Δ peak position observed in inclusive spectra of the charge-exchange (p,n)¹, (^3He ,t)², and (d,2p)³ reactions for nuclear targets compared to the proton target has been one of the most important issues in medium energy physics over the last several years. The Δ peak in the proton target spectrum appears at $\omega_L \approx 370$ MeV, where ω_L is the energy transfer in the laboratory system. On the other hand, for a nuclear target (^{12}C) the peak appears at $\omega_L \approx 300$ MeV. The peak position of $\omega_L \approx 370$ MeV for the proton target can be explained by the sum of the Δ resonance energy $\omega_R = 294$ MeV, the Δ recoil energy of 95 MeV received by the Δ in the (p,n) excitation process, and an additional shift (-19 MeV) due to momentum-transfer-dependent effects in the $t_{NN,N\Delta}$ transition operator. The downward energy shift of about 70 MeV in nuclear targets has been a mystery to be solved.

Clearly the shift observed is due to nuclear medium effects, and recently we have made⁴ a fairly realistic calculation of the energy spectra of the charge exchange reactions taking into account nuclear mean field and two-body type delta-hole interaction effects. The result has indicated that about 40 MeV out of the 70 MeV shift are due to Fermi motion (binding) of the target nucleons and the attractive one-body potential for a Δ (the nucleus mean field effect). A nuclear correlation effect produced by the Δ -particle-nucleon-hole (ΔN^{-1}) residual interaction in the medium gives rise to an overall shift of about 25 MeV. The interaction is strongly attractive in the spin-longitudinal (LO) channel due to the attractive π -exchange interaction, leading to a lowering of the Δ mass in that channel by about 50 MeV. No

significant shift is, however, observed in the spin-transverse part of the spectrum. Other effects contributing to the shift (~ 15 MeV) come from the spreading potential which describes the Δ conversion processes, such as $\Delta + N \rightarrow N + N$. A similar result has also been obtained by other authors.⁵

The calculations are based on the isobar model^{6,7} and DWIA. Denoting the charge-exchange process as $A + a \rightarrow (B + \Delta) + b$, where A (B) and a (b) denote the target (residual nucleus) and projectile (ejectile), respectively, the inclusive cross section is given by

$$\frac{d^2\sigma}{dJ_b d\Omega_b} = \frac{E_a E_b E_A E_{B+\Delta}}{(2\pi\hbar^2 c^2 W)^2} \frac{k_b}{k_a} R \quad (1)$$

where R is the strength function defined by

$$R = \text{Im}[-\langle \rho | G | \rho \rangle / \pi] = \text{Im}[-\langle \rho | \psi \rangle / \pi] \quad (|\psi\rangle \equiv G | \rho \rangle). \quad (2)$$

In Eq. (1), E_i is the total energy of particle i ($i = a, b, A$, and $B + \Delta$), k_a (k_b) is the wave number of a (b), and W is the total energy in the center of momentum system. $|\rho\rangle$ in Eq. (2) is the doorway state defined as $|\rho\rangle = (\chi_b^{(-)} \varphi_b | t_{NN,N\Delta} | \chi_a^{(+)} \varphi_a \varphi_A \rangle$, where $\chi_a^{(+)}$ ($\chi_b^{(-)}$) is the projectile distorted wave function in the incident (exit) channel, φ_a (φ_b) is the intrinsic wave functions of a (b), and φ_A is the initial target wave function, and $t_{NN,N\Delta}$ is the effective $NN \rightarrow N\Delta$ transition operator. The round bra $(| \rangle$ denotes integration with respect to the projectile coordinates only. Further, G in Eq. (2) is the Green's function that describes the propagation of the $(B + \Delta)$ system and contains all nuclear medium effects, while $|\psi\rangle$ is the wave function of the final $B + \Delta$ system. In the isobar model,^{6,7} G is given as

$$G = \frac{1}{\omega + i\Gamma_\Delta/2 - H_B - T_\Delta - U_\Delta - V_{N\Delta,N\Delta}}. \quad (3)$$

Here, ω is the excitation energy of the $(B + \Delta)$ system, $\Gamma_\Delta(\omega)$ is the energy dependent free decay width of the Δ , while T_Δ and $U_\Delta (= V_\Delta + iW_\Delta)$ are the kinetic energy operator and the Δ -nucleus potential, respectively. The Δ -nucleus potential U_Δ includes, as mentioned earlier, an imaginary part that represents the conversion (spreading) effect. H_B is the Hamiltonian of nucleus B , and $V_{N\Delta,N\Delta}$ is the residual interaction describing the ΔN^{-1} correlations.

In the numerical calculations, use was made of U_Δ determined from analyses of π -nucleus scattering.⁷ The $V_{N\Delta,N\Delta}$ was assumed to consist of π and ρ -exchange potentials with an

additional short range interaction.⁸ Further, the $t_{NN,N\Delta}$ -amplitude was assumed to be simple δ -function type force given as $t_{NN,N\Delta} = g'_{N\Delta} J_{\pi N\Delta} \left(\frac{\Lambda_\pi^2 - m_\pi^2}{\Lambda_\pi^2 - t} \right)^2 (\vec{\sigma}_1 \cdot \vec{S}_2^\dagger)(\vec{\tau}_1 \cdot \vec{T}_2^\dagger)$ with $J_{\pi N\Delta} = 4\pi\hbar c f_{\pi NN} f_{\pi N\Delta} / m_\pi^2 \approx 800 \text{ MeV fm}^3$, $g'_{N\Delta} = 0.335$. We made this choice since studies of the $t_{NN,N\Delta}$ -interaction by means of the $(\vec{d}, 2p)$ -reaction⁹ suggested that $t_{NN,N\Delta}$ is nearly constant in the $(\omega, \vec{q})$ -range relevant to the Δ -resonance region, except for the four momentum transfer, t , dependence coming from the form factor, and also that $t_{NN,N\Delta}$ should have a simple spin structure as assumed above.

In Fig. 6a, we reproduce our calculated results⁴ for the 0-degree spectra of the $^{12}\text{C}(p,n)$ reaction at 800 MeV incident energy, in comparison with the experimental data. The theoretical curves were calculated either with or without inclusion of Δ -hole correlations (full and dashed curves, respectively). Also the correlated spin-LO and spin-transverse (TR) cross sections are shown separately. It is seen that the peak position of the spin-LO spectrum is lowered considerably in energy in comparison with the spin-TR spectrum or with the uncorrelated result. This is due to the attractive π -exchange interaction in the spin-LO channel. The peak of the uncorrelated spectrum appears at $\omega \approx 320 \text{ MeV}$, somewhat lower than the 330 (=370-40) MeV that are expected from the binding energy effect on the Δ and the N . This small shift is due to the conversion (spreading) effect. Fig. 6b shows cross section contributions of various spin states. It is seen that the spectra of lower spin states have their peak at lower ω , showing that the ΔN^{-1} correlations are most important for these lower spin states.

Similar calculations were also done for the $(^3\text{He}, t)$ reaction, obtaining a similar fit to the data as obtained in the (p, n) reaction. The work reported here was presented at various international meetings.¹⁰⁻¹⁶

1. C. G. Cassapakis et al., Phys. Lett. **63B**, 35 (1976)
 B. E. Bonner et al., Phys. Rev. C **18**, 1418 (1978).
 D. A. Lind, Can. J. Phys. **65**, 637 (1987).
2. V. G. Ableev et al., Phys. JETP Lett. **40**, 763 (1984).
 D. Contardo et al., Phys. Lett. **168B**, 331 (1986).
3. C. Ellegaard et al., Phys. Rev. Lett. **59**, 974 (1987).
4. T. Udagawa, S. W. Hong and F. Osterfeld, Phys. Lett. **B245**, 1 (1990). (See also 3-14.)
5. J. Delorme and P. A. Guichon, Phys. Lett. **B263**, 157 (1991).
6. L. S. Kisslinger and W. Wang, Ann. of Phys. **108**, 116 (1977)
7. M. Hirata, J. H. Koch, F. Lenz and E. J. Moniz, Ann. of Phys. **120**, 205 (1979).
 E. Oset, H. Toki and W. Weise, Phys. Rep. **83**, 281 (1982).
8. M. R. Anastasio and G. E. Brown, Nucl. Phys. **A285**, 516 (1977).
9. C. Ellegaard et al., Phys. Lett. B **231**, 365 (1989).
10. F. Osterfeld, S.-W. Hong, and T. Udagawa, *Proceeding of the International Conference on Selected Topics in Nuclear Structure*, Dubna (1989) p.86-94. (See also 3-15.)
11. T. Udagawa, S.-W. Hong, and F. Osterfeld, *Proc. of Second LAMPF International Workshop on Pion-Nucleus Double Charge-Exchange*, eds. by W. R. Gibbs and M. J. Leitch (World Scientific, 1989), p. 439. (See also 3-16.)
12. S.-W. Hong, F. Osterfeld, and T. Udagawa, *Invited talk given at International Workshop on Nuclear Structure and Nuclear Reactions*, RIKEN, Japan, eds. by K.-I. Kubo, M. Ishihara, M. Ichimura, and S. Yamaji (World Scientific, 1991), p. 261. (See also 3-17.)
13. F. Osterfeld, S.-W. Hong, and T. Udagawa, Nucl. Phys. **A527**, (1991) 405c. (See also 3-18.)
14. T. Udagawa, P. Oltmanns, and F. Osterfeld, *Proc. of International Conference on Spin and Isospin in Nuclear Interactions*, eds. by S. W. Wissink, C. D. Goodman, and G. E. Walker (Plenum Press, New York, 1991), p. 69. (See also 3-19.)
15. T. Udagawa, F. Osterfeld, P. Oltmanns, and S.-W. Hong, *Proc. of the 6-th International Conference on Nuclear Reaction Mechanisms*, eds. by E. Gadioli (Universita Deglistudi di Milano, Milano, 1991) p.632. (See also 3-20.)
16. F. Osterfeld, B. Krfggen, P. Oltmanns, and T. Udagawa, *Proc. of International Workshop on Meson Production, Interaction and Decay*, Krakow, May 7-11, 1991. (See Also 3-21.)

2.2.2 Exclusive Cross Sections

The inclusive cross section given by Eqs. (1)-(2) in the previous subsection 2.2.1 includes contributions from three different processes characterized by the mode of the subsequent decay of the Δ after its creation. The three different decay modes in question are the quasi-free (qf) decay $\Delta \rightarrow \pi + N$, the decay due to the conversion (spreading) process $\Delta + N \rightarrow N + N$, and the decay resulting from single pion production, $\Delta + N^{-1} \rightarrow \pi$

(coherent π production). Correspondingly, the strength function R given by Eq. (2) can be split into a sum of three terms

$$R = [\langle \psi | \frac{\Gamma_{\Delta}}{2} | \psi \rangle + \langle \psi | -W_{\Delta} | \psi \rangle + \langle \psi | -Im(V_{N\Delta, N\Delta}) | \psi \rangle] / \pi. \quad (1)$$

Fig. 7 shows the exclusive cross section^{1,2} for the qf, conversion, and coherent π -production events. Note that W_{Δ} and $V_{N\Delta, N\Delta}$, involved in the conversion and coherent π -production components in the above expression, respectively, are ingredients of the nuclear medium that make an important contribution to the shift discussed in the previous subsection 2.2.1. Therefore, comparisons of the predicted exclusive cross sections with the data are expected to provide further tests of our theoretical calculations reported in 2.2.1.

Exclusive cross section data that we can use for such comparisons recently became available^{3,4} through measurements of the $^{12}\text{C}(^3\text{He}, t)$ and $^{12}\text{C}(p, n)$ reactions in coincidence with $\pi + p$, $2p$ and (single) π events. Such data were taken for incident ^3He and p energies of 2 GeV and 800 MeV, respectively, with the triton and neutron emitted at forward angles. These coincidence cross sections are compared with calculations in Fig. 8, taking the (p,n) reaction case.⁴ In doing this, however, we introduced overall normalization factors of 0.25, 0.38 and 1.67 for the qf, conversion, and coherent π production cross sections, respectively. With these normalization factors, the theoretical curves fit the experimental data fairly well. It is particularly remarkable that the different peak positions for the three cross sections are very well reproduced by the calculations. (A similar result has been obtained also for the $^{12}\text{C}(^3\text{He}, t)$ reaction. See Ref. 2.)

The normalization factors 0.25 and 0.38 introduced for the qf and conversion cross sections, respectively, account partially for the final state absorption of π and/or p by the target. (The absorption of π in the coherent π production process is automatically taken into account in the calculation.) Another source for the normalization factors is that the calculated qf and conversion cross sections include besides the πp and $2p$ events also the πn and pn production processes. The measured πp and $2p$ production cross sections should therefore be smaller than the cross sections predicted from the first and second term in Eq. (1). Another source for the need of a renormalization of the theoretical cross sections comes from the fact that the measurements of the coincidence cross sections are made with a certain finite *acceptance*

angle and energy. (See Refs. 3 and 4 for the details.) In comparing the calculated cross sections with the measured ones, we have to renormalize the former appropriately. This is done by the normalization factors in Fig. 8.

In contrast to the πp and $2p$ coincidence cross sections, the normalization factor 1.67 for the coherent π production cross section is larger than unity. This may reflect the fact that the experimental cross section includes not only the contribution from real coherent π production events, but also contributions from the qf decay processes in which only the π is detected, but the proton is not detected because of the “acceptance” problem of the detection system. The experimental cross section may also include contributions from the qf $\Delta \rightarrow \pi + n$ process. Considering all these factors, the fit obtained in Fig. 8 with the assumed normalization factors seems to be quite significant.

The result of the calculations has indicated that the qf πp and the conversion $2p$ events are dominated by the spin-TR channel, while the coherent π events are dominated by the spin-LO channel. Thus coherent π production events are a good source of information on the π correlation. A more detailed analysis of the data, i.e., the analyses of the energy and angular distributions of the emitted π , would be both interesting and important.

-
1. T. Udagawa, F. Osterfeld, and P. Oltmanns, Proceeding of the LAMPF Workshop on *N-N and N-Nucleus Scattering* held at Los Alamos, to be published. (See also 4-3.)
 15. T. Udagawa, F. Osterfeld, P. Oltmanns, and S.-W. Hong, *Proc. of the 6-th International Conference on Nuclear Reaction Mechanisms*, eds. by E. Gadioli (Universita Deglistudi di Milano, Milano, 1991) p.632. (See also 3-20.)
 3. T. Hennino *et al.*, Phys. Lett. B **283**, 42 (1992)
 4. J. Chiba *et al.*, Phys. Rev. Lett. **67**, 1982 (1991).

2.2.3 Spin-Observables

Another source of information on the π correlations discussed in the previous subsection is in the spin transfer coefficients C_{XX} and C_{ZZ} , since these quantities are quite sensitive to the

correlation. Defining the spin-LO and spin-TR strength functions, R_L and R_T , respectively, as

$$R_i = \text{Im}[-\langle \rho_i | G | \rho_i \rangle / \pi] \quad (i = L \text{ or } T), \quad (1)$$

where

$$|\rho_i\rangle = \hat{O}_i |\varphi_A\rangle, \quad (2)$$

with

$$\hat{O}_L = e^{i\vec{q}\cdot\vec{r}}(\vec{S} \cdot \hat{q})\hat{T}_-, \quad (3)$$

and

$$\hat{O}_T = e^{i\vec{q}\cdot\vec{r}}(\vec{S} \times \hat{q})\hat{T}_-, \quad (4)$$

the spin transfer coefficients can be evaluated as

$$C_{XX} = \frac{-|R_L|^2}{|R_L|^2 + 2|R_T|^2}, \quad (5)$$

$$C_{ZZ} = \frac{|R_L|^2 - 2|R_T|^2}{|R_L|^2 + 2|R_T|^2}. \quad (6)$$

From these equations, the sensitivity of the spin transfer coefficients to the π correlations in R_L is obvious.

In Fig. 9, we present the calculated spin transfer coefficients C_{XX} and C_{ZZ} for the $p(\vec{p}, \vec{n})\Delta^{++}$ reaction at the proton incident energy of 800 MeV in comparison with the data.¹ These data are found to be well described by our calculations using the $t_{NN,N\Delta}$ -operator given by Eq. (5). This confirms that the spin structure of the $t_{NN,N\Delta}$ -operator used in our study² of the medium effects on Δ is realistic.

The calculated spin transfer coefficients for the 800 MeV $^{12}\text{C}(p,n)$ reaction are shown in Fig. 10. Here the observables C_{XX} and C_{ZZ} calculated with inclusion of ΔN^{-1} correlations (full curves) are compared with the uncorrelated results (broken curves). A large effect due to the correlations is seen in both quantities. It is thus highly desirable to measure the coefficients C_{XX} and C_{ZZ} .

We remark that at Saturne in Saclay an experiment with similar motivation for seeing the spin-LO correlation effect in nuclei was performed. In this experiment the tensor analyzing power in the $^{12}\text{C}(\vec{d}, 2p)$ reaction at $E_{lab}=2$ GeV was measured. We have analyzed these data,

finding that values measured at $3-6^\circ$ could be explained fairly well by the calculations, but the data at the extreme forward angles (0° data particularly) were not reproduced very well, particularly in the excitation energy region where the pion correlations play an important role, i.e., at the lower excitation energy region. In Fig. 11, we present the results of these calculations, including the fit to the inclusive cross sections. We believe that the problems we find in describing the data at forward angles and lower excitation energies are due to cross section contributions from other reaction mechanisms, e.g., one- and two-step nucleon-knockout and various other two-step mechanisms. The fact that both tensor analyzing power and inclusive cross section data consistently defy explanation seems to support our view.

Just as in the $(\vec{d}, 2p)$ reaction, contributions from different reaction mechanisms other than one-step Δ excitation might also contribute to the (p,n) reaction. It seems likely, though, that these processes should have, if any at all, less effect in the (p,n) than in the $(\vec{d}, 2p)$ reaction. This is particularly true for the lower ω energy side of the Δ resonance, as the observed (p,n) spectra seem to show that the reaction is dominated here by the one-step Δ excitation process. The work described above was presented³ at the Telluride Conference held on 11-15, March 1991.

-
1. G. Glass et al., Phys. Lett. **B129**, 27 (1983).
 2. T. Udagawa, S. W. Hong and F. Osterfeld, Phys. Lett. **B245**, 1 (1990). (See also 3-14.)
 3. T. Udagawa, P. Oltmanns, and F. Osterfeld, *Spin and Isospin in Nuclear Interactions*, eds. by S. W. Wissink, C. D. Goodman, and G. E. Walker (Plenum Press, New York, 1991), p. 111. (See also 3-19.)

2.3 Nuclear Structure

We have proposed recently a new approach to continuum RPA calculations. In this approach, we construct a coupled-channel (CC) integral equation for a set of radial wave functions $\lambda_{ph}(r)$ for the particle p excited from a hole orbit h . We call $\lambda_{ph}(r)$ the correlated source functions. An important feature of $\lambda_{ph}(r)$ is that they are localized. We are thus able to solve the CC integral equation very efficiently by applying the Lanczos method. For solving the CC equation for the 2^+ channel in ^{208}Pb , which has a dimension of 149, for instance, we needed only 10 iterations to reach convergence. This means that we were able to reduce the computation time by at least an order of magnitude. The method has thus enabled us to perform such studies of the nuclear medium effects on Δ as described in 2.2.

Another advantage of the method is that it enables us to calculate contributions from particle-hole (ph) components involved in the total strength function. This means that with this method we are able to calculate the particle emission decay probabilities and also a variety of coincidence cross sections. In fact, we have successfully applied the method to calculate the particle decay widths for the giant monopole response states. In what follows, we describe the work done on the application to the giant resonances and improvement of the method.

2.3.1 Random-Phase-Approximation Calculations of Nuclear Response in the Continuum

In this subsection, we first present the results of the application to the 2^+ strength function, S , in the giant quadrupole resonance (GQR) region.¹ In Ref. 1, the formulation of the method of the calculation has been given. As already noted in the introductory remark, the essence of the approach is the use of the Lanczos method.²

Fig. 12 shows the calculated distributions $F(E)$ of the energy-weighted quadrupole transition strength in unit of the total sum rule, i.e.,

$$F(E) = [ES(E) / \int_0^\infty E' S(E') dE'] \times 100. \quad (1)$$

The corresponding experimental data are also presented by a histogram with a 1 MeV bin. These experimental data are those deduced from hadronic inelastic scattering experiments.³⁻⁴

As seen, the calculations reproduce the observed distributions fairly well. The calculated peak energies (E_R) and widths (Γ) of the GQR are $E_R=21.2, 16.8$ and 11.1 MeV and $\Gamma=6.4, 4.4$ and 1.8 MeV for ^{16}O , ^{40}Ca and ^{208}Pb , respectively. These are compared with the experimental values of $E_{R,exp}=21, 18$ and 11 MeV, and $\Gamma_{exp}=7.5, 4.0$ and 2.0 MeV, respectively.

The above calculations included effects of the damping of the excited particle by means of the imaginary part included in its mean field, but ignored the effects for the hole states as well as the interference between the particle and hole states. In spite of this, the calculation could reproduce fairly well the observed width Γ_{exp} . This suggests that the neglected effects are small as expected from the fact that the dominant contribution to S comes from ph components, whose hole states are in the last occupied shell, and that the contribution to the spreading from such hole states may be small. It might also be possible that this is achieved because the two contributions neglected are cancelling one another.⁵

The calculations predicted S^I/S at the peak energy to be 12%, 8% and 5%, respectively, for ^{16}O , ^{40}Ca and ^{208}Pb . This result shows that the decay should be dominated by the damping, which agrees with the data taken so far, although there are no data which would permit a quantitative comparison with the calculated results.

The calculations reported above were all done by assuming the mass multipole field as the external force. We recently extended the calculations to use a more realistic force, i.e., the M3Y force of Bertsch *et al.*⁶ for (p,p') reactions. With the M3Y force, the source function was calculated by means of DWIA. Such calculations were reported at two international meetings.^{7,8}

-
1. T. Udagawa and B. T. Kim, Phys. Rev. C **40**, 2271 (1989). (See also 3-21.)
 2. R. R. Whitehead, A. Watt, B. J. Cole and I. Morrison, *Advances in Nuclear Physics*, Vol.9, p.123 (1977).
 3. K. T. Knoopfle et al., Phys. Rev. Lett. **35**, 779 (1975).
 4. Y.-W. Lui et al., Phys. Rev. C **24**, 884 (1981).
 5. J. Lisantti et al., to be published.
 6. G. F. Bertsch, P. F. Bortignon and R. A. Broglia, Rev. Mod. Phys. **55**, 287 (1983).
 7. G. Co, K. F. Quader, R. D. Smith, and J. Wambach, Nucl. Phys. **A485**, 61 (1988).
 8. C. F. Bertsch, J. Borysowicz, H. Mcmanus and W. G. Love, Nucl. Phys. **A284**, 399 (1977).

7. B. T. Kim and T. Udagawa, *Proceedings of the 5th French-Japanese Symposium on Nuclear Physics*, ed. by K. Shimizu and O. Hashimoto, p.270 (1989). (See also 3-23.)
8. B. T. Kim and T. Udagawa, *Chin. Jour. of Phys.* **29**, 431 (1991). (See also 3-24.)

2.3.2 Continuum Random-Phase-Approximation Study of Direct Decay of Giant Monopole Resonances in ^{90}Zr and ^{208}Pb

As a continuation, such continuum RPA calculations as described in 2.3.1 were made¹ of particle emission widths from the giant monopole resonances (GMR) in ^{90}Zr and ^{208}Pb . As emphasized in the introductory remarks, our method can retain the particle-hole (ph) information, which makes it possible to evaluate contributions from various particle-hole (ph) components. In fact, the strength function $S(E)$ can be decomposed into a sum of the ph components $S_{ph}(E)$, as

$$S = \sum_{ph} S_{ph}. \quad (1)$$

S_{ph} can further be decomposed into two parts S_{ph}^I and S_{ph}^D , i.e., the damping (spreading) and direct particle decay parts, respectively. In terms of S_{ph}^I , the partial width Γ_h^I of the GMR may be expressed as

$$\Gamma_h^I = \Gamma \int_{E_R - \Delta E}^{E_R + \Delta E} \sum_p S_{ph}^I(E) dE / \int_{E_R - \Delta E}^{E_R + \Delta E} S(E) dE, \quad (2)$$

where E_R and Γ are the energy and the width of the resonance, respectively. The values of E_R and Γ may be extracted from numerically calculated spectra of $S(E)$. ΔE defines the range of the integration involved, for which we choose as $\Delta E=2$ MeV. With this choice for ΔE chosen value, the integration in Eq. (2) can cover the entire region of the resonance.

Numerical calculations were performed² for S , S_{ph}^I and Γ_h^I with two different ph interactions, i.e., those of Shlomo and Bertsch³ (SB) and of Ring and Speth (RS).⁴ In all the calculations, the strengths of the ph interaction were adjusted so as to reproduce the observed resonance energy E_R .

Fig.13 shows the calculated distributions $F(E)$ of the energy-weighted monopole transition strength defined as Eq. (1) of 2.3.1. The corresponding experimental data, deduced

Table 1: Calculated and experimental Γ_h^I -values

Nucleus	hole	Γ_h^{Iexp} (keV)	Γ_h^{ISB} (keV)	Γ_h^{IRS} (keV)
^{208}Pb	$p_{1/2}$	-	5	19
	$f_{5/2}$	<35	10	63
	$p_{3/2}$	75	16	43
	$i_{13/2}$	75	8	18
	$f_{7/2}$	<140	20	102
	$h_{9/2}$	-	6	1
^{90}Zr	$g_{9/2}$	150	45	56
	$p_{1/2}$		20	59

from the measured α -particle inelastic scattering cross sections^{5,6} are also presented there by histograms with a 1 MeV bin.

As seen, the calculated $F(E)$ fits the experimental data fairly well. From the calculated spectra, one can deduce the widths of $\Gamma^{cal}=3.0$ and 2.8 MeV for ^{90}Zr and ^{208}Pb , respectively. These values agree very well with the experimental values of $\Gamma^{exp}=3.0$ and 2.5 MeV.^{5,6} It should be remarked at this stage that the calculated F 's with two different ph interaction are almost identical. This is not the case, however, for Γ_h^I , as will be seen below.

In Table 1, we summarize the calculated and experimental values of Γ_h^I for neutron emission from various hole states h in ^{207}Pb and ^{89}Zr . Γ_h^{ISB} and Γ_h^{IRS} shown there are those obtained by using the SB and RS ph interactions, respectively. Although a large effort has been made in extracting the experimental values, the best values available now, particularly for ^{208}Pb , are still upper limit values. This results from the fact that to extract these values from the measured (coincidence cross section) data is extremely difficult, because in doing so one has to subtract from the measured data large background and statistical components, neither of which are very well known. The values listed in Table 1 are those determined by assuming the lower limit of the statistical components, and therefore should be viewed as an upper limit.

As seen in Table 1, the two theoretical values are fairly different from each other, indicating the sensitivity of Γ_h^{cal} to the ph interaction. Such a sensitivity was noticed earlier in Ref. 9.

The result that the theoretical calculations are sensitive to the ph interaction implies that we may use the data for Γ_h^{exp} in testing the ph interaction. At this moment, however, it is not possible to make such a test, since the data available now are only the upper limit values. It is thus highly desirable that more accurate data be taken. Note that Γ_h^{cal} is generally smaller than Γ_h^{exp} . This is particularly true for the case for Γ_h^{SB} ; Γ_h^{SB} is smaller by a factor of 4-8 in ^{208}Pb and 3 in ^{90}Zr than is Γ_h^{exp} . The situation is better in Γ_h^{RS} , but still underestimates Γ_h^{exp} (except the $f_{7/2}$ case).

In the study reported above, we included the damping of the excited particles, but not of the holes. We also ignored the interference effect, whose importance has been indicated in recent studies.¹⁰ The neglect of the damping of the hole states may be justified, since the holes of the main ph-components in GMR are those in the uppermost occupied shells, whose spreading width (damping) is rather small.

Microscopic calculations done in the past have indicated the importance of interference effects, but such calculations still significantly underestimate the width of the GMR.¹⁰ One possible source may be traced to the fact that the calculations introduce a rather dramatic approximation to replace real final nuclear states by pure 2p-2h states, into which the collective monopole state damps. Such an approximation leads to a large interference effect, which cancels out contributions from the damping of the particle and the hole. It is highly probable that the interference effects will be reduced if the approximation is removed; a strong mixing of 2p-2h and more complicated states with a random phase nature will then be induced, which in turn causes cancellations between contributions from various 2p-2h components, thereby reducing the net contribution. At this moment, it is very difficult to confirm what was argued above by carrying out a realistic calculation. In the present study, we have adopted a phenomenological approach, neglecting the effect of the hole damping and the interference for the reason, discussed above, but treating the effect of the damping of the excited particle in a realistic manner. The successes obtained in the present study and also in our previous study² support our view, though much has to be done before reaching a more concrete understanding of the problem.

The work described in this and the previous subsections was summarized at the international workshop on spectroscopy and structure of molecules and nuclei, held at Tallahassee,

March, 1992.¹¹

-
1. T. Udagawa and B. T. Kim, Phys. Lett. B **230**, 6 (1989). (See also 3-25.)
 2. T. Udagawa and B. T. Kim, Phys. Rev. C **40**, 2271 (1989). (See also 3-22.)
 3. S. Shlomo and G. F. Bertsch, Nucl. Phys. **A243**, 507 (1975).
 4. P. Ring and J. Speth, Phys. Lett. **B44**, 477 (1973).
 5. S. Brandenburg *et al.*, Nucl. Phys. **A466**, 29 (1987).
 6. C. M. Rozen, D. H. Youngblood, J. D. Bronson, Y.-W. Lui, and U. Garg, Phys. Rev. C **21**, 1252 (1980).
 7. Wim Borghols, Ph. D. Thesis (Groningen, 1988). (Unpublished)
 8. S. Brandenburg *et al.*, to be published in Phys. Rev. C **39**, 2448 (1989).
 9. A. Bracco, J. R. Beene, N. Van Giai, P. F. Bortignon, F. Zardi and R. A. Broglia, Phys. Rev. Lett. **60**, 2603 (1988).
 10. See, for instance, the following review articles;
G. F. Bertsch, P. F. Bortignon, and R. A. Broglia, Rev. Mod. Phys. **55**, 287 (1983);
J. Wambach, Rep. Prog. Phys. **51**, 989 (1988).
 11. T. Udagawa, to be published. (See 4-4.)

2.3.3 Random Phase Approximation Calculations of the Nuclear Response in the Continuum Using a Nuclear Matter G-Matrix

Most calculations of the nuclear response which use the continuum random phase approximation (CRPA), which is the subject of our work described in the previous subsections 2.3.1 and 2.3.2, have thus far used a δ -force, for instance the Skyrme force, for the residual particle-hole(ph) interaction, since zero-range interactions greatly simplify the numerical calculations. It is natural, however, to try to include realistic finite range residual interactions in CRPA calculations for the same reasons that extensive efforts have been made to use realistic effective interactions in calculating the properties of the low-lying collective states of nuclei. An effort¹ has thus recently been devoted to extend our CRPA approach² to handle the case where in addition to the direct part, the exchange part of the effective ph interaction is finite range in character. Shigehara, *et al.*³, recently computed the spin-isospin response in the quasielastic region using a nuclear matter finite-range G-matrix, including a full finite range treatment for both the direct and exchange contributions to the ph residual interaction. A limitation to their formalism is that because they directly compute the nuclear

response function, as opposed to solving the CRPA equations, they can only calculate the singles cross-section of the scattered particle.

We first applied the method to the lowest 3^- state for ^{16}O . This case is interesting since it is well known that a discrete RPA calculation with the G-matrix results in an excitation energy and $B(E3)$ value that agrees nicely with the experimental data⁴. We found that the use of the G-matrix in the finite range CRPA calculation caused the lowest 3^- state to become unstable. In order to obtain a stable RPA result with an excitation energy near the experimental energy (6.13 MeV) and $B(E3)$ value ($1470\text{ e}^2\text{fm}^6$) we found it necessary to reduce the overall strength of the G-matrix by 19% for the exchange part. This suggests that the coupling to the continuum plays a very important role. In addition we also applied the method to the 3^- in ^{40}Ca , finding it necessary to reduce the ph strength for the exchange term by 37%.

We then applied the method to calculate the 2^+ quadrupole strength functions for ^{16}O and ^{40}Ca in the giant resonance region. We used the same renormalized ph interaction that was obtained in the 3^- calculation for each nucleus. The necessity to introduce a renormalized ph interaction is consistent with the observation by previous researchers that using only the standard RPA in a discrete basis with the G-matrix for the low-lying collective states is inadequate because the ph correlations become unphysically large and requires that either one modifies the effective interaction(say by the addition of screening terms) or go to higher RPA⁵⁻⁶.

We studied the validity of the zero-range approximation of the exchange matrix element; use of the approximation pushed the quadrupole strength far below 10 MeV. As pointed out by Shigehara, *et al.*⁴, the exchange part of the Nakayama nuclear matter G-matrix is strongly momentum dependent, particularly for the $T=S=0$ channel, thus the delta-function approximation is poor. The final remark is that the ratio of S^1 to S at the peak resonance energy is .26 and .11 for ^{16}O and ^{40}Ca respectively. Comparing these results to those obtained in Ref. 2 (.12 and .08), we observe a significant difference for the ^{16}O case. The large emission strength for ^{16}O is probably due to the peak resonance occurring at an excitation energy 2.5 MeV greater than the peak resonance energy obtained in Ref. 2, although this point needs further clarification.

-
1. D. Knobles and T. Udagawa, Nucl. Phys. **A533**, 189 (1991) (See also 3-26.)
 2. T. Udagawa and B.T. Kim, Phys. Rev. C **40**, 271 (1989).
 3. T. Shigehara, K. Shimizu, and A. Arima, Nucl. Phys. **A492**, 388 (1989).
 4. K. Nakayama, S. Krewald, J. Speth, and W.G. Love, Nucl. Phys. **A431**, 419 (1984).
 5. M.G.E. Brand, K. Allaart, and W.H. Dickhoff, Phys. Lett. B **214**, 483 (1988).
 6. S. Drozd, S. Nishizaki, J. Speth, and J. Wambach, To be published.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 12

Fig. 13

*reprints + preprints
removed,
ds*

List of Papers (October 1991 - September 1992)

“Random-Phase Approximation Calculations of the Nuclear Response in the Continuum Using a Finite-Range Interaction”, D. Knobles and T. Udagawa, Nucl. Phys. **A533**, 189 (1991).

“Dynamical Fusion and Direct Reaction Polarization Potentials for Heavy-Ion Scattering and Fusion at Sub-Coulomb-Barrier Energies”, B. T. Kim and T. Udagawa, Phys. Lett. B **273**, 37 (1991).

“RPA Calculations of Nuclear Response in the Continuum for the Hadronic Inelastic Scattering”, B. T. Kim and T. Udagawa, Chinese Journal of Physics, **29**, 431 (1991).

“Delta Excitations in Nuclei and Its Decay”, T. Udagawa, F. Osterfeld, P. Oltmanns, and S. W. Hong, Proceedings of 6-th International Conference on *Nuclear Reaction Mechanisms*, ed. by E. Gadioli (Universita Delistudi di Milano, Milano, 1991), p.632.

“Systematics of Sub- and Near-Barrier Fusion Cross Sections and the Threshold Anomaly”, T. Udagawa, M. Naito, and B. T. Kim, Phys. Rev. C **45**, 876 (1992).

“ Δ -Excitations in Nuclei”, F. Osterfeld, B. Krfgn, P. Oltmanns, and T. Udagawa, Proceedings of the Workshop on *Meson Production, Interaction and Decay*, ed. by A. Magiera, W. Oelert, and E. Grosse (World Scientific, 1991), p.116.

“ Δ Excitations in Nuclei”, T. Udagawa, P. Oltmanns, and F. Osterfeld, Proceedings of the International Conference on *Spin and Isospin in Nuclear Interactions*, ed. by S. W. Wissink, C. D. Goodman, and G. E. Walker (Plenum Press, New York, 1991), p. 69.

“A Unified Approach for Low-Energy Nuclear Reactions”, B. T. Kim and T. Udagawa, Proceedings of International Conference on *Towards a Unified Picture of Nuclear Dynamics*, ed. by Y. Abe, S. M. Lee, and F. Sakata, (American Institute of Physics, New York, 1991), p.133.

“Breakup-Fusion Analyses of the $^{40}\text{Ca}(^6\text{Li},d)^{44}\text{Ti}$ Reactions and α -Cluster Structure in ^{44}Ti ”, Chong-Yeal Kim and T. Udagawa, Phys. Rev. C **46**, 532 (1992).

5. PERSONNEL

Faculty

Dr. Takeshi Udagawa, Professor

Post-Doctoral Appointments

Dr. D. Knobles (- February, 1992)

Dr. C. Y. Kim (January, 1992 -)

Pre-Doctoral Appointments and Graduate Students

K.-P. Cheng (Fully supported by Chinese government, - August, 1990)

A. Gomez-Chamacho (- August, 1991)

M. Naito (- August, 1992)

C.-Y. Kim (- December, 1991)

J. Young (December, 1991 -)

Visitors

Professor S. W. Hong (July - August, 1991 and July, 1992); (Department of Physics,
Sung Kyung Kwan University, Seoul, Korea)

Professor B. T. Kim (December, 1989 - December 1990 and July - August, 1991);
(Department of Physics, Sung Kyung Kwan University, Seoul, Korea)

Dr. R. C. Mastroleo (December, 1989 - February, 1990 and September, 1991 -
August, 1992);(Instituto de Estudos Avancados, Sao Jose dos Campos, Brazil)

END

**DATE
FILMED**

12 / 14 / 92