DOE/ER/40343--6

DE93 004375

Investigation of Pion-Nucleus Interactions

Progress Report

April 1, 1991 — March 31, 1992

C. Fred Moore
Department of Physics
University of Texas at Austin
Austin, Texas 78712-1081

RECEIVED OSTI

DEC 14 1992

September 1992

PREPARED FOR THE U. S. DEPARTMENT OF ENERGY
UNDER GRANT NUMBER DE-FG05-87ER40343

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Contents

1	Introduction	2
2	Overview	3
3	Double Giant Resonances in Pion Double Charge Exchange on ^{51}V, ^{115}In, and ^{197}Au	5
3.1	Abstract	5
3.2	Introduction	6
3.3	Experiment	6
3.4	Results and Analysis	7
3.4.1	Discussion of the data	7
3.4.2	Energies and widths of the double resonances	13
3.4.3	Splitting of the GDR built on the IAS of ^{51}V	15
3.4.4	Mass dependence of double-resonance cross sections and sum-rule analysis	17
3.5	Summary	21
4	Publications	23
5	Abstracts	26
6	Invited Talks	27
7	Proposal Defenses	28
8	Approved New Research Proposals	29
9	Personnel	30

List of Tables

1	Target compositions	7
2	DCX results	13
3	Intensity ratios	14
4	Giant-dipole energies	14
5	Giant dipole resonances built on the isobaric analog state	17
6	CCIA calculations	20

List of Figures

1	Doubly differential cross-section spectrum for ^{51}V	8

2	Doubly differential cross-section spectrum for ^{115}In	9
3	Angular distributions	10
4	Angular distributions for the double dipole	11
5	Missing mass spectrum	16
6	Cross sections	18
7	Double dipole cross sections	19

1 Introduction

This report summarizes the work carried out by personnel from the University of Texas at Austin at the Los Alamos Clinton P. Anderson Meson Physics Facility (LAMPF) during the calendar years 1991-1992 under grant DE-FG05-87ER40343 between the University of Texas at Austin and the United States Department of Energy.

One of our faculty and a research faculty member has been on site (at LAMPF) during five months or more of each year. An average of eight graduate students have been associated with this program, most being in residence at LAMPF.

The research activities involved experiments done with the Energetic Pion Channel and Spectrometer (EPICS), the Low Energy Pion Channel (LEP), the Pion and Particle Physics Channel (P^3), the High Resolution Spectrometer (HRS), and planning a new experimental program associated with the new high-resolution Neutral Meson Spectrometer (NMS) at LAMPF.

A brief overview of work supported by this grant is given in Section 2. An account of the study (Experiment #1154) of the double giant resonances in pion double charge exchange on ^{51}V , ^{115}In , and ^{197}Au forms Section 3. This report contains a list of published papers and preprints in Section 4, abstracts in Section 5, and invited talks in Section 6. These papers summarize experiments involving participants supported by this grant and indicate the work accomplished by these participants in this program of medium energy nuclear physics research. Section 7 contains a list of the most recent proposals on which we have participation at LAMPF, while Section 8 lists proposals which have been approved this past year to run as experiments. Section 9 contains a list of personnel who have participated in this research program.

The research cited in this report is, in many cases, the collaborative effort of many groups associated with research at LAMPF. Collaboration with the Los Alamos National Laboratory, the University of Minnesota, the University of Colorado, the Louisiana State University, the Ben Gurion University of the Negev, the King Fahd University of Petroleum and Minerals, the Lawrence Livermore Laboratory, the New Mexico State University, Argonne National Laboratory, the University of Colorado, and the University of Pennsylvania is acknowledged.

One Ph.D. degree was granted by the University of Texas at Austin during 1992, based on experimental research done at LAMPF, bringing the total number of Ph.D.'s granted to Texas students supported by this grant in meson-nuclear physics to twelve.

2 Overview

Our program concentrates on pion physics experimental results obtained using the Energetic Pion Channel and Spectrometer (EPICS), Pion and Particle Physics channel (P³), and the Low Energy Pion physics channel (LEP). These facilities are unique in the world in their intensity and resolution. Two classes of experiments can be done best with this equipment: scattering (elastic and inelastic) and double charge exchange (DCX). Several coincidence experiments are in progress and several are proposed.

Inelastic pion scattering proposal #1025, which has been approved with A⁺ priority, deals with measurements of pion scattering from a polarized ¹³C target. The asymmetry parameter A_y for $\pi^\pm$ elastic and inelastic scattering has been measured over an angular range of 30° to 100° from a polarized ¹³C target at EPICS. The elastic-scattering results should be sensitive to spin-dependent effects in the pion-nucleus interaction, such as the Δ -nucleus spin-orbit interaction. The inelastic-scattering results will be sensitive to the relative mixture of spin-flip and non-spin-flip contributions to the transitions. These polarized target experiments will give new information about the pion-nucleus interaction that is not available from experiments with unpolarized targets, because the former measures the interference between amplitudes and the latter measures squares of amplitudes. The target system was assembled principally from polarized-target elements available at LAMPF. The experimental beam time requested is one running cycle, taken to be 1,200 hours.

In experiment #1104 cross-sections to the GQR for the nuclei ⁹⁰Zr and ²⁰⁸Pb have been measured with π^+ and π^- at energies from 350 to 550 MeV, *i.e.*, above the $\Delta_{3/2,3/2}$ resonance region. The goal of this experiment is more accurate evaluation of neutron and proton transition amplitudes (M_n and M_p) extracted for GQR in those neutron excess nuclei. Those values of M_n and M_p obtained near the $\Delta_{3/2,3/2}$ resonance region have uncertainties in the wave functions involved and the shapes of the neutron and proton transition densities at the nuclear surface due to the short mean free path of the pion near the Δ resonance. The longer mean free path of the pion in these higher energy region may help to eliminate those uncertainties in extraction of the matrix elements.

Regarding proposal #1018, we have analyzed large-angle pion elastic data on ¹²C and ¹⁶O with standard first-order optical models using matter distributions deduced from electron scattering. We show that these models reproduce only the qualitative features of the data. However, much better agreement can be achieved with the same models provided the matter distributions are modified in a suitable way. At this stage it is not clear whether these corrections are applicable to all nuclei or whether they apply at energies other than those on resonance. To address these questions we propose to measure elastic excitation functions (at 175°) on ⁶Li, ²⁸Si, and ⁵⁸Ni in the energy range from 100 to 250 MeV.

Pion absorption: The LEP and P³ channels at LAMPF will be used. In the low energy pion channel, we propose to measure the cross sections for π^+ and π^- absorption on ¹²C, ⁵⁸Ni, and ²⁰⁸Pb in which the final state includes two or more free protons. Data will be taken for π energies of 50, 100, 150, and 200 MeV. The energy and angular distributions of the outgoing protons are measured using a large solid-angle bismuth germanate BGO detector

described in the section of last year's proposal entitled "BGO BALL".

In the pion particle physics (P^3) channel, we measure the cross sections for π^+ and π^- absorption on ^{12}C , ^{58}Ni , and ^{208}Pb in which the final state includes two or more free protons. Data are taken for π energies of 200, 300, 400, and 500 MeV. The energy and angular distributions of the outgoing protons are measured using a large solid-angle BGO detector.

Among the most exciting results obtained recently at LAMPF in pion physics is the observation of resolved giant-dipole resonances built on the isobaric analog states and the isovector giant dipole resonance built on another giant dipole. The isobaric analog, and the giant dipole were both observed in (π^+, π^-) single charge exchange reaction (with a typical maximum cross section around $0.5 \rightarrow 1.0$ mb/sr). Therefore a combination of these resonances could, in principle, be reached in double charge exchange. However, until recently, none of these double resonances had been found.

The results obtained so far are exciting and have a potential of new and interesting physics. Experiments #1050 and #1118 have demonstrated that pion DCX provides a unique reaction in which one can study the existence of double giant resonances in nuclei. The results indicate that two new different modes of nuclear excitations are observable in pion double charge exchange reactions. These are the giant dipole built on the isobaric analog state and the giant dipole resonance built on the giant dipole resonance. Due to the selectivity of the DCX reaction these exotic states are observable although they are located at high excitation energy where background from DCX to the continuum is large. The data show that the newly discovered resonances are general collective features of all nuclei for the double-dipole, and of all nuclei with $N-Z \geq 1$ for the GDR built on the IAS. In experiment #1118 at the energetic pion channel and spectrometer (EPICS) at LAMPF, we observed a clear splitting of the giant dipole built on the isobaric analog state ($\text{GDR} \otimes \text{IAS}$) in the $^{93}\text{Nb}(\pi^+, \pi^-)^{93}\text{Tc}$ reaction at $T_\pi = 295$ MeV. The results obtained in the first two research years of this program have been published (or submitted) in several scientific reports and invited talks at international conferences.

Experiment #1154 (which ran in two phases) was designed to accurately measure the width of the giant dipole built on the isobaric analog state on a wide range of mass and with higher statistics than previously obtained. The results indicate that the width of the GDR built on the IAS increases significantly with mass. For example, the resonance for ^{13}C has a width of 3.0 ± 0.6 MeV, but that for ^{59}Co has a width of 7.0 ± 1.0 MeV (more than twice as wide as for ^{13}C). The increase of the width with mass is not yet fully understood, and is one of the puzzles arising from this study. An interesting implication was also discovered during this study. The resonance on the $T = 1/2$ targets has a single isospin value ($T = 3/2$) and is the analog of the $T_>$ GDR state in the target nucleus. Thus pion DCX provides a new general method for an accurate determination of the $T_> = T + 1$ giant dipole state on any $T = 1/2$ nucleus available as a target for DCX measurements.

Multiparticle final state in pion interactions: We studied the decay protons in the reaction $(\pi, \pi' p)$ as a test of charge symmetry. Comparisons of photo proton and photo neutron breakup of ^4He have provided evidence for a large violation of charge symmetry, caused by forces other than the Coulomb force. However, pion inelastic scattering data show that the deviation from charge symmetry can be explained by the Coulomb potential only. In an

attempt to resolve this argument on the existence of charge symmetry breaking forces other than Coulomb interaction, a series of $(\pi, \pi' p)$ coincidence experiments on ^4He (#998), ^{12}C and ^{208}Pb (#1039) were conducted during the 1986-91 time period. The conclusion of these experiments is that there is no other force than Coulomb that breaks up charge symmetry, and the discrepancy between the result using electromagnetic probes and that using pions can be explained by direct-knock-out and semi-direct components, using the widely used Direct-Semidirect Model or Doorway-state Model.

Experiment #1103 aims to measure the ratio of $(\pi^+, \pi^{+'})$ vs $(\pi^-, \pi^{-'})$ on various nuclei to provide information on proton and neutron transition densities at energies around the Roper resonance where the relative strength of pure proton and pure neutron transitions should be reversed as compared to the $\Delta_{3/2,3/2}(1232)$ resonance case.

Experiment #1210 ran recently at LEP (July 1991). The experiment aimed to study the mechanism for DCX to excite analogs of anti-analog states in heavy nuclei. A previous, high statistics DCX experiment on ^{93}Nb showed for the first time the existence of a resonance at high excitation energy below the double isobaric analog state. The observation of the resonance with such a large cross section was unexpected. In a recent article we suggested a few possible identifications for this resonance. Experiment #1210 confirms the previous observation on ^{93}Nb and gives new results on ^{56}Fe and ^{44}Ca at a lower beam energy. The results are still under study, and additional beam time was approved for this experiment by the PAC.

Experiment #1209, run recently at EPICS, is designed to measure the energy dependence of the double giant resonance on ^{13}C and ^{51}V . The measurements provide the first attempt to determine the excitation function of a double resonance state. The data is currently being replayed for a subsequent analysis, however, on-line results show that the cross section of the giant dipole built on the isobaric analog state on ^{13}C and ^{51}V increases with energy as expected from simple sequential model calculations. The results indicate that pion double charge exchange involving the excitation of the closed-shell core proceeds predominantly by a two-step sequential $(\pi^+, \pi^0)(\pi^0, \pi^-)$ mechanism, and therefore should be a unique probe for exciting isotensor double giant resonances in nuclei.

In Section 3 we discuss in detail the results from Experiment #1154 which ran at EPICS in Sep-Oct 1990.

3 Double Giant Resonances in Pion Double Charge Exchange on ^{51}V , ^{115}In , and ^{197}Au

3.1 Abstract

Resonances were observed in the continuum in pion induced double charge exchange (at $T_\pi=295$ MeV) on ^{51}V at $Q=-14.2$, -32.9 , and -48.1 MeV, on ^{115}In at $Q=-25.2$, -37.4 and -49.7 MeV and on ^{197}Au at $Q=-35.8$, -45.1 and -54.9 MeV. The three resonances have different characteristic angular distributions. Partial angular distributions were measured on ^{51}V and ^{115}In and observed to have $J^\pi=0^+$, 1^- and 2^+ shapes respectively. Based on

their centroid energies, angular distributions and observed cross sections we identify the resonances as the double isobaric analog state, the giant dipole built on the isobaric analog and the double giant dipole resonance. The new observations are in close agreement with the gross features reported recently for the double giant resonances, and provide some of the best signals obtained to date for the double resonance states in nuclei.

3.2 Introduction

A long-standing issue in nuclear physics is whether multipole excitations of giant dipole resonances or of other giant resonances exist. If they do exist, then what are their fundamental features (*e.g.* energies, widths, strengths, mass dependence, etc.) and do they confirm broad aspects of nuclear structure theory[1, 2, 3, 4, 5]? In the last decade a large amount of effort in heavy-ion physics[6, 7, 8] has been devoted to the search for the possible existence of multiphonon states. The difficulty in observing double giant resonances arises mainly from the fact that these states are located high in the continuum in a region of very high density of states and also have a large width. In recent works we reported a first observation of resolved giant-dipole resonances built on the isobaric analog states[9] and the double isovector giant dipole state[10] in pion double charge exchange. More recent studies[11, 12] to search for double resonances in nuclei have confirmed the first observations. The data indicate that the giant dipole built on the isobaric analog state (GDR $\otimes$ IAS) is a general collective feature of all nuclei with at least one-neutron excess[11]. The double dipole is a general double-collective excitation of all nuclei[12], and is excited in both the $\Delta T_z = -2$ and the $\Delta T_z = +2$ double charge exchange (DCX) modes[13]. The observed double resonances behave, to some extent, as a superposition of the ingredient “ordinary” resonances excited in a sequential process. However, one of the puzzles arising from these studies is the width of the GDR $\otimes$ IAS which increases rapidly with mass, and does not follow the systematics of the widths of the ordinary single dipole built on the ground state (GS). Deformed nuclei such as ^{51}V and ^{59}Co do not follow a simple pattern and should be treated separately. The width of the double GDR state has been found to be close to 1.5 times the width of the single GDR and is somewhat in disagreement with the simple theoretical predictions that the width of the double GDR, as a two-phonon state of independent phonons, should be twice the GDR width[2]. The present study was aimed to give more information with high statistics on the width of the double resonance states in medium and heavy nuclei.

3.3 Experiment

The measurements were performed with the energetic pion channel and spectrometer (EPICS) at the Clinton P. Anderson Meson Physics Facility (LAMPF) with the standard pion double-charge-exchange setup[14]. Table 1 lists the pertinent information on the targets. Measurements were taken at $T_\pi = 295$ MeV and scattering angles of 5° , 11° and 22.5° for ^{51}V , 5° and 9° for ^{115}In , and at a single angle of 9° for ^{197}Au . Electrons were eliminated using an iso-butane velocity-threshold Cherenkov detector in the focal plane. A scintillator placed behind a series of aluminum and graphite wedges was used to detect and veto muon

Table 1: Target compositions, areal densities, and laboratory angles of measured data.

Target	Isoic purity (%)	Areal density (g/cm ²)	Scattering Angle (lab°)
⁵¹ V	99.75	1.94	5, 11, 22.5°
¹¹⁵ In	95.7	0.928	5, 9°
¹⁹⁷ Au	100.	1.966	9°

events[15]. The remaining background was pions resulting from continuum DCX in the target. The choice of the highest beam energy available at EPICS ($T_\pi=295$ MeV) for the present measurements has several advantages: a) the background level from the continuum is lower since the excitation energy region of interest is pushed away from the inclusive DCX peak, b) a larger outgoing energy range (≈ 50 MeV) is covered by the acceptance of the spectrometer in a single setting, c) cross sections are expected to increase as k^2 or faster, where k is the incident pion momentum, and d) the highest energy was chosen because it is here that sequential processes dominate DCX. This is evidenced by the dominance of the DIAS at $\theta=5^\circ$.

The acceptance of the spectrometer was measured with a ¹²C target using inelastic scattering to the 4.44-MeV state at an energy $T_\pi=180$ MeV and a scattering angle of 35° which corresponds to the peak cross section of the 2^+ state. The spectrometer field was varied to move the 4.44-MeV peak across the focal plane, covering approximately $\pm 10\%$ of the central momentum of the spectrometer. Absolute normalization was obtained using a polyethylene (CH₂) target of areal density 0.025 g/cm² to measure the hydrogen elastic cross section and comparing the yields with cross sections calculated from π -nucleon phase shifts[16].

3.4 Results and Analysis

3.4.1 Discussion of the data

Figure 1 contains the ⁵¹V Q -value spectra at three different angles: 5° , 11° and 22.5° . The spectra exhibit a clear signal of three resonances. The DIAS is observed at $Q = -14.2 \pm 0.1$ MeV ($E_x = 11.26$ MeV) and is most strongly populated at 5° . At larger angles the DIAS is weak, whereas the upper resonances are relatively stronger. The middle resonance is identified as the giant dipole built on the isobaric analog state (GDR $\otimes$ IAS) and the upper as the double giant dipole (GDR $\otimes$ GDR). We will return to the identification of these resonances later. The peaks were fit with a Gaussian shape of variable width for the DIAS and a Lorentzian shape for the giant resonances (GR). In Fig. 1 the fit to the spectra has $\Gamma_{\text{DIAS}}=0.9$ MeV, $\Gamma_{\text{GDR}\otimes\text{IAS}}=8.1$ MeV, and $\Gamma_{\text{GDR}\otimes\text{GDR}}=10.0$ MeV. The width of the DIAS presumably arises solely from resolution energy due to the thick (1.94 g/cm²) ⁵¹V target. The spectra have been corrected for the spectrometer acceptance as a function of momentum. The background (dashed line), which arises from DCX to discrete low-lying states and to the continuum was

FIG1_E1154.GRF

Figure 1: Doubly differential cross-section spectra for the $^{51}\text{V}(\pi^+, \pi^-)^{51}\text{Mn}$ reaction at $T_\pi=295$ MeV and $\theta_{\text{lab}}=5^\circ$, 11° , and 22.5° . The arrows indicate the fitted location of the DIAS, the $\text{GDR} \otimes \text{IAS}$ and the $\text{GDR} \otimes \text{GDR}$. Short vertical lines represent statistical uncertainty of the data. The dashed line is the background fitted with a polynomial shape and the solid line is the fit to the spectrum using NEWFIT.

FIG2_E115.GRF

Figure 2: Same as Fig. 1 except for ^{115}In target, and (a) $\theta_{\text{lab}}=5^\circ$ and (b) $\theta_{\text{lab}}=9^\circ$.

FIG3_E1154.GRF

Figure 3: Angular distributions for the resonances identified as the GDR $\otimes$ IAS in the (a) $^{51}\text{V}(\pi^+, \pi^-)^{51}\text{Mn}$ reaction and (b) $^{115}\text{In}(\pi^+, \pi^-)^{115}\text{Sb}$ reaction at $T_\pi=295$ MeV. The solid and chain-dashed lines are sequential-model calculations for the GDR $\otimes$ IAS using surface and volume transition densities, respectively, without any normalization factor (see text).

Figure 4: (a) Partial angular distribution for the resonance identified as the double dipole in the $^{51}\text{V}(\pi^+, \pi^-)^{51}\text{Mn}$ reaction at $T_\pi=295$ MeV. The curves are sequential-model calculations for the double dipole using NEWCHOP. The dashed curve is the 2^+ distribution arbitrarily normalized to fit the 22.5° data point. The dotted curve is the 0^+ distribution normalized to fit the forward-angle data, and the solid curve is the sum of the 0^+ and 2^+ cross sections. (b) Same as (a) except for the $^{115}\text{In}(\pi^+, \pi^-)^{115}\text{Sb}$ reaction, and a 2^+ curve normalized to the 9° data point.

fit using a third-order polynomial shape of the form $R_0 + R_1E + R_2E^2 + R_3E^3$ where E is the absolute Q value. The solid line is the resulting fit to the spectrum using the computer code NEWFIT[17].

Figure 2 presents the measured DCX spectra on ^{115}In at scattering angles of 5° and 9° . Again, in addition to the narrow transition to the DIAS at $E_x = 23.7$ MeV ($Q = -25.2 \pm 0.1$ MeV), both spectra contain a wider peak labeled GDR $\otimes$ IAS which shows up more clearly at 9° . The higher resonance is the double giant dipole labeled GDR $\otimes$ GDR.

Figure 3 presents the angular distributions extracted for the peaks labeled GDR $\otimes$ IAS on both targets. In ^{51}V the maximum cross section is observed at 11° with a smaller cross section at 5° and 22.5° . The curves in Fig. 3 are simple calculations in which sequential single charge exchange through the isobaric analog state to the giant dipole resonance is evaluated using the pion coupled-channel impulse-approximation (CCIA) code NEWCHOP[18]. The calculations include four states: the GS, the GDR, the IAS, and the GDR $\otimes$ IAS. The strength of each SCX transition has been adjusted to reproduce the measured (or extrapolated) cross section to the GDR and the IAS[19]. A widely accepted collective Tassie form factor for the transition density, $r^{L-1}\rho'(r)\mathcal{Y}_{LM}(\hat{r})$, with $L=1$ was used for the GDR[20], where $\rho(r)$ is the nuclear GS density. For the IAS we used two different types of transition densities. The volume transition density (solid curve) was assumed for simplicity to have the same radial distribution as the GS density but normalized to the neutron excess in the target nucleus:

$$\rho_{\text{volume}} = \rho(r)(N - Z)/A . \quad (1)$$

A surface transition density (chain-dashed curve) obtained by taking a derivative of the GS density distribution was also used. The calculated curves are shown with no additional normalization.

Figure 4 exhibits the angular distributions extracted for the peaks labeled GDR $\otimes$ GDR in Figs. 1 and 2. The calculations are sequential-model calculations for the double dipole via the single GDR using NEWCHOP. A coupling of two $L = 1$ multipoles gives a total angular momentum transfer $\Delta L=0, 1$, or 2 . However from symmetry considerations[2] only $\Delta L=0, 2$ are allowed. This will also be the total J transfer for the double dipole state[12]. The long-dashed line represents the calculations for $J^\pi=0^+$ and the dotted lines for $J^\pi=2^+$. The solid curves are the sum of the $L=0$ and $L=2$ cross sections. The 2^+ curves were arbitrarily normalized to fit the data. The normalization factors for the 0^+ curves were chosen to account for the forward-angle rises of the 5° cross sections when added to the 2^+ curves. Both angular distributions show a forward-angle rise which may indicate a small contribution of the $J^\pi=0^+$ double-dipole state. The figure indicates that the observed double dipole resonances are primarily the $J^\pi=2^+$ double-dipole states, but may include some strength of the unresolved $J^\pi=0^+$ component. A similar feature was observed earlier experimentally on ^{40}Ca and heavier nuclei[12]. The fact that the double dipole excitation favors an angular momentum transfer of $J=2$ rather than $J=0$ seems to be a more general feature of identical two-phonon excitation. In a recent theoretical work[21, 22] of sum rule analysis it was found that the double Gamow-Teller excitation is also dominated by an angular momentum transfer of $J=2$ and the allowed $J=0$ component is weaker than the $J^\pi = 2^+$. It is not yet clear why one should expect this behavior, and whether in fact this is a general feature of multiphonon excitation.

Table 2: Results from the double-charge-exchange reaction on ^{51}V , ^{115}In , ^{197}Au at $T_\pi=295$ MeV.

Target	DIAS	GDR $\otimes$ IAS			GDR $\otimes$ GDR		
	$-Q$ (MeV)	$-Q$ (MeV)	Γ (MeV)	$d\sigma/d\Omega^{peak}$ ($\mu\text{b}/\text{sr}$)	$-Q$ (MeV)	Γ (MeV)	$d\sigma/d\Omega^{peak}$ ($\mu\text{b}/\text{sr}$)
^{51}V	14.2 ± 0.1	32.9 ± 0.2^a	8.1 ± 0.6	3.4 ± 0.2	48.1 ± 0.3	10.0 ± 4.3	6.8 ± 0.5
^{115}In	25.2 ± 0.1	37.4 ± 0.3	4.5 ± 0.7	2.1 ± 0.2	49.7 ± 0.7	8.5 ± 3.2	4.1 ± 0.8
^{197}Au	35.8 ± 0.1^b	45.1 ± 0.8	8.5 ± 1.5	4.0 ± 0.5	54.9 ± 0.4	10.0 ± 3.2	7.7 ± 1.6

^a Fitting the GDR $\otimes$ IAS on ^{51}V with three peaks gives the results listed in Table V.

^b The value for the DIAS is from a previous measurement at $T_\pi=295$ MeV and $\theta_{lab}=5^\circ$, see Ref. [11]. The listed values for the GR are from the present study.

3.4.2 Energies and widths of the double resonances

Table 2 summarizes the experimental results obtained for the DIAS, the giant dipole built on the IAS and the double dipole from the present study. For $T\geq 1$ target nuclei, the $\Delta T_z=-1$ charge exchange GDR, and the $\Delta T_z=-2$ GDR $\otimes$ IAS are split into three components with isospin $T-1$, T , and $T+1$ arising from coupling of one unit of isospin to a state with isospin T . The strength ratios for the GDR $\otimes$ IAS turn out to be very different from the intensity ratios predicted for the $\Delta T_z=-1$ GDR in SCX. Simple isospin geometry arguments predict the following strength ratio for the GDR $\otimes$ IAS components[2, 11]:

$$\begin{aligned} \sigma(T-1) : \sigma(T) : \sigma(T+1) = \\ (2T-1)(T-1)/T(2T+1) : \\ (2T-1)/T(T+1) : 3/(2T+1)(T+1) \end{aligned} \quad (2)$$

The different ratios between the $\Delta T_z=-1$ GDR and the $\Delta T_z=-2$ GDR $\otimes$ IAS arise from the fact that the transition induced by the isospin lowering operator are T dependent. Table 3 lists the expected ratios for the GDR $\otimes$ IAS assuming equal reduced matrix elements for the three states. The Pauli blocking effect will further suppress the T and $T+1$ states giving a larger strength to the $T-1$ component. For heavy nuclei with $T\gg 1$ (e.g. ^{115}In and ^{197}Au), the $T+1$ component is negligible, and most of the strength lies in the lowest component. However, for ^{51}V ($T=5/2$) the middle member could have a strength as large as that of the $T-1$ member. The use of different reduced matrix elements (see footnote of Table 3) suppresses the upper components but still predicts about equal strengths for the $T-1$ and T members. We will return to the ^{51}V splitting in the next section.

Table 4 lists the giant dipole energies derived from DCX compared with results from photonuclear reactions. For ^{115}In and ^{197}Au the GDR $\otimes$ IAS is predominantly $T-1$, whereas the “normal” GDR in the target nucleus is predominantly T . Therefore the energy of the GDR $\otimes$ IAS relative to the DIAS is related via the symmetry energy ΔE^- (listed in Table 4)

Table 3: Intensity ratios for isospin components of the GDR built on the IAS in pion double charge exchange reaction, assuming equal reduced matrix elements.

Target	$ T, T_z\rangle_i$	$ T, T_z\rangle_f$	Isospin members ^a		
			$ T_i - 1\rangle$	$ T_i\rangle$	$ T_i + 1\rangle$
⁵¹ V ^b	(5/2, 5/2)	(1/2, 1/2)	0.40	0.46	0.14
¹¹⁵ In	(17/2, 17/2)	(13/2, 13/2)	0.78	0.20	0.02
¹⁹⁷ Au	(39/2, 39/2)	(35/2, 35/2)	0.90	0.10	0.00

^a Cross section ratios have been calculated using the relation, $\sigma(T-1) : \sigma(T) : \sigma(T+1) = (2T-1)(T-1)/T(2T+1) : (2T-1)/T(T+1) : 3/(2T+1)(T+1)$. The T and $T+1$ members are further suppressed by Pauli blocking, not included in the listed ratios (Ref. [2, 11]).

^b Using different reduced matrix elements taken from SCX on ⁶⁰Ni (Ref. [19]), gives for ⁵¹V intensity ratios $|T_i-1\rangle : |T_i\rangle : |T_i+1\rangle = 0.48 : 0.44 : 0.08$

Table 4: Giant-dipole energies derived from DCX on ⁵¹V, ¹¹⁵In, and ¹⁹⁷Au compared with results from photonuclear reactions.

Target	δE_x^a (MeV)	ΔE^-^b (MeV)	$E1^{DCX}^c$ (MeV)	$E1^{DCX} A^{1/3}$ (MeV)	$E1^{DCX} A^{1/6}$ (MeV)	$E1^{photo}^d$ (MeV)
⁵¹ V	18.9±0.4	2.5	18.9±0.4	70.1±0.7	36.4±0.8	18.98 ^e
¹¹⁵ In	12.2±0.4	3.2	15.4±0.4	74.9±1.9	34.0±0.9	15.63
¹⁹⁷ Au	9.3±1.5	2.5	11.8±1.5	79.1±2.9	32.8±1.2	13.8

^a $\delta E_x = Q_{DIAS} - Q_{GDR\otimes IAS}^{middle}$ for ⁵¹V and $\delta E_x = Q_{DIAS} - Q_{GDR\otimes IAS}^{wholepeak}$ for ¹¹⁵In and ¹⁹⁷Au.

^b $\Delta E^- = T[E_v - (2T+3)E_t]$; $E_v = 50/A$; $E_t = -0.003$ for $A < 80$ and $E_t = +0.003$ for $A > 80$, (Ref. [28]).

^c $E1^{DCX} = \delta E_x + \Delta E^-$ for ¹¹⁵In and ¹⁹⁷Au, and $E1^{DCX} = \delta E_x$ for ⁵¹V.

^d Ref. [23].

^e ⁵¹V is an oblate nucleus. The photonuclear spectrum has two peaks at $E_< = 17.86$ and $E_> = 21.22$ MeV. The listed value is the weighted average, Ref. [23].

to the E1 energies from photonuclear data[23]. Table 4 shows that the E1 energies deduced from the present DCX experiment are surprisingly close to the photonuclear data. We also list in Table 4 $E1^{DCX} A^{1/3}$ and $E1^{DCX} A^{1/6}$. The results indicate that the E1 energies deduced from DCX are consistent with the $A^{-1/3}$ or $A^{-1/6}$ dependence as observed for the “normal” GDR.

For heavy nuclei a simple relation is expected between the three double resonances observed in the present study, namely[2]:

$$Q_{GDR \otimes IAS} = (1/2)(Q_{GDR^2} + Q_{DIAS}) \quad (3)$$

The experimental energies (Table 3) are in close agreement with the above simple relation, which gives an additional support for the proper identification of the resonances.

3.4.3 Splitting of the GDR built on the IAS of ^{51}V

The frequency of the GDR vibration is known to scale inversely with the size of the nucleus (going as $1/R$ or $1/R^2$). For axially symmetric deformed nuclei the GDR is split into two components. The low-frequency component arises from oscillation along the major axis of the nucleus[23]. For prolate deformation a strength ratio of lower : upper = 1 : 2 is expected, and for oblate deformed nuclei a ratio of lower : upper = 2 : 1 should prevail[23]. The nucleus ^{51}V is well known to be oblate. The photonuclear cross section on ^{51}V exhibits a clear double-peak structure in the GDR region, which is well fitted with a sum of two Lorentzian functions with relative strengths $\sigma_{lower} : \sigma_{upper} = 1.8 : 1$ in close agreement with the expectation for an oblate-shape nucleus[23]. The deformation found in the ground state of ^{51}V is not changed much when one or two Coulomb energies are added to the nucleus. Thus, the charge exchange GDR (in the $T_z=T-1$ nucleus) and the GDR $\otimes$ IAS (in the $T_z=T-2$) should have a similar deformation splitting as observed in the normal GDR in the $T_z=T$ target nucleus.

Figure 5 shows again the DCX spectrum of ^{51}V at 11° displayed earlier in Fig. 1, but with the peak labeled GDR $\otimes$ IAS fitted with three Lorentzian peaks. The energies and the widths of the three components were varied to obtain the least χ^2 fit for the entire spectrum. Table V summarizes the deduced energies, widths and cross sections of the three components of the GDR $\otimes$ IAS on ^{51}V at $\theta_{lab} = 11^\circ$. The resulting fit is represented by the solid curve in Fig. 5. The deduced cross section ratios are $\sigma_{lower} : \sigma_{middle} : \sigma_{upper} = 0.67 : 0.90 : 0.45$. The energy spacing among the three components is $\Delta E_1 = \Delta E_2 = 3.3$ MeV.

One can imagine the following scenario. The “normal” GDR in a $T > 0$ nucleus splits into two isospin components with T and $T+1$ and strength ratio of $\sigma_{lower} : \sigma_{upper} = T : 1$, respectively. The charge exchange GDR (in the $\Delta T_z = -1$ mode) splits into three isospin components $T-1$, T , and $T+1$. The predicted intensity ratios based on isospin coupling coefficients only are listed in Table 3. The Pauli blocking effect will further suppress the T and $T+1$ components and enhances the $T-1$ strength. When the difference in the reduced matrix elements is included (estimated from pion SCX data on $^{60}\text{Ni}(T=2)$ [19]) one obtains for ^{51}V the following ratios $|T-1\rangle : |T\rangle : |T+1\rangle = 0.48 : 0.44 : 0.08$. Thus the GDR built on the IAS is predicted to have about equal strength for the $|T-1\rangle$ and the $|T\rangle$ components whereas the $|T+1\rangle$ is negligible. The energy difference between $|T-1\rangle$ and $|T\rangle$ is $\Delta E^- = 2.5$ MeV

Figure 5: Missing mass spectrum for the $^{51}\text{V}(\pi^+, \pi^-)^{51}\text{Mn}$ reaction at $T_\pi=295 \text{ MeV}$ and $\theta_{\text{lab}}=11^\circ$ presented in Fig. 1, but with the GDR $\otimes$ IAS fitted with three peaks labeled GR1, GR2, and GR3.

Table 5: Results of the three-Lorentzian-peaks fit for the giant dipole resonance built on the isobaric analog state in the DCX reaction on ^{51}V at $\theta_{lab} = 11^\circ$ and $T_\pi = 295$ MeV.

Component	$-Q$ (MeV)	Γ (MeV)	$d\sigma/d\Omega$ ($\mu\text{b/sr}$)
GR1	29.8 ± 0.2	3.0 ± 0.5	0.67 ± 0.06
GR2	33.1 ± 0.4	4.0 ± 1.2	0.90 ± 0.08
GR3	36.4 ± 0.5	2.5 ± 1.5	0.45 ± 0.06

(Table 4). In addition, each of the two components has a deformation splitting as discussed earlier. Since the isospin splitting energy ($\Delta E^- = 2.5$ MeV) and the deformation splitting energy ($\Delta E = 3.3$ MeV) are quite close, one would expect a three-component structure in the GDR $\otimes$ IAS with the middle member being the sum of two overlapping components of different isospin. In the ideal case we expect a strength ratio of $\sigma_{lower} : \sigma_{middle} : \sigma_{upper} = 2 : 3 : 1$. The observed structure is close to the above simple estimate for the GDR $\otimes$ IAS on ^{51}V . Other possibilities, however, cannot be ruled out.

An additional source for splitting of both the GDR $\otimes$ GS and the GDR $\otimes$ IAS is the dependence on the spin coupling. For nuclei with $J_i \geq 1$ the GDR splits into three members with $J_f = J_i - 1$, J_i , and $J_i + 1$. However, the splitting due to spin is quite negligible relative to deformation and isospin splittings. This is evidenced by the single narrow peak of the GDR in many odd A spherical heavy nuclei (*e.g.* ^{91}Zr , ^{93}Nb , ^{117}Sn , ^{119}Sn , ^{197}Au , *etc*). In all the above cases there is no observable effect of spin splitting.

3.4.4 Mass dependence of double-resonance cross sections and sum-rule analysis

We now address the A dependence of the cross section of the GDR $\otimes$ IAS and the GDR $\otimes$ GDR. Figure 6 contains the previous and present data for the giant dipole built on the isobaric analog state. Following the discussion of Ref. [11], the figure displays the normalized peak cross section for the GDR $\otimes$ IAS as a function of A in a log-log scale. Including the present data on ^{51}V , ^{115}In and ^{197}Au , the experimental peak cross sections are well represented by the function:

$$d\sigma/d\Omega_{\text{GDR}\otimes\text{IAS}}^{\text{peak}} = 26,470(N - Z)NZ/A^{2.74} \text{ (nb/sr)}. \quad (4)$$

The present fit includes also the ^{13}C and ^{27}Al data points[24]. The nuclei ^{13}C and ^{27}Al are $T = 1/2$ nuclei and therefore the GDR $\otimes$ IAS is expected to be suppressed due to isospin considerations as illustrated in Fig. 2 of Ref. [24]. However, the measured cross sections for the above nuclei[24] are quite large and are well represented by the global power-law fit displayed in Fig. 6. It is likely that for $T = 1/2$ nuclei the GDR $\otimes$ IAS may gain some extra strength through the IAS coupling since in these cases the IAS is also the ground state of the intermediate nuclei. The new fit to the GDR $\otimes$ IAS (solid line) which includes the ^{13}C , ^{27}Al and the new data points yields a somewhat stronger A dependence of the cross sections in comparison with the previous fit (dashed line). In Ref. [11] we give the arguments for

Figure 6: Plot of the GDR⊗IAS cross sections at the peak divided by $(NZ)(N-Z)$ vs A for the (π^+, π^-) reactions on a log-log scale. The filled circles are the data points from the present study and the X's are from Ref. [11]. The dashed line is a power-law fit to the previous data with the parameters indicated (dashed arrow). The solid line includes the present data except for data from the deformed nucleus ^{51}V .

FIG7_E1154.GRF

Figure 7: Plot of the double dipole cross sections at 5° divided by $(NZ)^2$ vs A for the (π^+, π^-) reactions. The filled circles are the present results and the X's are from Ref. [12]. The dashed line is a power-law fit to the previous data[12], and the solid line is a refit including the new data. The ^{12}C and ^{32}S data are not included in either fit[12].

Table 6: Details of the CCIA calculations normalized to the energy weighted sum rule.

Target	E1 ^a (MeV)	Matrix element ^b (efm)	β^c	$d\sigma/d\Omega_{DD}^{5^\circ}$			
				Exp.	Theory		Sum
				($\mu\text{b/sr}$)	2 ⁺ ($\mu\text{b/sr}$)	0 ⁺ ($\mu\text{b/sr}$)	($\mu\text{b/sr}$)
⁵¹ V	18.51	3.177	0.134	6.8 $\pm$ 0.5	1.33	2.28	3.81
¹¹⁵ In	15.79	5.135	0.072	4.1 $\pm$ 0.8	1.28	6.73	8.01
¹⁹⁷ Au	13.79	7.127	0.053	7.7 $\pm$ 1.6	1.37	8.08	9.45

^a E1 energies were obtained using the relation: $E1 = 77.9A^{-1/3}[1 - \exp(-A/238)] + 34.5A^{-1/6}\exp(-A/238)$ MeV, Ref. [23].

^b This is the integral in Eq. (7).

^c Strength for the first-step GDR. For the second step, we used $\sqrt{5/3}\beta$, for the $J^\pi=2^+$, and $\sqrt{2/3}\beta$ for the $J^\pi=0^+$ double dipole states, Ref. [27].

decomposing the cross section into the above products in terms of the first and second step form factors times an overall A -dependent attenuation factor. Equation (4) gives an adequate overall behavior of the cross section of the GDR $\otimes$ IAS as a function of N , Z , and A .

Figure 7 displays the normalized 5° cross sections for the double dipole *vs* A . The data includes all previous and present measurements. The solid line is the best power-law fit to the data:

$$d\sigma/d\Omega_{GDR\otimes GDR}^{5^\circ} = 10,480(NZ)^2/A^{3.60} \text{ (nb/sr)}. \quad (5)$$

The fit in Fig. 7 does not include the ¹²C and ³²S deformed nuclei[12]. One can understand the above expression for the GDR $\otimes$ GDR as a product of the strength of the first and second step times an overall attenuation factor for DCX[12]. The present fit with the new data is in agreement with the previous fit[12]. The above analysis of the cross sections of the GDR $\otimes$ IAS and the GDR $\otimes$ GDR in terms of the two ingredient resonances indicates that pion double charge exchange involving the excitations of the closed-shell cores proceeds at this energy predominantly by a two-step sequential (π^+ , π^0)(π^0 , π^-) mechanism, and therefore should be a unique probe for exciting isoscalar double giant resonances in nuclei.

Table 5 summarizes the sum rule analysis for the double dipole. We used the energy weighted sum rule[23, 25, 26] to extract the deformation parameter β for the GDR:

$$\Delta E \left[\int \beta r \rho'(r) r^2 dr \right]^2 = (9/4\pi)(\hbar/2M)(NZ/A)e^2 \quad (6)$$

or equivalently,

$$\Delta E |M(E1)|^2 = (9/4\pi)(\hbar/2M)(NZ/A)e^2 \quad (7)$$

where $M(E1)$ is the isovector transition strength and ΔE is the E1 energy obtained from the

following parameterization:[23]

$$\Delta E(E1) = 77.9A^{-1/3} \left[1 - e^{-A/238} \right] + 34.5A^{-1/6}e^{-A/238} \text{ MeV} \quad (8)$$

Table 5 gives the details of these calculations and a comparison between the measured cross sections for the double dipole from the present study and the theoretical calculations for 0^+ and 2^+ double dipole states using NEWCHOP and the prescription of Ref. [27].

3.5 Summary

The (π^+, π^-) reaction was used to populate isotensor ($\Delta T=2$) double giant resonances on ^{51}V , ^{115}In , and ^{197}Au . The E1 energies derived from DCX in the present study are consistent with an $A^{-1/3}$ or $A^{-1/6}$ dependence, and agree (after correcting for the symmetry energy) with the results from photonuclear reactions. Partial angular distributions were measured on ^{51}V and ^{115}In and found to have a dipole shape for the GDR built on the IAS and predominantly a quadrupole shape for the double dipole. The width of the double GDR is about 1.5 times the width of the “single” dipole. The GDR built on the IAS on ^{51}V has a three-peak structure in the resonance region, which may arise from the deformed (oblate) shape of ^{51}V and from the isospin splitting into the $T=3/2$ and $T=5/2$ components. The observed intensity ratios and energy splitting of the three components are in qualitative agreement with the above simple arguments.

References

- [1] N. Auerbach, Proc. Workshop on Pion-Nucleus Physics: Future directions and New Facilities at LAMPF, AIP Conf. Proc. **163**, 34-35 (1987).
- [2] N. Auerbach, Annuals of Physics, 197, 376–395 (1990).
- [3] F. G. Scholz and F. J. W. Hahne, Z. Physical A – Atomic Nuclei **336**, 145 (1990).
- [4] G. Lauritsch and P. G. Reinhard, Nuclear Physics **A509**, 287 (1990).
- [5] G. Baur and C. A. Bertulani, Physics Letters B **174**, 23 (1986).
- [6] N. Francaria, Nuclear Physics **A482**, 245C (1988) and references therein.
- [7] G. Baur and Carlos Bertulani, Physical Review C **34**, 1654 (1986).
- [8] F. Catara, Nuclear Physics **A482**, 271C (1988).
- [9] S. Mordechai *et al.*, Physical Review Letters **60**, 408 (1988).
- [10] S. Mordechai *et al.*, Physical Review Letters **61**, 531(1988).

- [11] S. Mordechai *et al.*, Physical Review C **40**, 850 (1989).
- [12] S. Mordechai *et al.*, Physical Review C, **41**, 202 (1990); and *ibid.* **43**, R1509 (1991).
- [13] H. Ward *et al.*, Physical Review C **45**, 2723 (1992).
- [14] H.A.Thiessen *et al.*, Los Alamos Scientific Laboratory Report No. LA-6663-MS, 1977 (unpublished); S. J. Greene *et al.*, Physics Letters **88B**, 62 (1979).
- [15] C. L. Morris, J. F. Amann, R. L. Boudrie, N. Tanaka, S. J. Seestrom-Morris, L. C. Bland, P. A. Seidl, R. Kiziah, and Steven J. Greene, Nuclear Instruments and Methods A **238**, 94 (1985).
- [16] G. Rowe, M. Salomon, and R. H. Landau, Physical Review C **18**, 584 (1978).
- [17] C. L. Morris, unpublished.
- [18] E. Rost, computer code CHOPIN (unpublished). The code has been modified by C. L. Morris to calculate pion charge-exchange reactions and renamed NEWCHOP.
- [19] A. Erell, J. Alster, J. Lichtenstadt, M. A. Moinester, J. D. Bowman, M. D. Cooper, F. Irom, H. S. Matis, E. Piasetzky, and U. Sennhauser, Physical Review C **34**, 1822 (1986).
- [20] L. J. Tassie, Australian Journal of Physics **A305**, 509 (1978).
- [21] K. Muto, Physics Letters B **277**, 13 (1992).
- [22] D. C. Zheng, L. Zamick, and N. Auerbach, Annals of Physics **197**, 343 (1990).
- [23] B. L. Berman and S. C. Fultz, Reviews of Modern Physics **47**, 713 (1975).
- [24] S. Mordechai, *et al.*, Physical Review C **43**, 1111 (1991).
- [25] A. Bohr and B. Mottelson, Nuclear Physics, Volume II, (Benjamin, New York), (1975).
- [26] A. Gal, Physical Review C **25**, 2680 (1982).
- [27] J. M. O'Donnell, H. T. Fortune and E. Rost, Physical Review C **44**, 2426 (1991).
- [28] N. Auerbach, A. Klein and N. V. Giai, Physics Letters **106B**, 347 (1981).

4 Publications

Since 1991

1. D. L. Watson, R. Gilman, C. L. Morris, J. D. Zumbro, S. Mordechai, M. Burlein, J. M. O'Donnell, H. T. Fortune, A. Kotwal, J. Puttz, J. D. Silk, J. S. McDonald, C. Fred Moore, A. L. Williams, and S. Y. Yoo, "Pion-induced double charge exchange on ^{12}C , ^{24}Mg , ^{32}S , and ^{40}Ca ," *Physical Review C* **43**, 1318-1320 (1991).
2. A. L. Williams, J. A. McGill, C. L. Morris, G. R. Burleson, J. A. Faucett, D. S. Oakley, M. Burlein, H. T. Fortune, J. M. O'Donnell, G. P. Kahrmanis, and C. Fred Moore, "Pion double charge exchange on $^{42,44,48}\text{Ca}$ for $300 \leq T_\pi \leq 550$ MeV," *Physical Review C* **43**, 766-770 (1991).
3. S. Mordechai, C. L. Morris, J. M. O'Donnell, M. A. Kagarlis, D. Fink, H. T. Fortune, D. L. Watson, R. Gilman, H. Ward, A. Williams, Sung Hoon Yoo, and C. Fred Moore, "Analog of the $T_{>}$ giant dipole resonance in light nuclei," *Physical Review C* **43**, 1111-1115 (1991).
4. S. Mordechai, H. Ward, K. Johnson, G. Kahrmanis, David Saunders, C. Fred Moore, J. M. O'Donnell, M. A. Kagarlis, D. Smith, H. T. Fortune, C. L. Morris, "Comparison of the double giant-dipole states observed in (π^-, π^+) and (π^+, π^-) reactions on ^{40}Ca and ^{27}Al ," *Physical Review C* **43**, R1509-R1512 (1991).
5. S. Mordechai and C. Fred Moore, "Giant resonances in single and double charge exchange," *Proceedings of the International Workshop on Pions in Nuclei*, Edited by E. Oset, M. J. Vicente Vacas, and C. G. Recio, World Scientific, Peniscola, Spain, 22-31 (1991).
6. C. L. Morris, R. D. Ransome, B. G. Ritchie, J. D. Zumbro, D. L. Watson, and C. Fred Moore, "Evidence for physics beyond the quasi-deuteron model in pion absorption," *Proceedings of the International Workshop on Pions in Nuclei*, Edited by E. Oset, M. J. Vicente Vacas, and C. G. Recio, World Scientific, Peniscola, Spain, 22-31 (1991).
7. Herb Ward, C. Fred Moore, P. Susie Maloney, and Jimmy D. Martin, "Probabilistic Neural Nets in Analysis of Magnetic Spectrometer Data," *Nuclear Instruments & Methods* in press (1992).
8. S. Mordechai and C. Fred Moore, "Double giant resonances in atomic nuclei," *NATURE* **352**, 393-397 (1991).
9. C. Fred Moore, K. Johnson, G. P. Kahrmanis, James McDonald, M. Snell, H. J. Ward, Sung Hoon Yoo, C. L. Morris, S. Mordechai, M. Burlein, N. Claytor, H. T. Fortune, R. Ivie, G. B. Liu, J. M. O'Donnell, D. Smith, N. Auerbach, and D. Robson, "Angular distributions for the double isobaric analog and a $T_{<}$ state at high excitation in pion double charge exchange on ^{93}Nb ," *Physical Review C* **44**, 2209 (1991).

10. H. Ward, K. Johnson, G. Kahrimanis, D. Saunders, C. Fred Moore, S. Mordechai, C. L. Morris, J. M. O'Donnell, N. Auerbach, M. A. Kagarlis, H. T. Fortune, and D. Smith, "Double giant dipole resonance in the (π^-, π^+) reaction," *Physical Review C* **45**, 2723 (1992).
11. R. D. Ransome, C. L. Morris, V. R. Cupps, B. W. Ferguson, J. A. McGill, D. L. Watson, J. D. Zumbro, B. G. Ritchie, J. R. Comfort, J. R. Tinsley, R. A. Loveman, S. Dawson, S. Green, P. C. Gugelot, and C. Fred Moore, "Pion absorption in heavy nuclei," *Physical Review C* **45**, R509-R513 (1992).
12. B. Brinkmüller, C. L. Bili, D. Dehnhard, M. Jones, M. Martines, S. Nanda, S. M. Sterbenz, Yi-Fen Yen, L. Atencio, S. Greene, C. L. Morris, S. J. Seestrom, G. Burleson, K. Dhuga, J. Faucett, R. Garnett, K. Maeda, C. Fred Moore, S. Mordechai, A. Williams, S. H. Yoo, and L. C. Bland, "Elastic Scattering of π^+ and π^- from ^4He between 90 and 240 MeV," *Physical Review C* **44**, 2031 (1991).
13. D. S. Oakley, R. J. Peterson, S. J. Seestrom, C. L. Morris, M. A. Plum, J. D. Zumbro, A. L. Williams, M. A. Bryan, J. W. McDonald, and C. Fred Moore "Energy dependence of pion inelastic scattering from ^{208}Pb ," *Physical Review C* **44**, 2058 (1991).
14. A. L. Williams, K. W. Johnson, G. P. Kahrimanis, H. Ward, C. Fred Moore, J. A. McGill, C. L. Morris, G. R. Burleson, J. A. Faucett, M. Rawool-Sullivan, D. S. Oakley, M. Burlein, H. T. Fortuen, E. Insko, R. Ivie, J. M. O'Donnell, and D. Smith, "Mass dependence of high-energy pion double charge exchange," *Physical Review C* **44**, 2025 (1991).
15. M. K. Jones, D. Dehnhard, S. K. Nanda, S. M. Sterbenz, C. L. Morris, M. Plum, J. D. Zumbro, A. Hussein, D. Oakley, M. J. Smithson, A. Williams, J. McDonald, M. Bryan, A. Fuentes, M. Lynker, D. Crockett, M. Machuca, S. Mordechai, and C. Fred Moore, " $^4\text{He}(\pi, \pi'p)^3\text{H}$ reaction near the $P_{3,3}$ π -nucleon resonance," *Physical Review C* **46**, 52 (1992).
16. J. L. Langenbrunner, M. K. Jones, D. Dehnhard, S. K. Nanda, S. M. Sterbenz, C. L. Morris, M. Plum, J. D. Zumbro, A. Hussein, D. S. Oakley, M. Smithson, A. Williams, J. McDonald, M. Bryan, A. Fuentes, M. Lynker, D. Crockett, M. Machuca, S. Mordechai, and C. Fred Moore, "Coincidence Measurement of $^4\text{He}(\pi, \pi'p)$ and $^4\text{H}(\pi, \pi't)$," *Physical Review C* submitted (1991).
17. E. Gülmez, D. L. Adams, S. Beedoe, M. Bleszynski, J. Bystricky, V. Ghazikhanian, G. Igo, T. Jaroszewicz, A. G. Ling, M. Nasser, M. W. McNaughton, S. Penttilä, G. Glass, G. Kahrimanis, W. F. Kielhorn, K. H. McNaughton, P. J. Riley, S. Sen, D. Wolf, "Elastic scattering of N, L, and S-Type polarized 794 MeV protons from an ND_3 target polarized in the S-L Plane," *Physical Review C* submitted (1991).
18. J. Bar-Touv and S. Mordechai, "Systematics of the double giant dipole resonances in nuclei," *Physical Review C* **45**, 197 (1992).

19. D. A. Smith, H. T. Fortune, G. B. Lui, J. M. O'Donnell, M. Burlein, S. Mordechai, and A. R. Fazely, "Pion double charge exchange on $^{128,130}\text{Te}$," *Physical Review C* **46**, 477 (1992).
20. S. Mordechai, C. Fred Moore, H. T. Fortune, et al., "Pion double charge exchange on ^{24}Mg at 165 MeV," in preparation.
21. M. K. Jones, R. D. Ransome, V. R. Cupps, R. W. Fergerson, C. L. Morris, J. A. McGill, J. D. Zumbro, J. R. Comfort, B. G. Ritchie, J. R. Tinsley, P. C. Gugelot, C. Fred Moore, "Pion absorption above the $\Delta(1232)$ resonance," *Physical Review C* **46**, 52 (1992).??????
22. N. M. Hintz, X. H. Yang, M. Gazzaly, S. J. Seestrom-Morris, C. L. Morris, D. C. Cook, A. M. Mack, J. W. McDonald, S. Mordechai, D. S. Oakley, C. F. Moore, M. Lynker, J. D. Zumbro, "Determination of neutron and proton multipole matrix elements in ^{208}Pb from π^- and π^+ scattering at 180 MeV," *Physical Review C* **45**, 601 (1992).
23. J. O'Donnell, R. Lomax, B. McCloud, J. McGill, C. L. Morris, J. Davis, R. A. DeHaven, R. Johnson, G. Tubb, J. Novak, E. Gray, B. Rusnak, J. M. Applegate, T. D. Averett, J. Beck, B. G. Ritchie, E. Haebe, D. Kiehlmann, U. Klein, M. Peiniger, P. Schafer, H. Vogel, H. Ward, and C. Fred Moore, "A superconducting radio-frequency cavity for manipulating the phase space of pion beams at LAMPF," *Nuclear Instruments and Methods* **A317**, 445 (1992).
24. R. A. Lindgren, B. L. Clausen, G. S. Blanpied, J. Hernandez, C. S. Mishra, W. K. Mize, C. S. Shisnant, B. G. Ritchie, C. L. Morris, S. J. Seestrom, C. Fred Moore, P. A. Seidl, B. H. Wildenthal, R. Gilman, and J. Carr, "Pion scattering to 6^- stretched states in ^{24}Mg and ^{26}Mg ," *Physical Review C* **44**, 2413-2418 (1991).
25. H. Ward, J. Applegate, N. Auerbach, J. Beck, J. Johnson, K. Koch, C. Fred Moore, S. Mordechai, C. L. Morris, J. M. O'Donnell, M. Rawool-Sullivan, B. G. Ritchie, D. L. Watson, and C. Whitley "Systematics of the DIAS cross section at 50 MeV," in preparation (1992).
26. J. Bar-Touv and S. Mordechai, "Soft giant dipole resonance in the framework of open-shell linear response method," in preparation (1992).
27. C. L. Morris, R. Gilman, G. P. Gilfoyle, J. D. Zumbro, M. G. Burlein, S. Mordechai, T. Fortune, K. Maeda, G. S. Blanpied, B. A. Brown, L. C. Bland, M. Brown, D. S. Saunders, P. A. Seidl, and C. Fred Moore, "On the relative sign of the neutron- and proton-quadrupole matrix element for 2_2^+ transition in ^{34}S ," *Physics Letters B* submitted (1991).
28. J. M. O'Donnell, H. T. Fortune, J. D. Silk, C. L. Morris, J. D. Zumbro, S. Mordechai, C. Fred Moore, and S. H. Yoo, "The ^{138}Ce double isobaric analog state studied by pion induced double charge exchange," *Physics Review C* submitted (1991).

29. Y. Grof, N. Auerbach, D. Benton, H. T. Fortune, K. Johnson, M. A. Kagarlis, G. Kahrimanis, S. Mordechai, C. L. Morris, J. M. O'Donnell, I. Orion, D. Saunders, D. A. Smith, H. Ward, and C. Fred Moore, "Double giant resonances in pion double charge exchange on ^{51}V , ^{115}In , and ^{197}Au ," *Physical Review C* submitted (1992).
30. I. Orion, N. Auerbach, D. Benton, H. T. Fortune, Y. Grof, K. Johnson, M. A. Kagarlis, G. Kahrimanis, S. Mordechai, C. L. Morris, J. M. O'Donnell, D. Saunders, D. A. Smith, H. Ward, and C. Fred Moore, "Properties of the double isobaric analog state on ^{51}V , ^{115}In , and ^{197}Au ," in preparation (1992).
31. A. Salman, N. Auerbach, D. Benton, H. T. Fortune, Y. Grof, K. Johnson, M. A. Kagarlis, G. Kahrimanis, S. Mordechai, C. L. Morris, J. M. O'Donnell, I. Orion, D. Saunders, D. A. Smith, H. Ward, and C. Fred Moore, "The lowest three 0^+ states in ^{12}O ," *Physical Review C* submitted (1992).
32. R. D. Ransome, C. L. Morris, M. Martinez, V. R. Cupps, R. W. Fergerson, J. A. McGill, D. L. Watson, J. D. Zumbro, J. R. Comfort, B. G. Ritchie, J. R. Tinsley, R. A. Loveman, S. Dawson, A. Green, P. C. Gugelot, C. Fred Moore, "Inclusive pion single charge exchange on nuclei," *Physical Review C* submitted (1991).
33. M. K. Jones, B. G. Ritchie, R. D. Ransome, D. Clayton, K. Pujara, C. L. Morris, J. A. McGill, D. L. Watson, C. Fred Moore, I. Brown, and P. Campbell, "Energy dependence of the total $^{12}\text{C}(\pi^+, 2p)$ cross section," *Physics Letters B* submitted (1992).
34. R. D. Ransome, C. L. Morris, M. K. Jones, B. G. Ritchie, D. L. Watson, J. A. McGill, K. Pujara, D. B. Clayton, I. Brown, P. Campbell, C. Fred Moore, "Pion absorption in light nuclei," *Physical Review C* submitted (1992).
35. H. Ward *et al.* "Energy dependence of the GDR built on the IAS in pion double charge exchange on ^{13}C ," in preparation (1992).

5 Abstracts

Since 1991

1. H. J. Ward, K. Johnson, G. P. Kahrimanis, D. Saunders, C. Fred Moore, Shaul Mordechai, C. L. Morris, J. M. O'Donnell, H. T. Fortune, M. A. Kagarlis, D. Smith, "Observation of the double giant dipole resonance in (π^-, π^+) double charge exchange," *Bulletin of American Physical Society* **36**, 99 (1991).
2. S. Mordechai, C. L. Morris, J. M. O'Donnell, H. T. Fortune, M. A. Kagarlis, D. Smith, H. J. Ward, A. Williams, Sung Hoon Yoo, C. Fred Moore, and D. L. Watson, "Observation of the T_{γ} giant dipole resonance in light nuclei," *Bulletin of American Physical Society* **36**, 99 (1991).

3. K. Johnson, M. Purcell, H. Ward, A. Williams, S. Worm, B. Brinkmüller, D. Dehnhard, J. Langenbrunner, M. Palarczyk, C. Reidel, Y. Yen, B. Burleson, M. Ghossain, A. Klein, G. Kyle, S. Mukhopadhyay, M. Wang, R. Ivie, M. Kagarlis, S. Loe, M. Mackenzie, J. O'Donnell, D. Smith, M. Khandaker, M. Khayat, C. Allgower, J. Comfort, S. Greene, J. Jarmer, S. Penttilä, and N. Tanaka, "Pion elastic and inelastic scattering on polarized ^{13}C ," *Bulletin of American Physical Society* **36**, 99 (1991).
4. J. L. McDonald, M. A. Bryan, C. Fred Moore, J. W. McDonald, Thomas E. Cowan, C. L. Bennett, M. Clark, R. H. Howell, R. R. Rohatgi, D. H. G. Schneider, J. Schweppe, J. Fajans, "Positron-electron spectroscopy in heavy-ion collisions," *Bulletin of American Physical Society* **36**, 98 (1991).
5. C. L. Morris, John McGill, Ronald Ransome, Mark Jones, David Clayton, Junal Pujara, Barry Ritchie, Ian Brown, Paul Campbell, Doug Watson, and C. Fred Moore, "Excitation function of partial cross sections for pion absorption on ^{12}C ," *Bulletin of American Physical Society* **36**, 1299 (1991).
6. B. G. Ritchie, J. M. Applegate, T. D. Averett, J. Beck, D. S. Oakley, C. Kormanyos, R. J. Peterson, J. A. McGill, C. L. Morris, C. F. Moore, M. Snell, and H. Ward, "50 and 65 MeV $\pi^\pm$ scattering from ^{12}C with the SCRUNCHER," *Bulletin of American Physical Society* **36**, 1299 (1991).
7. S. Mordechai and C. Fred Moore, "Double resonance excitations in pion double charge exchange," Notre Dame Workshop on Giant Resonances and Related Phenomena, Notre Dame, Indiana, Oct. 21-23 (1991).
8. J. E. Wise, S. Høibråten, R. J. Peterson, J. A. McGill, C. L. Morris, S. J. Seestrom, R. M. Whitton, J. D. Zumbro, and A. L. Williams, "Y-Scaling in 500 MeV pion quasielastic scattering," *Bulletin of American Physical Society* **36**, 1299 (1991).
9. Y. Grof, I. Orion, S. Mordechai, C. Fred Moore *et al.*, "Double Resonances in Pion Double Charge Exchange on ^{51}V , ^{115}In , and ^{197}Au ," *Bulletin of the Israel Physical Society* **38**, 118 (1992).
10. S. Mordechai, C. L. Morris, Y. Grof, I. Orion, D. Binton, M. S. Kagarlis, H. T. Fortune, J. M. O'Donnell, H. Ward, K. Johnson, G. Kahrmanis, D. Saunders, and C. Fred Moore, "Double giant resonances in pion double charge exchange on ^{51}V , ^{115}In , and ^{197}Au ," *Bulletin of American Physical Society* submitted (1992).

6 Invited Talks

Since 1991

1. S. Mordechai, "Two-step giant dipole excitations in nuclei," Annual Meeting of the Israel Physical Society, Beer Sheva, Israel (1991).

2. S. Mordechai and C. Fred Moore, "Giant resonances in single and double charge exchange," International Workshop on Pions in Nuclei, Peniscola, Spain (1991).
3. S. Mordechai and C. Fred Moore, "Giant resonances," Pion Double Charge Exchange Workshop, DCX Working Group, Los Alamos National Laboratory, Los Alamos, New Mexico (1991)
4. S. Mordechai and C. Fred Moore, "Double resonance excitations in pion double charge exchange," Notre Dame, Indiana (1991).
5. S. Mordechai and C. Fred Moore, 'Giant resonances in single and double charge exchange,' *Proceedings of the International Workshop on Pions in Nuclei*, Ed., E. Oset, M. J. Vicente Vacas, and C. Garcia Recio, World Scientific, Penyscola (Castellón), Spain 22-23 (1991).
6. S. Mordechai and C. Fred Moore, "Double resonance excitations in pion double charge exchange," *Notre Dame University Workshop on Giant Resonances and Related Phenomena*, Notre Dame, Indiana, Oct. 21-23, No. 4 (1991).
7. S. Mordechai and C. Fred Moore, "Double Resonance Excitations at PILAC Energies," *PILAC Users Group Report on the Physics with PILAC*, Los Alamos National Laboratory, Los Alamos, NM 249-263 (1991).
8. R. D. Ransome, C. L. Morris, D. L. Watson, M. K. Jones, B. G. Ritchie, J. A. McGill, and C. Fred Moore, "Pion absorption on light nuclei," *Proceedings of the International Workshop on Pions in Nuclei*, Ed., E. Oset, M. J. Vicente Vacas, and C. Garcia Recio, World Scientific, Penyscola (Castellón), Spain 486-495 (1991).
9. S. Mordechai and C. Fred Moore, "New results on Double Giant Resonances in Nuclei," International Conference on Nuclear Structure and Nuclear Reactions at Low and Intermediate Energies," Dubna, Russia (1992).

7 Proposal Defenses

Since 1991

1. "Measuring neutron-neutron scattering length and effective range using the $^2\text{H}(\pi^-, 2n)\gamma$ reaction," LAMPF Proposal 1182.
2. "Energy dependence of double resonances in pion DCX," LAMPF Proposal 1209.
3. "Study of the mechanism for DCX to excite analogs of anti-analogs in heavy nuclei," LAMPF Proposal 1210.
4. " $(\pi, \pi'X)$ coincidence measurement on and below the $\Delta_{3,3}$ resonance," LAMPF Proposal 1211.

5. " ${}^6\text{Li}(\pi, \pi'X)$ coincidence measurements near the $\Delta_{3,3}$ resonance," LAMPF Experiment 1216.
6. "Accurate determination of the $T_{\geq}$ giant dipole state in light $T = 1/2$ nuclei using pion DCX," LAMPF Proposal 1217.
7. "Elastic scattering of π^+ and π^- from ${}^4\text{He}$ and ${}^3\text{He}$ above the $\Delta_{3,3}$ resonance," LAMPF Proposal 1219.
8. "A study of the pion-nucleus isovector spin-flip mechanism at low energies using ${}^{10}\text{B}$," LAMPF Proposal 1226.
9. "The two-proton component of the pion absorption reaction on nuclei at low energies," LAMPF Proposal 1227.
10. "Feasibility of the direct production of pionic atoms at LAMPF," LAMPF Proposal 1239.
11. "Isospin structure of the giant dipole resonance in pion single charge exchange," LAMPF Proposal 1241.
12. "The double giant dipole resonance in deformed nuclei," LAMPF Proposal 1242.
13. "Giant resonance excitations by low energy pion scattering," LAMPF Proposal 1253.
14. "Search for double Gamow-Teller resonances using pion double charge exchange," LAMPF Proposal 1186U.
15. "The double giant dipole resonance in deformed nuclei," LAMPF Proposal 1262U.
16. "Accurate Measurement of the nuclear isovector monopole resonance using the NMS," LAMPF Proposal 1281

8 Approved New Research Proposals

Since 1991

1. "Measuring neutron-neutron scattering length and effective range using the ${}^2\text{H}(\pi^-, 2n)\gamma$ reaction," LAMPF Experiment #1182.
2. "Energy dependence of double resonances in pion DCX," LAMPF Experiment #1209.
3. "Study of the mechanism for DCX to excite analogs of anti-analogs in heavy nuclei," LAMPF Experiment #1210.
4. " ${}^6\text{Li}(\pi, \pi'X)$ coincidence measurements near the $\Delta_{3,3}$ resonance," LAMPF Experiment #1216.

5. "Accurate determination of the $T_{>}$ giant dipole state in light $T = 1/2$ nuclei using pion DCX," LAMPF Experiment #1217.
6. "Elastic scattering of π^+ and π^- from ^4He and ^3He above the $\Delta_{3,3}$ resonance," LAMPF Experiment #1219.
7. "A study of the pion-nucleus isovector spin-flip mechanism at low energies using ^{10}B ," LAMPF Experiment #1226.
8. "The two-proton component of the pion absorption reaction on nuclei at low energies," LAMPF Experiment #1227.
9. "Feasibility of the direct production of pionic atoms at LAMPF," LAMPF Experiment #1239.
10. "Isospin structure of the giant dipole resonance in pion single charge exchange," LAMPF Experiment #1241.
11. "Giant resonance excitations by low energy pion scattering," LAMPF Experiment #1253.
12. "Search for double Gamow-Teller resonances using pion double charge exchange," LAMPF Proposal 1186U.
13. "The double giant dipole resonance in deformed nuclei," LAMPF Proposal 1262U.
14. "Accurate Measurement of the nuclear isovector monopole resonance using the NMS," LAMPF Proposal 1281.

9 Personnel

Since 1991

William T. Buttler	graduate student
Steve Hamilton	graduate student
John Johnson	graduate student
Kevin Johnson	graduate student
George P. Kahrmanis	graduate student
James L. McDonald	graduate student
Joseph W. McDonald	graduate student
C. Fred Moore	faculty
Shaul Mordechai	faculty (Ben Gurion University of the Negev)
Arturo Morosoff	graduate student
Mike Snell	graduate student (Air Force Academy)
Steve Brent Thomas	graduate student
Herbert J. Ward	graduate student

Charles Whitley

graduate student

Institutions in parentheses indicate either current employment or a permanent place of employment.

END

**DATE
FILMED**

1 / 13 / 93