ANNUAL PROGRESS REPORT
ON
DEEXCITATION PROCESSES IN NUCLEAR REACTIONS

Prepared by
Norbert T. Porile
Department of Chemistry
Purdue University
W. Lafayette, Indiana

September 1992

Grant Number DE-FG02-88ER40408

U.S. Department of Energy

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

During the past year, our research program has involved continuing analysis of Fermilab E-735, search for quark-gluon plasma (QGP) in $\bar{p}$ -p collisions; continuing study of target fragments produced in the interaction of copper with intermediate-energy heavy ions; an exclusive study of multifragmentation using reverse kinematics at the Bevalac; and detector development for the STAR detector at RHIC.

A. Search for QGP in $\bar{p}$ -p Collisions at the Tevatron

Fermilab E-735, a Duke-Fermilab-Iowa State-Notre Dame-Purdue-Wisconsin collaboration, is designed to search for the formation of QGP in $\bar{p}$ -p collisions at $\sqrt{s} = 1.8$ TeV. The experimental observables are the transverse momenta ($p_t \leq 1.5$ GeV/c) of centrally produced mass-identified particles ($\pi^\pm$, $K^\pm$, $\bar{p}$, Λ^0) as a function of charged particle multiplicity. Continuing effort has been devoted to the analysis of the data obtained in a year-long experiment performed in 1988-89.

The collaboration has completed a study of source size using pion interferometry. The experiment detected approximately 10^7 pions in a magnetic spectrometer located at $\sim 90^\circ$ with respect to the colliding $p\bar{p}$ beams. Approximately 10^6 like-sign pion pairs were observed and these constituted the raw data for the analysis. Assorted selection criteria reduced this number to $\sim 3 \times 10^5$ pairs.

The data were analyzed on the basis of several models. In one type of analysis the correlation function can be expressed as [1]

$$C_2 = 1 + \lambda e^{-q_t^2 R^2} \quad q_0 \leq 0.2 \text{ GeV} \quad (1)$$

where q_t is the relative transverse momentum of the $\pi\pi$ pair, R is the source

radius, and λ is the coherence parameter. Eq. 1 requires that the $\pi\pi$ energy difference q_0 be zero and in practice only events for which $q_0 \leq 0.2$ GeV were used. The correlation function must be evaluated with respect to a reference distribution. After considerable investigation of various possible reference distributions we chose pions from different events with the same multiplicity and vertex location. Thus the experimental correlation function was evaluated as

$$C_2 = (\pi^+\pi^+)_{\text{S}}/(\pi^+\pi^+)_{\text{D}} \quad (2)$$

where the denominator is the number of pion pairs at a point in a different event reference distribution, and the numerator is the number at a corresponding point of a same-event distribution.

Figure 1 shows the experimental correlation function and the fit of Eq. 1, which corresponds to $R = 1.06 \pm 0.07$ fm and $\lambda \sim 0.23$. This value of λ is substantially smaller than the maximum theoretical value of 1.0, which should be obtained when the production phases of the two pions are completely random. A completely coherent source would lead to zero enhancement at low q_t [2].

The number of pion pairs in our final data set was sufficiently large to permit a limited investigation of source size as a function of charged particle multiplicity N_c . The resulting values of R as a function of multiplicity density $dN_c/d\eta$, where η is the pseudorapidity, are shown in Fig. 2. A significant increase in source size is seen to accompany increasing $dN_c/d\eta$. Also plotted in Fig. 2 are lower energy results from the ISR [3] and UA1 [4]. These results are not strictly comparable to the present values because of differences in pseudorapidity and azimuthal angle intervals. However, they show similar trends with $dN_c/d\eta$. The coherence parameter λ decreases from ~ 0.4 to ~ 0.2 with increasing $dN_c/d\eta$, suggesting a concomitant increase in the relative number of

Fig. 1. Experimental ratio of like-sign pion pairs from the same event to like-sign pion pairs from different events vs q_t . The curve represents a fit of Eq. 1.

Fig. 2. Interaction radius vs $dN_c/d\eta$. Note that the experimental conditions for the ISR and UA1 points differ from those of the present experiment in a number of significant respects.

phase-correlated pion pairs within our acceptance.

The results can be used to estimate the energy density as a function of $dN_c/d\eta$ using an approach similar to that of Bjorken [5]. Preliminary results indicate that the energy density is approximately independent of multiplicity. Although the energy deposited increases with multiplicity, as evidenced by our previous results showing an increase in mean transverse momentum with $dN_c/d\eta$ [6], this increase is matched by that in source size (Fig. 2), thereby leading to an approximately constant energy density. Searches for QGP such as the present one are predicated on the belief that the QGP can only be made at high energy density and consequently at the highest multiplicities attainable [7]. Confirmation of our preliminary result would suggest that energy densities sufficiently high to result in QGP formation may not be attainable in high multiplicity events.

B. Target Fragments from the Interaction of Copper with Intermediate Energy Heavy Ions

We have extended our previous [8,9] studies of target fragments of the interaction of copper with 15-45 MeV/nucleon ^{12}C ions to 90 MeV/nucleon in order to study the evolution of the reaction dynamics in the intermediate energy regime. An article describing this research has been submitted for publication [10].

The evolution of the mass yield distribution is displayed in Fig. 3. Although the basic shape of the distribution remains the same, the details do change with energy. Thus, the distribution becomes flatter as the energy increases and the minimum observed between spallation and fragmentation products moves to lower mass number. The variation with energy of the mean mass loss from the target is plotted in Fig. 4. The mass loss rises steeply between 15 and 45 MeV/nucleon and then only slightly between 45 and 90 MeV/nucleon.

Fig. 3. Mass-yield distributions for the interaction of copper with 24, 45, and 90 MeV/nucleon ^{12}C . The curves represent an empirical parametrization and the points are experimental cross sections.

Fig. 4. Energy dependence of the mean mass loss from the target for Cu + 90 MeV/nucleon ^{12}C .

The results at the highest two energies have been compared with cascade-evaporation calculations. We used the intranuclear cascade code ISABEL [11] and the evaporation code EVA [12]. The comparison is shown in Fig. 5. At 90 MeV/nucleon the calculation predicts the general shape of the distribution quite well. Although the cross sections of near-target products are somewhat overpredicted and those of low-mass products are underpredicted, the exponential spallation region is well represented by the cascade-evaporation model. The under-estimation of the yields of low mass products is a consequence of the fact that these products are formed in binary fragmentation, a process not incorporated into the cascade-evaporation model. The calculation cannot account for the 45 MeV/nucleon data indicating that the model breaks down between 45 and 90 MeV/nucleon. Evidently, mean field effects are still important at the lower of these energies.

The forward component of velocity of the composite nucleus leading to a particular product, v_1 , was obtained from the measured product recoil range. The results for 90 MeV/nucleon ^{12}C plus copper are compared with the cascade-evaporation calculations in Fig. 6. Both sets of velocities are expressed in terms of the velocity of the putative compound nucleus, v_{CN} . The experimental remnant velocity increases with increasing mass loss to a plateau for those products farthest removed from the target. The fractional longitudinal momentum transfer in the plateau is ~ 0.3 . Products in this region are presumably the result of central collisions while those involving much smaller fractional velocity transfer are formed in peripheral interactions [13]. The cascade-evaporation calculation correctly predicts the v_1/v_{CN} values for $\Delta A \sim 10-30$. However, the velocity transfer in peripheral interactions, leading to products close to the target, is severely underestimated. Note in Fig. 5 that the calculation also fails to reproduce the cross sections of the products formed in

Fig. 5. Comparison of the experimental and calculated (o) mass yield distribution for (a) 90 MeV/nucleon $^{12}\text{C} + \text{Cu}$ and (b) 45 MeV/nucleon $^{12}\text{C} + \text{Cu}$.

Fig. 6. Comparison of the experimental (•) and calculated (◊) fractional velocity transfer for Cu + 90 MeV/nucleon ^{12}C as function of mass loss from the target, ΔA .

these peripheral interactions.

C. Exclusive Multifragmentation Studies at the Bevalac

The EOS collaboration (Catania-GSI-Kent State-LBL-Purdue-Texas A&M-UC Davis) is working on an exclusive study of heavy ion reactions at the Bevalac using the EOS time projection chamber as the central detector in a multi-detector system. The experiments include studies of flow effects in symmetric collisions and multifragmentation of heavy projectiles using reverse kinematics. Our first data run took place during two weekends last March and April; a six-week run was completed in early September, and an additional eight-week run is scheduled in late 1992. Depending on the fate of the Bevalac, there may be additional runs in 1993.

Figure 7 shows an integrated display of a typical interaction and the response of the various detector systems. The target (not shown) is located just upstream of the TPC. Light fragments ($Z = 1 \sim 6$) can be identified in the TPC on the basis of dE/dx vs. rigidity (see Fig. 8). The insert in Fig. 7 shows a projection onto the pad plane of all the hits recorded in the TPC in this particular event. As shown on the main diagram, some 30 mass-identified tracks can be associated with these hits.

The TPC is followed by the MUSIC II detector (multiple sampling ionization chamber). This detector consists of three separate ionization chambers each of which performs some fourteen independent dE/dx measurements and provides tracking in either the horizontal or vertical plane. MUSIC II is sensitive to fragments with $Z \geq 8$ and can resolve all charges up to at least gold, the heaviest projectile that we have run so far. Figure 9 shows the remarkable charge resolution of MUSIC II for the fragmentation of 1 GeV/nucleon lanthanum ($Z = 57$) in a plot of number of detected fragments vs fragment Z .

Fig. 7. Integrated display of EOS experiment. The main detectors are shown in plan view and charged reaction products from a typical interaction are tracked. The insert shows a projection of hits in the TPC onto the pad plane.

1 A GeV Au + Au

Fig. 8. Scatter plot of dE/dx vs rigidity in the TPC for 1 GeV/nucleon Au + Au.

92/07/23 23.35

tope169f1-4

z c=2

z c=2

Fig. 9. Charge distribution of fragments from 1 GeV/nucleon La + C detected in MUSIC II. The bottom graph shows an expanded view of Z = 30-40 fragments.

The MUSIC II detector is followed by the TOF wall. The purpose of this array is to bridge the gap between the TPC and MUSIC II by the detection of fragments with $Z \approx 3-10$. The central portion of the array consists of 100 2 m long by 2 cm wide vertical slats arranged into a 50 slat upstream wall backed up by an overlapping 50-slat downstream wall, thereby providing 100% acceptance for projectile fragments. Each slat is coupled to two phototubes located at the top and bottom of the array. The TOF array provides both time-of-flight and pulse height information. In principle, this will provide both fragment mass and charge distributions in the above range, the latter on the basis of dE/dx .

The final detector shown in Fig. 7 is MUFFINS, a multi-segmented neutron detector. This detector is placed behind the TPC at 0° to the beam and detects neutrons from projectile fragmentation. It is operated as a stand-alone detector by the Catania group.

Our chief interest in the various experiments being performed by the EOS collaboration is in exclusive measurements of multifragmentation. We measure projectile breakup by use of reverse kinematics. The kinematic focussing and energy boost provided by the projectile motion ensures that we can detect virtually every charged fragment emitted in a given interaction. As detailed in our last 3-year proposal and proposed originally by Campi [14], these data will permit us to first determine the moments of the charge distribution and then extract the critical exponents associated with the phase transition model of multifragmentation. We have so far measured the breakup of 1 GeV/nucleon Kr, La, and Au projectiles interacting with carbon, and in some instances also beryllium and aluminum. (The probability of multifragmentation depends on the target mass. While carbon should give a reasonable yield of fragmentation events, it may not be the optimal target for all projectiles.) We have obtained close to 3×10^5 events for each of the above beams and are in the process of determining the

number of additional events that will be required to obtain adequate statistics.

D. Time-of-Flight System for STAR

During the past year we participated with other members of the STAR collaboration (from Purdue, Rice, and LBL) in simulations and design of a possible TOF system for the STAR detector at RHIC. The purpose of the TOF detector is to extend the momentum measurements for mass-identified particles to higher momenta than can be achieved in the central TPC [15]. Thus, the π -K and K-p separations will be extended by the TOF detector from an upper limit of 0.65 GeV/c to one of 1.3 GeV/c and from 1 GeV/c to 2.4 GeV/c, respectively. This more extensive momentum interval will permit the determination of more complete transverse momentum spectra and more accurate mean momenta for π , K, and p, as well as more accurate particle ratios.

All the configurations under consideration involved the use of plastic scintillators. The essential difference between them was the placement of the phototube, either inside or outside the 5 kG field of the TPC magnet. The recent availability of mesh dynode tubes [16], which can operate in such a field, makes the more compact in-field phototube configuration a more attractive choice. After extensive simulations we settled on a shingle design, in which the cylindrical outer field cage of the TPC is tiled with 7776 single ended scintillators arranged in 216 trays. Each tray is 2.25 m long (half the length of the TPC), 11.8 cm wide, and supports 36 scintillators. Each shingle scintillator will consist of a 1 inch diameter proximity mesh dynode phototube, e.g., Hamamatsu R3432-01 optically coupled to a $1.5 \times 2.6 \times 23.5$ cm³ Bicron BC404 scintillator. The 36 scintillators will be mounted four abreast in a tray. Figure 10 shows a cross section of the shingle counters and trays. Further details are given in the STAR technical design report [15].

Fig. 10. A cross section of the TOF shingle counters and trays. The radii are measured relative to the center of the beam line.

The estimated cost of the STAR detector relative to the availability of funds indicates that the TOF system cannot be incorporated in the Phase I detector. It will hopefully be added to the detector as funds become available for an upgrade.

References

1. S. Y. Fung, et al., Phys. Rev. Lett. 41, 1592 (1978).
2. W. A. Zajc in Hadronic Multiparticle Production, p. 235, P. Carruthers, Ed., World Scientific Press (1988).
3. T. Akesson, et al., Phys. Lett. B187, 420 (1987).
4. J. D. Dowell, UA1 Collaboration, Proceedings of the VIIth Topical Workshop of Proton-Antiproton Physics, p. 115, R. Raja, A. Tollestrup, and J. Yoh, Ed., World Scientific Press (1988).
5. J. D. Bjorken, Phys. Rev. D27, 140 (1983).
6. T. Alexopoulos et al., Phys. Rev. Lett. 64, 91 (1990).
7. L. Van Hove, Phys. Lett. B118, 138 (1982).
8. S. Y. Cho et al., Phys. Rev. C36, 2349 (1987).
9. S. Y. Cho et al., Phys. Rev. C39, 2227 (1989).
10. J. P. Whitfield and N. T. Porile, submitted to Phys. Rev. C.
11. Y. Yariv and Z. Fraenkel, Phys. Rev. C20, 2227 (1979); Phys. Rev. C24, 488 (1981).
12. I. Dostrovsky, et al., Phys. Rev. 116, 683 (1959).
13. L. Pienkowski, et al., Phys. Rev. C43, 1331 (1991).
14. X. Campi, J. Phys. A19, L917 (1986); Phys. Lett. B208, 351 (1988).
15. Conceptual Design Report for the Solenoidal Tracker at RHIC, The Star Collaboration, Pub.-5347 (1992).
16. G. Finset, et al., NIM A290, 450 (1990).

Publications (September 1991-August 1992)

Target Residues from the Interaction of Silver with 15-45 MeV/nucleon ^{12}C Ions. Y. H. Chung, S. Y. Cho, and N. T. Porile, Nucl. Phys. A533, 170 (1991).

Target Fragmentation of Silver by 14.6 GeV/nucleon ^{16}O Ions. M. Bronikowski and N. T. Porile, Phys. Rev. C44, 1661 (1991).

Comment on "On the Anomalous Interpretation of $^{40}\text{Ar} + \text{Cu}$ Collisions at 0.9 and 1.8 A GeV". G. Dersch, R. Beckmann, G. Feige, T. Lund, P. Vater, R. Brandt, E. Ganssauge, K. Aleklett, E. M. Friedlander, P. L. McGaughey, G. T. Seaborg, W. Loveland, J. Herrmann, N. T. Porile, G. Haase, V. A. Leskin, and K. D. Tolstov, Isotopenpraxis 27, 6 (1991).

A Quark-Gluon Plasma Search in $\bar{p}$ -p at $\sqrt{s} = 1.8$ TeV. T. Alexopoulos, C. Allen, E. W. Anderson, H. Areti, S. Banerjee, P. D. Beery, P. Bhat, J. Bishop, N. N. Biswas, A. Bujak, D. D. Carmony, T. Carter, P. Cole, Y. Choi, R. De Bonte, V. De Carlo, A. R. Erwin, C. Findeisen, A. T. Goshaw, L. J. Gutay, A. S. Hirsch, C. Hojvat, V. P. Kenney, D. Koltick, C. S. Lindsey, C. Loomis, J. M. Lo Secco, T. McMahon, A. P. McManus, N. K. Morgan, K. Nelson, S. H. Oh, J. Pickarz, N. T. Porile, D. Reeves, W. R. Robertson, R. P. Scharenberg, S. R. Stampke, B. C. Stringfellow, M. A. Thompson, F. Turkot, W. D. Walker, C. H. Wang, D. K. Wesson, and H. Zhong, Nucl. Phys. A525, 165c (1991).

Recoil Properties of Target Fragments from the Interaction of Silver with 218 GeV ^{16}O Ions. M. Bronikowski and N. T. Porile, Phys. Rev. C45, 1389 (1992).

Hyperon Production from p - $\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV. T. Alexopoulos et al., submitted to Phys. Rev. D.

Longitudinal Momentum Transfer from Thick-Target Recoil Studies of Intermediate Energy Heavy Ion Reactions. J. P. Whitfield and N. T. Porile, submitted to Nucl. Phys. A.

Evolution of the Reaction Dynamics in the Interaction of Copper with 15-90 MeV/nucleon ^{12}C Ions. J. P. Whitfield and N. T. Porile, submitted to Phys. Rev. C.

Personnel

Professor N. T. Porile, Principal Investigator

Dr. B. Srivastava, Assistant Research Scientist

Working on multifragmentation experiment at the Bevalac and on STAR detector development.

A. Rimai, Graduate Research Assistant

Supported on this grant during past four years. Working on analysis of Fermilab E-735.

J. Whitfield, Graduate Research Assistant

Supported on this grant during past three years. Working on target fragmentation at intermediate energies.

END

**DATE
FILMED**

12 / 1 / 92