2 cyp.

UCRL-20242

MASTER

THE STRUCTURE OF BACTERIOCHLOROPHYLL B

Dea Linda Baumgarten
(Ph. D. Thesis)

December 18, 1970

AEC Contract No. W-7405-eng-48

UCRL

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

LAWRENCE RADIATION LABORATORY
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-20242

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

-ii-

THE STRUCTURE OF BACTERIOCHLOROPHYLL B

Contents

Abstract	iii
List of Abbreviations	iv
I. Introduction	1
II. Experimental	
A. Preparation of the Bacteriochlorophylls	12
B. Instrumentation	14
III. Data and Interpretation	
A. Implications of the Visible Spectrum	16
B. The Mass Spectrum	19
C. The Infrared Spectrum	23
D. Nuclear Magnetic Resonance	27
E. Hydrogenation	38
F. Thin Layer Chromatography of Phytol	39
IV. Discussion	
A. General Considerations of Chlorophyll Structure	41
B. The Structure of Bacteriochlorophyll <u>b</u>	45
C. Suggestions for Further Work	49
V. Conclusion	52
Acknowledgments.	54
References	55

THE STRUCTURE OF BACTERIOCHLOROPHYLL B

Dea Linda Baumgarten

University of California
Berkeley, California

December 18, 1970

ABSTRACT

A bacteriochlorophyll absorbing at longer wavelength than any other known chlorophyll molecule occurs in the photosynthetic bacterium, Rhodopseudomonas viridis. We have developed a procedure for isolating this molecule, called bacteriochlorophyll b, and have measured its visible and infrared absorption spectra, its mass spectrum, and its nuclear magnetic resonance spectrum. The spectral data show that the structure of bacteriochlorophyll b must be very closely related to that of the well-known bacteriochlorophyll a. Our experimental results, when combined with a study of the basic similarities among chlorophyll structures, enable us to propose a structure for the bacteriochlorophyll b molecule; i. e., an unsaturated (4-ethylidene) bacteriochlorophyll a.

LIST OF ABBREVIATIONS

Chl	chlorophyll
Bchl	bacteriochlorophyll
<u>R. viridis</u>	<u>Rhodopseudomonas viridis</u>
<u>R. spheroides</u>	<u>Rhodopseudomonas spheroides</u>
<u>R. rubrum</u>	<u>Rhodospirillum rubrum</u>
MS	mass spectrometry
IR	infrared, or infrared spectroscopy
CD	circular dichroism
NMR	nuclear magnetic resonance

For the numbering of the carbon atoms on the porphyrin ring, see Fig. I-5, p. 10. Side-chain carbon atoms are given primed numbers, i. e., an ethyl group at position 4 on the ring would be numbered -CH₂-CH₃.

4' 4"

I. INTRODUCTION

In the blue-green algae and photosynthetic bacteria one finds quite a number of different chlorophylls whose ability to absorb light ranges from about 300 nm (the blue band of Bchl a) to 1014 nm (the red band of Bchl b), or nearly the whole region in which the transmittance of water is appreciable.¹ The variety of chlorophyll structures found in these organisms, in contrast to the consistent use of chlorophyll a by green plants, probably evolved as a result of brisk competition for sunlight in the primitive aqueous environment to which they were confined.

The first organism discovered to absorb light of wavelength longer than 1000 nm was reported by Eimhjellen² in 1963 and later named Rhodopseudomonas viridis.³ The bacteriochlorophyll was isolated from this organism and found to have a visible spectrum which was similar to that of the well-known Bchl a but shifted about 20 nm to longer wavelength. The newly discovered chlorophyll was called bacteriochlorophyll b. Later, Bchl b was also found in a sulfur bacterium.⁴ In Table I-1, the in vivo absorption of R. viridis is compared with the closely related Rhodopseudomonas spheroides, and the absorption spectra of the two bacteriochlorophylls are compared in acetone. Complete spectra are shown on the following pages (Figs. I-1 and I-2).

	<u>In vivo</u> absorption of the bacteriochl.	Absorption maxima in acetone			
<u>R. spheroides</u> (Bchl <u>a</u>)	875 nm	771	576	390	358
<u>R. viridis</u> (Bchl <u>b</u>)	1014 nm	794	580	407	368

Table I-1

In following the path of the absorbed light energy as it is converted to chemical energy in the bacterium, the study of light-induced absorbance changes of the chlorophylls in the reaction center has been useful. The chromatophores of bacteria contain large amounts of chlorophyll for efficient capture of light energy, but only a small percentage of the molecules are involved in the energy conversion which occurs in the reaction center. In the R. spheroides reaction center, pigments are found which absorb at 870 and 800 nm and are thought to be specialized forms of Bchl, differing from the light-harvesting Bchl perhaps only because of the nature of their immediate environment.⁵ On illumination one observes a reversible bleaching at 870 nm and a shift of the 800 nm band toward the blue. Analogous changes occur in R. viridis, but at longer wavelength.⁶ Table I-2 compares some of the in vivo absorptions of the two systems. From thermodynamic considerations, the maximum chemical potential which an organism can obtain in photosynthesis is a function of the energy of the absorbed light. Thus the R. viridis system, in shifting to longer wavelength than

THE VISIBLE ABSORPTION SPECTRUM OF BCHL. a IN ACETONE

Fig. I-1

XBL 7012-4947

THE VISIBLE ABSORPTION SPECTRUM OF Bchl B IN ACETONE

Fig. I-2

XBL 7012-4943

	Absorption by light-harvesting Bchl	Reaction center pigment subject to reversible photobleaching	Reaction center pigment subject to blue shift
<u>R. spheroides</u>	875 nm	870	800
<u>R. viridis</u>	1014 nm	985	830

Table I-2

R. spheroides, has done so at the sacrifice of about ten percent of the maximum chemical energy.⁷

The R. viridis system has contributed to our understanding of how energy is trapped at the reaction center. Because the photo-bleachable pigment in the reaction center is at comparable or lower energy than the light-harvesting pigment in most organisms, it was thought to act as an energy sink for light absorbed by the bulk of the chlorophyll.⁸ In R. viridis this reaction center pigment absorbs at considerably higher energy than the light-harvesting pigment, and a more complex mechanism must be proposed to explain the high efficiency with which the light energy is retained.⁶

R. viridis may also prove useful for investigating the molecular arrangement of the chlorophylls in reaction centers. Most of our knowledge of this subject comes from circular dichroism measurements,⁵ and greater energy separation between the pigments of the reaction center may allow the CD components to be more clearly resolved.

R. viridis is of potential interest because of its ability to absorb light of very long wavelength, and because of the spectrum

of its reaction center. We propose to determine the structure of the chlorophyll found in this organism. Since the structure of chlorophyll has been one of the great accomplishments of classical organic chemistry as well as the subject of investigation by modern techniques, we will give a brief historical account of the methods used in the determination of chlorophyll structure.

Our understanding of the chemical structure of chlorophyll a, the most extensively studied chlorophyll, evolved over the first half of the twentieth century. It was first studied by the traditional chemical procedure of stepwise degradation, and finally by comparison with a synthetic compound. Although we recognize R. Willstätter to be the pioneer in chlorophyll chemistry, he makes reference to F. Hoppe-Seyler for recognizing that chlorophyll and heme are closely related.⁹ The contemporary structures for heme greatly influenced Willstätter's formulation of chlorophyll. In the work of Willstätter and Stoll¹⁰ in the first decade of this century are summarized the following accomplishments: (1) devising of a method for separating chlorophylls a and b and their derivatives on the basis of partition between ether and hydrochloric acid of varying concentration; (2) determination of the correct empirical formulae for chlorophyll a and for phytol; (3) confirmation of the similarity between chlorophyll and heme on the basis of their degradation products; (4) identification of four different pyrroles derived from chlorophyll, although the more labile substituents did not survive the degradation process;

(5) realization that magnesium is complexed to the nitrogen of the pyrroles; (6) description of phytol as having a hydrocarbon chain containing one double bond and numerous branches, and the suggestion that it might be related to isoprene. Willstätter and Stoll realized that the porphyrin structure must be highly unsaturated and made up of several rings, on the basis of its low hydrogen to carbon ratio. The structure of one chlorophyll derivative and the structure of phytol which Willstatter proposed is shown in Fig. I-3 so that the reader may appreciate the state of knowledge at that time.

Fig. I-3

The actual elucidation of the porphyrin ring structure awaited the work of H. Fischer about twenty-five years later.¹¹

Several proposals for the ring structure were being discussed at that time, and Fischer turned to synthesis to distinguish the correct one. The first synthetic compound to have the characteristic porphyrin spectrum was Etioporphyrin II, shown in Fig. I-4. The ordering of the side groups in the chlorophyll molecule could only be solved by comparison of a chlorophyll derivative with a synthetic compound, and for this step the preparation and resolution of a formidable number of isomers was necessary. The first synthetic compound to be linked to a derivative of chlorophyll was Phylloerythrin, shown in Fig. I-4 below. The location of phytol and the carboxymethyl

Fig. I-4

group quickly followed. The presence of a vinyl group was determined by stepwise conversion to acetyl, and it was located at position 2 by

comparison with several synthetic compounds. The extra hydrogens in chlorophyll were shown to be in ring IV by identification of the optically active imide produced by oxidation. Fischer established the basic structure of bacteriochlorophyll a by dehydration to 2-acetyl-chlorophyll a,¹² and he used degradation to the maleimides to locate the second optically active ring.¹³

The total synthesis of chlorophyll a which was begun by R. B. Woodward in 1956¹⁴ made use of modern methods to obtain high yields, and the structure determination by Fischer was entirely substantiated. The structures of Chl a and Bchl a are shown in Fig. I-5, p. 10.

The use of the various types of spectroscopy in determining chlorophyll structure grew as instrumentation improved and data on organic molecules was collected. Visible spectroscopy has long been a common tool for studying the relationships among porphyrin compounds. The literature on the visible spectroscopy of porphyrins was systematized and added to by A. Stern in a series of papers published between 1934 and 1936 and referred to by G. R. Seely in The Chlorophylls.¹⁵ The infrared spectra of Chl a and numerous derivatives were published by Weigl and Livingston in 1953¹⁶ and by Holt and Jacobs in 1955.¹⁷ Katz and coworkers¹⁸ continued the interpretation of the infrared spectra and did a great deal of work on solvent effects and aggregation. The remarkable effect of the porphyrin ring current on NMR spectra was reported by Becker and Bradley¹⁹ and by Ellis, et al.²⁰ about 1960. Further studies on porphyrin NMR and calculation of

CHLOROPHYLL-A

BACTERIOCHLOROPHYLL-A

PHYTOL

the ring current were done by Abraham.²¹ J. J. Katz made use of the decoupling technique to assign proton resonances in the pheophorbides, and studied the aggregation shifts which occur as a function of concentration.²² Mass spectroscopy is still seldom used for chlorophyll structures, and the most extensive work in the area of porphyrin mass spectroscopy is that of Jackson.²³

In our study of the structure of Bchl b, we will rely almost entirely on spectroscopic methods. These techniques give good and mutually supporting information about the nature of most of the substituents on the porphyrin ring. In the Discussion we will look at the biosynthetic pathway and survey the variations commonly found among chlorophylls in order to determine the positions of these substituents and propose a structure for bacteriochlorophyll b.

II. EXPERIMENTAL

A. Preparation of the Bacteriochlorophylls

The bacteriochlorophyll from Rhodopseudomonas viridis was isolated in small amounts by Eimhjellen et. al. in 1963 by means of circular paper chromatography.² We undertook larger scale preparation with the goal of obtaining 40 mg for NMR experiments. R. viridis was grown in 11.5 liter batches in the medium described by Eimhjellen. After 10 days growth, the bacteria were collected by centrifugation, and 40 to 50 ml of paste was usually obtained. To slow the oxidation process, Bchl b solutions were exposed only to a dim green light, and nitrogen was bubbled through the solutions whenever they were left standing. The following extraction procedure seems to maximize the yield. Add seven parts acetone and one part water to the paste, and blend for about 3 minutes. Allow to stand for half an hour, then filter the dark green acetone extract through paper. Add five percent more water, and filter through fine sintered glass.

Our first attempt at chromatography followed the procedure of Sauer, Lindsay-Smith and Schultz for Bchl a.²⁴ Although polyethylene chromatography with acetone-water provides good resolution of Bchl a, the R. viridis extract yielded only an oily uncrystallizable product by this procedure. Professor Sauer suggested that this bacterium may contain more lipids or colorless materials which are not resolved with polyethylene. Sugar chromatography, a method used by Houssier and

Sauer²⁵ for preparation of protochlorophylls, reverses the order of elution, allowing carotenoids and lipids to be washed out before the chlorophyll. The sugar column, 3.5 cm in diameter and 30 cm in height, is packed dry with powdered sugar (C and H, containing 3% cornstarch) and washed with 1 liter of isooctane. The bacteriochlorophyll extract is transferred to isooctane in a separatory funnel. Then it is washed twice with water to remove most of the acetone, dried over Na_2SO_4 , and concentrated to about 150 ml. The extract is added to the column, and about 10 hours (at about 150 ml per hour) is required to elute the carotenoids. When the eluate is colorless, 0.5% n-propanol is added to the isooctane, and elution of the bacteriochlorophyll proceeds for several hours.

The one serious disadvantage of the sugar column is the long period of time required for elution of the bacteriochlorophyll and the consequent possibility of oxidation. To hinder oxidation, all solvents are degassed with N_2 , and the column is of course kept in the dark. The oxidized material moves very slowly with 0.5% propanol, and only the last fractions show an oxidation peak in the spectrum. The best fractions are washed with water and give a blue-green solid material which is collected by centrifugation and dried. The purity of the sample is judged by the absorbance ratios in the visible spectrum which are most sensitive to the amount of oxidation or the presence of carotenoids. Paper chromatography on cellulose (Eastman Chromagram Sheets, No. 6064) with 20% acetone in petroleum ether was also used to

check the purity. The blue-green Bchl b moves faster than the oxidation products. In impure samples, four or five other bands ranging from bright green to gray could be seen. Dry Bchl b samples oxidize completely in one to two weeks if stored in the dark in a vacuum dessicator. Bchl b fractions in isooctane taken directly from the chromatography column and not crystallized may be kept at -20°C for a month or more.

Bacteriochlorophyll a was prepared in the usual way²⁴ by extraction with acetone and chromatography on polyethylene. It is eluted with 70:30 acetone-water; the pigment crystallizes as the acetone is evaporated. It may be recrystallized from isooctane, and it is stable for many months if stored in the dark and away from air.

B. Instrumentation

The visible spectra were taken on a Cary 14R spectrophotometer in the near infrared mode. The absorbances were measured on freshly prepared acetone solutions with an optical density close to 1.0, or about 10^{-5}M .

Circular dichroism spectra of the Q_y absorption band were taken by Mr. Ken Philipson on a Durrum-JASCO spectrophotometer. The samples were dissolved in ether and had an optical density of about 1.0.

Infrared spectra were measured on a Beckman IR-7 spectrophotometer using a matched pair of 0.1 mm cells. The cells had a capacity of 0.1 ml and contained approximately 4 mg of the pigment dissolved in tetrahydrofuran.

Mass spectra were measured by Dr. Gerald Han on an Associated Electronics Industries model MS 12 mass spectrometer using a direct inlet system at about 270°C.

NMR spectra were recorded on the 220 megacycle instrument manufactured by Varian Associates to which a field sweep decoupling device was attached. The Varian A-60 NMR spectrometer was used for preliminary examination of the samples. Three separate preparations of Bchl b were made for NMR experiments. About 40 mg of the pigment was dissolved in 0.5 to 1 ml of deuterioacetone. Some of the samples contained T.M.S. as an internal standard. In others, the position of the solvent peak, which had previously been measured relative to T.M.S., was used as the standard. The solubility of Bchl b in pure deuteriochloroform is too low for NMR measurements, but we obtained one spectrum in DCCl_3 containing a few percent of deuteropyridine.

III. DATA AND INTERPRETATION

In this section we will compare the data we have obtained for Bchl b with data on Bchl a, which can be interpreted in terms of its known structure, shown in Fig. I-5.

A. Implications of the Visible Spectrum

From Figs. I-1 and I-2 (pp. 3 and 4) one can see that the visible spectrum of Bchl b is very similar to that of Bchl a but shifted approximately 20 nm to the red. The spectrum consists of a long wavelength band around 794 nm, a smaller middle band about 580 nm, and a complex blue band of height comparable to the red band. This pattern is characteristic of a porphyrin which has two saturated rings. In Bchl a, opposite rings are saturated, but it is conceivable that Bchl b might have adjacent rings saturated. However, calculations by Seely²⁶ and by Bernard and Jackman²⁷ indicate that an adjacent tetrahydroporphyrin would not have a band so far in the red. Their arguments are based on symmetry: the opposite tetrahydroporphyrin is more symmetric and lower in energy than the adjacent tetrahydroporphyrin. If the calculations are correct, Bchl b must have opposite rings saturated.

On oxidation, the red band of Bchl b shifts to the shorter wavelength typical of a chlorin. Both Eimhjellen² and Brockmann²⁸ have suggested that the oxidation product of Bchl b is similar, maybe identical, to that of Bchl a. We were unable to confirm the identity

because the reaction is complicated by the presence of at least five oxidation products. The spectrum of one of these products after chromatography is shown in Fig. III-1. A comparison of this spectrum with that of 2-desvinyl-2-acetylchlorophyll a²⁹, which is obtained from oxidation of Bchl a, is made in Table III-1 below. It indicates that the two oxidation products are certainly very similar.

2-Desvinyl-2-acetylchlorophyll <u>a</u>							
$\lambda_{\max}$	677	628	591	538	505	436	388
A_{λ}/A_{677}		.196	.124	.065	.040	1.17	.788
Oxidation product of Bchl <u>b</u>							
$\lambda_{\max}$	677	628	586			436	386
A_{λ}/A_{677}		.152	.095			1.22	.752

Table III-1

The fact that the visible spectrum of Bchl b is red shifted relative to that of Bchl a would be most satisfactorily explained by a change in the nature of one or more of the substituents on the ring. It is also possible that substitution of another metal for Mg could cause such a shift, but Brockmann²⁸ reported that he did find Mg in Bchl b. The possibilities for substituents are considerably limited by the results of the infrared and mass spectra, which follow.

Our conclusions from the visible spectrum are (1) that Bchl b is more reduced than a chlorin, and is probably a bacteriochlorin, with opposite rings reduced, and (2) upon oxidation, Bchl b forms products with absorption spectra characteristic of a chlorin.

THE VISIBLE ABSORPTION SPECTRUM OF AN OXIDATION PRODUCT OF BCHL B

Fig. III-1

XBL 7012-4946

We had hoped to obtain circular dichroism spectra for the two bacteriochlorophylls, but the signal-to-noise ratio was too low for quantitative measurement. Qualitatively, we can say that the rotational strength of the Q_y (red) band of Bchl b is less than half that of Bchl a. Houssier and Sauer³⁰ have reported the CD spectra of a number of chlorophylls and chlorophyll derivatives, and made calculations of the rotational strengths based on an independent contribution from each asymmetric center in the molecule. Their calculations are qualitatively in agreement with experimental data and could help us to select the correct stereoisomer of Bchl b once the substitution pattern is known.

B. The Mass Spectrum

Jackson, et al.²³ have reported mass spectra of a large number of natural and synthetic porphyrin compounds. Some generalizations about the fragmentation patterns of porphin and chlorin free bases which arose from their work are (1) the molecular ion is almost always the most intense peak in the spectrum; (2) side chain fragmentations from porphins are benzylic in character, that is, fission occurs at a bond once removed from the conjugated macrocycle; (3) chlorins lose substituents directly from the reduced pyrrole ring, however; (4) there is very little fragmentation of the porphyrin nucleus itself; (5) doubly charged ions form a prominent series of peaks. Among the specific features mentioned by Jackson was loss of the complete propionate ester side-chain. The $-CO_2CH_3$ group at

position 10 is also easily lost, and the loss of both 10-CO₂CH₃ and 7-CH₂CH₂CO₂CH₃ together produces a strong peak. The 10-CO₂CH₃ group

appears to be able to lose methanol to form $O=C=$. Jackson

and co-workers could not obtain a mass spectrum of chlorophyll a or b, but they did report a weak spectrum for pheophytin a. In addition to the molecular ion, pheophytin a showed a weak ion corresponding to fission of the 10-CO₂CH₃ group, and a somewhat stronger ion at m/e = 278, resulting from fission of the phytol ester. Phytol itself has a molecular weight of 296, but it readily loses the elements of water and forms the fragment with m/e = 278 which is presumably phytadiene³¹.

Mass spectrometry on the intact chlorophyll molecules is hindered by their low volatility. Probably the high temperatures (270 to 300°C) needed to vaporize the molecules cause considerable decomposition, and confused fragmentation patterns result. The mass spectra which we obtained for the two bacteriochlorophylls and their pheophytins, though not entirely rationalizable, did allow us to compare the molecular weights of the porphyrins and also their hydrocarbon tails*. Table III-2, p. 21, shows the major peaks in the mass spectra and their intensities. In the Bchl a and Bchl b compounds, clusters of peaks occur around the same mass numbers, and the major peak in each cluster almost always occurs at the same mass number for the two molecules.

The two pheophytin molecules show a few major differences and frequent

* My thanks to Dr. Gerald Han for obtaining the mass spectra.

m/e	Bchl <u>a</u> (two runs)	Bchl <u>b</u>	Bpheo <u>a</u>	Bpheo <u>b</u>
627	24 - 3	12	----	----
625	34 - 6	25	----	----
623	14 - 2	11	----	----
592	8 - 35	8	40	3
590	3 - 20	5	3	18
575	----	----	----	40
574	----	----	----	100 base
567	50	50	----	----
566	100 base	100 base	5	----
564	29 - 50	46	5	6
552	5	6	40	5
551	7	8	100 base	9
550	7	6	16	21
549	6	5	21	14
548	7	6	6	30
535	16 - 25	18	15	8
534	30 - 50	29	27	11
533	5 - 19	13	70	9
532	8 - 36	25	18	21
516	4	----	----	26
515	2	----	----	40
506	7	6	17	----
505	11	9	30	5
504	6	----	42	11
503	6	----	17	22
488	----	----	6	20
487	----	----	----	40
486	----	----	----	29
479	50 - 21	27	21	----
478	10	10	50	5
477	11	15	34	----
390	34 - 4	12	70	----
389	----	5	150	----
372	0 - 115	----	----	----
311	21 - 5	17	200	----
279	100 - 6 base	74	18	50
278	0 - 8	100 base	100 base	100 base
274	5 - 100 base	26	----	----

Table III-2

discrepancies of one or two mass numbers. The visible spectrum of bacteriopheophytin b, taken just before the mass spectrum, indicated some oxidation of the sample, and that could account for the differences. The data on the bacteriochlorophylls suggests that the molecular weights of the two molecules are identical, at least as far as the heavy atoms are concerned. It is possible, though, for a couple of protons to be gained or lost during ionization³². The presence of a peak at $m/e = 278$ indicates that the esterifying group of Bchl b is phytol. In Table III-3, the major peaks in the bacteriochlorophyll spectra, which are the same for both molecules, are assigned so as to agree as much as possible with the work of Jackson, et. al.²³. The parent peak is assumed to be methyl bacteriopheophorbide, mol. wt. 624, produced after cleavage of phytol. We cannot explain the fact that the intensity of the $m/e = 566$ base peak is very high in both

Major peak	Side groups present	Calc. mass no.
$m/e = 625$	$\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_3$, CO_2CH_3	+ 1 proton
592	$\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_3$, >C=O	+ 1
566	$\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_3$, -----	+ 1
534	-----, CO_2CH_3	- 3
479	-----, -----	+ 1

Table III-3

compounds. The loss of 59 mass units ($-\text{CO}_2\text{CH}_3$) and 146 mass units ($-\text{CO}_2\text{CH}_3$ and $-\text{CH}_2\text{CH}_2\text{CO}_2\text{CH}_3$) makes a fairly strong case for the presence

of a $10\text{-CO}_2\text{CH}_3$ group in the Bchl b molecule. There is no indication in the mass spectra of substituents which might be present in the Bchl b molecule but not in Bchl a. The small fragments such as acetyl, methyl, or ethyl are apparently too weak to show up in our spectra, so it is not surprising that a subtle difference between the two molecules should go undetected.

From the mass spectra we conclude that (1) Bchl a and Bchl b have the same molecular weight to within a couple of hydrogen atoms, (2) both molecules cleave so as to form the phytadiene fragment, and (3) Bchl b contains the C-10 carboxymethyl group and the C-7 propionic ester. Whatever differences there are between the two molecules do not change the basic fragmentation pattern.

C. The Infrared Spectrum

The infrared spectra of the two compounds, Bchl a and Bchl b, are very much alike, as can be seen from Figs. III-2 and III-3. The similarity in the carbonyl region of the spectrum makes a major contribution to our knowledge of the Bchl b structure. The bands in this region are listed in Table III-4 for both molecules in the solvent tetrahydrofuran. The interpretation follows that of Katz for Bchl a¹⁸.

Bchl <u>a</u>	Bchl <u>b</u>
1735 cm^{-1}	1735-6 cm^{-1}
1685	1688-90
1655	1656-7

Table III-4

-24-

THE INFRARED SPECTRUM OF Bchl a IN TETRAHYDROFURAN

XBL 7012-4944

Fig. III-2

THE INFRARED SPECTRUM OF BCHL B IN TETRAHYDROFURAN

Fig. III-3

XBL 7012-4945

The 1735 cm^{-1} peak is the carbonyl stretching vibration of the two esters at positions 7 and 10. If only one ester were present, as in pyrobacteriochlorophyll for example³³, the intensity of this band would be noticeably reduced relative to the other bands. The 1688 cm^{-1} band is assigned to the ketone of ring V. The 1656 cm^{-1} absorption, which typically has about half the intensity of the other two, is assigned to the acetyl group at position 2. An aldehyde function, though it also absorbs at approximately 1656 cm^{-1} in chlorophyll b, is ruled out because it has an extinction coefficient about twice that of acetyl and thus would be comparable in size to the other two bands. The absorptions in the carbonyl region strongly indicate the presence of two esters, a ketone on an intact ring V, and an acetyl group somewhere in the molecule. If the above interpretation is correct, the IR results also exclude any additional carbonyl groups.

The small band shifts in the fingerprint region are uninterpretable except perhaps after the fact, and will not be discussed here. It would be helpful if the region where vinyl group stretching vibrations occur, about $900\text{ to }1000\text{ cm}^{-1}$, would offer some hint of a possible vinyl function, but solvent absorption makes the small changes which occur here unreliable.

It might be useful at this point to sum up the ways in which the infrared and mass spectra work together to determine what substituents are present in the Bchl b molecule. Neither spectrum allows any conclusions to be drawn about the arrangement of these substituents around the porphyrin ring, and that subject will be left for the Discussion. The

mass spectrum tells us that the molecular weights of the two molecules are practically identical, and the infrared spectrum implies that they have the same carbonyl functions. Specifically, the mass spectrum suggests that Bchl b has both a labile $\text{-CO}_2\text{CH}_3$ group and a phytol ester. The infrared supports the presence of the two esters and also indicates a ketone in a position comparable to that of ring V in Bchl a. If the infrared data is accepted as evidence for the acetyl group, there are only a few substituents which can still be considered under the molecular weight restrictions. If we assume that Bchl b is a close relative of Bchl a, some of the most probable substituents are $\text{-CH}_2\text{OH}$, -CH=CH_2 , and >C=CHCH_3 . Thus the structures one might propose for Bchl b are practically limited to the removal of two protons or to simple geometric isomers of Bchl a.

D. Nuclear Magnetic Resonance

Because the NMR spectrum of Bchl b will be of central importance in the Discussion, we will begin this section with some general remarks about chlorophyll NMR. In spite of the large number of protons, the spectrum of a chlorophyll molecule consists mainly of well-resolved lines apart from those associated with the phytol ester. Most of the protons are isolated and do not split each other, and the groups are spread out over a large range of chemical shifts. The spreading out of chemical shifts is partly due to the porphyrin ring current¹⁹, which acts much like the benzene ring current in shifting peripheral substituent group protons to low field. Before we make specific

assignments, it may be helpful to discuss the NMR spectrum in a general way by dividing it into several regions based on chemical shift. In the lowest field region, from 8 to 9 ppm, we expect to find the methine protons, which are bonded directly to the aromatic ring and also experience the largest ring current. The second region, 8 ppm to approximately 3.9 ppm, includes the protons directly attached to the saturated rings II and IV, proton 10, and the vinyl protons of phytol. In the low field methyl region, approximately 3.9 to 2.7 ppm, are found several methyl groups which either show a substantial shift due to ring current or occur downfield because they are part of an acetyl or ester function. Between 2.7 and 1.6 ppm are found the resonances of methyl and methylene protons attached to the saturated rings II and IV. Farthest upfield are the methyl and methylene protons of the saturated part of the phytol chain and protons well removed from the effect of the ring current, such as the 4" methyl of Bchl a.

The 220 megacycle NMR spectra of the two bacteriochlorophylls in deuterioacetone appear in Figs. III-4 and III-5, and the integrated areas are shown in Table III-5. The assignment of Bchl resonances will be based on the work of Katz on methyl bacteriopheophorbide a²² and on the bacteriochlorophyll a spectrum published by Saucer²⁴. The resonances in the methine region are summarized in Table III-6. Integration shows them to have nearly equal areas. In Bchl a, the α proton is assigned to the lowest field resonance on the basis of its proximity to the acetyl function on carbon 2. The β and δ protons occur quite close together.

-29-

THE 220 MEGACYCLE NMR SPECTRUM OF Bchl a IN DEUTEROACETONE

Fig. III-4

XBL 711-5111

-30-

XBL 711-5010

THE 220 MEGACYCLE NMR SPECTRUM OF BCHL B IN DEUTEROACETONE

Fig. III-5a

-30 a-

THE 60 MEGACYCLE NMR SPECTRUM OF AN INDEPENDENT PREPARATION OF BCHL B IN DEUTEROACETONE

Fig. III-5b

-30b-

THE 60 MEGACYCLE NMR SPECTRUM OF AN INDEPENDENT PREPARATION OF BCHL B IN DEUTEROACETONE

Fig. III-5b

Areas are normalized to 12 protons in the low-field methyl region.

Bchl <u>a</u>		Bchl <u>b</u>	
ppm from T.M.S.	area, protons	ppm from T.M.S.	area, protons
8.80	0.81	8.90	2.28
8.38 } 8.36 }	1.48	8.88 } 8.42 }	1.36
5.85	0.46	5.86	1.16
5.04	3.22	5.20 } 5.11 }	1.72
4.73	0.51	4.32	2.39
4.34	3.83	3.93	0.73
4.02 } 3.91 }	1.58		
3.75	3.32	3.77	2.41
3.38	3.32	3.43 } 3.32 }	6.8
3.25	2.55	3.02	2.76
2.96	2.81		
2.80	4.6	2.73	5.6
water		water	
2.30	4.85	2.48 } 2.32 }	6.0
section deleted for acetone peak	(21 protons of phytol)	1.84, 1.81	3.13
		1.68, 1.65	2.94
1.72, 1.69 } 1.60 } 1.54 } 1.25 } 1.08 }	19.7	1.58	3.36
		1.30	
		1.15, 1.12 } 1.08 }	25.3
	5.10	0.88, 0.85	14

Table III-5

Bchl <u>a</u>	Bchl <u>b</u>	Assignment
8.80 ppm	8.90 ppm	α
8.38	8.88	β
8.36	8.42	δ

Table III-6

In Bchl b, one of these latter protons appears to be shifted downfield. The appearance of all three methine protons in the Bchl b spectrum rules out the possibility of alkyl substituents at these positions, as is found in Chlorobium chlorophyll³⁴.

Discussion of peaks in the second region will be postponed until we come to the high field methyl resonances because coupling occurs between protons of those two regions. The peaks that occur in the third region, 3.9 to 2.7 ppm, are summarized in Table III-7. The protons absorbing in this region belong to four isolated methyl groups (see p. 31 for integration), and the assignments are given in the table. Their positions in

Bchl <u>a</u>	Bchl <u>b</u>	Assignment
3.77 ppm	3.77 ppm	C-11 methyl
3.38	3.43	C-1 and C-5
3.37	3.32	methyls
3.00	3.02	C-2 methyl
2.65	2.73	water
(position varies in different samples)		

Table III-7

the two molecules are close enough to confirm the conclusion from IR and MS that Bchl b has an acetyl group and a methyl ester. The assignment for water was based on its temperature dependence and on the integrated areas, which are included in Table III-5 on p. 31.

The general approach we have been using, the comparison of Bchl a and Bchl b spectra, runs into difficulties in the high field region. Since we are working with the chlorophylls rather than their pheophorbides, our spectra include the 15 methyl and 21 methylene protons of phytol. These protons occur upfield from the significant chlorophyll resonances except in the case of Bchl a. In Bchl a, for unknown reasons,* the phytol peaks are shifted downfield by approximately 0.6 ppm. Table III-8 compares the two bacteriochlorophylls and phytol. Because the

Bchl <u>a</u>	Bchl <u>b</u>	Phytol	Assignment
	1.58 ppm	1.68 ppm	vinyl -CH ₃
~ 2.0	1.22	1.23	methylenes of phytol
1.57	0.90 0.82	0.917 0.833	methyls of phytol

Table III-8

phytol band obscures several of the important methyl resonances of Bchl a, we will have to abandon the direct comparison approach, and

* The phytol chain of Bchl a appears to have a different configuration which allows it to experience the ring current more strongly. We do not know what the structural basis for this difference might be. It may be an artifact of our preparation procedure, or associated with the R. rubrum strain which we grow. The shift of the phytol peaks has also been seen by Sauer, Lindsay-Smith and Schultz in our laboratories²⁴. It is surprising that all the upfield protons of phytol experience a uniform shift.

base our assignments on the pheophorbide spectra of Katz, and on our own double resonance experiments.

There are two regions of the NMR spectrum which remain to be discussed. The peaks which occur in the high field methyl region of the Bchl b spectrum and their integrated areas are given in Table III-9. The same information is given for the low field protons, and successful decoupling experiments between protons of the two groups are indicated.

ppm	area		ppm	area
2.48	6.0	coupled	5.86	1.72
2.32			5.20	
2.05			5.11	
1.84	3.13	coupled	4.32	2.39
1.81			3.93	0.73
1.68	2.94	coupled		
1.65				

Table III-9

The assignments which can be made with some certainty in these two regions are the following. The absorptions at 2.48 and 2.32 ppm must include the four methylene protons at positions 7' and 7". The peak at 5.86 ppm is the proton at position 10. The absorptions at 5.20 and 5.11 ppm must include the vinyl proton of the phytol chain. The complex peak at 4.32 ppm must include the methylene of phytol which is bonded to both the oxygen atom and the double bond. These two phytol resonances should be coupled, but since the peaks are complex and

close together, it is not surprising that the coupling could not be demonstrated.

In Bchl a, the upfield region would include the two methyl groups at positions 3 and 8. The methylene part of the ethyl group would be found in the same region as the 7' and 7'' methylenes, and the methyl part of that group would be well upfield around 1.1 ppm. One would predict that the two methyl groups and the 4' and 7' methylenes would be coupled to single protons in the low field region. Decoupling of the 8' methyl and the C-7 proton has in fact been shown by Katz for the methyl pheophorbides a and b^{22b}.

In our Bchl b NMR spectrum, there are two upfield doublets, each of which are coupled to low field protons, and each doublet has roughly the area of a methyl group. The doublet at 1.68, 1.65 ppm is coupled to a proton at 4.32 ppm. Both the coupling constant (7.0 ± 0.5 cps) and the relative shift of the two resonances (2.66 ppm) are about right for a methyl group split by a proton on one of the saturated pyrrole rings. Although the coupling constant for the second doublet at 1.84, 1.81 ppm is the same, the proton to which it is coupled, at 5.20 ppm, is considerably farther downfield; that is, the separation in chemical shift between the coupled protons is 3.38 ppm. This separation is outside the range reported by Katz, 2.49 to 2.70 ppm, for the methyl to proton coupling in the saturated rings of various chlorophyll derivatives^{22a}. Since we have been assuming that the two bacteriochlorophylls are closely related, we will now assume that Bchl b has a methyl substituent at position 8 and assign the normal

1.68, 1.65 ppm doublet to that methyl group. The assumptions we have made will be more fully explained in the Discussion.

The conclusions which we have drawn from NMR are the following:

- (1) the bridge positions α , β , and δ are unsubstituted in Bchl b;
- (2) one C-10 proton is present; (3) Bchl b has the low field methyl substituents at positions 1 and 5; (4) the presence of the C-2 acetyl group and the C-10 methyl ester, suggested by IR and MS, are confirmed in the NMR; (5) the high field bands of phytol occur in the Bchl b spectrum, and their chemical shifts are close to those of free phytol;
- (6) the 7' and 7'' methylenes are found in the Bchl b spectrum; (7) the presence of a methyl group at position 8 is indicated by a doublet around 1.66 ppm coupled to a low field proton. The Bchl b molecule is apparently identical to Bchl a except for ring II.

In assigning the doublet at 1.84, 1.81 ppm, we will consider two possible structures for Bchl b. One is the suggestion made by Brockmann²⁸ that Bchl b retains the methyl and ethyl groups at positions 3 and 4 but has a cis configuration of the protons in ring II. The other suggestion, which originated with Holt³⁵, is that Bchl b has an ethylidene group, >CHCH_3 , at position 4. Neither of these two hypotheses has been previously tested using NMR. If we followed Brockmann's hypothesis, we would assign the doublet at 1.84, 1.81 ppm to a methyl group at position 3, and the low field proton to which it is coupled would differ from the other chlorophylls in that it is cis rather than trans to the proton on carbon 4. The CH_3 protons of the

ethyl group would appear as a triplet about 1.1 ppm, and they would be coupled to a multiplet around 2.2 ppm. We have two arguments against this assignment scheme. First, it is unlikely that the cis configuration would cause a downfield shift of the C-3 proton. In fact, Inhoffen reports that the cis-trans effect on the NMR spectrum of dihydro-octaethylporphyrin is very small and in the opposite direction³⁶. Second, we feel that the triplet at 1.1 ppm which appears in the Bchl b spectrum shown on p. 30 is an artifact or an impurity. The sharp triplet was reduced to a shoulder in the other two Bchl b spectra which we ran on independent preparations (see spectrum on p. 30b). Further, we were unable to decouple this triplet by irradiation in the vicinity of 2.2 ppm, though it is possible that the failure of the decoupling experiment was due to instrumental problems. The presence of an ethyl group received some support from the integration of the peaks around 2.32 ppm and above 1.30 ppm (see table III-5) but because of the large absorption by phytol, the areas in the high field region may be uncertain by several protons.

The hypothesis of the ethylidene group provides a very good explanation for the coupling of the 1.84, 1.81 ppm doublet to a proton at 5.20 ppm, for that proton would now be part of a vinyl group, and the shift to low field would be reasonable. The presence of a double bond at carbon 4 would also provide a nice explanation for the low field position of two of the three methine resonances at 8.90 and 8.88 ppm which we would now assign to protons α and β . The major weakness

in this assignment scheme is that we have no sharp doublet in the spectrum which we can assign to the C-3 methyl group. A methyl group at carbon 3 is of course necessary because of the molecular weight results. Possibly this methyl group has experienced a downfield shift from its usual position due to the nearby double bond, and is located under the acetone peak at 2.05 ppm. Transfer of the sample to DCCl_3 - D-pyridine revealed a broad multiplet around 2.02 ppm which we attempted but failed to decouple. Since the broad peak could well be residual acetone, this experiment was inconclusive. It is also possible that this methyl group doublet could be obscured by one of the phytol bands. The two hypotheses for the structure of Bchl b will be looked at in broader perspective in the Discussion.

E. Hydrogenation

An attempt was made to determine whether or not Bchl b possesses an ethylidene group by hydrogenation. Positive evidence for such a group would be the blue shift of the visible spectrum after hydrogenation. If indeed Bchl b differs from Bchl a only by the presence of this additional double bond, hydrogenation should yield a product which will co-chromatograph with Bchl a.

Chl a was used as a control for the microhydrogenation experiment using palladium on carbon in dioxane. It took up $2/3$ mole of H_2 , and the red band of the visible spectrum shifted from 657 to 650 nm. Under the same conditions, Bchl b took up a small amount of H_2 but showed no change in its visible spectrum. After continuation of the hydrogenation

procedure for 1 hour with PtO_2 catalyst under 3 atm. of H_2 , the spectrum of Bchl b was still unchanged. Dilution to 50% methanol and hydrogenation under 3 atm. for 1 hour longer resulted in a shift of the red band to 610 nm. The absorption at 610 nm is probably due to a compound in which ring V has been opened.

The hydrogenation experiment does support our earlier results that Bchl b does not have a free vinyl group. But regarding the possibility of an ethylidene group it is inconclusive, for such a group is more sterically hindered than a vinyl, and may resist hydrogenation. This was apparently the case for the similarly hindered double bond in phytol, which was not reduced in the micro-hydrogenation of either Chl a or Bchl b.

F. Thin Layer Chromatography of Phytol

Phytol can be removed from chlorophyll molecules by dissolving in 2.5% KOH in methanol and heating for 20 minutes at 66°C . The phytol may be recovered by extraction into ether followed by evaporation of the solvent. Bacteriochlorophyll b is oxidized by this procedure and the products were not recovered.

The procedure of Shimizu, Fukushima, and Tamaki was used for the detection of phytol³⁷. The phytol is dissolved in a small amount of petroleum ether and spotted on a silica gel sheet (Eastman Chromagram Sheet no. K301R). The chromatogram is developed with 19:1 benzene-ethyl acetate. After drying, the sheet is sprayed with 0.25% aqueous KMnO_4 . The phytol shows up as a brown spot on a purple background.

The alcohol which was obtained from Bchl b was run in parallel with a solution of 1% commercial phytol in petroleum ether. Both compounds had an R.F. value of 0.45 ± 0.05 when the chromatogram was developed with 19:1 benzene - ethyl acetate. Chromatograms were also developed with 85:15 isooctane - ethyl acetate, and both compounds had an R.F. between 0.3 and 0.4.

This experiment supports the hypothesis that Bchl b contains phytol, though isomers of phytol would probably not be detected by this procedure. It does not contain farnesol. The experiment does prove that the alcohol from Bchl b definitely contains a carbon-carbon double bond, for the development of the spot by KMnO_4 relies on the oxidation of a double bond.

IV. DISCUSSION

A. General Considerations of Chlorophyll Structure

The discussion will begin with a look at the biosynthesis of chlorophyll and at the variations in chlorophyll structures which are commonly found. These considerations will allow us to discount the likelihood of many of the conceivable isomers of the chlorophyll ring system and locate the positions where changes in the substitution pattern are most probable.

The biosynthetic pathway to chlorophyll a in plants has been reviewed by L. Borograd in The Chlorophylls³⁸. The porphyrin ring is formed by enzymatic joining of four monopyrroles. It has been shown that only one kind of pyrrole is used in the formation of porphyrin, and this molecule has one acetic and one propionic acid side chain³⁹. The condensation of this pyrrole, porphobilinogen, to form Urogen III is shown below (Fig. IV-1). Logically, four porphobilinogen molecules

Fig. IV-1

can condense to urogen in four different ways. The symmetrically substituted molecule, Urogen I, which has a four-fold axis of rotation, is known to exist in plants, but it is only the very unsymmetric Urogen III which is useful in the later steps of chlorophyll synthesis. After formation of the urogens, decarboxylation of the acetic acid substituents to methyl groups takes place. The resulting compounds are called Coprogen I and III. In the next step, Coprogen III is converted to Protoporphyrinogen IX by an enzyme which selects against the symmetric Coprogen I isomer. This molecule contains two vinyl and two propionic acid groups, and is shown in Fig. IV-2. Before the esterification steps can occur, Protoporphyrinogen IX must be oxidized to the extensively conjugated Protoporphyrin IX, which is capable of chelating a magnesium ion. Magnesium Protoporphyrin IX is the substrate for formation of Mg Protoporphyrin IX monomethyl ester, shown in Fig. IV-2.

Fig. IV-2

At this step, the esterification of just one propionic acid group is selected enzymatically. The esterified side group is then involved in the formation of the cyclopentanone ring, ring V. Reduction of the vinyl group at position 4 probably occurs after closure of ring V⁴⁰. Thus, the last steps in the synthesis of Chl a are reduction of one vinyl group, esterification with phytol, and reduction of the double bond between carbons 7 and 8. Since pigments related to Chl a are found in R. spheroides, it has been suggested that Bchl a is formed from Chl a⁴¹. This process would have to involve the oxidation of the 2-vinyl group, possibly through the intermediate, α -hydroxyethyl, and reduction of ring II.

From the point of view of evolution, any change in the structure of a chlorophyll becomes progressively less probable as it occurs in earlier stages of the biosynthesis. All chlorophylls are likely to have the first stages in common. In fact, we wish to assume that the only changes which are probable enough to be considered are (1) opening of ring V, (2) substitution of another metal for magnesium, (3) a free acid at position 7, or esterification with an alcohol other than phytol, (4) altered pattern of reduction, or a change in the asymmetry at rings II and IV, (5) altered substituents at carbon atoms 2 and 4, and possibly oxidation of some of the methyl groups. Let us see how well this assumption bears up in examination of naturally occurring chlorophylls.

The reader is referred to Table IV-1 on p.44 which compares the basic structure and substituents of known chlorophylls. It should be

Chlorophyll	Presence of Ring V	Mg	Esterifying Group	Reduction of Ring IV	Ring II	C-10 -CO ₂ CH ₃	Substituents at C-2	C-4	Other	Reference
Chl <u>a</u>	Yes	Yes	Phytol	Yes	No	Yes	vinyl	ethyl		15
Chl <u>b</u>	Yes	Yes	Phytol	Yes	No	Yes	vinyl	ethyl	formyl group at carbon 3	15
Bchl <u>a</u>	Yes	Yes	Phytol	Yes	Yes	Yes	acetyl	ethyl		15
Chlorobium chls	Yes	Yes	Farnesol	Yes	No	No	α- hydroxy -ethyl	ethyl, n-propyl, or iso- butyl	methyl or ethyl group at C-5; methyl or ethyl group at bridge position 8	43
Chl <u>d</u>	Yes	Yes	Phytol	Yes	No	Yes	formyl	ethyl		43
Chl <u>c</u>	Yes	Yes	none: free acid	No	No	Yes	vinyl	vinyl or ethyl		42
Protochls	Yes	Yes	Phytol	No	No	Yes	vinyl	vinyl or ethyl		25

Table IV-1

pointed out that only the structure of chlorophyll a is known by total synthesis. The substitution patterns of several other chlorophylls have been definitely established by chemical conversion to Chl a or its derivatives, and some are based on analogy to the common chlorophyll structure. From Table IV-1 it can be seen that the known chlorophylls all contain ring V and magnesium. These apparently are not points of variation in chlorophyll structure. Chlorophyll c contains a free acid group, and the Chlorobium chlorophylls are esterified with farnesol. Hence, variation in the hydrocarbon tail has to be considered. In the level of reduction of rings II and IV, all combinations occur except for the saturation of ring II alone. The substituents at carbons 2 and 4 vary, but with a couple of exceptions, they contain two carbon atoms. Chl b has a formyl group instead of a methyl at carbon 3. The Chlorobium chlorophylls have several exceptional changes which occur as a group: (1) a saturated hydrocarbon substituent at position 8; (2) lack of the C-10 carboxymethyl group; (3) esterification with farnesol. Thus in arriving at a proposal for the structure of Bchl b, we can feel safe in ignoring fundamental changes in the porphyrin structure, and concentrate on possibilities 1 to 5 in the paragraph above.

B. The structure of Bacteriochlorophyll b

We have inferred that Bchl b has the carbocyclic ring V on the basis that (1) proton 10 is present in the NMR, (2) the $-CO_2CH_3$ group is easily split off in the mass spectrum, (3) the C-9 carbonyl resonance occurs at the expected frequency in the infrared. A. S. Holt reports that the

molecule gives a positive phase test, for which an intact ring V is essential⁴³. We have assumed that magnesium is present in Bchl b as it is in other chlorophylls. Brockmann confirmed the presence of Mg by precipitating it with titanium yellow²⁸. Evidence from the mass spectrum strongly suggests esterification by phytol. The presence of phytol is supported by the high-field peaks in the NMR spectrum and, by thin layer chromatography. The visible spectrum of the Bchl b molecule makes necessary the reduction of two opposite pyrrole rings. We assume that these are the customary rings, II and IV. The stereochemistry of the reduced rings, or of position 10, is not established by our data, but we will assume that the carbons in ring IV and position 10 have their common configuration. Bchl b does not appear to possess any of the unique structural characteristics of the Chlorobium chlorophylls. NMR tells us that the bridge positions are unsubstituted, and that the C-10 carboxymethyl group is present. The NMR spectrum also indicates the presence of the two low-field methyl groups which are regularly found at carbon atoms 1 and 5. Evidence for an acetyl group is found in both the IR and NMR. The acetyl group is most likely to occur at carbon 2 because of the overall similarity of this molecule to Bchl a. There are coupled resonances in the NMR spectrum which may be assigned to proton 8 and to the methyl protons 8'. Thus the substituents and stereochemistry in ring II are the only variables which remain.

We know from the mass spectrum that the remaining substituents must add up to a molecular weight of 44 ± 2 ; further, the infrared spectrum will not support an additional carbonyl group in the molecule. Consider-

ation of the biosynthetic pathway and the basic similarity of most chlorophyll structures leads us to expect a methyl group at position 3 and a two-carbon substituent at position 4, or else groups which can be easily derived from the above by oxidation. Combinations which are allowed under the above restrictions are (1) $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$ (cis arrangement), (2) $-\text{CH}_3$, $-\text{CH}=\text{CH}_2$, (3) >CH_2 , $-\text{CH}_2\text{CH}_3$, (4) $-\text{CH}_3$, >CHCH_3 , and possibly (5) $-\text{CH}_3$, CH_2OH . Of these possibilities, we can rule out (2) on the basis that a vinyl group would have been detected in the hydrogenation experiment, and additional low field protons would be expected in the NMR. We rule out (3) because its NMR spectrum would not show an upfield doublet coupled to a low-field proton. The hydroxyl group in (5) would be expected to show an additional low field proton in the NMR. Structure (5) is also rejected on the basis of the low field position in the NMR of the proton coupling the methyl group at position 3, which we will discuss in connection with structure (1).

Of the two hypotheses which remain, Brockmann and Kleber²⁸ proposed the first one: that Bchl b is identical to Bchl a except in having the cis configuration of protons in ring II. The central point of Brockmann's work was the crystallization of dehydro-bacteriomethylpheophorbide b and establishment of its identity with 2-acetyl-2-desvinyl-methylbacteriopheophorbide a on the basis of melting point. That is, both Bchl a and Bchl b yield the identical compound after oxidation to the chlorin and removal of Mg and phytol. These authors further observed that on treatment of Bchl b with 0.1 N HCl, 20% bacteriopheophytin a is produced along with bacteriopheophytin b. Brockmann suggested that the cis configuration in

ring II could explain the red shift of the visible spectrum of Bchl b over that of Bchl a³⁶ and also the relatively easy oxidation of the Bchl b molecule.

We would like to suggest that the alternate hypothesis, the presence of an ethylidene group at position 4, is also consistent with Brockmann's observations. With the ethylidene structure, the "oxidation" of Bchl b to the chlorin would indeed be easy. It would be a tautomerism, rather than a real oxidation, involving only the relocation of a carbon-carbon double bond. It may be catalyzed by oxygen and/or light. Since formation of the chlorin cannot occur without the loss of the double bond between carbons 4 and 4', Brockmann, in working with the oxidized molecules, would not have been able to detect the difference between the ethylidene structure and the one he proposed. In the preparation of bacteriopheophytin b, the acidic conditions might facilitate partial reduction of the ethylidene double bond, perhaps by disproportionation, to form bacteriopheophytin a.

We also suggest that the NMR data on Bchl b fits better the hypothesis of an ethylidene group than the isomerized structure of Bchl a. This conclusion is based on the large chemical shift between the doublet at 1.84, 1.81 ppm and the proton at 5.20 ppm to which it is coupled. The doublet would be assigned to the 4" methyl group of the ethylidene function, and the single proton would appear at low field because it is next to a double bond. The ethylidene hypothesis accounts for the red shift of the visible spectrum for it contributes another site of unsaturation one carbon removed from the porphyrin skeleton. It also explains

the easy "oxidation" of the Bchl b molecule, and the observation that the oxidation products of the two bacteriochlorophylls may well be identical. We would like to propose the structure shown in Fig. IV-3 below for the bacteriochlorophyll b molecule.

Fig. IV-3

C. Suggestions for Further Work

We have not tried to determine the stereochemistry at carbon atom 3. Since the classical method of degradation to the optically active imide is quite likely to destroy the ethylidene function, precise measurement of the circular dichroism spectrum may prove to be the best way of obtaining this information. Mr. Ken Philipson has extended the comparison of calculated and experimental rotational strengths made by Houssier and

Sauer³⁰ to Bchl a⁴⁴. Using his estimates of the contribution which each asymmetry center makes to the total rotational strength, we can calculate the rotational strength expected for both configurations at C-3. The results are shown in Table IV-2. Both configurations are consistent with our qualitative CD measurement, but should be easily distinguishable if we can obtain quantitative results.

(Q _y band)	C-3	C-4	C-7	C-8	C-10	Calculated sum, (x10 ⁴⁰ cgs)	Experimental R _A (x10 ⁴⁰ cgs)
Bchl <u>a</u>	1.9	1.9	-3.3	-2.3	5.2	3.4	5.0
Bchl <u>b</u> , normal config. at C-3	1.9	0	-3.3	-2.3	5.2	1.5	absolute value less than 1.7
Bchl <u>b</u> , reversed config. at C-3	-1.9	0	-3.3	-2.3	5.2	-2.3	

Table IV-2

Successful hydrogenation of the ethylidene group and observation of a blue shift in the visible spectrum would be the most direct way of confirming our proposed structure for Bchl b. If the asymmetry at carbon 3 is the same as for Bchl a, as we might expect, the two molecules should co-chromatograph after hydrogenation. Hydrogenation of the ethylidene double bond is not so easy as for a vinyl group, but we might be able to accomplish it after exploration of various solvents and experimental conditions.

Preparation of the pheophorbide of Bchl b, and location of the additional upfield doublet due to the C-3 methyl group in the NMR would be satisfying evidence for our proposed structure.

Finally, we would like to be able to prepare the pheophorbide derivative of Bchl b so as to confirm Brockmann's result that it is identical to 2-desvinyl-2-acetyl-pheophorbide a.

V. CONCLUSION

The purpose of this work was to study by physical methods the structural features of the Bchl b molecule and finally to propose a structure for this molecule which is consistent with our data and with what we know about the structure of chlorophylls in general. Since Bchl b is found in a bacterium which is closely related to those which normally contain Bchl a, and since the visible spectra of the two bacteriochlorophylls are very similar, we took the approach of comparing data on the two molecules whenever it was possible.

We inferred from the visible spectrum that Bchl b has the bacteriochlorin skeleton. The molecular weight of the molecule was set within narrow limits by measurement of its mass spectrum. The mass spectrum also helped to establish the presence of two ester groups and the identity of the long-chain alcohol. The infrared spectrum defined the number and the nature of the carbonyl groups, and the protons adjacent to these carbonyl groups were then easily assigned in the NMR spectrum. In the NMR spectra, all of the protons which are found in Bchl a could be assigned in Bchl b except for those on ring II. One of the protons in the Bchl b spectrum which was coupled to a methyl group occurred at a lower field than was observed for any group in Bchl a. We proposed that Bchl b contain an ethylidene function to account for this low-field resonance.

Assignment of the various substituent groups to specific locations on the bacteriochlorin skeleton was made on the assumption that basic changes in the biosynthetic pathway to chlorophyll are very unlikely to

↓

occur. Brockmann's success in relating the dehydro-bacteriomethyl-pheophorbide b derivative to a known derivative in the Bchl a series supported the analogy which we made between the structures of the two bacteriochlorophylls.

ACKNOWLEDGMENTS

I would like to thank Professor Kenneth Sauer, for without his guidance and encouragement, much of this work would not have been completed. My thanks go also to Dr. Gerald Han for making time in a busy schedule to do mass spectrometry of the bacteriochlorophylls, to Mr. Ken Philipson for several battles against instrumental deficiencies to obtain circular dichroism measurements, and to Dr. Ercole Cavaliere for his spontaneous interest in our nuclear magnetic resonance studies. This work was supported by a National Science Foundation grant, GB-6738, and by the United States Atomic Energy Commission.

REFERENCES

1. J. M. Olson, Science 168, 438 (1970).
2. K. E. Eimhjellen, O. Aasmundrud, and A. Jensen, Biochem. Biophys. Res. Comm. 10, 232 (1963).
3. J. M. Olson, J. W. Carroll, M. L. Clayton, G. M. Gardner, A. E. Linkins, III, and C. M. C. Moreth, Biochem. Biophys. Acta, 172, 338 (1969).
4. K. E. Eimhjellen, H. Steensland, and J. Traetteberg, Arch. Mikrobiol. 59, 82 (1967).
5. K. Sauer, E. A. Dratz, and L. Coyne, Proc. Nat. Acad. Sci. 61, 17 (1968).
6. A. S. Holt and R. K. Clayton, Photochem. Photobiol. 4, 829 (1965).
7. R. T. Ross and M. Calvin, Biophys. J. 7, 595 (1967).
8. W. L. Vredenberg and L. N. M. Duysens, Nature 197, 355 (1963).
9. F. Hoppe-Seyler, Zeitschr. f. Physiol. Chem. 3, 339 (1879).
10. R. Willstätter and A. Stoll, Investigations on Chlorophyll, transl. by F. M. Schertz and A. R. Merz, The Science Press Printing Company, Lancaster, 1928.
11. H. Fischer and H. Orth, Die Chemie des Pyrrols, Vol. II, part 1, 1937, and H. Fischer and A. Stern, Vol. II, part 2, Akademische Verlagsgesellschaft M. B. H., Leipzig, 1940.
12. H. Fischer, W. Lautsch, and H. K. Lin, Liebigs Ann. Chem. 534, 1 (1938).
13. H. Fischer and H. Wenderoth, Liebigs Ann. Chem. 637, 179 (1939).
14. R. B. Woodward, Pure Appl. Chem. 2, 383 (1961); Angew. Chem. 72, 651 (1960).

15. G. R. Seely in The Chlorophylls, ed. by L. P. Vernon and G. R. Seely, Academic Press, New York, 1966, Ch. 3.
16. J. W. Weigl and R. Livingston, J. Am. Chem. Soc. 75, 2173 (1953).
17. A. S. Holt and E. E. Jacobs, Plant Physiol. 30, 553 (1955).
18. a. J. J. Katz, R. C. Dougherty, and L. J. Boucher in The Chlorophylls, ed. by L. P. Vernon and G. R. Seely, Academic Press, New York, 1966, Ch. 7.
b. J. J. Katz, G. L. Closs, F. C. Pennington, M. R. Thomas, and H. H. Strain, J. Am. Chem. Soc. 85, 3801 (1963).
19. E. D. Becker and R. B. Bradley, J. Chem. Phys. 31, 1413 (1959).
20. J. Ellis, A. H. Jackson, G. W. Kenner, and J. Lee, Tetrahedron Letters 2, 23 (1960).
21. a. R. J. Abraham, A. H. Jackson, and G. W. Kenner, J. Chem. Soc. p. 3468 (1961).
b. R. J. Abraham, Mol. Phys. 4, 145 (1961).
22. a. J. J. Katz, R. C. Dougherty, and L. J. Boucher in The Chlorophylls, ed. by L. P. Vernon and G. R. Seely, Academic Press, New York, 1966, Ch. 7.
b. G. L. Closs, J. J. Katz, F. C. Pennington, M. R. Thomas, and H. H. Strain, J. Am. Chem. Soc. 85, 3809 (1963).
23. A. H. Jackson, G. W. Kenner, K. M. Smith, R. T. Alpin, H. Budzikiewicz, and C. Djerassi, Tetrahedron 21, 2913 (1965).
24. K. Sauer, J. R. Lindsay-Smith, and A. J. Schultz, J. Am. Chem. Soc. 88, 2681 (1966).

25. C. Houssier and K. Sauer, *Biochem. Biophys. Acta*, 172, 476 (1969).
26. G. R. Seely, *J. Chem. Phys.* 27, 125 (1957).
27. J. R. Barnard and L. M. Jackman, *J. Chem. Soc.* p. 1172 (1956).
28. H. Brockmann, Jr. and I. Kleber, *Tetrahedron Letters* 25, 2195 (1970).
29. J. R. Lindsay-Smith and M. Calvin, *J. Am Chem. Soc.* 88, 4500 (1966).
30. C. Houssier and K. Sauer, *J. Am. Chem. Soc.* 92, 779 (1970).
31. C. Liljenberg and G. Odham, *Physiologia Plantarum* 22, 686 (1969).
32. H. Budzikiewicz and S. E. Drewes, *Liebigs Ann. Chem.* 716, 222 (1968).
33. Unpublished work of Dr. Shirley Tsai.
34. R. B. Woodward and V. Škarić, *J. Am. Chem. Soc.* 83, 4676 (1961).
35. A. S. Holt, private communication.
36. H. H. Inhoffen, J. W. Bucher, and R. Thomas, *Tetrahedron Letters* 14, 1145 (1969).
37. S. Shimizu, H. Fukushima, and E. Tamaki, *Phytochem.* 3, 641 (1964).
38. L. Borograd in *The Chlorophylls*, ed. by L. P. Vernon and G. R. Seely, Academic Press, New York, 1966, Ch. 15.
39. D. Shemin and J. Wittenberg, *J. Biol. Chem.* 192, 315 (1951).
40. O. T. G. Jones, *Biochem. J.* 89, 100 (1963).
41. a. W. R. Sistrom, M. Griffiths, and R. Y. Stanier, *J. Cellular Comp. Physiol.* 48, 459 (1956).
b. M. Griffiths, *J. Gen. Microbiol.* 27, 427 (1962).
42. R. C. Dougherty, H. H. Strain, W. A. Svec, R. A. Uphaus, and J. J. Katz, *J. Am. Chem. Soc.* 88, 5037 (1966).

43. A. S. Holt in The Chlorophylls, ed. by L. P. Vernon and G. R. Seely, Academic Press, New York, 1966, Ch. 4.
44. K. Philipson, S. Tsai, and K. Sauer, J. Phys, Chem., submitted, 1970.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE RADIATION LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720