Technical Progress Report

Substantial progress has been achieved on question I.1 about the dynamics of predator-prey systems in a patchy environment described on page 20 of the grant proposal. The approaches used have been the ones described in the proposal and some new avenues that have been found during the year. Work on question I.2 on competition will begin shortly.

First, a new phenomenon that might contribute to outbreaks in systems of discrete patches has been determined using a discrete time model with both spatial and age structure. As described in the attached preprint, differences in dispersal rates and density dependence at different ages may lead to chaotic behavior. This discovery implies the importance of coupling age and spatial structure, and is described completely in the preprint.

I have also written a short note which will appear in the journal *ECOLOGICAL* describing the difference between different modelling approaches for including spatial structure. Appreciation of these differences is the first step for understanding the factors leading to local outbreaks of insects.

A model for a single species in a patchy environment with migration, local population growth and disasters within patches, including the possibility of disasters which remove only some of the individuals within a patch has been formulated. Additionally, the equilibrium states of this model have been determined, and sufficient conditions for stability have been found. A manuscript reporting these findings is under preparation. Currently, simulations are being performed to study this model further, and to extend this model to include the case of a host and parasitoid in a patchy environment. Also, the distributions of numbers within patches generated by the model will be compared to data from entomological studies. A brief description of the model follows.

The model could be phrased in terms of a density function for the fraction of patches with n individuals in them. However, it is simpler to use the following device to change the model into one where patches are defined in terms of *age*, and not *size*. I will define the *age* of a patch with n individuals to be the time it takes for a patch settled by the initial propagule size to grow to have n individuals in it, given that the patch has suffered no disasters. Under the assumptions about local growth dynamics given above, this device provides a unique way to assign ages and sizes of patches. Let $p(t, a)$ be the

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

fraction of patches of 'age' a , $\mu(a) = \mu_t(a) + \mu_p(a)$ be the rate of all disasters in a patch of age a , with μ_t the rate of 'total' disasters, disasters which remove all individuals within a patch, and μ_p the rate of partial disasters, disasters which remove only some proportion of the individuals. Partial disasters need not be of a fixed size, so let $\gamma(\alpha, a)$ be the probability that a disaster in a patch of 'age' α leads to a patch of 'age' a . Consequently,

$$\int_0^\infty \gamma(\alpha, a) da = 1 \quad (1)$$

for all α . Finally, define a total 'size' of the population by $s(t) = \int_0^\infty \sigma(a)p(t, a)da$ where $\sigma(a)$ is a measure of the 'size' of patches of 'age' a , which enters into the model in defining the rate at which empty patches are colonized:

$$p(t, 0) = \int_0^\infty p(t, a)M(a, s(p(t, \cdot)))da \quad (2)$$

Here, $M(a, s(p(t, \cdot)))$ is the rate at which empty patches are colonized due to colonists from a patch of 'age' a when the total 'size' of the population is given by $s(t)$.

The dynamics within patches are thus governed by two features: local aging and disasters. These two events are incorporated within the following equation:

$$\frac{\partial p}{\partial t} + \frac{\partial p}{\partial a} = -\mu(a, p(t, \cdot))p + \int_0^\infty \mu_p(\alpha, p(t, \cdot))p(t, \alpha)\gamma(\alpha, a)d\alpha \quad (3)$$

The solution of (3) along characteristics leads to the following equation,

$$p(t, a) = p(t-a, 0)e^{-\int_0^a \mu(\alpha)d\alpha} + \int_0^a \left[\int_0^\infty \mu_p(\alpha, p(t, \cdot))p(t-v, \alpha)\gamma(\alpha, a-v)e^{-\int_{a-v}^a \mu(\beta)d\beta}d\alpha \right] dv \quad (4)$$

Denote the equilibrium, i.e. time independent solution of (4), by $\hat{p}$. This equilibrium satisfies the equation

$$\hat{p}(a) = \hat{p}(0)e^{-\int_0^a \mu(\alpha)d\alpha} + \int_0^a \left[\int_0^\infty \mu_p(\alpha, \hat{p}(\cdot))\hat{p}(\alpha)\gamma(\alpha, a-v)e^{-\int_{a-v}^a \mu(\beta)d\beta}d\alpha \right] dv \quad (5)$$

Now, I will define the stable patch distribution, $f(a)$, by

$$f(a) = \frac{\hat{p}(a)}{\hat{p}(0)}, \quad (6)$$

and I will define the probability of survival to age a as

$$l(a) = e^{-\int_0^a \mu(\alpha) d\alpha}. \quad (7)$$

Then f satisfies the equation:

$$f(a) = l(a) + \int_0^a \int_0^\infty \mu_p(\alpha, \hat{p}(\cdot)) f(\alpha) \gamma(\alpha, a-v) \frac{l(a)}{l(a-v)} d\alpha dv \quad (8)$$

This equation is linear if μ and μ_p are independent of $p(\cdot)$, so I now make that assumption, and indicate this by dropping the second variable in the functions μ and μ_p .

Finally, define

$$g(a) = \mu_p(a) f(a) \quad (9)$$

so

$$g(a) = \mu_p(a) l(a) + \int_0^a \int_0^\infty \mu_p(a) g(\alpha) \gamma(\alpha, a-v) \frac{l(a)}{l(a-v)} d\alpha dv. \quad (10)$$

Rewrite this by changing the order of integration as:

$$g(a) = \mu_p(a) l(a) + \int_0^\infty \mu_p(a) g(\alpha) \int_0^a \gamma(\alpha, a-v) \frac{l(a)}{l(a-v)} dv d\alpha. \quad (11)$$

and then by letting $z = a - v$ as:

$$g(a) = \mu_p(a) l(a) + \int_0^\infty \mu_p(a) g(\alpha) l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz d\alpha, \quad (12)$$

which is a Fredholm integral equation of the form

$$g(a) = q(a) + \int_0^\infty g(\alpha) K(\alpha, a) d\alpha \quad (13)$$

where

$$K(\alpha, a) = \mu_p(a) l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz. \quad (14)$$

$$q(a) = \mu_p(a) l(a) \quad (15)$$

The analysis of (13) is facilitated by the fact that the kernel $K(\alpha, a)$ is nonnegative and continuous. Define an eigenvalue λ and eigenfunction u as a nontrivial solution of the homogeneous equation:

$$u(a) = \lambda \int_0^\infty u(\alpha) K(\alpha, a) d\alpha \quad (16)$$

Since the kernel $K(\alpha, a)$ is nonnegative and continuous, there exists a smallest positive eigenvalue, denoted Λ (Zabreyko et. al., 1975). I will make use of the following theorem (Zabreyko et. al., 1975):

Theorem 1 *Consider the non-homogenous equation*

$$u(a) = q(a) + \lambda \int_0^\infty u(\alpha) K(\alpha, a) d\alpha \quad (17)$$

where $K(\alpha, a)$ is continuous. Let $\lambda < \Lambda$. Then for any nonnegative function $q(a)$, the equation (17) has a unique nonnegative solution u^ which can be obtained by the method of successive approximations:*

$$u_{n+1}(x) = q(a) + \lambda \int_0^\infty u_n(\alpha) K(\alpha, a) d\alpha, \quad (18)$$

where the initial approximation u_0 is arbitrary. The rate of convergence of the successive approximations is described by:

$$\sup_a |u^*(a) - u_n(a)| \leq C(\lambda) \left(\frac{\lambda}{\Lambda} \right)^n \quad (19)$$

where $C(\lambda)$ is a constant.

Now note that (13) is an equation of the form (17) with $\lambda = 1$. I will now show that for the homogenous equation corresponding to (13) that the smallest positive eigenvalue $\Lambda > 1$, so I can apply the previous theorem to find the 'stable patch distribution'. First note that

$$\begin{aligned} \int_0^\infty K(\alpha, a) da &= \int_0^\infty \mu(a) l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz da. \\ &+ \int_0^\infty [\mu_p(a) - \mu(a)] l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz da. \end{aligned} \quad (20)$$

Using integration by parts, one finds

$$\begin{aligned}
\int_0^\infty \mu(a) l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz da &= \left[-l(a) \int_0^a \frac{\gamma(\alpha, z)}{l(z)} dz \right]_0^\infty \\
&\quad + \int_0^\infty \gamma(\alpha, a) da \\
&= \int_0^\infty \gamma(\alpha, a) da \\
&= 1
\end{aligned} \tag{21}$$

Equations (20), (21) imply that

$$\int_0^\infty K(\alpha, a) da \leq \nu < 1 \tag{22}$$

for all values of α .

Since the spectrum of $K(\alpha, a)$ and $K(a, \alpha)$ are the same, (22) provides the estimate (Zabreyko et. al., 1975) that

$$1 < \frac{1}{\nu} \leq \Lambda. \tag{23}$$

It is clear from the preceding that the size of Λ is governed by how large the fraction of partial disasters is relative to total disasters.

The theorem and (23) now show that the 'stable patch distribution' is unique, and provide a means of computing the 'stable patch distribution'. This can then be substituted into equation (2) and solved for $p(0)$ and thus the system has a unique equilibrium if (2) has a unique solution in this case.

Proceeding as in Hastings and Wolin (1989), we can demonstrate one simple sufficient condition for stability. If older (true age) patches are less likely to suffer extinction than younger patches, then the unique equilibrium found above is locally stable.

Proposed research

There are no changes from the original three year proposal.

Final Report-- DE-FG03-89ER60886/A000 7-15-89 to 7-14-93 (Alan Hastings,
Inst. for Theoretical Dynamics, Univ. of California, Davis, CA 95616)

"Development of Paradigms for the Dynamics of Structured Populations"

Many of the technical details have been described in earlier reports, so here I will provide an overview of the results, referring to the published papers attached. In particular, I will focus on the extent to which I have answered the following two questions, which were the focus of the grant proposal.

Question I.1. Can local 'outbreaks' of pests be explained by predator prey systems in 'patchy environment'? What role do local dynamics play in the interaction between a predator and a patchily distributed prey?

Question I.2 What are the patterns produced by competition in a patchy environment, and can they be used to infer the presence of competition?

Regarding the first question, the answer turns out to be very affirmative -- complex local dynamics can be generated in a variety of ways (Hastings, 1990, 1991; Hastings et al., 1993, Hastings, 1995). First, as shown in Hastings (1992, 1993) interactions between space and dynamics in a single species can lead to complex local dynamics. This was demonstrated in a spectacular fashion in Hastings and Higgins (1994), where the vital role of space in generating very long transient dynamics was emphasized.

A second theme that emerged was the role of local interactions alone in generating complex dynamics (Hastings et al., 1993). This was emphasized in a variety of investigations of simple ecological models (Hastings and Powell, 1991; Klebanoff and Hastings, 1994a,b).

A brief answer to the second question on competition was discussed in Hastings (1991b). Work on this problem is ongoing.

Finally, as indicated in the original proposal (but excluded from emphasis based on reviews and funding) a similar approach was used to look at dynamics of the flour beetle (Hastings and Costantino, 1991).

Publications resulting from grant DE-FG03-89ER60886/A000
(Attached as appendices)

Hastings, A. (1990). Spatial heterogeneity and ecological models. Ecology, 71, 426-428.

Hastings, A. (1991a). Structured models of metapopulation dynamics. Biological Journal of the Linnean Society, 42, 57-71.

Hastings, A., & Costantino, R. F. (1991). Oscillations in population numbers: Age dependent cannibalism. Journal of Animal Ecology, 60, 471-482.

Reprints / Preprints - removed

Hastings, A., & Powell, T. (1991). Chaos in a three species food chain. Ecology, 72, 896-903.

Hastings, A. (1991b). McKendrick-Von Foerster Models for Patch Dynamics. in Lecture Notes in Biomathematics, v. 92

Hastings, A. (1992). Age dependent dispersal is not a simple process: density dependence, stability and chaos. Theoretical Population Biology, 41, 388-400.

Hastings, A. (1993) Complex interactions between dispersal and dynamics: Lessons from coupled logistic equations. Ecology, 74, 1362-1372.

Hastings, A., Hom, C., Ellner, S., Turchin, P., & Godfray, H.C.J. (1993) Chaos in ecology: Is mother nature a strange attractor? Annual Reviews of Ecology and Systematics, 24, 1-33.

Hastings, A. & Higgins, K. (1994) Persistence of transients in spatially structured ecological models, Science, 263, 1133-1136.

Klebanoff, A. & Hastings, A. (1994a). Chaos in three species food chains. Journal of Mathematical Biology 32:427-451

Klebanoff, A. & Hastings, A. (1994b) Chaos in one-predator, two-prey models: general results from bifurcation theory. Mathematical Biosciences, in press.

Hastings, A. (1995) Conservation and spatial structure: Theoretical approaches. in Lecture Notes in Biomathematics. v. 100, in press.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.