Excitation of Large- k_θ Ion-Bernstein Waves in Tokamaks

E. J. Valeo and N. J. Fisch

Plasma Physics Laboratory, Princeton University

P. O. Box 451, Princeton, NJ 08543

Abstract

The mode-converted ion-Bernstein wave excited in tokamaks is shown to exhibit certain very interesting behavior, including the attainment of very small poloidal phase velocities, the reversal of poloidal direction, and up-down asymmetries in propagation and damping. Because of these effects, this wave holds promise for channeling α -particle power to ions, something that would make a tokamak fusion reactor far more attractive than presently envisioned.

One way to achieve significantly improved tokamak performance is to direct fusion α -particle power into either electrons to sustain the plasma current[1] or into ions to increase the plasma reactivity[2]. Such a channeling of the α -particle power could make possible the operation of fusion reactors in regimes in which there is a significant hot, nonmaxwellian component to the fuel ions, or in which the fuel ion temperature can be much greater than the electron temperature[3, 4]. The fusion reactivity is then greatly enhanced, making possible a much more economical reactor.

The free energy in the α -particles, namely, the free energy associated with the gradient in α -particle pressure, may be tapped through a resonant quasi-linear interaction with a poloidally propagating short wavelength electrostatic wave[1]. Such a wave diffuses the α -particles both in space and energy, rather than just in energy. The free energy may be accessed if $k_\perp \rho_\alpha \gg 1$, where $\rho_\alpha \equiv v_\alpha / \Omega_\alpha$ is the α -particle gyroradius, and where, for radial diffusion, $k_\perp \simeq k_\theta$. (The confining magnetic field defines the $\parallel$ and $\perp$ directions; in cylindrical coordinates, θ is the azimuthal or poloidal direction.)

Importantly, the change in the radial guiding center displacement, $\Delta r_{gc} = \Delta v_\theta / \Omega_\alpha$, is related directly to the energy change, $\Delta E = m_\alpha(v_\theta \Delta v_\theta + v_\parallel \Delta v_\parallel)$. This occurs because particles in a magnetic field are diffused by waves along the paths

$$v_\perp \Delta v_\perp + (v_\parallel - \omega/k_\parallel) \Delta v_\parallel = 0, \quad (1)$$

whereas, there are two resonance constraints: One, since $k_\perp \rho_\alpha \gg 1$, α -particles are unmagnetized in the wave, and, hence, Landau resonant if

$$v_\theta = (\omega - k_\parallel v_\parallel) / k_\theta. \quad (2)$$

Two, so that successive kicks from the wave add,

$$v_\parallel = (\omega - n\Omega_\alpha) / k_\parallel. \quad (3)$$

It follows from Eqs. (1) – (3) that

$$\Delta r_{gc} = \Delta E k_\theta / m_\alpha \Omega_\alpha \omega. \quad (4)$$

It is this relation that leads to the quest for large- k_θ waves. Clearly, in order to extract successfully α -particle power, the key ratio $\Delta r_{gc} / \Delta E$ must neither be too large nor too small. In the former case, the α -particle leaves the tokamak before giving up its energy. In the latter case, it is difficult to establish a population inversion. The ideal condition is for this ratio to be such that Δr_{gc} is on the order of the minor radius for extracting the full α -particle energy. From Eq. (4), this maximum excursion is

$$\Delta r_{\max} = (1/2)(k_\theta \rho_{\alpha 0}) \rho_{\alpha 0} (\Omega_\alpha / \omega), \quad (5)$$

where $\rho_{\alpha 0} \simeq 4\text{cm}$ is the initial gyroradius for 3.5 MeV α -particles. For $1 < \Omega_\alpha / \omega < 3/2$, corresponding to wave frequencies between the deuterium and tritium resonances (such as the IBW), then for large k_θ sufficiently large, say $k_\theta \simeq 4\text{cm}^{-1}$, one obtains $32\text{cm} < \Delta r_{\max} < 48\text{cm}$, which is a range suitable for extraction, although toroidal effects can be expected to modify somewhat this range. The required diversion of power occurs if the wave then damps on fuel ions.

To achieve large k_θ , one might consider the lower hybrid wave[1], but the wave may be difficult to excite in the proper regime, or there may be other difficulties[5], The mode-converted ion Bernstein wave[6], however,

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

might be excited more easily with the requisite poloidal phase velocity and high wave number. This can occur as follows[7]: In a D-T (or other multi-species) plasma, the ion Bernstein wave (IBW) may be excited by launching a fast wave that mode-converts at the ion-ion resonance surface into an IBW[8]. This mode-converted ion Bernstein wave propagates essentially in the poloidal direction if two choices are made: one, the ion-ion hybrid resonance layer, which is more or less a vertical plane, is chosen to intersect near the tokamak magnetic axis; and, two, the fast wave is launched so as to intersect this vertical plane somewhat off the magnetic axis. Thus, horizontal is equivalent to poloidal propagation.

Of course, coupling to an IBW is possible through a variety of other methods[9]. One possibility would be to launch a slow electrostatic wave (electron plasma wave), which will subsequently convert to an IBW at the lower hybrid resonance. It would be difficult, however, in such a manner, to achieve the high poloidal wave numbers that are necessary, and that are achieved here, through mode conversion at a vertical ion-ion hybrid resonance. The method of coupling analyzed here, in contrast, exploits the production of the short wavelengths characteristic of a resonance layer that is aligned precisely perpendicular to the poloidal direction.

The location of this ion-ion hybrid resonance is determined by[10]

$$\omega^2 = (\Omega_D \Omega_T) \frac{f_D \Omega_T + f_T \Omega_D}{f_D \Omega_D + f_T \Omega_T}, \quad (6)$$

where ω is the wave frequency, $\Omega_D \equiv eB/m_D$ is the deuterium gyrofrequency, Ω_T is the tritium gyrofrequency, and f_D and f_T are their respective fractional densities. The horizontal location of this resonance, which may be chosen so as to intersect the magnetic axis, may be controlled through the toroidal magnetic field, B , through the wave frequency, ω , or through the relative concentration of the different ionic species.

Note that, since the gradient of the magnetic field strength in the major radius direction ($\hat{R}$) is the principal inhomogeneity intrinsic to the mode conversion process, the $\hat{R}$ component of the wavevector of the outgoing wave will increase most rapidly as the IBW propagates away from the mode conversion layer, thus achieving essentially poloidal propagation with high poloidal wavenumber.

In our analysis of the propagation of such a wave in a reactor-grade plasma, several things have been discovered: One, through a suitable choice

of wave and tokamak parameters, and under conditions[4] important for channeling α -particle power, namely ion temperatures about 20 KeV, and electron temperatures about 10 KeV, high k_θ can in fact be reached. Two, in this regime, the IBW can be made to reverse direction, so that the fields near high k_θ will be enhanced, as will the channeling effect. Three, it is possible to augment and exploit certain up-down asymmetries (because of the poloidal magnetic field) both for off-axis current-drive and for channeling α -particle power. Four, a deuterium rich mixture achieves simultaneously both the high- k_θ channeling effect and the diverting of power to ions.

The dispersion relation, which describes the ion Bernstein wave, may be written in the form[9]

$$K_{xx} = \frac{|K_{xy}|^2}{n^2} - n_{||}^2 \frac{K_{zz}}{n_{\perp}^2 - K_{zz}}, \quad (7)$$

where the K_{jj} are elements of the susceptibility tensor, and $n \equiv kc/\omega$ is the index of refraction. In the regime important for channeling, it is not possible[11] to employ conventional (small $k_{\perp}\rho$) approximations[12, 13, 14] of these susceptibilities. In what follows, the wave propagates horizontally ($\hat{x}$) and toroidally ($\hat{z}$), but not vertically ($k_y = 0$).

Because our interest is primarily in off-axis effects, the poloidal field importantly relates the parallel to the perpendicular wavenumber through $k_{||} \simeq k_z + \theta k_x$, where $\theta \equiv B_\theta/B$. The first term, k_z , is the launched parallel wavenumber or the parallel offset. The second term represents the projection of the horizontal wavenumber, $k_x \simeq k_\theta$, on the magnetic field in the presence of a small poloidal magnetic field, B_θ , which is assumed horizontal in the regime of interest. Suppose that the antenna launching the fast wave is up-down symmetric, and situated on the low-field periphery of the tokamak. Then one might envision two rays of fast waves propagating from the antenna, one that undergoes mode conversion to the IBW in the top half of the tokamak and the other in the bottom half. The resulting IBW spectrum, however, is not symmetric for nonzero poloidal field, since B_θ is up-down antisymmetric.

In Fig. 1, we trace such rays after the mode conversion for a 50:50 DT mixture. Note that for the case of cancellation of $k_{||}$ ($\theta < 0$, or, say, for the IBW propagating in the top half of the tokamak), there is a tendency for the wave to reach sufficiently high $k_{\perp}$ to be of interest for channeling.

Note also that, particularly for $k_z = 0.1$, the wave in fact turns around prior to damping, returning to the mode conversion layer at an upshifted wavenumber[15]. A caveat to the results here, and to be explored further, is that the ray tracing approach may not be valid, particularly for $\theta > 0$, where the damping is so intense as to occur in a distance short compared to a wavelength.

For channeling alpha power, it is significant both that the horizontal group velocity passes through zero and that this can be made to occur at high k_θ . There is field enhancement where the group velocity vanishes, so that less power is required for the waves to diffuse α -particles in a time short compared to their slowing down time.

Note also that there are considerable up-down asymmetries. First, clearly, the sense of poloidal propagation is opposite. Second, for $\theta > 0$, high- k_θ are not reached, since this wave damps immediately. Third, the damping for $\theta > 0$ is on ions, whereas, for $\theta < 0$, it is on electrons.

These asymmetries lead to different but not necessarily canceling effects. If the fast wave antenna is poloidally phased, then it would be possible to launch just the wave that produces the more desirable effect. However, this phasing is not easily achieved, and, in any event, present tokamaks are generally not equipped for such phasing. Hence, it is of interest, therefore, to examine whether, for example, extraction of α -particle power can occur by the mode-converted IBW with toroidal phasing only.

For the extraction process, it is clear that the large disparity in k_θ leads to significant movement in the α -particles from the top interaction with insignificant cancellation from the bottom one. In addition, there would be amplification of the wave by α -particles at the top, but damping at the bottom. The only loss in not having poloidal phasing is that half the power is wasted with respect to extraction.

The current drive effect in the midplane[16], i.e. for $\theta = 0$, should not be affected by poloidal fields. The current drive effect off axis, however, is more complicated. Since the wave travels in opposite directions on top and bottom, and in both cases is completely damped, the current drive effect might be thought to vanish off axis, and, in fact, has been shown to vanish at least for very small $k_\parallel$ [17]. However, for realistic $k_\parallel$, there are interesting asymmetries: First, in the top case, the wave is 90% damped on electrons, whereas in the bottom case it is 2/3 damped on deuterium, 1/3 on tritium. Hence, the electrons and ions absorb momentum in opposite directions, but

their currents from this effect *add*. Second, the parallel velocity of electrons and tritium ions that absorb wave power must be of the same sign as $\omega/k_{\parallel}$, whereas the parallel velocity of the resonant deuterium ions must be of opposite sign. Hence, to the extent that the resonant ions absorb power, there is the minority species heating current drive effect (MSHCD) that could be in opposite directions for these ions[18]. Depending on the effective bulk ion charge state, the trapped particle fraction, and other parameters, the MSHCD currents can actually be made to flow in either direction[19, 20]. Thus, it is clear that sufficient asymmetry exists for the current drive effect even off axis, and that here the effects of the oppositely traveling waves can even be additive.

Note, however, that while extraction of α -particle power appears to be in place, and while this power may in fact be channeled to produce current drive even with toroidal phasing only, it follows from Fig. 1 that this power can not be channeled to ions for increased reactivity. To accomplish this channeling effect, it will be necessary to explore further the parameter space. In Fig. 2, we vary the fuel mix, showing the damping on each species for (a) bottom case (i.e. $\theta > 0$), (b) $\theta = 0$, and (c) top case. (Note that the on-axis case has only 50% electron damping, which limits the mode-converted current drive, but the electrons for this case are relatively cold.)

Of particular interest is that, for a 70:30 DT mix, there is the opportunity for both high k_{θ} and damping on tritium ions. Here, most of the power can be channeled to tritium ions at near maximum k_{θ} . Note that this is quite relevant to reactor optimization, since at 70:30, there is 85% of the maxwellian fusion power, plus increased opportunity for an enhanced reactivity of a nonmaxwellian tritium tail[4].

In Fig. 3, we show the ray tracing for the 70:30 case, which indicates that here, as in the 50:50 case, even with toroidal phasing only, the channeling effect remains. However, the current drive effect on top and bottom largely cancels, since in both cases primarily tritium absorbs the power. As in the 50:50 case, the horizontal group velocity vanishes, and the dissipation, especially in the bottom trajectory, is over a very short distance. Also, note that the mode has shifted very much nearer to the tritium resonance.

Since the high- k_{θ} now occurs at only $\omega \simeq 0.74 \Omega_{\alpha}$, it is of interest to check whether there is sufficient Doppler shift for α -particles to interact with the wave. From Eq. (3), using $\rho_{\alpha} \simeq 4$ cm, and, from Fig. 3, taking $k_{\perp} \simeq 3$ cm $^{-1}$ and $k_{\parallel} \simeq -.075$ cm $^{-1}$, we find that $v_{\parallel}/v_{\perp} \simeq 1$, indicating that the resonance

still falls within the bulk of the α -particle distribution.

Although the effect reported is neither fully explored nor optimized here, it can be concluded from this study that large k_θ waves can be excited in tokamaks, off-axis, so as to divert α -particle power to ions. At some sacrifice in optimization, both the diversion of power and current drive can be expected, even without poloidally asymmetric antenna phasing.

The authors are indebted to C. F. F. Karney and R. Majeski for very useful comments. This work was supported by the United States Department of Energy under contract number DE-AC02-76-CHO3073.

References

- [1] N. J. Fisch and J. M. Rax, Phys. Rev. Lett. **69**, 612 (1992).
- [2] N. J. Fisch and J. M. Rax, in Plasma Physics and Controlled Nuclear Fusion Research (Proc. 14th Int. Conf. on Plasma Phys. and Controlled Thermonuclear Fusion, Würzburg, 1992) IAEA, Vienna, Vol. 1, 769 (1993).
- [3] J. F. Clarke, Nucl. Fusion **20**, 563 (1980).
- [4] N. J. Fisch and M. C. Herrmann, Princeton Plasma Physics Laboratory Report 2989 (1994).
- [5] K. Kupfer, S. C. Chiu and V. S. Chan, GA Report GA-A21683 (1994).
- [6] I. B. Bernstein, Phys. Rev. **109**, 10 (1958).
- [7] N. J. Fisch, E. J. Valeo, C. F. F. Karney, and R. Majeski, to appear in Proc. 21st Int. Conf. on Plasma Phys. and Controlled Nucl. Fusion, Montpellier, France (1994).
- [8] F. W. Perkins, Nucl. Fusion **17**, 1197 (1977).
- [9] M. Ono, Phys. Fluids B **5**, 241 (1993).
- [10] T. H. Stix, *The Theory of Plasma Waves*, McGraw-Hill, NY (1962).
- [11] R. Koch, et al., Report ERM/KMS-89, Ass. Euratom-Belgium, Ecole Royale Militaire, Brussels, Belgium (1989)

- [12] M. Brambilla, Nucl. Fusion **28**, 549 (1988).
- [13] P. L. Colestock and R. J. Kashuba, Nucl. Fusion **23**, 763 (1983).
- [14] T. H. Stix, Phys. Rev. Lett. **15**, 878 (1965).
- [15] This effect, which would vanish for very small $k_{||}$, does not appear to have been remarked upon in the literature, and would be missed numerically if the susceptibilities are approximated by the usual finite Larmor radius expansion.
- [16] R. Majeski et al., in Proc. 10th Top. Conf. on rf Heating, ed. M. Porkolab and J. Hosea (AIP, NY) AIP Conf. Proc. Vol. 289, 401 (1994).
- [17] A. K. Ram and A. Bers, Phys. Fluids B **3**, 1059 (1991).
- [18] N. J. Fisch, Nucl. Fusion **21**, 15 (1981).
- [19] S. C. Chiu, V. S. Chan, J. Y. Hsu, G. E. Guest, R. Prater, and J. M. Rawls, Nucl. Fusion **23**, 499 (1983).
- [20] N. J. Fisch, Rev. Mod. Phys. **59**, 175 (1987).

Figure Captions

Fig. 1: Trajectory of IBW $k_x(x)$ from dispersion relation of Eq. (7), with parameters: $\omega = 1.69 \cdot 10^9 \text{ rad sec}^{-1}$, $n_e = 10^{14} \text{ cm}^{-3}$, $T_e = 10 \text{ keV}$, $T_D = T_T = 20 \text{ keV}$, with fractional densities, $f_D = f_T = 0.5$. The plasma inhomogeneity in the $\hat{x}$ direction is due entirely to the variation in magnetic field $\mathbf{B} = 4.8 \cdot 10^4 (\hat{z} + \theta \hat{x}) R/x$ gauss, with $\theta = \pm 0.05$, and "major radius" $R = 260 \text{ cm}$. The broken curves are trajectories for the values .05, .75, and .1 cm^{-1} of the toroidal wavevector k_z . The solid lines connect the coordinates at which .75, .5, and .2 of the initial power remains.

Fig. 2: Left axis displays fractional power absorbed by electrons (long dashes), deuterium (dots), and tritium (short dashes). Right axis displays k_x for which 50% of power remains (solid line). All are plotted as a function of fractional deuterium density f_D , with $\theta = .05, 0, -.05$ in (a), (b), and (c), respectively.

Fig. 3: Ray trajectories for $f_D = .7$. Other parameters are as in Fig. 1.

FIGURE 1

FIGURE 2

FIGURE 3

EXTERNAL DISTRIBUTION IN ADDITION TO UC-420

Dr. F. Paoloni, Univ. of Wollongong, AUSTRALIA	Dr. H. Yamato, Toshiba Res & Devel Center, JAPAN
Prof. R.C. Cross, Univ. of Sydney, AUSTRALIA	Prof. I. Kawakami, Hiroshima Univ., JAPAN
Plasma Research Lab., Australian Nat. Univ., AUSTRALIA	Prof. K. Nishikawa, Hiroshima Univ., JAPAN
Prof. I.R. Jones, Flinders Univ, AUSTRALIA	Librarian, Naka Fusion Research Establishment, JAERI, JAPAN
Prof. F. Cap, Inst. for Theoretical Physics, AUSTRIA	Director, Japan Atomic Energy Research Inst., JAPAN
Prof. M. Heindler, Institut für Theoretische Physik, AUSTRIA	Prof. S. Itoh, Kyushu Univ., JAPAN
Prof. M. Goossens, Astronomisch Instituut, BELGIUM	Research Info. Ctr., National Instit. for Fusion Science, JAPAN
Ecole Royale Militaire, Lab. de Phy. Plasmas, BELGIUM	Prof. S. Tanaka, Kyoto Univ., JAPAN
Commission-European, DG. XII-Fusion Prog., BELGIUM	Library, Kyoto Univ., JAPAN
Prof. R. Bouciqué, Rijksuniversiteit Gent, BELGIUM	Prof. N. Inoue, Univ. of Tokyo, JAPAN
Dr. P.H. Sakanaka, Instituto Fisica, BRAZIL	Secretary, Plasma Section, Electrotechnical Lab., JAPAN
Prof. Dr. I.C. Nascimento, Instituto Fisica, Sao Paulo, BRAZIL	Dr. O. Mitarai, Kumamoto Inst. of Technology, JAPAN
Instituto Nacional De Pesquisas Espaciais-INPE, BRAZIL	Dr. G.S. Lee, Korea Basic Sci. Ctr., KOREA
Documents Office, Atomic Energy of Canada Ltd., CANADA	J. Hyeon-Sook, Korea Atomic Energy Research Inst., KOREA
Ms. M. Morin, CCFM/Tokamak de Varennes, CANADA	D.I. Choi, The Korea Adv. Inst. of Sci. & Tech., KOREA
Dr. M.P. Bachynski, MPB Technologies, Inc., CANADA	Prof. B.S. Liley, Univ. of Waikato, NEW ZEALAND
Dr. H.M. Skarsgard, Univ. of Saskatchewan, CANADA	Inst of Physics, Chinese Acad Sci PEOPLE'S REP. OF CHINA
Prof. J. Teichmann, Univ. of Montreal, CANADA	Library, Inst. of Plasma Physics, PEOPLE'S REP. OF CHINA
Prof. S.R. Sreenivasan, Univ. of Calgary, CANADA	Tsinghua Univ. Library, PEOPLE'S REPUBLIC OF CHINA
Prof. T.W. Johnston, INRS-Energie, CANADA	Z. Li, S.W. Inst Physics, PEOPLE'S REPUBLIC OF CHINA
Dr. R. Bolton, Centre canadien de fusion magnétique, CANADA	Prof. J.A.C. Cabral, Instituto Superior Tecnico, PORTUGAL
Dr. C.R. James,, Univ. of Alberta, CANADA	Prof. M.A. Hellberg, Univ. of Natal, S. AFRICA
Dr. P. Lukác, Komenského Universzita, CZECHO-SLOVAKIA	Prof. D.E. Kim, Pohang Inst. of Sci. & Tech., SO. KOREA
The Librarian, Culham Laboratory, ENGLAND	Prof. C.I.E.M.A.T, Fusion Division Library, SPAIN
Library, R61, Rutherford Appleton Laboratory, ENGLAND	Dr. L. Stenflo, Univ. of UMEA, SWEDEN
Mrs. S.A. Hutchinson, JET Library, ENGLAND	Library, Royal Inst. of Technology, SWEDEN
Dr. S.C. Sharma, Univ. of South Pacific, FIJI ISLANDS	Prof. H. Wilhelmson, Chalmers Univ. of Tech., SWEDEN
P. Mähönen, Univ. of Helsinki, FINLAND	Centre Phys. Des Plasmas, Ecole Polytech, SWITZERLAND
Prof. M.N. Bussac, Ecole Polytechnique,, FRANCE	Bibliotheek, Inst. Voor Plasma-Fysica, THE NETHERLANDS
C. Mouttet, Lab. de Physique des Milieux Ionisés, FRANCE	Asst. Prof. Dr. S. Cakir, Middle East Tech. Univ., TURKEY
J. Radet, CEN/CADARACHE - Bat 506, FRANCE	Dr. V.A. Glukhikh, Sci. Res. Inst. Electrophys. I Apparatus, USSR
Prof. E. Economou, Univ. of Crete, GREECE	Dr. D.D. Ryutov, Siberian Branch of Academy of Sci., USSR
Ms. C. Rinni, Univ. of Ioannina, GREECE	Dr. G.A. Eliseev, I.V. Kurchatov Inst., USSR
Preprint Library, Hungarian Academy of Sci., HUNGARY	Librarian, The Ukr.SSR Academy of Sciences, USSR
Dr. B. DasGupta, Saha Inst. of Nuclear Physics, INDIA	Dr. L.M. Kovrizhnykh, Inst. of General Physics, USSR
Dr. P. Kaw, Inst. for Plasma Research, INDIA	Kernforschungsanlage GmbH, Zentralbibliothek, W. GERMANY
Dr. P. Rosenau, Israel Inst. of Technology, ISRAEL	Bibliothek, Inst. Für Plasmaforschung, W. GERMANY
Librarian, International Center for Theo Physics, ITALY	Prof. K. Schindler, Ruhr-Universität Bochum, W. GERMANY
Miss C. De Palo, Associazione EURATOM-ENEA , ITALY	Dr. F. Wagner, (ASDEX), Max-Planck-Institut, W. GERMANY
Dr. G. Grosso, Istituto di Fisica del Plasma, ITALY	Librarian, Max-Planck-Institut, W. GERMANY
Prof. G. Rostangni, Istituto Gas Ionizzati Del Cnr, ITALY	