ADVANCED SEQUENCING TECHNOLOGY

2nd Year Progress Report

Grant # DE-FG02-88ER60700

Awarded to the University of Utah

Development of multiplex sequencing with a transposon front end, chemiluminescent detection and automated reading of the patterns, has progressed to the point where we will work at integrating this into a simple, self contained system during the next year.

Bob Weiss and Diane Dunn have perfected transposon insertion and mapping to the point where robustness of the system is being tested on pilot projects. The informatics group of our genome center has worked with Weiss and Dunn to implement a "transposon mapper" system that automates analysis of data from combinatorial pools to select the minimally overlapping set of inserts, which when sequenced, will give double strand coverage with minimal redundancy. It now seems clear that the mini gamma delta transposon system

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

JP

has the needed randomness of insertion to be useful for large DNA sequencing projects.

Considerable experience has been obtained over the last year in using the direct blot sequencing gels for producing membranes for probing. Longer gels are better, and careful construction of the crucial gel - membrane interface has improved the quality and robustness so that reads of 600-700 are routinely obtained.

An automated probing chamber has been engineered and constructed. Early tests show that uniform probing and washing can be obtained with minimal volumes of reagents. A pumping system, including temperature control of buffer reservoirs, is just being tested out.

Achim Karger has shown that direct reading of chemiluminescent sequence patterns on membranes can be done with a CCD camera. The camera scans the film and the CCD readout is synchronized with the movement so that the data is integrated as the bands cross the array of pixels. High quality images are obtained. Sensitivity and background are being investigated with the

hope of reaching the point where deep levels of multiplexing can be used with direct chemiluminescent detection.

Development of the sequence reading algorithm by Tom Stockham and Jeff Ives has proceeded well with demonstrations of reads to 600 bases with very low error rates. The system is now being tested quantitatively on pilot projects where manual reading will be compared with the automated system. Also, a friendly and convenient front end is being added to the reader to make it uniformly useful.

In addition to continuing the development of each of these components during the next year, we will work on ways to fold the pieces into an integrated and self-contained system. Direct reading of the chemiluminescent membranes makes this possible.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Publications:

Chu, T.J., Caldwell, K.D., Weiss, R.B., Gesteland, R.F., and Pitt, W.G., 1991. Low fluorescence background electroblotting membrane for DNA sequencing. *Electrophoresis* 13:105-114.

Karger, A., Harris, J.M. and Gesteland, R.F. 1991. Multiwavelength fluorescence detection for DNA sequencing using capillary electrophoresis. *Nucleic Acids Res.* 19:4955-4962.

Swerdlow, H. 1991. Capillary gel electrophoresis for DNA sequencing. The University of Utah. (Ph.D. dissertation).

Swerdlow, H., Zhang, J.Z., Chen, D.Y., Harke, H.R., Grey, R., Wu, S., Fuller, C. and Dovichi, N.J. 1991. Three DNA sequencing methods using capillary gel electrophoresis and laser-induced fluorescence. *Anal. Chem.* 63:2835-2841.

Presentation Abstract for: "Genome Mapping and Sequencing" conference
at Cold Spring Harbor Laboratory on May 6-10,
1992

LARGE-SCALE TEST OF COMBINATORIAL MAPPING AND SEQUENCING
WITH MICRO- $\gamma\delta$ TRANSPOSONS.

R. Weiss, D. Dunn, W. Wheatley, L. DiSera, A. Kimball, B. Milner, M. Ferguson, B. Duval, R. Lee, C. Rote and R. Gesteland, Department of Human Genetics, University of Utah, Salt Lake City, UT 84112.

An integrated front-end approach for processing sets of large plasmid clones into high quality sequence ladders is in an initial test phase using a 100 kb contiguous stretch of human DNA . 5000 individual inserts of a sequencing vector, composed of a family of 260 nt. micro- $\gamma\delta$ transposons, that will serve as bi-directional, multiplex priming sites for dideoxy-sequencing, have been isolated in 12 large plasmid targets. These plasmids range in size from 6 to 20 kb, and collectively span 100 kb of the neurofibromatosis 1 gene from chromosome 17. A mapping scheme based on restriction fragment sizing from combinatorial pools allows culling of a minimal spanning set of these inserts, while facilitating batch processing of clones. A pooling strategy, relying on robotic pipetting, reduces the 5000 clones into 168 DNA pools, from which the map positions of each micro- $\gamma\delta$ insert is detected by multiple probings of high-resolution Southern transfers. These high-resolution blots are generated by a direct blotting agarose gel electrophoresis apparatus we have designed, that integrates the separation and transfer steps in one device. The resulting films are scanned and interpreted within an X-windows application that records and stores the fragment sizes, as well as organizing and tracking the pooling process. The derived map position then allows computer prediction of a minimal spanning set of inserts required to complete both strands of the target sequence with minimal redundancy. Templates are prepared in multiplex pools and converted to single strands following attachment to a solid support via a biotin-streptavidin linkage. Multiplex sequence ladders are generated using Mn^{++} -Sequenase conditions; the electrophoretic separation and transfer to nylon membranes is accomplished using direct blotting electrophoresis. Direct blotting conditions are allowing ascertainment of > 600 nts., resolved with constant band-spacing, from each end of the micro- $\gamma\delta$ insert. We will discuss the design and construction of an automated probe chamber for processing membranes through the hybridization and washing procedures, and we will discuss the development of graphical interfaces, written using X windows on top of a relational database, that will allow data organization, monitoring and analysis throughout the sequencing process.

Presentation Abstracts for: "High Performance Capillary Electrophoresis"
international symposium at Amsterdam, The
Netherlands, on February 9-13, 1992

ACHIM E. KARGER*, JOEL M. HARRIS, RAYMOND F. GESTELAND

University of Utah, Dept. of Human Genetics, Eccles Institute for
Human Genetics, Salt Lake City, UT 84112, USA

Multiwavelength Fluorescence Detection for Capillary DNA Sequencing

Multiwavelength detection of laser induced fluorescence for DNA sequencing with different fluorophores is described. A N₂-cooled 2-dimensional charge-coupled device (CCD) camera is used for on-column capillary electrophoresis detection by recording emission spectra over a range of 160nm. The minimum concentration of fluorescent labeled primer solution required for selective detection is 26pM (Joe-A fluorescent labeled primer, Applied Biosystems Inc., 10nm bandwidth, S/N=2). The optical system, fluorescence spectroscopy with a 2-D array, the operation of the CCD camera in the 'time delayed integration mode' (TDI) and the processing of spectroscopic raw data are demonstrated. Increased selectivity of multiwavelength detection over single sensor detectors can be verified by Monte-Carlo as well as analytical calculations. The optical system has no moving parts, is not limited to a certain set of sequencing fluorophores and could function in multiple fluorophore applications other than DNA sequencing without modification.

HAROLD SWERDLOW*, KERRY DEW-JAGER and RAYMOND GESTELAND

Human Genetics Dept., 6160 Eccles Genetics Bldg. Univ. of Utah SLC, UT 84112, USA

Studies on Stability of Capillary Gels for Electrophoresis of DNA Sequencing Reaction Products.

Recent interest in capillary gel electrophoresis (CGE) has been fueled by the Human Genome Project and other large-scale sequencing projects. Advances in gel polymerization, handling technology and low-level detector design have enabled sequencing of DNA directly in capillaries. Efforts to exploit this technology on a widespread basis have been hampered by problems with the reproducibility and stability of gels, and by the formation of bubbles during electrophoresis. Various capillary pre-treatments and alternative gel recipes can improve the situation, but probably do not completely eliminate bubbles. The use of CGE as an instrumental technique, has proven problematic, when actual DNA sequencing samples are employed.

Instability of gels during electrophoresis can be visualized by the formation of different classes of bubbles and a decrease in the current passing through the capillary under a constant voltage. These aberrations all seem to express themselves at the injection (cathodic) end of the capillary gel. In order to improve CGE, we sought to determine the mechanisms underlying the disruptive processes in a simplified system (i.e. a standard polyacrylamide gel which was not covalently attached to the silica capillary wall). Experimental evidence for amelioration of the most prominent sources of gel breakdown will be shown. Improvements in sample preparation, allowing multiple DNA sequencing runs on the same capillary, will also be discussed.

GRANT NUMBER: _____

BIOGRAPHICAL SKETCH

Give the following information for all new key personnel, consultants, and collaborators.
Copy this page for each person.

NAME Raymond F. GESTELAND		POSITION TITLE Professor of Human Genetics, Biology, and Howard Hughes Med. Inst.	
EDUCATION (Begin with baccalaureate or other initial professional education, such as nursing, and include postdoctoral training.)			
INSTITUTION AND LOCATION	DEGREE	YEAR CONFERRED	FIELD OF STUDY
University of Wisconsin	B.S.	1960	Chemistry
University of Wisconsin	M.S.	1961	Biochemistry
Harvard University	Ph.D.	1965	Biochemistry
Institut de Biologie Moleculaire, Univ. de Geneve	Post-doc	1966-67	

RESEARCH AND PROFESSIONAL EXPERIENCE: Concluding with present position, list, in chronological order, previous employment, experience, and honors. Key personnel include the principal investigator and any other individual who participate in the scientific development or execution of the project. Key personnel typically will include all individuals with doctoral or other professional degrees but in some projects will include individuals at the masters or baccalaureate level provided they contribute in a substantive way to the scientific development or execution of the project. Include present membership on any Federal Government public advisory committee. List, in chronological order, the titles, all authors, and complete references to all publications during the past three years and to representative earlier publications pertinent to this application. DO NOT EXCEED TWO PAGES.

Experience:

- 1967-73 Assistant Professor (adjunct), Dept. of Biochemistry, State University of New York at Stony Brook
- 1967-78 Assistant Director for Research and Senior Staff Investigator, Cold Spring Harbor Laboratory
- 1971-74 Member of the National Science Foundation Advisory Panel on Molecular Biology
- 1973-78 Associate Professor (adjunct), Dept. of Biochemistry, State University of New York at Stony Brook
- 1977-78 Member of National Institutes of Health-AID Task Force on Virology
- 1978-pres Investigator, Howard Hughes Medical Institute, Laboratory for Genetic Studies, University of Utah
- 1978-pres Professor of Biology, University of Utah
- 1982-88 Member, National Science Foundation Advisory Panel on Biochemistry
- 1984-pres Co-Chairman, Human Genetics Dept., University of Utah
- 1985-pres Professor of Human Genetics, University of Utah
- 1987-pres Member, Selection Committee for the Pediatric Scientist Training Program.
- 1988-90 Associate Editor, "The New Biologist," a new journal formed in 1988.
- 1989-pres. Board of Trustees (Research Committee), Primary Children's Medical Center Foundation, Salt Lake City, Utah.
- 1989-pres. Adjunct Professor of Bioengineering, University of Utah

Publications from 1989 to present:

- Karger, A., J.T. Ives, R.B. Weiss, J.M. Harris, and R.F. Gesteland. 1990. Imaging of fluorescent and chemiluminescent DNA hybrids using a 2-D CCD camera. Meeting: SPIE Proceedings (Los Angeles, CA) Vol. 1206.
- Milton, D.L., Casper, M.L., Wills, N.M., and Gesteland, R.F. 1990. Guanine tracts enhance sequence directed DNA bends. NAR 18:817-820.
- Swerdlow, H. and Gesteland, R.F. 1990. Capillary gel electrophoresis for rapid, high resolution DNA sequencing. NAR 18:1415-1419.

- Milton, D.L., Casper, M.L. and Gesteland, R.F. 1990. Saturation mutagenesis of a DNA region of bend: Base steps other than ApA influence the bend. *J. of Mol. Biol.* 213:135-140.
- Weiss, R.B., Dunn, D.M., Atkins, J.F. and Gesteland, R.F. 1990. The Ribosome's Rubbish, in "Ribosomes: Structure, Function and Genetics", W.Hill, P.Moore, R.Garrett, J.Warner, A.Dahlberg and D.Schlessinger eds.,chapter 46, ASM publications, pp.534-540.
- Wagner, L.A., Weiss, R.B., Driscoll, R., Dunn, D.S., and Gesteland, R.F. 1990. Transcriptional slippage occurs during elongation at runs of adenine or thymine in *Escherichia coli*. *Nucleic Acids Research* 12:3529-3535.
- Cawthon, R. Weiss, R., Xu, G., Viskochil, D., Culver, M., Stevens, J., Robertson, M., Dunn, D., Gesteland, R., O'Connell, P. and White, R. 1990. A major segment of the neurofibromatosis I gene: cDNA sequence, genomic structure and point mutations. *Cell* 62:193-201.
- Xu, Gangfeng, O'Connell, P., Viskochil, D., Cawthon, R., Robertson, M. Culver, M., Dunn, D., Stevens, J., Gesteland, R.F., White, R.L., and Weiss, R. 1990. The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell* 62:599-608.
- Skuzeski, J.M., Nichols, L.M., and Gesteland, R.F. 1990. Analysis of leaky viral translation termination codons in vivo by transient expression of improved B-glucuronidase vectors. *Plant Molecular Biology* 15:65-79.
- Tuohy, T.M.F., Thompson, S., Hughes, D., Gesteland, R.F., and Atkins, J.F. 1990. The role of EF-Tu and other translation components in determining translocation step size. *Biophys. Acta* 1050:274-278. (Special issue on protein synthesis).
- Weiss, R.B., Dunn, D.M., Atkins, J.F. and Gesteland, R.F. 1990. Ribosomal Frameshifting from -2 to +50 nucleotides, in "Progress in Nucleic Acids Research and Molecular Biology", W.Cohn and K.Moldave eds., Vol. 39:159-183.
- Atkins, J.F., Weiss, R.B., and Gesteland R.F. 1990. Ribosome gymnastics - Degree of difficulty 9.5, Style 10.0. *Cell* (review) 62: 413-423.
- Skuzeski, J.M., Nichols, L.M., Gesteland, R.F., and Atkins, J.F. The signal for a leaky UAG stop codon in several plant viruses includes the two downstream codons. *J. Mol. Biol.* 218:365-373.
- Xu, G, Lins, B., Tanaka, K., Dunn, D., Wood, D., Gesteland, R., White, R., Weiss, R., and Tamanoi, F. 1990. The catalytic domain of the neurofibromatosis type 1 gene stimulates the ras GTPase and complements ira mutants of *S. cerevisiae*. *Cell* 63: 835-841.
- Wills, N.W., Gesteland, R.F. and Atkins, J.F. Evidence that a downstream pseudoknot is required for translational read-through of the Moloney Murine Leukemia Virus gag stop codon. 1991. *Proc. Natl. Acad. Sci. USA* 88:6991-6995 USA.
- Atkins, J.F., Weiss, R.B., Thompson, S., and Gesteland, R.F. 1991. Towards a genetic dissection of the basis of triplet decoding, and its natural subversion: programmed reading frame shifts and hops. *Annu. Rev. Genet.* 25: 201-228.
- Swerdlow, H., Gesteland, R. Wu, S. Harke, H., and Dovichi, N.J. 1990. Capillary gel electrophoresis for DNA sequencing: Laser-induced fluorescence detection with the sheath flow cuvette. *J. of Chrom.* 516: 61-67.
- Karger, A., Harris, J.M. and Gesteland, R.F. 1991. Multiwavelength fluorescence detection for DNA sequencing using capillary electrophoresis. *Nucleic Acids Research* 19:4955-4962
- Condon, B. G., Gesteland, R. F. and Atkins J. F. 1991. An analysis of sequences stimulating frameshifting in the decoding of gene 10 of bacteriophage T7. *Nucleic Acids Research* 19:5607-5612.
- Condon, B.G., Atkins, J.F., and Gesteland, R.F. 1991. Frameshifting in gene 10 of bacteriophage T7. *Journal of Bacteriology* 173:6998-7003.

BIOGRAPHICAL SKETCH

Give the following information for all new key personnel, consultants, and collaborators.
Copy this page for each person.

NAME Robert B. WEISS		POSITION TITLE Research Assistant Professor	
EDUCATION (Begin with baccalaureate or other initial professional education, such as nursing, and include postdoctoral training.)			
INSTITUTION AND LOCATION	DEGREE	YEAR CONFERRED	FIELD OF STUDY
Massachusetts Inst. of Technology Cambridge, Massachusetts	B.S.	1978	Life Sciences
Univ. of Washington, Seattle, WA	Ph.D.	1983	Genetics

RESEARCH AND PROFESSIONAL EXPERIENCE: Concluding with present position, list, in chronological order, previous employment, experience, and honors. Key personnel include the principal investigator and any other individual who participate in the scientific development or execution of the project. Key personnel typically will include all individuals with doctoral or other professional degrees but in some projects will include individuals at the masters or baccalaureate level provided they contribute in a substantive way to the scientific development or execution of the project. Include present membership on any Federal Government public advisory committee. List, in chronological order, the titles, all authors, and complete references to all publications during the past three years and to representative earlier publications pertinent to this application. DO NOT EXCEED TWO PAGES.

Experience:

1978 - 1983

Graduate student, Department of Genetics
University of Washington, Seattle, Washington
Funding: N.I.H. Pre-doctoral Training Grant
Advisor: Jonathan A. Gallant

1983 - Sept. 1984

Post-doctoral fellow, Cold Spring Harbor Laboratory
Funding: Exxon Research Foundation Grant for Anaerobic Genetics
Advisor: A.I. Bukhari

Oct. 1984 - 1988

Post-doctoral fellow, Howard Hughes Medical Institute, University of Utah
Funding: N.S.F. and Helen Hay Whitney Foundation Fellowship
Advisor: Raymond F. Gesteland

July 1988 - August 1991

Senior Associate, Howard Hughes Medical Institute, University of Utah
Funding: Howard Hughes Medical Institute
Advisor: Raymond F. Gesteland

Oct. 1988 - present

Research Assistant Professor, University of Utah, Human Genetics Dept.

Representative Publications:

Huang, W.M., S. Ao, S. Casjens, R. Orlandi, R. Zeikus, R. Weiss, D. Winge and M. Fang. 1988.
A novel untranslated sequence within gene 60 of the bacteriophage T4 DNA topoisomerase.
Science 239:1005-1012.

Weiss, R., D. Lindsley, B. Falahee, and J. Gallant. 1988. On the mechanism of ribosomal frameshifting at hungry codons. J. Mol. Biol. 203:403-410.

Falahee, M.B., R.B. Weiss, M. O'Connor, S. Doonan, R.F. Gesteland, and J.F. Atkins. 1988.
Mutants of translational components that alter reading frame by two steps forward or one step

- back. *J. Biological Chemistry* 263:18099-18103.
- Weiss, R.B., Dunn, D.M., Shuh, M., Atkins, J.A. and Gesteland, R.F. (1989) E.coli ribosomes rephase on retroviral frameshift signals at rates ranging from 2 to 50 percent. *The New Biologist* 1, 159-170.
- Karger, A., Ives, J., Weiss, R., Harris, J. and Gesteland, R.F. (1990) Imaging of fluorescent and chemiluminescent DNA hybrids using a 2-D CCD camera. In "SPIE Proceedings, Vol. 1206.
- Weiss, R.B., Dunn, D.M., Atkins, J.F. and Gesteland, R.F. (1989). tRNA hopping during translation. in "Ribosomes: Structure, Function and Genetics", W.Hill, P.Moore, R.Garrett, J.Warner, A.Dahlberg and D.Schlessinger eds., ASM Publications, pp. 534-540.
- Weiss, R.B., Dunn, D.M., Atkins, J.F. and Gesteland, R.F., Ribosomal Frameshifting from -2 to +50 nucleotides, in "Progress in Nucleic Acids Research and Molecular Biology", W.Cohn and K.Moldave eds., 39:159-183.
- Wagner, L., Weiss, R., Driscoll, R., Dunn, D. and Gesteland R. F. (1990) Transcriptional slippage occurs during elongation at runs of adenine or thymine in *E. coli*. *Nuc. Acids Res.* 18, 3529-3535.
- Weiss, R.B., Huang, W.M. and Dunn, D.M. (1990) A nascent peptide is required for ribosomal bypass of the coding gap in bacteriophage T4 gene 60. *Cell* 62, 117-126.
- Cawthon, R., Weiss, R., Xu, G., Viskochil, D., Culver, M., Stevens, J., Robertson, M., Dunn, D., Gesteland, R., O'Connell, P. and White, R. A major segment of the neurofibromatosis I gene: cDNA sequence, genomic structure and point mutations. *Cell* 62:193-201.
- Atkins, J.F., Weiss, R.B., and Gesteland, R.F. (1990) Ribosome gymnastics - Degree of difficulty 9.5, style 10.0. *Cell* 62, 413-423.
- Xu, G., O'Connell, P., Viskochil, D., Cawthon, R., Robertson, M., Culver, M., Dunn, D., Stevens, J., Gesteland, R., White, R., and Weiss, R. (1990) The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell* 62, 599-608.
- Cawthon, R.M., O'Connell, P., Buchberg, A.M., Viskochil, D., Weiss, R.B., Culver, M., Stevens, J., Jenkins, N.A., Copeland, N.G., and White, R.L. (1990) Identification and characterization of transcripts from the neurofibromatosis 1 region: the sequence and genomic structure of *EVI2* and mapping of other transcripts. *Genomics* 7, 555-565.
- Xu, G., Lin, B., Tanaka, K., Dunn, D., Wood, D., Gesteland, R., White, R., Weiss, R., and Tamanoi, F. (1990) The catalytic domain of the neurofibromatosis type 1 gene stimulates the ras GTPase and complements *ira* mutants of *S. cerevisiae*. *Cell* 63, 835-841.
- Weiss, R.B., Dunn, D., Atkins, J. and Gesteland, R. 1990. "The Ribosome's Rubbish," chap. 46, pp. 534-540, in "The Ribosome: Structure, Function, and Evolution." ASM Publication.
- Weiss, R.B. (1990) Frameshift and frame-jump sites as control points during elongation. in: "Post-transcriptional Control of Gene Expression," NATO ASI Series, Vol.H 49., pp. 579-590. J.E.G. McCarthy and M.F. Tuite (eds.), Springer-Verlag Berlin Heidelberg.
- Cawthon, R.M., Andersen, L.B., Buchberg, A.M., Xu, G., O'Connell, P., Viskochil, D., Weiss, R.B., Wallace, M.R., Marchuk, D.A., Culver, M., Stevens, J., Jenkins, N.A., Copeland, N.G., Collins, F.S., and White, R. (1991) cDNA sequence and genomic structure of *EVI2B*, a gene lying within an intron of the neurofibromatosis type 1 gene. *Genomics* 9, 446-460.
- Atkins, J.F., Weiss, R.B., Thompson, S., and Gesteland, R.F. (1991) Towards a genetic dissection of the basis of triplet decoding, and its natural subversion: programmed reading frame shifts and hops. *Annu.Rev.Genet.* 25, 201-228.
- Weiss, R.B. (1991) Ribosomal frameshifting, jumping and read-through. *Current Opinion in Cell Biology* 3(6), 1051-1055.
- Chu, T.J., Caldwell, K.D., Weiss, R.B., Gesteland, R.F., and Pit, W.G. (1991) Low fluorescence background electroblotting membrane for DNA sequencing. *Electrophoresis* 13: 105-114.

END

**DATE
FILMED**

12 / 2 / 92