APR 23 1962

BNL 5994

MASTER

THE SPECIFICITY OF INCORPORATION OF GLUTAMIC ACID
AND GLUTAMINE INTO PROTEIN*

By G. Zubay

Biology Department, Brookhaven National Laboratory, Upton, New York

Communicated by Milislav Demerec

The sequence of most amino acids in proteins is believed to be determined by a nucleotide coding mechanism.¹ There is evidence to suggest that not all amino acids are coded for. Thus we know that prolines are hydroxylated after incorporation into the polypeptide chain to produce hydroxyproline;² and that cystines are formed by the interaction of two peptide linked cysteines. In both these cases the final chemical form of the amino acid is partly determined after it has been incorporated into the polypeptide chain. Quite clearly hydroxyproline and cystine have no mutual representation in nucleic acids. Instead proline and cysteine are coded for, and some of these are converted into hydroxyprolines and cystines respectively. Although it has been assumed³ that the acid and amide forms of glutamate and aspartate are separately coded for, no convincing experimental support has been obtained. Experiments described here have been designed to test these alternate possibilities.

Materials and Methods.--E. coli S-RNA was used. The preparation of this as well as the preparation of E. coli enzyme extract and the routine procedure used to assay amino acid incorporation has been described in detail elsewhere.⁴ Isolation of amino acid from amino acid labelled S-RNA and paper electrophoresis: after incubation with C¹⁴-glutamic acid and the standard incubation mixture, amino acid S-RNA was reisolated by phenol extraction. The coupled amino acid was hydrolyzed from the S-RNA by incubation for three

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

hours in 0.2 M ammonium formate at pH 8.0 and 37°C. The RNA is then removed by ethanolic precipitation. The amino acid is concentrated by evacuation which removes the alcohol, water and finally the ammonium formate. The dried residue is resuspended in a minimum amount of 0.05 M sodium phosphate, pH 8.0 + 2 mM C-glutamic acid + 2mM C-glutamine. 100 μ l of this solution are electrophoresed at pH 8.0 for 40 minutes at 16 volts/cm on Whatmann 3 MM paper. The positions of the glutamine and glutamic acid markers was detected by ninhydrin. The radioactivity of the glutamine and glutamic acid was determined by cutting samples from the paper and counting them in a gas flow counter.

Results and Discussion.--Amino acids form specific complexes with S-RNAs which serve as immediate precursors in protein synthesis.⁵ In an incubation mixture containing purified Escherichia coli S-RNA and crude Escherichia coli enzyme extract it has been found that uniformly labelled C¹⁴-glutamic acid is slowly incorporated into the S-RNA.⁴ In contrast uniformly labelled C¹⁴-glutamine is rapidly incorporated into the S-RNA (Fig. 1). The maximum incorporation in both cases amounts to about 0.3 μ M amino acids per μ M S-RNA nucleotides. In parallel experiments an equimolar amount of unlabelled glutamine inhibits 90% of the incorporation of C¹⁴-glutamic acid into S-RNA, whereas the same amount of unlabelled glutamic acid results in less than 10% inhibition of C¹⁴-glutamine incorporation (Table 1). These results strongly suggest that the observed glutamic acid incorporation into S-RNA results from its prior conversion into glutamine and the subsequent incorporation of glutamine into the S-RNA. This was confirmed by the isolation of S-RNA which had been incubated with C¹⁴-glutamic acid. Separation of the amino acid from the S-RNA and paper electrophoresis at pH 8.0 demonstrated that the radioactivity which had become incorporated

into the S-RNA was entirely accountable as glutamine (Fig. 2). It would appear that an S-RNA exists for glutamine but not for glutamic acid. There is a possibility that the activating enzyme for the direct incorporation of glutamic acid into S-RNA has been destroyed in the preparation of the enzyme extract. However, this is improbable since in thirteen out of fifteen amino acids tested, sufficient activating enzyme is present to give appreciable incorporation of the amino acid into S-RNA.⁶ It is concluded that in protein synthesis glutamine is coded for while glutamic acid probably is not. The asparagine-aspartic acid system has not been thoroughly studied, but a similar inhibition of labelled aspartic acid incorporation into S-RNA by unlabelled asparagine has been observed.

If the glutamic acid in protein arises by the hydrolysis of the γ -amino group of glutamine, there must be a mechanism which determines when this hydrolysis occurs. It is a possibility that only glutamines on the exposed surface of the protein accessible to glutaminase are converted to the acid form. If this is correct, knowledge of the distribution of glutamines and glutamates might be of some value in the determination of the tertiary structure of proteins.

The author is indebted to Drs. D. E. Koshland, Jr., and C. H. W. Hirs for stimulating discussion.

*Research carried out at Brookhaven National Laboratory under the auspices of the U. S. Atomic Energy Commission.

¹v.g., G. Gamov, A. Rich and M. Vcas, in Advances in Biological and Physics, Vol. IV (New York: Academic Press, 1956) p. 23.

²Stetten, M. R., J. Biol. Chem., 181, 31 (1949).

³Crick, F. H. C., J. S. Griffith and L. E. Orgel, these PROCEEDINGS, 43, 416 (1956).

⁴Experimental details of the assay system used here may be found in Zubay, G., J. Mol. Biol., 4, 000 (1962).

⁵Hoagland, M. B., in The Nucleic Acids, ed. E. Chargaff and J. N. Davidson (New York: Academic Press, 1960), vol. 3, chapter 37.

⁶Zubay, G., unpublished results.

TABLE 1

The Incorporation of Glutamic Acid and Glutamic Acid Into S-RNA. In the control experiments the S-RNA was added after the reaction was stopped by addition of TCA

Incubation Mixture	cpm
Control	43
80 μ M C ¹⁴ -glutamic	340
80 μ M C ¹⁴ -glutamic + 80 μ M C ¹² -glutamine	70
Control	20
80 μ M C ¹⁴ -glutamine	96
80 μ M C ¹⁴ -glutamine + 80 μ M C ¹² -glutamic	92

Fig. 1.--Rate of incorporation of glutamine and glutamic acid into S-RNA at the same concentration of added Escherichia coli enzyme extract. Specific activities of the amino acids were the same.

Fig. 2.--Electrophoresis of radioactive amino acid hydrolysate from S-RNA which has been enzymatically coupled with C^{14} -glutamic acid. Nonradioactive glutamine and glutamic acid markers have been added. Only the glutamine spot is radioactive. Electrophoresis done at pH 8.0 in 0.05 M sodium phosphate, 16 volts/cm for 40 minutes on Whatman 3 MM paper.

C^{12} & C^{14} - GLUTAMINE

C^{12} - GLUTAMIC
ACID