	<u>Page</u>
I. INTRODUCTION.	1
II. BRIEF SUMMARY OF ACCOMPLISHMENTS DURING THE PRIOR YEAR GRANT PERIOD	2
III. PUBLICATIONS DURING THE GRANT PERIOD SPONSORED IN FULL OR IN PART THROUGH THE DEPARTMENT OF ENERGY	15
IV. STUDIES TO BE UNDERTAKEN DURING THE COMING YEAR . <i>Removed.</i>	18
BUDGET. <i>Removed.</i>	21

*Studies of the Formation Chemical Reactivity
+ Properties of Small Clusters Application to
an Understanding of Aerosol Formation and
Heterogeneous Chemistry;*

*Pennsylvania State Univ.
University Park, PA*

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

I. INTRODUCTION

The overall intent of our continuing program is to provide basic underpinning research that will contribute to a quantification of the behavior of radionuclides and pollutants associated with advanced energy activities after these materials emanate from their source and are transferred through the environment to the biota and human receptor. The results of studies on the interaction of electromagnetic radiation with matter provide important data in the area of radiation biology and are also of value in assessing photoinduced transformation of pollutants.

Chemical reactions that proceed following either a photophysical or ionizing event, are directly influenced by the mechanisms of energy transfer and dissipation away from the primary site of absorption. Neighboring solvent or solute molecules can affect this by collisional deactivation (removal of energy), through effects in which dissociating molecules are kept in relatively close proximity for comparatively long periods of time due to the presence of the solvent, and in other ways where the solvent influences the energetics of the reaction coordinate. Research on clusters offers one of the most viable ways of elucidating the molecular details of these processes.

A unique aspect of clusters is that they are comprised of a distribution of subunits that can be shifted by making suitable changes in the conditions employed in their preparation, e.g., temperature and pressure. Femtosecond techniques are especially amenable to investigating processes associated with energy dissipation, and the course of reactions in real time at the site of an ionizing event following the absorption of electromagnetic radiation. Significant insight can be gained through such studies, especially by comparing the findings with the mechanisms of isolated ion-molecule reactions which have a close parallel to many of those of importance in the condensed phase. Work in clusters of considerable value in unraveling structure-function interrelationships in the area of radiation biology where the findings serve to reveal the details of reactions and elucidate the effects due to energy transfer to neighboring bound ligands and effects of solvation due to caging phenomena, or alternatively influence on the energy barriers along the reaction coordinate.

The current program is comprised of three general areas of research: 1) Investigation of the dynamics of ionization and the mechanisms of the early-time reactions following the interaction of ionizing electromagnetic radiation with matter; 2) Measurement of the kinetics of ensuing ion reactions with effort focused on the influence of solvation effects and identifying similarities and differences between gas and condensed phase reactions; and 3) Determination of the structure of solvated reaction centers via spectroscopy, dynamics and thermochemistry.

II. BRIEF SUMMARY OF ACCOMPLISHMENTS DURING THE PRIOR YEAR GRANT PERIOD

During the last year, we have published and/or submitted for publication 13 papers in the open scientific literature that deal with research sponsored in full or in part by this grant. The first four listed in Section III were in press at the date of submission of last years continuation, and have now appeared in print; the relevant publication information is shown. The other 9 represent new publications which were written and submitted after the continuing proposal submission was made. As is evident from our publications, we have had a very productive year; indeed, we have made a number of significant scientific findings, especially the discovery of metallo-carbohedrenes (Met-Cars) which has led to considerable publicity in both the scientific and popular press and some important new findings in the area of ionization in the femtosecond time regime and the suppression of dissociative channels upon the ionization of solvated biological molecules.

The research has proceeded in accordance with the objectives set forth in the proposal for this program which commenced March 1, 1991. During the last report period, we undertook work in all areas. The details are given in the papers to which reference is made in Section III.

One major thrust of the work during the past year has been to studies of the role of ionization on reactions and energetics. Particular attention has been paid to systems having varying proton affinities. From the data, we can determine the influence of solvation on these values and ascertain

the ultimate chemical nature of the ion center formed as a result of the molecular rearrangement reactions. (See papers III-323, 329, 331, 336, 337, 338 and 349.)

Recently, we have demonstrated our capabilities to obtain ultrafast light pulses in time regimes of interest in terms of processes related to radiation effects on matter (i.e., ~ 100 femtoseconds) and have succeeded in ionizing molecules and clusters in this time domain. The completed system we constructed is shown in Figure 3. These include ammonia clusters, water clusters, methanol clusters, trimethylamine clusters, and tryptamine clusters, as well as a number of mixed clusters such as mixed water-ammonia clusters, methanol-water clusters, tryptamine-water clusters, and tryptamine-ammonia clusters, etc. Figure 1 is a typical mass spectrum of tryptamine (TA) molecules ionized by the femtosecond laser pulses (wavelength at 620 nm, pulse duration ~ 150 fs, and pulse energy $\sim 800 \mu\text{j/pulse}$). It is evident that ionization of tryptamine molecules generally initiates fragmentation of the molecular ions, occurring through the loss of NH_2 and CH_2NH_2 fragments. However, when tryptamine is solvated by water (W) molecules, the processes which lead to the fragmentation of tryptamine are quenched since neither $(\text{TA}-\text{NH}_2)(\text{W})_n$ nor $(\text{TA}-\text{CH}_2\text{NH}_2)(\text{W})_n$ are observable in the spectrum. This effect is demonstrated in Figure 2. What processes are responsible for dissipating the energy away from the site of initial deposition of ionizing radiation? In what time domains do these processes take place? Studies of these solvated biological molecules on the femtosecond time scale have great promise for elucidating the roles of solvation on the energy transfer processes and represent an important objective of our continuing program.

Recently, we also commenced limited theoretical studies of the solvation structure around the biological molecules, and their functional groups, for several of the molecules proposed for study herein. The methods include quantum mechanical SCF techniques for local structure determination, and computations using atom-atom potentials for deducing the solvation structures. See Figure 4

Figure 1. Mass spectrum showing the extensive fragmentation of isolated tryptamine upon ionization; see text for experimental conditions.

Figure 2. Mass spectrum of tryptamine solvated with water demonstrating the lack of fragmentation due to solvation. For simplicity of presentation, only a few peaks are labelled.

Femtosecond Pulse Generation and Amplification

Femtosecond Pulse Amplification, Pump-Probe Delay, and Recompression

Colliding Pulse Mode-Locked
Ring Dye Laser (CPM) Bowtie
Amplifier

Figure 3. Femtosecond laser facility in operation in our laboratory for use in the proposed pump-probe experiments.

which shows the calculated solvation structure that accounts for our experimental findings in the context of the suppression of the dissociation of tryptamine.

A particularly interesting and instructive observation which has come from recent femtosecond laser experiments demonstrates that the time domain over which energy is absorbed can often influence the structure at the site of absorption. In particular, using the femtosecond apparatus, we have commenced a study of ammonia and the generation of NH_2 and NH_4^+ ; upon its ionization the former is a major functional group of many biological molecules. The value of detailed experimental findings viewed in conjunction with theoretical calculations, will serve to elucidate structural changes and ultimate mechanisms important in the field of radiation biology.

Studies of proton transfer events following energy absorption enable an investigation of the influence of the adsorption of ionizing radiation on other groups in amino acids such as NH_2 which is present in all of them, and NH_3 which is an important group in glycylglycine. Likewise, study of proton transfer can be investigated following energy absorption in certain other functional groups of amino acids such as in phenol, which is the chromophore of tyrosine. The detailed chemistry of the OH group on tryosine has also been elucidated through investigations of the behavior and reactions of alcohols.

A typical mass spectrum of ammonia clusters ionized by nanosecond laser pulses produces protonated clusters that usually completely dominate the mass spectrum through the formation of NH_2 during the course of the reactions. By contrast, upon ionization in the femtosecond time regime, intact unprotonated clusters are also observed, revealing the differences which can arise due to ionization in different time regimes and hence assessing different regions of the potential energy surfaces. See Figures 5(a) and 5(b).

Studies of the influence of solvation on the absorption of photons in the chromophore of tyrosine, namely phenol, also continued during the report period. These experiments were done to investigate the mechanisms of energy disposal following the absorption of electromagnetic radiation

Figure 5. (a) Mass spectrum of protonated ammonia clusters. (b) Mass spectrum showing the co-existence of intact ammonia clusters upon ionization with the femtosecond time domain.

at a chromophoric site. To accomplish this, we extended the stimulated emission pumping method to a cluster-ion dip spectroscopy technique employing a two-color laser scheme in combination with time-of-flight mass spectrometry. Cluster-ion-dip spectrometry (CIDS), which has been pioneered in our laboratory, is mass selective and is a powerful method for probing the ground-state vibrational spectra of clusters. It offers the potential of obtaining high spectral resolution data. In this method, the frequency of one laser is tuned to a desired transition between the ground and first electronic state, and the excited molecule is then subjected to photons of a second frequency. This results in two-color resonant-enhanced multiphoton ionization from the S_1 state. Tuning the second laser leads to a dip in the ion-cluster spectrum only when its energy coincides with that corresponding to vibrational states in the ground electronic state. The findings have also led to new insight of the influence of solvation on proton transfer reactions effected through the absorption of electromagnetic radiation. (See publications III-329 and 351).

The results of recent ion dip spectroscopy studies of the phenol-OD [PhOD] molecule and phenol-OD(D_2O)₁ [PhOD(D_2O)₁] from 500-1300 cm^{-1} are shown in Figure 6(a) and (b), and the spectra compared with the undeuterated molecule and cluster. The greater modulation of the PhOD ion dip spectrum compared to that of phenol-OH [PhOH] is the result of a longer S_1 lifetime for the deuterated molecule due to quenching of internal conversion. The PhOD(D_2O)₁ cluster spectrum allows us to identify the ground state intermolecular PhOD- D_2O stretch as 144 cm^{-1} compared to 151 cm^{-1} for the undeuterated cluster. Importantly, the blue shift of the 7a mode (C-O stretch) provides evidence that the water acceptor molecule donates charge into the C-O bond.

A fast-flow reactor was also used to produce and investigate ionic water clusters with the general composition $M^+(H_2O)_n$; where M is one of the alkali metals Li, Na, K or Cs and n is up to 45. A series of magic numbers were observed, especially pronounced for $K^+(H_2O)_{20}$ and $Cs^+(H_2O)_{20}$ and to some degree for lithium, and for some other cluster sizes. Evidence was obtained that these stable cluster ions are gas phase hydrates which have direct analogy to the well-known

Figure 6(a). Ion dip spectra of PhOH and PhOD from 500-1300 cm⁻¹ taken through the O⁰ intermediate state. The greater modulation of the PhOD spectrum is due to the increased S₁ lifetime due to the quenching of internal conversion by deuteration. Notice that the intensities of the 12₁⁰/a_{6a}₂⁰ Fermi doublet bands are approximately the same for both isotopomers.

Figure 6(b). Ion dip spectra of the $\text{PhOH}(\text{H}_2\text{O})_1$ and $\text{PhOD}(\text{D}_2\text{O})_1$ clusters from 500-1300 cm⁻¹ taken through the 0_0^0 intermediate state. The intermolecular phenol-water stretch (σ_1^0) is indicated by the dashed bars, built off of the phenol intramolecular vibrations. The intensity of the $12_1^0/16a_2^0$ Fermi doublet bands is quite different for $\text{PhOH}(\text{H}_2\text{O})_1$ through the $\text{PhOD}(\text{D}_2\text{O})_1$ cluster follows that of the PhOD molecule. The $7a_1^0$ bands for both clusters are blue-shifted by $< 8 \text{ cm}^{-1}$.

clathrate hydrates in the condensed phase. (See publications III-331, 336 and 338.) Interestingly, analogous species are thought to play a role in conduction processes in biological systems.

In order to reveal the role which ionization can play on the reactions of molecules in the condensed phase, we commenced studies of simple alkali metal ones generally thought to be inert. Alkali metal-methanol cluster ions $A^+(\text{CH}_3\text{OH})_n$ ($A = \text{Li, Na, K, Rb, Cs}$; n up to 20) were produced and their subsequent reactions investigated using a fast-flow reactor operated under the usual conditions. Surprisingly, dehydration reactions of methanol clusters are observed to occur at degrees of solvation beyond certain critical sizes. The critical size is found to be dependent on the specific alkali metal ion in the clusters, with the observed product ions being $A^+(\text{CH}_3\text{OH})_{n-1}\text{H}_2\text{O}$, where $n \geq 8$ for Li, Na, and K, and $n \geq 9$ for Rb and Cs. We have interpreted the size dependent dehydration reactions on the basis of charge, and solvation effects, and on the reaction energies. (See publication III-349.)

Recently, we discovered a new class of cluster compounds termed metallo-carbohedrenes which are comprised of metal atoms bound within a carbon-cage lattice. (See publications III-353, 356, 358, 360 and 364.) The new molecular clusters are generated through reactions of early transition metals with vapors of small hydrocarbon molecules, utilizing a new laser-based plasma reactor concept. Figure 7 displays a typical ionic cluster distribution of Ti_mC_n^+ using CH_4 as the reactant gas. It is evident that $\text{Ti}_8\text{C}_{12}^+$ is "super" magic in the mass spectrum. We accounted for our observations through suggestions that this species, as well as its corresponding neutral molecule discovered later, has a pentagonal dodecahedron structure, where the Ti atoms occupy eight unique positions that are identically coordinated. Thereafter, findings of magic peaks corresponding to $\text{M}_8\text{C}_{12}^+$ (where M is V, Zr, and Hf) along with prior observations of $\text{Ti}_8\text{C}_{12}^+$, established metallo-carbohedrenes as a general class of stable molecular cluster ions. Following the foregoing work, we thereafter commenced investigations of larger cluster systems. Interestingly, these

Figure 7. Mass spectrum for the metallo-carbohedrene Ti_8C_{12} showing the super magic peak.

revealed a unique structural growth pattern found in the system of Zr_mC_n , i.e., the formation of the multicage structures.

Studies in our laboratory reveal that these species are easily ionized, evidently have delocalized electron character, and may easily bind to the functional groups of biological molecules. If their low ionization potentials are confirmed, these species may be useful as a radiopharmaceuticals under some circumstances, and all may be good scavengers of free radicals. Several research organizations have contacted us to also suggest their possible utility as agents to delivery heavy metals to cancer cells for use in radiotherapy, and also as cages for delivering antiviral agents to selected cells through bonding these to various antibodies.

III. PUBLICATIONS DURING THE GRANT PERIOD SPONSORED IN FULL OR IN PART THROUGH THE DEPARTMENT OF ENERGY, GRANT NO. DE-FGO2-88ER60648 ARE MARKED WITH **.

323. "Evaporative Dissociation of Ammonia Cluster Ions: Quantification of Decay Fractions and Isotope Effects," S. Wei, K. Kilgore, W. B. Tzeng, and A. W. Castleman, Jr., *J. Phys. Chem.* **95**, 8306-8309 (1991). **
324. "Chemistry of Large Hydrated Anion Clusters $X^-(H_2O)_N$, $N=0-59$ and $X=OH$, O, O_2 , and O_3 , III: Reaction of SO_2 ," X. Yang and A. W. Castleman, Jr., *J. Phys. Chem.* **95**, 6182-86 (1991).
325. "Chemistry of Large Hydrated Anion Clusters $X^-(H_2O)_N$, $N=0-59$ and $X=OH$, O, O_2 , and O_3 , II: Reaction of CH_3CN ," X. Yang, X. Zhang and A. W. Castleman, Jr., *J. Phys. Chem.* **95**, 8520-8524 (1991).
326. "Elucidating the Origin of Magic Numbers: Trends in the Relative Binding Energies of Xenon Cluster Ions and Their Implications," S. Wei, Z. Shi, and A. W. Castleman, Jr., *J. Chem. Phys.* **94**, 8604-07 (1991).
327. "Mass Spectrometric Study of the Formation, Evaporation, and Structural Properties of Doubly Charged MgO Clusters," P. J. Ziemann and A. W. Castleman, Jr., *Phys. Rev. B* **44**, 6488-99 (1991).
328. "Evidence for the Polymerization of Ethylene Catalyzed by Ti^+ in the Gas Phase," B. C. Guo and A. W. Castleman, Jr., *J. Am. Soc. for Mass Spectrom.* **3**, 464-466 (1992).

329. "Solvation Effects on the Structure and Reactivity of Clusters," A. W. Castleman, Jr., *Int. J. Quantum Chemistry* **25**, 527-544 (1991). **
330. "Kinetic Study of Neutral Lead Cluster Reactions," R. W. Farley, P. J. Ziemann, R. G. Keesee, and A. W. Castleman, Jr., *Z. Phys. D.*, submitted.
331. "Magic Numbers in Atomic Clusters: Energetic Electronic and Structural Effects," A. W. Castleman, Jr., in *Clustering Phenomena in Atoms and Nuclei* (M. Brenner, T. Lönnroth, and F. B. Malik, Eds.), Springer-Verlag, pp 99-109 (1992). **
332. "Effects of Ligands on the Reactivity of V^+ and Ti^+ Toward Dehydrogenation of Ethylene," B. C. Guo and A. W. Castleman, Jr., *Int. J. Mass Spectrom. Ion Proc.* **113**, R1-R5 (1992).
333. "Laboratory Studies of Large Protonated Water Clusters Under the Conditions of Formation of Noctilucent Clouds (NLC) in the Summer Mesopause," X. Yang and A. W. Castleman, Jr., *J. Geophys. Res.* **96**, 22,573-22,578 (1991).
334. "An Ab Initio Calculation of Potential Energy Surface of Ar_3^+ with MP4 Correlation Correction," Z. Y. Chen, B. D. May, and A. W. Castleman, Jr., *Z. Phys. D.*, in press.
335. "Study of Protonated Methanol Cluster Ions Under Thermal Conditions," X. Zhang, X. Yang, and A. W. Castleman, Jr., *Chem. Phys. Lett.* **185**, 298-302 (1992).
336. "Evidence for the Encagement of Alkali Metal Ions Through the Formation of Gas-Phase Clathrates: Cs^+ in Water Clusters," A. Selinger and A. W. Castleman, Jr., *J. Phys. Chem.* **95**, 8442-8444 (1991). **
337. "Reactions and Properties of Clusters," A. W. Castleman, Jr., to be published in *Advances in Mass Spectrometry*, Vol. 12, Elsevier Science Publishers, in press. **
338. "Encagement of Alkali Metal Ions in Gas-Phase Water Clusters," A. Selinger and A. W. Castleman, Jr., *Proc.*, International Symposium on the Physics and Chemistry of Finite Systems: From Clusters to Crystals, in press. **
339. "Magic Numbers of Metal and Metal Alloy Clusters and Their Chemical Reactivity," Y. Yamada and A. W. Castleman, Jr., *Proc.*, International Symposium on the Physics and Chemistry of Finite Systems: From Clusters to Crystals, in press.
340. "Studies Reactions of Small Titanium Oxide Cluster Cations Toward Oxygen at Thermal Energies," B. C. Guo, K. P. Kerns, and A. W. Castleman, Jr., *Int. J. Mass Spectrom. Ion Proc.*, in press.
341. "Chemistry and Kinetics of Size Selected Cobalt Cation Clusters at Thermal Energies: 1. Reactions with CO", B. C. Guo, K. P. Kerns and A. W. Castleman, Jr., *J. Chem. Phys.* **96**, 8177-8186 (1992).
342. "Chemistry and Kinetics of Primary Reactions of Ti^+ with H_2O , NH_3 , CH_3OH , C_2H_4 , and C_3H_6 at Thermal Energies," B. C. Guo, K. P. Kerns and A. W. Castleman, Jr., *J. Phys. Chem.* **96**, 4879-4833 (1992).

343. "Mass-Spectrometric Investigation of the Stabilities and Structures of Mn-O and Mn-Mg-O Clusters," P. J. Ziemann and A. W. Castleman, Jr., *Phys. Rev. B*, in press.
344. "Structures and Bonding Properties of Ca-O Clusters Inferred from Mass Spectral Abundance Patterns," P. J. Ziemann and A. W. Castleman, Jr., *J. Phys. Chem.* **96**, 4271-4276 (1992).
345. "Chemistry and Kinetics of Size Selected Cobalt Cation Clusters at Thermal Energies: 2. Reactions with O₂," B. C. Guo, K. P. Kerns, and A. W. Castleman, Jr., *J. Phys. Chem.*, in press.
346. "Dehydrogenation of Ethylene and Propylene and Ethylene Polymerization Induced by Ti⁺ in the Gas Phase," B. C. Guo and A. W. Castleman, Jr., *J. Am. Chem. Soc.*, in press.
347. "Magic Numbers of Sodium/Antimony-Clusters Produced by Gas Aggregation and Subsequent Photoionization," A. Hartmann and A. W. Castleman, Jr., *Proc.*, International Symposium on the Physics and Chemistry of Finite Systems: From Clusters to Crystals, Richmond, VA, Oct. 8-12, 1991, in press.
348. "Ti₈C₁₂⁺ -- Metallo-carbohedrenes: A New Class of Molecular Clusters?," B. C. Guo, K. P. Kerns and A. W. Castleman, Jr., *Science* **255**, 1411-1413 (1992).
349. "Influence of Solvation of Dehydration Reactions of Alkali Metal Methanol Cluster Ions," X. Zhang and A. W. Castleman, Jr., *J. Am. Chem. Soc.*, in press. **
350. "The Formation, Growth and Nucleation of Water Cluster Ions," A. W. Castleman, Jr., *Proc. 9th International Conference on Atmospheric Electricity*, St. Petersburg, USSR, June 15-19, 1992, in press.
351. "Ion Dip Spectroscopy of Phenol-OD and Phenol-OD(D₂O)₁," R. J. Stanley and A. W. Castleman, Jr., *J. Chem. Phys.*, submitted. **
352. "Metallo-Carbohedrenes [M₈C₁₂⁺ (M=V, Zr, Hf and Ti)]: A Class of Stable Molecular Cluster Ions," B. C. Guo, S. Wei, J. Purnell, S. Buzza and A. W. Castleman, Jr., *Science* **256**, 515-516 (1992).
353. "Metallo-Carbohedrenes as a Class of Stable Neutral Clusters: Formation Mechanism of M₈C₁₂ (M = Ti and V)," S. Wei, B. C. Guo, J. Purnell, S. Buzza, and A. W. Castleman, Jr., *J. Phys. Chem.* **96**, 4166-4168 (1992). **
354. "Metallo-Carbohedrenes: Formation of Multi-Cage Structures," S. Wei, B. C. Guo, J. Purnell, S. Buzza and A. W. Castleman, Jr., *Science* **256**, 818-820 (1992).
355. "The Magic Numbers of Metal and Metal Alloy Clusters," Y. Yamada and A. W. Castleman, Jr., *J. Chem. Phys.*, in press.
356. "Generation of Metal-Carbon and Metal-Nitrogen Clusters with a Laser Induced Plasma Technique," B. C. Guo, S. Wei, Z. Chen, J. Purnell, S. Buzza, K. P. Kerns and A. W. Castleman, Jr., *J. Chem. Phys.*, in press. **

357. "On Reaction Channels in a Plasma Reactor-Laser Vaporization Source: formation of Carbon Clusters and Metal-Carbon Clusters," Z. Y. Chen, B. C. Guo, B. May, S. Cartier and A. W. Castleman, Jr., *Chem. Phys. Lett.*, submitted.
358. "Metallo-Carbohedrenes -- A New Class of Molecular Clusters," A. W. Castleman, Jr., B. C. Guo, and S. Wei, *Proc. Adriatico Research Conference on Clusters and Fullerenes*, Trieste, Italy, June 23-26, 1992, in press. **
359. "Reactions of Metal Oxide Anions," R. G. Keesee, B. Chen, A. C. Harms and A. W. Castleman, Jr., *Int. J. Mass Spectrom. Ion Proc.*, submitted.
360. "Laser Plasma Reactors: Forming Met-Cars -- A New Class of Cluster Materials," A. W. Castleman, Jr., B. C. Guo, S. Wei and Z. Y. Chen, in *Plasma Physics and Controlled Fusion*, Pergamon Press, in press. **
361. "A New Technique for Producing Binary Clusters: Transition Metal-Silicides," Z. Y. Chen, S. F. Cartier, B. D. May, and A. W. Castleman, Jr., *Int. J. Mass Spectrom. Ion Proc.*, submitted.
362. "Growth of Titanium Nitride: From Cluster to Microcrystal," Z. Y. Chen and A. W. Castleman, Jr., *J. Chem. Phys.*, submitted.
363. "Clathrate Structures in Water-Methanol Mixed Clusters," Z. Shi, S. Wei, J. V. Ford, and A. W. Castleman, Jr., *J. Chem. Phys.*, submitted.
364. "Metallo-Carbohedrenes--A New Class of Molecular Clusters," B. C. Guo and A. W. Castleman, Jr., in *Advances in Metal and Semiconductor Clusters* (M. Duncan, Ed.), JAI Press, invited. **

END

**DATE
FILMED**

2 / 23 / 93