Proceedings of RIKEN BNL Research Center Workshop

Volume 58

RHIC Spin Collaboration Meeting XX

November 18, 2003

Organizer:

Les Bland

RIKEN BNL Research Center

Building 510A, Brookhaven National Laboratory, Upton, NY 11973-5000, USA

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any employees, nor any of their contractors, subcontractors or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Available electronically at-

<http://www.doe.gov/bridge>

Available to U.S. Department of Energy and its contractors in paper from-

U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831
(423) 576-8401

Available to the public from-

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22131
(703) 487-4650

Printed on recycled paper

Preface to the Series

The RIKEN BNL Research Center (RBRC) was established in April 1997 at Brookhaven National Laboratory. It is funded by the "Rikagaku Kenkyusho" (RIKEN, The Institute of Physical and Chemical Research) of Japan. The Center is dedicated to the study of strong interactions, including spin physics, lattice QCD, and RHIC physics through the nurturing of a new generation of young physicists.

During the first year, the Center had only a Theory Group. In the second year, an Experimental Group was also established at the Center. At present, there are three Fellows and seven Research Associates in these two groups. During the third year, we started a new Tenure Track Strong Interaction Theory RHIC Physics Fellow Program, with six positions in the first academic year, 1999-2000. This program had increased to include ten theorists and one experimentalist in academic year, 2001-2002. With five fellows having already graduated, the program presently has ten theorists and two experimentalists. Five of the ten RHIC Physics Fellows have been awarded/offered tenured positions, and this will be their final year in the program.

Beginning in 2001 a new RIKEN Spin Program (RSP) category was implemented at RBRC, presently comprising six RSP Researchers, and six RSP Research Associates. In addition, RBRC has four RIKEN Jr. Research Associates and one RBRC Young Researcher. A number of Visiting Scientists also contribute to the physics program at the Center.

RBRC also has an active workshop program on strong interaction physics with each workshop focused on a specific physics problem. Each workshop speaker is encouraged to select a few of the most important transparencies from his or her presentation, accompanied by a page of explanation. This material is collected at the end of the workshop by the organizer to form proceedings, which can therefore be available within a short time. To date there are fifty-eight proceeding volumes available.

The construction of a 0.6 teraflops parallel processor, dedicated to lattice QCD, begun at the Center on February 19, 1998, was completed on August 28, 1998. A 10 teraflops QCDOC computer is under development and expected to be completed in JFY 2004.

**N. P. Samios, Director
November 2003**

***Work performed under the auspices of U.S.D.O.E. Contract No. DE-AC02-98CH10886.**

CONTENTS

Preface to the Series.....	i
RHIC Spin Collaboration Meeting XX ~ November 18, 2003	
RHIC Working Point	
<i>W. Fischer & R. Tomas</i>	1
Quest for a New Working Point – Status Report	
<i>R. Tomas</i>	5
Helical Dipole Partial Snake for AGS	
<i>J. Takano</i>	13
2 Partial Siberian Snakes in AGS	
<i>W. MacKay</i>	41
RHIC Spin Tune Measurement & Polarization Reversal	
<i>M. Bai</i>	43
Status of PLL	
<i>P. Cameron</i>	49
AGS Polarization – Comments from the 2003 Run	
<i>L. Ahrens</i>	55
Polarized Proton Ramps	
<i>J. van Zeijts</i>	63
Update on the H-Jet	
<i>A. Nass</i>	73
RHIC Polarimeter Status	
<i>H. Huang</i>	87
CNI Polarimetry Status	
<i>O. Jinnouchi & S. Bravar</i>	93

CONTENTS

New Theory Studies for pp Scattering	
<i>W. Vogelsang</i>	105
π^0 ALL Analysis in PHENIX	
<i>K. Tanida</i>	149
STAR Spin Analysis Update	
<i>S. Trentalange</i>	177
List of Registered Participants.....	194
Agenda.....	199
Additional RIKEN BNL Research Center Proceeding Volumes.....	200
Contact Information	

RHIC Working Point

Wolfram Fischer
and
Rogelio Tomas

BROOKHAVEN
NATIONAL LABORATORY

RHIC Spin Collaboration Meeting 2003
17 November 2003

- Working Point
 - = choice of transverse oscillation frequencies
 - **Need to minimize effect of resonances**
- Resonances are driven by
 - Non-linear magnetic fields in triplets
(single beam, independent of intensity)
 - Beam-beam interaction
(two beams, dependent of intensity)
- Collider should run dominated by beam-beam
 - Can correct triplet errors
 - Cannot compensate beam-beam
 - **Need to correct triplet errors first
(IR Correction Task Force)**

- Triplet error correction
 - Low order errors (octupole and sextupole) can be corrected with beam measurements
 - Technique commissioned last Run (F. Pilat), work not yet finished
 - High order errors (12-poles) correction better based on magnet measurements
 - Robust correction being develop (Y. Luo)
 - Need better machine model for corrector calculation
 - Being developed (R. Tomas),
DX, D0, splits, end effects, data base driven
- Goal: Finish IR correction by end of Run-4

Quest for a new working point - status report

Contents

- Background measurements vs. Dynamic Aperture
- Beam Beam effect on Dynamic Aperture
- Possible working points
- IR Task Force preliminary study

Spin Collaboration meeting _____ R. Tomás

Background measurements

dAu, January 2003

Spin Collaboration meeting _____ R. Tomás

Background measurements vs. DA

The model contains beam-beam, triplet errors, 5% σ low frequency orbit modulation, and power supply ripples (no correctors).

Spin Collaboration meeting _____ R. Tomás

Problem: Beam-Beam does not dominate

Next year lattice - DA

β^* 's = 1 1 4 5 3 5 and 1 1 3 5 3 5

⇒ IR non-linear correction is needed

Spin Collaboration meeting _____ R. Tomás

Possible working points - 113535

⇒ SPS working point seems a good candidate
(ISR w.p. ≈ 0.95 will be considered too)

Spin Collaboration meeting _____ R. Tomás

IR Task Force preliminary study

Improvement of the model by a simple slicing of the triplet quadrupoles leads to changes:

⇒ A more refined model of the IR non-linearities is needed and is presently under construction.

Spin Collaboration meeting _____ R. Tomás

Summary

- Simulations and observations show limited agreement
- Beam Beam does not dominate in simulations

⇒ Improved model under construction

- DA could be enlarged in two ways:
 - by non-linear correction in the IRs
 - by changing the tunes to SPS ones, ≈ 0.68 . (ISR tunes, ≈ 0.95 , need further consideration too).

Helical Dipole Partial Snake for AGS

Junpei Takano / TITech

Masahiro Okamura / RIKEN

Koji Takano / Takano Original Magnet CO. LTD.

- Design study
- Optimization
- Fabrication
- Future plan

Design study

Double pitch structure

- spin rotation is 5% (9degree)
- Max B = 1.5 T
- Beam offset and diffraction angle = 0

Double helical pitch structure is very reasonable in space and cost.

Design study

Cross section of helical coil

Shape of helical coil is larger than normal window frame type coil.

Design study

Cross section of helical magnet

Slow pitch region

Rapid pitch region

Coil position is window frame type.
The yoke has pole piece like H type magnet.

Design study

1/5 model of the helical coil

This model is constructed to verify processes of assembling the magnet.

Design study

1/5 model of the helical magnet

This model is constructed to verify processes of assembling the magnet.

Design study

Property of the helical magnet

	Required property	Designed property
Water pressure	200–250 [PSI]	213.5 [PSI] (=15kg/cm2)
Temperature rise	17 [degree C]	14.97 [degree C]
Current	3000 [A]	2671.429 [A]
Voltage	300 [V]	237.44 [V]
Power	900 [kW]	634.3 [kW]
Ramp rate	0.5 [sec]	0.5 [sec]
Duty factor	100 [%]	100 [%]
Weight	18000 [kg]	15000 [kg]

Optimization of helical pitch

L2, P2 : parameter

Optimization of helical pitch

Example of beam trajectory

beam trajectory of x vs. y

beam trajectory of z vs. x, y

If dx and $dy = 0$ then the beam trajectory will be closed.

Optimization of helical pitch

The table of dx and dy when changing the parameter (L2, P2)

	L2	P2	dx	dy
pattern1(ref.)	130	200	-3.158	0.481
pattern2	130	240	-9.881 (-)	1.682 (+)
pattern3	125	200	-4.564 (-)	0.133 (-)

Matrix for dx, dy = 0

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \Delta L2 \\ \Delta P2 \end{pmatrix} = \begin{pmatrix} \Delta dx \\ \Delta dy \end{pmatrix}$$

Solving this matrix,

$$L2 = 131.8371575$$

$$P2 = 183.7732405$$

Optimization of helical pitch

Beam trajectory with optimized parameter

$L1 = 39.08142125[\text{cm}]$, $L2 = 131.8371575[\text{cm}]$

$P1 = 90[\text{cm/rev}]$, $P2 = 183.7732405[\text{cm/rev}]$

Beam trajectory is closed.

Offset and diffraction angle = 0.

Fabrication

Winding helical snake coil

Winding frame

1st layer is wound

Fabrication

Winding helical snake coil

Simulation 3D model

Completed winding

Fabrication

Helical snake coil

Fabrication

Alimentation of coil position

Reference plate is for alimentation of coil and pressing lamination steels.

captured snake in the yoke

Fabrication

Lamination steels

Notch

Transition point

Helical guide

Slow ← → Rapid

Fabrication

Helical pole face

Fabrication

Under construction

Upper side lamination is complete

Fabrication

Upside down

Crane track's work

Fabrication

50% and 100% lamination

After upside down (50% laminated)

Lamination is completed

Fabrication

Outer base frame

Fabrication

Complete the helical magnet

Fabrication

Helical magnet with pipe

148mm diameter pipe is installed.
The designed pole gap is 150mm.

Fabrication

Zoom up the helical pole face

Future plan

High order magnetic field measurement of this helical magnet
will be done at magnet division.

Cross checking the simulation and measurement data.

Adjusting the magnetic field based on the measurement data
before install to beam line.

After installation, the experiment for the effect of this helical magnet will be done.

Future plan

Adjusting

Current = 80% to 110%

We can find out the good operation current point.

Future plan

Operation pattern

Red snake	Blue snake	
On	Off	normal operation before installing the blue snake
Off	On	new operation after blue snake is set up
On	On	the pattern we can run before removing the red snake

After super conductor partial snake is completed,
we can have more combination operating pattern.

2 Partial Siberian Snakes in AGS

— Vertical component of stable spin
— Spin tune

BROOKHAVEN
NATIONAL LABORATORY

From T. Roser

Waldo Mackay

RHIC Spin Tune Measurement & Polarization Reversal

RHIC Spin Collaboration Meeting, Nov.
18, 2003

Mei Bai, BNL, Upton, NY 11973

□ Spin tune measurement

- In resonance rotating frame, the vertical and horizontal projections of the polarization are given by

$$P_y = P_{\vec{A}}^{\circ} \quad P_x = P_{\vec{A}}^{\gg}$$

$$\circ = \frac{(f_{drive} - f_{spin})}{f_{rev}} \quad \gg = \frac{(1/G') B_d L}{4\epsilon B_C} \quad \dot{A} = \sqrt{\circ^2 + \gg^2}$$

- Method 1: map P_y as a function of the spin flipper frequency

spin precession frequency =

spin flipper frequency at which P_y vanishes

- This requires no new development
- However, this is rather time consuming and is not practical for beam at store

□ Spin tune measurement

- Method 2: measure the amplitude of the coherent precession of the horizontal component. The spin tune is then given by

- This requires a modification of the existing polarimeter so that one can measure the turn by turn horizontal component. In principle, this is achievable using CNI as well as STAR polarimeter, according to Gerry and Les.
- However, the accuracy of this measurement highly depends on how close the driving frequency is to the intrinsic spin precession frequency. To achieve an accuracy of 0.001, a $\sim 1.0\%$ error bar on P_x is required assuming the total polarization is about 50% and δ is about 0.01.
- According to Gerry and HH, a 200 sec will yield a 2% P_x measurement provided the total polarization is 50%.

- Method 3: measure P_y while sweeping the spin flipper frequency.

□ Polarization reversal

- F-S formula

$$P_f = P_i \left(2 e^{-\frac{|\mathbf{A} \cdot \mathbf{p}|^2}{2}} - 1 \right)$$

⇒

- Assume total n times spin flipping are performed, let η be the efficiency of each spin flipping. The polarization at the end of the nth spin flipping is then $P_f = P_i \eta^n$. To achieve $P_f > 0.90$

$$\frac{1}{2} \dots \sqrt[n]{0.90}$$

- Say $n=20$, $\eta > 0.995$, with $\epsilon. = 4.6 \times 10^{-4}$, one needs to sweep the spin flipper tune in

0.02 in 57379 revolution turns.

□ Polarization reversal commissioning

. *Three sets of measurements:*

- a) measure the spin flipping efficiency vs. flipper strength. The flipper frequency sweep range as well as the spin tune remain constant.
- b) measure the spin flipping efficiency vs. flipper frequency sweep range with fixed flipper strength as well as the spin tune.
- c) repeat b) with different spin tune.
- with the current snake power supplies, we are able to move the spin tune
- to 0.45 the outer current of one snake should be 94Amps and the inner
- current should be 294.54Amps. The nominal current setting for the snake
- is (inner,outer)=(326.23,100). It takes about 2 minutes to ramp the snake
- current from 326.23 to 294.54.

Status of PLL

P. Cameron, BNL

**for
RHIC Spin Collaboration Meeting XX
November 18, 2003
RIKEN BNL Research Center**

PLL/TF Lessons Learned – in order?

- Humility
- *Stability and Loop Tuning - modeling*
- Signal to Noise, Dynamic Range
- Anomalous phase shifts
- Coupling
- Chromaticity control (not)

Peter Cameron - ENL

Instrumentation Update

1

PLL/Tune Feedback Block Diagram

Peter Cameron - ENL

Instrumentation Update

2

Stability

- Minimize phase shifts – two sources
 - Sampling delay – RHIC 2003 was 4 degrees/Hz with 90Hz update rate
 - BW limiting filter – RHIC 2003 was 40Hz BW, lots of phase shift (100 degrees at 20Hz)
 - Result was loop was twitchy, always close to instability, efforts went into derivative, auto loop gain,...
- For RHIC 2004 (many thanks to Carl Schultheiss)
 - Increasing sampling rate to ~1777Hz
 - Move filter to ~100Hz, switchable digital filters
 - Result is ~x10 improvement in loop bandwidth, gain margin,...

Peter Cameron - ENL

Instrumentation Update

3

Loop tuning with beam 1

- Simple tuning algorithm (if FFT not available) – apply square wave at 10% of desired BW
 - Adjust K_p until for ~10% overshoot
 - Raise K_d until critically damped
 - Raise K_{damp} for ~1% overshoot
- But
 - max dither of ~.001 limits gain to regime of no overshoot?
 - Magnet system BW not sufficient?
 - Questions remain about matrix?

Loop tuning with beam 2

- We do have FFT! Apply modified Ziegler-Nichols tuning (Rasmussen - www.control.auc.dk/~fur/adapt/anto.pdf)
 - Adjust $K_{prop, manual}$ for oscillation, measure frequency
 - Calculate $K_{der, auto}$ and $K_{integr, auto}$ from oscillation period
- But
 - max. dither of ~ 0.001 imposed by machine requirements again limits gains to regime of no oscillation?
 - Dither at oscillation often sufficient to cause excursions outside beam resonance (is this a problem?)
 - Experimental results are questionable

Peter Caspersen - ENL

Instrumentation Update

13

Almost random thoughts on loop tuning

- Loop BW is not an explicit function of S/N
 - Loop BW is a function of loop gain
 - Maximum loop gain is a function of phase and gain margins
 - Phase and gain margins are a function of S/N
- BTF portion of loop gain varies considerably during ramp
 - We can (and do) control signal amplitude with kicker power feedback - hopefully this corrects BTF variation of loop gain
- What's left? Tune and 'noise' fluctuations. We require sufficient gain and phase margin to accommodate tune and noise fluctuations during ramping.
- Beam has memory
 - increases effective loop gain? How do we model this?
- Magnet BW sufficient for tuning? Use beam-beam?

Peter Caspersen - ENL

Instrumentation Update

14

PLL Lessons Learned - in order?

- Humility
- Stability and Loop Tuning - modeling
- *Signal to Noise*
- Anomalous phase shifts
- Coupling
- Chromaticity control (not)

Peter Caspersen - ENL

Instrumentation Update

15

Signal to Noise (need ~ 20 dB minimum)

- Loop bandwidth (and filtering)
 - As low as possible to maximize S/N
 - As high as possible to minimize phase shift, maximize loop stability
 - minimum $\sim \times 10$ magnet system BW
- Revolution line - saturation and dynamic range
 - Moveable BPM, feedback to keep it centered
 - Hotter mixers
- Kick strength and emittance growth
 - Kicker power < 100 mW
 - Sensitive to linewidth
 - Feedback on kicker power essential

Peter Caspersen - ENL

Instrumentation Update

17

PLL Lessons Learned – in order?

- Humility
- Stability and Loop Tuning - modeling
- Signal to Noise
- *Anomalous phase shifts*
- Coupling
- Chromaticity control (not)

Peter Cascone - ENL

Instrumentation Update

20

PLL Lessons Learned – in order?

- Humility
- Stability and Loop Tuning - modeling
- Signal to Noise
- Anomalous phase shifts
- *Coupling*
- Chromaticity control (not)

Peter Cascone - ENL

Instrumentation Update

22

Coupling correction on the ramp

- During tune meter kicks
- Schottky - passive
 - Far free from LFS, more difficult from HFS
 - Need GPM, include in Schottky
 - Motion control, improved S/N
- PLL – skew quad modulation (Roser)
 - First efforts during last run's Beam Experiments
 - Small (~10-4) tune modulations give good signal
 - Supplement to Artus/BPM method when beam is kicked
 - Gives coupling when beam is not kicked

Peter Cascone - ENL

Instrumentation Update

24

PLL Lessons Learned – in order?

- Humility
- Stability and Loop Tuning - modeling
- Signal to Noise
- Anomalous phase shifts
- Coupling
- *Chromaticity control (not)*

Peter Cameron - DES

Instrumentation Update

25

Chromaticity Effect on PLL

- Conclusion from the chromaticity study is that PLL tune measurement can cope with a large range of chromaticity
- Chromaticity control is not a primary issue for PLL tune measurement and tune feedback
- Chromaticity control is an issue primarily in the usual operational sense

Peter Cameron - DES

Instrumentation Update

26

Conclusions

- Not obvious that much more can be done to improve S/N with the present approach (superconducting pickup?)
- Remaining improvements will be in loop tuning
- Testing on resonator
 - refine PID
 - improve model (include beam memory? How?)
 - open-loop measurements
- Startup - single ring, single bunch; then six bunches
 - tune PID (on background first, then modulate magnets) - one and six bunches
 - close magnet loop and modulate - six bunches
- Startup - both rings, six bunches
 - Use beam-beam for fast efficient tuning?
- Warm and Fuzzy? Not yet; still in a new realm
- Feedforward?

Peter Cameron - DES

Instrumentation Update

27

AGS Polarization Comments from the 2003 Run

L. Ahrens, BNL

for
RHIC Spin Collaboration Meeting XX
November 18, 2003
RIKEN BNL Research Center

1) The "Gauss Clock" timing framework seems adequate.

AC dipole timing scans give polarization dependence close to expectations.

Magnetic "flat-tops" at correct $G\gamma$. No tuning of the intermediate porches necessary. If Woody's sharp polarization loss spikes are at (.3 and .7) on 46.5 porch then our position is nearly perfect. Fig 1

However calibration measurements no so robust during the run. (gold work)

2) Intensity dependence of measured polarization in AGS (?) and the poor polarization over the last shift of the run - effects not yet cleanly understood.

Plot of some (Pol vs Int) from the last two weeks of the run. Fig 2

3) AC Dipole behavior and associated beam behavior.

Deterioration (over the run) of the beam coherent motion during the dipole pulse. That this structure was getting "worse" was not particularly obvious and does not particularly correlate with polarization loss. But it may be important. We need to quantify the structure and pay attention to it. Fig 3 & 4

The (magnet-power supply) system for the AC dipole was not healthy. Will be better next time.

4) Up-the-Ramp measurement clearly showed no polarization loss as the beam passed through the $G\gamma=48$ -v resonance region, where a 10 % loss is expected. Fig 5

The measurement looks good. Can our machine and model differ enough to explain this?

Maybe we can make some measurements of the AGS beta function around the machine using the response of the beam position monitors to dipole orbit distortions.

46.5 bumpless porch field scans
runs 238-252; 759-775

◆ ms 238-252
X ms 759-775

AGS Flat Top Magnetic Field Setting (kGauss)

Fig 1

21 March 03

fig 3 (early)
mm

mid-May '03

figure 4 (late in run)

http://www.cadops.bnl.gov/cgi-bin/machlog.pl?nb=rhic_pp_fy03&action=view&page=-6909 11/17/03

fig 5

Polarized Proton Ramps

Johannes van Zeijts
CAD/AP

Outline

- FY03 ramp as designed
- FY03 ramp optics analyzed
- FY04 proton ramp issues

FY03 Ramp Design

Blue Design Beta

FY03 model tunes- part I

Quadrupole Transfer Functions

- 13cm quadrupoles: Q 1/2/3
 - $dT/T = 0.0028$
- 8cm quadrupoles
 - $dT/T = -0.0026$

FY03 model tunes – part B

Lattice Functions at Store

Beta* at Store

- Not quite as designed:
 - STAR 1m $\rightarrow$ 0.93/0.86 (B) , 1.23/0.91 (Y)
 - Phenix 1m $\rightarrow$ 1.22/0.89 (B), 1.41/0.91 (Y)
 - Phobos 3m $\rightarrow$ 2.96/2.51 (B), 2.30/2.51 (Y)
 - Brahms 3m $\rightarrow$ 2.81/3,33 (B), 3.06/3.62 (Y)
- Effect on Luminosity
 - $6/8 = 1.12$, observed ratio = 1.14
 - $8/10 = 2.36$, observed ratio = 2.3

Proton Ramp Issues

- 'Regular' ramp issues
 - Minus transition
- Add path length changes from snake/rotators
- With new TF knowledge (to be confirmed..)
 - Can just squeeze on ramp
- How do we want to handle the rotators?
 - Too much time wasted (20 min.) at store
 - Always on at full field?

Conclusion

- FY03 pp ramp very useful for AP analysis

Update on the H-Jet

1. High frequency transitions installed and tested.
2. Holding field magnet powered up.
3. Running under PLC control.
4. Modifications to the IR completed.
5. Moved the jet to the IR.
6. Water cooling system completed.
7. Jet systems running in the IR.
8. Started to monitor the jet through the VME crate.
9. Results of the Safety Review.
10. Next steps.

The Jet

Achieving polarized atoms in the Jet

Hyperfine structure of hydrogen in an external magnetic field.

$$|1\rangle = \left| +\frac{1}{2}, +\frac{1}{2} \right\rangle$$

$$|2\rangle = \cos \theta \left| +\frac{1}{2}, -\frac{1}{2} \right\rangle + \sin \theta \left| -\frac{1}{2}, +\frac{1}{2} \right\rangle$$

$$|3\rangle = \left| -\frac{1}{2}, -\frac{1}{2} \right\rangle$$

$$|4\rangle = \cos \theta \left| -\frac{1}{2}, +\frac{1}{2} \right\rangle - \sin \theta \left| +\frac{1}{2}, -\frac{1}{2} \right\rangle$$

with $\theta = \frac{1}{2} \arctan \frac{B_c}{B}$

$B_c = 0.507 \text{ kGauss}$

Achieving polarized atoms in the Jet

Calculation of the nuclear polarization:

$$P_z = n_1 - n_3 + (n_4 - n_2) \cos 2\theta$$

- Focussing atoms with electron spin $+\frac{1}{2}$ into the target region while defocussing atoms with electron spin $-\frac{1}{2}$ by 2 sets of sextupole magnets using the Stern-Gerlach effect
- Exchange of occupation numbers of states $|1\rangle$ and $|3\rangle$ ($|2\rangle$ and $|4\rangle$) with the WFT (SFT) to obtain a negative (positive) nuclear polarization.
- $\Rightarrow$ 95% (96%) maximal polarization at 1kGauss (1.2 kGauss) holding field.

High frequency transitions and holding field magnet

- High frequency transitions installed in a shielding box to reduce z-field in the transition region.
- Installed water cooling to remove the heat from the coils.
- Measured efficiencies of all transitions to be $> 99.5\%$.
- Installed 2 beam blockers to remove states $|3\rangle$ and $|4\rangle$ and molecules completely from the BRP beam.
- Main holding field magnet powered up without problems. No polarization losses due to non adiabatic conditions of the field.
- $\Rightarrow > 94.5\%$ (95.5%) polarization at 1kGauss (1.2 kGauss) field.

PLC control

- Setup of a PLC system to completely interlock the Jet.
- Hardware now protected in case of failures (power, water, comp. air ...).
- PLC can be monitored and controlled by a PC.
- Additional interlocks for power supplies of holding field magnet, HFT coils and HFT RF (temperature, cooling water, vacuum).
- Readout of vacuum gauges, water flow meters, valve and for-pump status via ADC of PLC.

Modifications and moving to the IR

- Crane installed and tested in the IR.
 - Rails mounted on the floor.
 - Cable trays installed from the service building to the IR.
 - Cables pulled and terminated for the whole system.
-
- Moving started October 9th by splitting the jet in 3 parts:
 1. ABS stage (3 upper chambers)
 2. RHIC and BRP stage (6 chambers)
 3. Racks
 - Riggers moved all 3 parts out of Bldg. 930 to Bldg. 1012.
 - Items 1 and 2 $\Rightarrow$ IR, 3 $\Rightarrow$ service building.
 - Moved part 1 ontop of the rails with the crane.
 - Moved part 2 ontop of 1 using a forklift (only this time, modifications on the crane in progress).
 - Beamline not in place.
 - Connected cables between Jet and racks.
-
- $\Rightarrow$ whole moving will need 1 day of RHIC down.
+ an additional day for leak checking.

Water cooling system

- Heat exchange system located in the service building.
- Water lines running to the IR as well as to the racks.
- Cooling needed for the holding field magnet, the turbo pumps, the RF-coils, the nozzle cold head compressor and the RF-amplifier for the dissociator.
- Additional chiller in the IR for the dissociator tube cooling.
- Closed cycle system using tap water for heat exchange at the moment.
- Filters installed to clean up the system to prevent blocking of the small conductance holding field magnet lines.
- Decision to use cooling tower or chiller for heat exchange will follow.

Jet running in the IR

- After fixing some leaks start of all pumps. Pressures achieved $\approx 10^{-8}$ Torr without hydrogen beam (chamber in RHIC vacuum $7 \cdot 10^{-9}$ Torr without RHIC pumps running).
- Dissociator was running but intensity dropped after some hours running.
- Exchange of the glass tube (2 mm thickness $\Rightarrow$ 1mm thickness) $\Rightarrow$ stable operation at a high intensity level.
- Running the high frequency transitions without any big problems. Stable operation and high polarization.
- Holding field magnet powered up and running without problems (up to 1.2 kGauss).
- No problems found when running all devices with the 50 m cables instead of the short cables before.
- Replaced (50 m) BRP gauge cable with shielded one and improved power supply to increase the signal to noise ratio.
- Now testing for long term stability.
- Started to monitor the jet via VME crate.
- Using 'pet' pages and archiving to monitor the different parameters.
- Cabling on the way to fully monitor and digitally control the jet.

Next Steps

- When vacuum established in the beam lines adjoining the jet,
⇒ open the vacuum valves and measure the true jet beam gas load.
- Control the power supplies through the VME crate and PET control.
- Some minor things to fix after Safety Review Committee review and initial walkthrough.
- Move out of the IR on November 20th with beamline in place.
- Move back to Bldg. 930 and continue long term stability tests as well as control and measurement of the polarization via VME.
- Focus on preparations to install the silicon detectors on the jet (flanges, collimators, etc.).
- Developing and constructing the H_2 (H_1 , H_2O) monitor
- Check alignment of the whole system.

RHIC Polarimeter Status

H. Huang, BNL

for
RHIC Spin Collaboration Meeting XX
November 18, 2003
RIKEN BNL Research Center

RHIC Polarimeter Status

- All targets and Si detectors have been installed at the end of October. The targets survived the vacuum pump down. All Si detectors show reasonable signals from the Am source.
- The 90 degree Si detectors are in parallel with beam direction now: no angle information available anymore and distribute the events originally at central strip.
- ∞ • All electronics boards are out to test the oscillation problem. They will be put back before pp run.
- All electronics will be moved into the jet target control hut to share electronics with jet target and convenient for control purpose.
- Need a fast turn around time for the coming run. We estimate that a 12 hours dedicated time (spread over days) is needed for polarimeter commissioning. Expect the polarimeter to be ready within the first week.

AGS CNI Polarimeter

pC Analyzing Power

FY03 AGS Polarimeter Performance

About 10 min measurement to give 2.6% errorbar for 6×10^{10}

In the long run, the polarimeter is required to handle

$\sim 2 \times 10^{11}$ /bunch intensity

Two limitations in last run: oscillation in electronics induced by bunch passing (proportional to beam intensity) and data processing speed

• DAQ can only handle 600K/spill (1.3×10^{11} /spill for 2 sec.) with 5 sec. rep. rate. This was a limitation in multiple bunch running mode.

• Narrower targets below 200 μm broke before testing with beam.

• Si detectors are dying near the end of run.

AGS Polarimeter Modification

If don't care of measuring time, we can always reduce the target-in time to meet the DAQ constraint. For the oscillation problem, we can raise the t value (with lower A_N).

- Reduce target width by a factor 2. 250-300 μm target. Delivery schedule is under discussion.
- Increase distance by 1.3. Add spacer to increase distance between target and Si detector from 25 cm to 33 cm.
- Put two Si detectors in one flange. Flange design is almost done.
- New shaper board is in design stage and should be ready by Feb.
- Install peltier cooler to extend lifetime of the detector.
- Faster DAQ using USB 2.0 CAMAC controller (increase speed by about 4 times).
- Continue testing narrower target while modify motor running profile to smooth out target motion as much as we can.

AGS Polarimeter Modification (2)

- Acceptance increase by $2(\# \text{ of Si detectors})/1.3(\text{distance increase})=1.6$
- Target width reduces by $\frac{1}{2}$.
- Increase bunch intensity by $(2*10^{11}/0.6*10^{11})=3.3$.
- The event rate will be higher than last year by about 2.5 times and reduce measurement time from 10 min. to 4 minutes. This will be within the new DAQ system capability.
- For 6 bunch running mode, the event rate will be higher and may require further upgrade on DAQ system.

CNI polarimetry status

Osamu Jinnouchi (RIKEN BNL Research Center)
/ Sandro Bravar (BNL)

RHIC-Spin-Collaboration Meeting

November 18, 2003

Update of RHIC polarization status

- During the run, polarization from the scaler numbers was distributed to the experiments. The number depends on,
 - $T_{1/4}$, ADC_{max} determined by WFD algorithm
 - WFD look-up-table banana cut
- This online results were defined as tentative “official polarization” for run-03
- The calibration and correction were done at some level at the online stage, and the recoil Carbon energy is one of the most important value which determine the AN
- Energy calibration (Correction)
 - Scale – 241 Am source
 - Stable within a few % throughout the run (c.f. 20% in RUN-02)
 - Scale and shift –Energy loss in SSD dead layer
 - Estimated by kinetic fit to E-T banana curve
 - Would generate 50-80keV shift in detection energy

Comparison WFD algorithm vs. Waveform

- (A, T) values from the WFD FPGA algorithm were compared with the result from waveform data samples
- Clear correspondence in each event
- The Event mode data and scaler mode data were confirmed to be correct for the first time
- Storing the timing info at the maximum pulse height will be also useful

SSD dead layer estimation

- Ambiguity about dead layer width is one of the systematic error source in AN $\rightarrow$ polarization error
- The width could be different in each Si detector, and each strip
- It can be roughly (and indirectly) estimated by kinetic fit for banana distribution
- Direct measurement using the test beam was done during the last three weeks at Tandem in Kyoto (Basic data sets were collected for 3 SSDs) The run was very successful !!
- It is desirable to perform this test for all the detectors, each year – Since this is time consuming, we need to develop test bench system in Lab space

The radial component (at BLUE Flattop)

False asymmetry (radial component) had been seen throughout the Run-03
The effect was significant and there are many possible candidates for the source

O.Jinnouchi @ RHIC-Spin-J

Nov. 14, 2003

Limitation of the square root formula

- Square root formula (actually any other formula) can generate false asymmetry with 45 degree SSDs, when A_N of each detector is different
- Also it is possible to generate false asymmetry by just eliminating the suspicious S_i strip channels
- A_N for each strip would be the useful parameter to look at so that we can visually understand the anomalous behavior of asymmetries

$$\text{The Ratio}(i) = (U_i - R D_i) / (U_i + R D_i)$$

U_i : Spin Up counts for i -th strip

D_i : Spin Down counts for i -th strip

R : luminosity ratio L_u/L_d

represents A_N for each strip

AN for each strip (in YELLOW)

Ratio (AN) is corrected with the sign (1-3 for plus and 4-6 for minus) and factor $\sqrt{2}$ for 45 degree SSDs

- Both energies in yellow can indicate the tilted spin from vertical
- Opposite direction btw injection and flattop
- Other things look to be normal

O.Jinnouchi @ RHIC-Spin-J

Nov. 14, 2003

A_N for each strip (in BLUE)

- A_N for BLUE Si-3 is extremely small and this has created the large asymmetry in radial direction
- The reduction cannot be explained by,
 - Wrong dead layer estimation
 - Wrong energy scale
 - Banana cut is not right
 because the size is too significant

Mass distribution

- One way to check the consistency in kinematics is to see the mass distribution
- The width has similar behavior, but centroid has different slope only in Si-3

Mass distribution for
Every strips in Si-3

Mass vs. C Energy

Beam position was off-centered

- The data from BPM and survey group indicates the target (and beam) was not centered w.r.t. polarimeter chamber
- The effective angle for each 45 degree detector (= effective A_N) was changed ($\cos 45 = 1/\sqrt{2} \rightarrow ??$)
- The size of this effect can even become $\pm 10\%$ according to the geometrical displacement
- Only Si-3 could be largely affected while the others get small effects

Summary

- Algorithm of WFD has been studies with waveform analysis
- SSD dead layer measurement in Kyoto successfully finished
- To understand the anomaly in Blue flattop is the top priority
- The mechanism has been understood, then we need to find out the source and how to avoid it in the run-04

New theory studies for pp scattering

Werner Vogelsang

RIKEN-BNL Research Center / BNL Nuclear Theory

RSC meeting, BNL, November 18, 2003

Outline :

- Near-term prospects for measuring Δg at RHIC
 - * high- p_T jet / hadron production
 - * sign of Δg ?
- Single transverse-spin asymmetries

Near-term prospects for measuring Δg

– High- p_T jets and hadrons –

$$p_T^3 \frac{d\Delta\sigma}{dp_T d\eta} =$$

$$+ \mathcal{O}\left(\frac{\lambda}{p_T}\right)^n$$

$$d\Delta\sigma^{pp \rightarrow hX} = \sum_{abc} \Delta f_a \otimes \Delta f_b \otimes d\Delta\hat{\sigma}^{ab \rightarrow cX'} \otimes D_c^h + \text{P.C.}$$

$$\begin{array}{ccc} \Delta\hat{\sigma}^{(0)} & + & \alpha_s \Delta\hat{\sigma}^{(1)} + \dots \text{pert.} \\ \text{LO} & & \text{NLO} \end{array}$$

• nucleon spin structure $\Delta f_{a,b}$

• for hadrons : fragmentation functions $D_c^h \rightsquigarrow e^+e^-$ annihilation

- NLO = state of the art

LO

NLO

- lowest order : good for qualitative descriptions
- however, precise predictions afford higher-order NLO calculations :
 - may be sizeable, in particular in polarized case
 - reduction in scale dependence

$$\mu \frac{d}{d\mu} d\sigma_{\text{phys}} = 0$$

$$\neq 0 \quad \text{in truncated perturbation theory}$$

- NLO corrections now known for all relevant reactions

Gordon, WV; Contogouris et al.; de Florian, Frixione, Signer, WV; Stratmann, Bojak;
de Florian; Jäger, Schäfer, Stratmann, WV; . . .

- jet and hadron calculations share many features
- however :

- final-state singularities
 - cancel for jets
 - imply additional non-perturbative input for incl. hadrons
→ fragmentation functions

- fragm. fct. analyses : Kniehl,Kramer,Pötter and Kretzer

Unpolarized cross section

$pp \rightarrow \pi^0 X$ by PHENIX

113

(Why so good . . . ?)

Apanasevich et al. (see also : Aurenche et al.)

- contributions by partonic scatterings : pions

- contributions by partonic scatterings : jets

- contributions by partonic scatterings :

- average x and z :

- average x and z in forward region :

Kretzer

Spin asymmetry

Polarized gluon densities : Glück,Reya,Stratmann,WV

$pp \rightarrow \pi^0 X, \sqrt{S} = 200 \text{ GeV}, |\eta| \leq 0.38$

Jäger, Schäfer, Stratmann, WV

- latest development : first preliminary data from PHENIX !

- $\Rightarrow$ can A_{LL}^{π} be negative ? How negative can it be ?

- partonic subprocess asymmetries $\hat{a}_{LL} \equiv \frac{\hat{\sigma}_{++} - \hat{\sigma}_{+-}}{\hat{\sigma}_{++} + \hat{\sigma}_{+-}}$

- $gg \rightarrow q\bar{q}$ has negative $\hat{a}_{LL}$
- however, $\hat{\sigma}_{gg \rightarrow gg} \approx 160 \times \hat{\sigma}_{gg \rightarrow q\bar{q}}$ (at $\hat{\eta} = 0$)

A lower bound on A_{LL}^π

Jäger, Kretzer, Stratmann, WV

- integrate LO cross section over all rapidities $|\eta| \leq \cosh^{-1}(x_T)$,
and take mellin moments in $x_T^2 \equiv 4p_\perp^2/S$:

$$\Delta\sigma^\pi(N) \equiv \int_0^1 dx_T^2 (x_T^2)^{N-1} \frac{p_\perp^3 d\Delta\sigma^\pi}{dp_\perp}$$

126

- obtain

$$\Delta\sigma^\pi(N) = \sum_{a,b,c} \Delta f_a^{N+1} \Delta f_b^{N+1} \Delta \hat{\sigma}_{ab}^{c,N} D_c^{\pi,2N+3}$$

- in terms of moments of Δg :

$$\Delta\sigma^\pi(N) = (\Delta g^{N+1})^2 \mathcal{A}^N + 2 \Delta g^{N+1} \mathcal{B}^N + \mathcal{C}^N$$

$\uparrow$
 gg scatt.

$\uparrow$
 qg

$\uparrow$
 $qq, qq', q\bar{q}, \dots$

- a parabola – with a minimum given by

$$\mathcal{A}^N \Delta g^{N+1} = -\mathcal{B}^N$$

$$\Delta\sigma^\pi(N)\Big|_{\min} = -\frac{(\mathcal{B}^N)^2}{\mathcal{A}^N} + \mathcal{C}^N$$

- Mellin inverse :

$$\frac{p_\perp^3 d\Delta\sigma^\pi}{dp_\perp}\Big|_{\min} = \frac{1}{2\pi i} \int_\Gamma dN (x_T^2)^{-N} \Delta\sigma^\pi(N)\Big|_{\min}$$

- resulting asymmetry : negative, but tiny

$$p_\perp = 1.5 \text{ GeV} :$$

$$A_{\text{LL}}^\pi \approx -10^{-4}$$

$$p_\perp = 4.5 \text{ GeV} :$$

$$A_{\text{LL}}^\pi \approx -10^{-3}$$

- the polarized gluon that minimizes A_{LL}^π :

- a node – helps by allowing $\Delta g(x_a) \times \Delta g(x_b) < 0$
- a full “global analysis” confirms :
pQCD at leading power predicts that a negative A_{LL}^π is small

$pp \rightarrow \text{jet} X, \sqrt{S} = 200 \text{ GeV}, 0 \leq \eta \leq 1$

de Florian, Frixione, Signer, WV

$$pp \rightarrow \pi^0 X, \quad \sqrt{S} = 200 \text{ GeV}, \quad \eta = 3.3$$

- dominance of $qg \rightarrow qg$ sets in

π^+ and π^-

positive Δg : $A_{LL}^{\pi^+} > A_{LL}^{\pi^0}$
 negative Δg : $A_{LL}^{\pi^+} < A_{LL}^{\pi^0}$

positive Δg : $A_{LL}^{\pi^-} < A_{LL}^{\pi^0}$
 negative Δg : $A_{LL}^{\pi^-} > A_{LL}^{\pi^0}$

... only at $p_T > 5$ GeV, good statistics required

Single-transverse spin asymmetry A_N

- correlation $\sim i \vec{s}_T \cdot (\vec{p}_{\text{beam}} \times \vec{l}_\pi)$

- * large A_N seen in fixed-target experiments at BNL,ANL,Fermilab,Serpukhov

- * E704 ('96) :

- A_N is power-suppressed as $1/p_T$...

STAR

* higher p_T : see expected $1/p_T$ behavior ?

- Sivers '90 :

$$f_{1T}^\perp = \text{[Diagram 1]} - \text{[Diagram 2]}$$

The diagram shows the difference between two parton distribution functions. The first diagram (left) represents $f_{1T}^\perp$ and shows a parton with momentum q and transverse momentum k_T inside a nucleon with momentum P and transverse spin S_T . The second diagram (right) shows the same parton and nucleon, but with the transverse spin S_T reversed. The difference between these two diagrams is the Sivers function $f_{1T}^\perp$.

asymmetry

$$q_{p\uparrow}(x, \vec{k}_T) \neq q_{p\uparrow}(x, -\vec{k}_T)$$

- mechanism is

$$A_N \propto f_{1T}^\perp(x_a, k_\perp) q(x_b) \hat{\sigma}^{qq \rightarrow qq}(k_\perp) D(z)$$

- however Collins '93 : for distribution functions correlation $\vec{S}_T \cdot (\vec{P} \times \vec{k}_T)$ ruled out by T invariance of QCD
- recent development : gauge links that make pdfs *gauge invariant* allow the “T-odd” structure

(Brodsky,Hwang,Schmidt; Collins; Belitsky,Ji,Yuan; Boer,Mulders,Pijlman)

- A_N for a single-inclusive observable :
 - * interpretation complicated : Sivers vs Collins vs Qiu/Sterman,Koike
 - * $p^\uparrow p \rightarrow \gamma X$, $p^\uparrow p \rightarrow \mu^+ \mu^- X$, $p^\uparrow p \rightarrow \text{jet } X \dots$
 - * factorization with $k_\perp$ an *assumption*
- there is a class of observables that depend directly on a measured k_T :
 - * azimuthal asymmetries in $ep^\uparrow \rightarrow e'\pi X$
 - * azimuthal asymmetries in $e^+e^- \rightarrow \pi\pi X$
 - * $p^\uparrow p$ scattering ?
- for these, $k_\perp$ effects are not power-suppressed,
and more experience with factorization is available

- example : $ep^\uparrow \rightarrow e\pi X$

Collins

$$\sin(\phi + \phi_S) \sum_q e_q^2 \delta q(x) H_1^{\perp,q}(z)$$

$$\sin(\phi - \phi_S) \sum_q e_q^2 f_{1T}^{\perp,q}(x) D_q(z)$$

- HERMES

- only moderate Q^2

What's a similar observable in pp ?

- example : near back-to-back jets in $p^\uparrow p \rightarrow \text{jet jet } X$

Boer, WV

STAR

- A simple estimate : Boer,WV
- Gaussian $k_{\perp}$ distribution for partons in unpolarized proton :

$$f(k_{\perp}) = \frac{e^{-(k_{\perp})^2 / \langle k_{\perp}^2 \rangle}}{\pi \langle k_{\perp}^2 \rangle} \quad (\text{mostly gluons})$$

- leads to distribution

$$\propto e^{-a (1 - \cos(\Delta\phi))}$$

- fit to STAR back-to-back data :

$$\sqrt{\langle k_{\perp}^2 \rangle} \approx 1.3 \text{ GeV}$$

- Sivers type correlation :

$$f_{1T}^{\perp}(k_{\perp}) \propto k_{\perp} e^{-(k_{\perp})^2 / \langle r_{\perp}^2 \rangle}$$

- leads to spin asymmetry

$$A_N \propto \left(|\vec{P}_c^{\perp}| \sin \phi_c + |\vec{P}_d^{\perp}| \sin \phi_d \right) \frac{2 \langle r_{\perp}^2 \rangle}{(\langle r_{\perp}^2 \rangle + \langle k_{\perp}^2 \rangle)^2} e^{-[\vec{P}_c^{\perp} + \vec{P}_d^{\perp}]^2 / (\langle r_{\perp}^2 \rangle + \langle k_{\perp}^2 \rangle)}$$

- for $\vec{P}_c^{\perp} \parallel \vec{S}_{\perp}$

$$\propto \sin(\Delta\phi) e^{-a(1-\cos(\Delta\phi))}$$

- use parameters of Anselmino, D'Alesio, Murgia
- very sensitive to gluon Sivers function
- future : perturbative tail, Sudakov effects

- d'Alesio, Murgia

$$\sqrt{S} = 200 \text{ GeV}, 8 \leq p_{T1,2} \leq 12 \text{ GeV}, -1 \leq \eta_{1,2} \leq 1$$

Boer, WV

Conclusions :

$$\Delta g$$

- a lot can be learned from jets and hadrons
- sign of Δg from forward region ?
- most theoretical tools in place – some challenges remain

146

$$A_N$$

- look for $1/p_T$ behavior of $p^\uparrow p \rightarrow \pi X$
- large potential for back-to-back observables

- physics importance of Sivers function :

- * interference of $J_z = +\frac{1}{2}$ and $J_z = -\frac{1}{2}$ amplitudes
→ orbital angular momentum

- * Burkardt : connection $q(x, \vec{k}_T) \leftrightarrow q(x, \vec{b}_T) \leftrightarrow H, E$ "GPD's"

- * universality ?

$$f_{1T}^\perp|_{\text{DY}} = -f_{1T}^\perp|_{\text{DIS}} \quad \text{Collins; Brodsky, Hwang, Schmidt;}$$

Belitsky, Ji, Yuan; Boer, Mulders, Pijlman; Anselmino, D'Alesio, Murgia

π^0 A_{LL} analysis in PHENIX

K. Tanida (RIKEN/RBRC)
RSC meeting 11/18/03

Outline

- Relative luminosity
 - estimation of systematic error
 - A_{LL} in ZDC/BBC
- LocalPol
- Counting number of π^0 's
- Result

A_{LL}

$$A_{LL} = \frac{\sigma_{++} - \sigma_{+-}}{\sigma_{++} + \sigma_{+-}} = \frac{1}{|P_B P_Y|} \frac{N_{++}/L_{++} - N_{+-}/L_{+-}}{N_{++}/L_{++} + N_{+-}/L_{+-}}$$

++ same helicity

+- opposite helicity

(P) Polarization

(L) Relative Luminosity

(N) Number of pi0s

How to get relative luminosity?

- Principle: $R = N_{++}/N_{+-}$ -- from any counter (trigger)
- BBCLL1 trigger
 - coincidence of two Beam-Beam Counters
 - small background ($\sim 10^{-4}$)
 - high statistics (detects $\sim 1/2$ of all collisions)
 - A_{LL}^{BBC} : expected to be small, need confirmation

Method of accuracy estimation

- Cross-check by ZDCCLL1 trigger
(Zero Degree Calorimeter)
 - low background ($\sim 10^{-4}$)
 - statistics: moderate ($\sim 3\%$ of BBCLL1)
- $N(\text{ZDCCLL1})/N(\text{BBCCLL1})$ must be constant
- Crossing-by-crossing, fill-by-fill ratio calculation
Blue: + - + - + - + -
Yellow: + + - - + + - -
→ We have all spin combinations in each fill
- GL1p scaler counts both BBCCLL1 and ZDCCLL1 for each crossing
- Required accuracy: constant to 10^{-3} level for each fill.

crossing-by-crossing ratio

Significant deviation from constant. Why?

Possible explanation – detector acceptance

- Extreme cases
 - Flat acceptance
→ N: area
 - Narrow acceptance
→ N: height

- ZDCLL1: Z-vertex resolution ~ 20 cm (wider acceptance)
 - BBCLL1: ~ 5 cm (narrower)
- (both have Z-vertex cut at ± 30 cm)

Correlation with Z-vertex width

We understand the reason now $\rightarrow$ correction

After correction

Relative luminosity accuracy $< 0.09\%$ (stat. limited)

Alternative correction

- Actual vertex width is not yet available in most runs (will come soon)
- Alternative width estimation
 - use ZDCout/ZDCin ratio
 - ZDCin: ZDCLL1 counts with Z-vertex cut of 30 cm
 - ZDCout: ZDCLL1 outside of ZDCin.
(both are counted crossing-by-crossing)
 - wider vertex distribution $\Leftrightarrow$ larger ratio

Actual accuracy estimation

-- Bunch fitting

- Crossing-by-crossing analysis, taking care of possible effect of A_{LL} in BBC and ZDC
- For i-th crossing, calculate

$$r(i) = N_{ZDC_{LL1}}(i)/N_{BBC_{LL1}}(i)$$
 corrected for vertex width
- Fit $r(i)$ by

$$r(i) = C[1 + A_{LL}^{BBC/ZDC} P_B(i) P_Y(i)]$$
 fit parameter: $C, A_{LL}^{BBC/ZDC}$
- You will get $C, A_{LL}^{BBC/ZDC}$, their errors, and χ^2
 - accuracy is obtained as $\delta A_{LL}^{BBC/ZDC} (\times P_B P_Y)$
 - χ^2 gives a good check for systematic errors.

χ^2 vs stat. error – before correction

Graph

correlating
 $\rightarrow \sigma_{\text{syst}} \sim 0.002$

χ^2 vs stat. error – after correction

Graph

correlation
is weak, but
still some
systematic
effect remains

Result & BBC/ZDC A_{LL}

- Systematic error seen $\rightarrow \chi^2$ correction required
- If χ^2/dof is larger than 1, then enlarge statistical error by $\sqrt{\chi^2/\text{dof}}$ – conservative correction
- Averaged result: $A_{LL}^{\text{BBC/ZDC}} = (1.8 \pm 1.8) \times 10^{-3}$
(without vertex width correction, δA_{LL} is ~ 2 larger)
- This corresponds to relative luminosity accuracy of $\delta R = 2.5 \times 10^{-4}$
 $\rightarrow$ Achieved the goal
- ZDC-BBC A_{LL} is 0 consistent
 $\rightarrow$ we can use both for reference of $A_{LL} = 0$

Some notes

- We reached $\delta A_{LL}^{BBC/ZDC} = 1.8 \times 10^{-3}$
 - this is dominated by statistics of ZDC, so it's rather conservative estimation
 - actual relative luminosity is given by BBC
- In π^0 analysis, vertex position dependence of acceptance is the same as BBCCLL1 because
 - BBC hits are required in the analysis
 - PHENIX central arm acceptance is almost flat for π^0 within the region we are using (± 30 cm)

Remaining tasks for rel. luminosity

- Apply offline Z-vertex cut
 - better resolution: $\sim 5 \text{ cm} \rightarrow \sim 2 \text{ cm}$
(effect is found to be small)
- Full accuracy estimation using real vertex distribution.
 - correction for other than vertex width?
- Study effect of multiple collisions
 - negligible in Run3, significant for higher luminosity.
- Z-vertex dependence of acceptance (efficiency) for BBCLL1, π^0 and other reaction channels.
 - reaction channel specific relative luminosity.
 - BBCLL1 itself is almost perfect for π^0 case, but may be not for other channels.

Vertex distribution (BBCLL1)

changes rel. lumi. by $< 0.04\%$ per fill (negligible)

PHENIX Local Polarimeter

- ✓ Forward neutron transverse asymmetry (A_N) measurements
- ✓ SMD (position) + ZDC (energy)

ϕ distribution

Vertical $\rightarrow \phi \sim \pm\pi/2$

Radial $\rightarrow \phi \sim 0$

Longitudinal $\rightarrow$ no asymmetry

Spin Longitudinal Component

$$S_L = \sqrt{1 - S_T^2}, \quad S_T = \sqrt{S_{T\text{-vertical}}^2 + S_{T\text{-radial}}^2}$$

S_T is measured with PHENIX Local Polarimeter

Left-Right asymmetry

Up-Down asymmetry

$$S_L(\text{blue}) = 99.3^{+0.5}_{-1.4} \quad {}^{+0.0}_{-0.9}$$

$$S_L(\text{yellow}) = 97.4^{+1.3}_{-3.2} \quad {}^{+0.1}_{-0.9}$$

π^0 counting

Data collected with high p_T photon trigger

Based on EMCal; Threshold ~ 1.4 GeV/c

Rejection factor ~ 110

Analyzed data sample: **42.7M events (~ 0.215 pb $^{-1}$)**

$\text{sqrt}(\langle P_b P_y \rangle) \sim 26\%$

Minimum Bias data

To obtain “unbiased” π^0 cross section at low p_T

For high p_T photon trigger efficiency study

π^0 reconstruction for A_{LL}

Results obtained for four pt bins from 1 to 5 GeV/c

π^0 peak width varies from 12 to 9.5 MeV/c^2 from lowest to highest pt bins

Background contribution under π^0 peak for $\pm 25 \text{ MeV}/c^2$ mass cut varies from 45% to 5% from lowest to highest pt bins

π^0 counting for A_{LL}

N_{π^0} :

$\pm 25 \text{ MeV}/c^2$ around π^0 peak (and also ± 15 and $\pm 35 \text{ MeV}/c^2$ for cross checks)

N_{bck1} :

Two $50 \text{ MeV}/c^2$ wide areas adjacent to π^0 peak

N_{bck2} :

$250 \text{ MeV}/c^2$ wide area between π^0 and η peaks

N_{π^0} and N_{bck} accumulated statistics

pt GeV/c	N_{π^0} 15 MeV/c ²	N_{π^0} 25 MeV/c ²	N_{π^0} 35 MeV/c ²	N_{bck1}	N_{bck2}
1-2	1278k	1777k	2129k	1470k	3478k
2-3	874k	1059k	1146k	335k	989k
3-4	176k	201k	208k	27k	83k
4-5	34k	38k	39k	3.9k	12k

N_{π^0} statistical error estimation

In multi- π^0 events:

$$\sigma_{N_{\pi^0}}^2 = \sigma_k^2 N_{ev} + \langle k \rangle^2 \sigma_{N_{ev}}^2 = \langle k^2 \rangle N_{ev}$$

Comparison to naïve approach: $\sigma_{N_{\pi^0}}^2 = N_{\pi^0}$

$$(k^{enhance})^2 = \frac{\langle k^2 \rangle N_{ev}}{N_{\pi^0}} = \frac{\langle k^2 \rangle}{\langle k \rangle}$$

Fluctuation enhancement (squared) due to multi-particle events

pt GeV/c	N_{π^0} 15 MeV/c ²	N_{π^0} 25 MeV/c ²	N_{π^0} 35 MeV/c ²	N_{bck1}	N_{bck2}
1-2	1.134	1.220	1.302	1.360	1.708
2-3	1.047	1.081	1.114	1.178	1.406
3-4	1.015	1.026	1.037	1.101	1.222
4-5	1.008	1.015	1.021	1.059	1.162

A_{LL} measurements

$$A_{LL} = \frac{\sigma_{++} - \sigma_{+-}}{\sigma_{++} + \sigma_{+-}} = \frac{1}{|P_B P_Y|} \frac{N_{++}/L_{++} - N_{+-}/L_{+-}}{N_{++}/L_{++} + N_{+-}/L_{+-}}, \quad \delta A_{LL} = \frac{1}{|P_B P_Y|} \frac{1}{\sqrt{N_{++} + N_{+-}}}$$

++ same helicity

+- opposite helicity

Procedure

1. Collect N and L for ++ and +- configurations (sum over all crossings) and calculate A_{LL} for each fill
2. Average A_{LL} over fills; use χ^2/NDF to control fit quality; use “bunch shuffling” to check syst. errors

A_{LL} Calculations

A_{LL} averaged over fills

1-2 GeV/c

$$A_{LL} = -2.8\% \pm 1.2\%$$

$$\chi^2/ndf = 64/48$$

2-3 GeV/c

$$A_{LL} = -2.2\% \pm 1.5\%$$

$$\chi^2/ndf = 34/48$$

3-4 GeV/c

$$A_{LL} = -0.2\% \pm 3.3\%$$

$$\chi^2/ndf = 49/48$$

4-5 GeV/c

$$A_{LL} = -2.3\% \pm 7.4\%$$

$$\chi^2/ndf = 39/48$$

A_L check for “yellow” beam

$$A_L = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-} = -\frac{1}{|P|} \frac{N_+/L_+ - N_-/L_-}{N_+/L_+ + N_-/L_-}$$

pt GeV/c	$A_L^{\pi^0+bck}$ 15 MeV/c ²	$A_L^{\pi^0+bck}$ 25 MeV/c ²	$A_L^{\pi^0+bck}$ 35 MeV/c ²	A_L^{bck1}	A_L^{bck2}
1-2	0.001±0.004	0.000±0.003	0.000±0.003	0.002±0.004	0.000±0.003
2-3	0.001±0.004	0.000±0.004	0.000±0.004	0.002±0.007	0.002±0.005
3-4	0.007±0.009	0.011±0.009	0.008±0.009	-0.033±0.025	-0.010±0.015
4-5	-0.001±0.021	0.004±0.020	0.008±0.020	0.020±0.064	0.050±0.039

All are zeros within 1.5σ

A_L check for “blue” beam

$$A_L = \frac{\sigma_+ - \sigma_-}{\sigma_+ + \sigma_-} = -\frac{1}{|P|} \frac{N_+/L_+ - N_-/L_-}{N_+/L_+ + N_-/L_-}$$

pt GeV/c	$A_L^{\pi^0+bck}$ 15 MeV/c ²	$A_L^{\pi^0+bck}$ 25 MeV/c ²	$A_L^{\pi^0+bck}$ 35 MeV/c ²	A_L^{bck1}	A_L^{bck2}
1-2	-0.001±0.004	0.001±0.003	0.000±0.003	-0.002±0.004	-0.000±0.003
2-3	0.001±0.004	0.000±0.004	0.002±0.004	0.009±0.007	0.000±0.005
3-4	0.004±0.009	0.006±0.009	0.006±0.008	-0.004±0.024	-0.036±0.015
4-5	-0.024±0.021	-0.016±0.020	-0.016±0.019	-0.011±0.062	0.013±0.038

All are zeros within 1.5σ , except

π^0 A_{LL} result

p_T GeV/c	$A_{LL}^{\pi^0+bck}$ (r_{bck})	A_{LL}^{bck}	$A_{LL}^{\pi^0}$ (Background subtracted)
1-2	-0.028 ± 0.012 (45%)	-0.006 ± 0.014	-0.046 ± 0.025
2-3	-0.022 ± 0.015 (17%)	-0.035 ± 0.027	-0.019 ± 0.019
3-4	-0.002 ± 0.033 (7%)	0.094 ± 0.092	-0.009 ± 0.036
4-5	-0.023 ± 0.074 (5%)	0.38 ± 0.24	-0.045 ± 0.079

Polarization scaling error $\delta P \sim 30\%$: is not included

- ✓ Enters the A_{LL} quadratically
- ✓ Analyzing power $A_N(100 \text{ GeV}) \sim A_N(22 \text{ GeV})$ is assumed
- ✓ $\delta P \sim 30\%$: combined stat. and syst. error for $A_N(22 \text{ GeV})$ (AGS E950)

p_T smearing correction is not included

Summary

- Relative luminosity – systematics is mostly due to crossing by crossing variation of Z-vertex width and acceptance difference
→ correction applied
- Relative luminosity accuracy
 $\delta A_{LL} = 1.8 \times 10^{-3}$ achieved
- Relative luminosity is reaction channel dependent
 - For π^0 analysis, BBCLL1 is good enough.
- Preliminary result for π^0 A_{LL} obtained
- Various null tests – all OK
statistical and systematic errors are well studied.

STAR Spin Analysis Update

Stephen Trentalange

Beam-Beam Counters

- **scintillator annulus** installed around the beam pipe, on the east and west poletips of STAR magnet at $\pm 3.74\text{m}$ from IR for detection of charged particles in forward region $2 < |\eta| < 5$

STAR Beam-Beam Counter Schematic
Front View

Small hexagonal annulus:

18 pixels (16 PMT) covering $3.3 < |\eta| < 5.0$ and $0 < \phi < 2\pi$
- **azimuthal segmentation !**

A coincidence condition between BBC East and West - suppresses beam-gas background and used for:

- triggering in pp (minimum bias, jet triggers),
- (relative) luminosity measurements
- local polarimetry

BBC cross section:

$\sigma(\text{BBC}) \sim 50\% \text{ of } \sigma_{\text{tot}}(\text{pp}) = 26 \text{ mb}$ Large!

- Van der Meer scans at RHIC (Run2 and Run3 data)
- are consistent with Geant simulations for STAR

Transverse Single Spin Asymmetries

Asymmetries measured for $p+p \rightarrow A + X$, where A – hit(s) in the BBC (small tiles, $3.3 < |\eta| < 5.0$)

$$\epsilon_{LR} = \frac{\sqrt{N_L^\uparrow N_R^\downarrow} - \sqrt{N_L^\downarrow N_R^\uparrow}}{\sqrt{N_L^\uparrow N_R^\downarrow} + \sqrt{N_L^\downarrow N_R^\uparrow}} \approx P_{\text{beam}}^{\text{vert}} \times \underline{A_N(\text{BBC})}$$

RHIC CNI polarimeter $\left(\epsilon_{\text{CNI}} = P_{\text{beam}} \times A_N(\text{CNI}) \right)$

$$\epsilon_{TB} = \frac{\sqrt{N_T^\uparrow N_B^\downarrow} - \sqrt{N_T^\downarrow N_B^\uparrow}}{\sqrt{N_T^\uparrow N_B^\downarrow} + \sqrt{N_T^\downarrow N_B^\uparrow}} \approx P_{\text{beam}}^{\text{rad}} \times A_N(\text{BBC})$$

**Zero at RHIC
(Vertical pol)**

$N_{L(R,T,B)}$ – nr of counts from BBC East or West in Left, Right, Top, Bottom part counted every bunch crossing by scaler system]

Number of counts from the BBC (L,R,T,B) sorted by beam polarization states
Example: Polarized Yellow beam (sum over Blue beam polarization states)
 heads towards **BBC East**

Local Polarimetry

Rotators	OFF	ON
CNI polarimeter	non-zero	non-zero
BBC Left-Right (vertical)	NON_ZERO → ZERO	
BBC Top-Bottom (radial)	zero	zero

Spin rotators worked as designed and stable,
BBC worked very well as a local polarimeter

- vertical and radial polarization components smaller than 3%

Longitudinal polarization for the first time!

Joanna Kiryluk

Transverse Single Spin Asymmetries – Preliminary Results

Top-Bottom BBC asymmetry (Radial P_{beam} component)
Left-Right BBC asymmetry (Vertical P_{beam} component)

Ratio: ϵ (BBC East)/ ϵ (CNI)
consistent with
 ϵ (BBC West)/ ϵ (CNI)

We get:
 $A_N(\text{BBC}) = 0.0066(8)$
 Stat. error only!

using: $A_N(\text{CNI}) = 0.0118(10)$
 from CNI online analysis

Joanna Kiryluk

Relative Luminosity - Motivation

Double spin longitudinal asymmetry (e.g. $p+p \rightarrow \text{jet} + X$) is defined as:

$$A_{LL} = \frac{1}{P_1 P_2} \times \frac{N_{++} - R \times N_{+-}}{N_{++} + R \times N_{+-}}$$

$P_{1,2}$ - beam polarization, from CNI polarimeter at RHIC

N_i where $i=++$, $+-$ are spin dependent yields

$R = \frac{L_{++}}{L_{+-}}$ - relative luminosity of bunch crossings with different spin directions

Assumption: $A_{LL} = 0.01$

Relative luminosity - one of the dominant systematic errors in A_{LL} measurement. It must be known at (least) 10^{-3} level

Requirements for a luminosity process/detector:

- high rates
- small background
- no (or small) spin dependence

$$\delta R \ll P_1 P_2 \times A_{LL}$$

Relative Luminosity - Measurement

- RHIC filled with 55 - transversely polarized - proton bunches,
- spin orientation related to the bunch crossing number 1-120, **polarization pattern at STAR:**

- integrated luminosity (number of BBC E.W counts) recorded every 5 minutes - scaler board system
- relative luminosities calculated for each STAR run, which typically lasts from a few minutes to $\sim$ hour
- statistical uncertainty $\delta R_{\text{stat}} \sim 10^{-4} - 10^{-3}$

Joanna Kiryluk

Relative Luminosity – Systematics

Beam-Gas Background Contribution

$$R(\text{BBC}) - R(\text{true}) = Bg \times [1 - R(\text{true})]$$

if $R(\text{true}) = 1$ then $R(\text{BBC}) = R(\text{true})$

$$R(\text{BBC}) = \frac{N_{++}^{\text{BBC}}}{N_{+-}^{\text{BBC}}} \cong \frac{L_{++} + N_{bg}}{L_{+-} + N_{bg}} = \Lambda \quad \Lambda \quad \text{simple math} \quad \Lambda \quad = \frac{L_{++}}{L_{+-}} + Bg \times \left(1 - \frac{L_{++}}{L_{+-}}\right)$$

Contribution to systematic uncertainty on R $\delta R(\text{sys}) \sim 0.004 \times 0.2 \sim 8 \times 10^{-4}$
- must correct for background

Joanna Kiryluk

Summary

1. **Single transverse spin asymmetry measured with the Beam Beam Counters** in 2003 were found to be $A_N(\text{BBC})=0.0066(8)$ using $A_N(\text{CNI})=0.0118(10)$ from CNI online analysis. Systematic studies in progress.
2. The BBC was used to tune spin rotators needed to get longitudinally polarized beams at STAR IP and to monitor on-line transverse and radial beam polarization components. They are found to be small: $P_{\text{rad(vert)}} < 3\%$.
3. The BBC in STAR is a good luminosity monitoring detector. **The relative luminosity is known at the level of $\sim 3 \times 10^{-3}$**
 - statistical uncertainty: $10^{-4} - 10^{-3}$
 - (conservative) systematic uncertainty (BBC and ZDC comparison) $\sim 3 \times 10^{-3}$

This needs further studies (background contribution)

Joanna Kiryluk

2003 Detector Configuration

Leading Jet Asymmetries are Sensitive to Gluon

- Simulation based on Pythia
- Use Fast TPC and EMC Simulators
- EMC+TPC Jets Reconstructed using Cone Algorithm
- Includes Jet-Patch Trigger ($E_T > 5$) covering the Full Half Barrel Calorimeter
- A_{LL} includes all jet processes (not a direct measure of ΔG)

Simulations by Les Bland

pp Longitudinal Data Set

- May 16- May 30 2003
- Ave Polarization / Beam ~ 0.25
- RHIC Delivered Integrated Luminosity $\sim 420 \text{ nb}^{-1}$
- Full STAR Detectors Sampled $\sim 330 \text{ nb}^{-1}$
- Recorded Luminosity $\sim 180 \text{ nb}^{-1}$
- Barrel EMC Covers
0-1 in Eta and the Full Azimuth
- High Tower Trigger
 - Requires Single Tower $\text{ADC} > 287$ ($E_T \sim > 2.4 \text{ GeV}$)
- Jet Patch Trigger
 - Requires Patch of 1×1 (η, ϕ) $E_T > 5 \text{ GeV}$

Jet Finder (Mike Miller/Thomas Henry)

For this Analysis, Jets are defined as a grouping of one or more (charged) Tracks measured in the TPC and satisfying the following requirements...

- Using an Iterative, Midpoint, Cone Algorithm
- Cone Angle = 0.7 in eta-phi
- Seed Energy = 0.5 GeV
- $p_T \text{ Track} > 0.1 \text{ GeV}$
- Track Range = ± 1.6
- $p_T \text{ Jet} > 5 \text{ GeV}$
- $Z = \pm 75 \text{ cm}$

TPC Jet Characteristics – EMC Trigger

- Strong Trigger Bias at low E_T

- Goal: Measured E_T --> Parton E_T

Longitudinal Spin Asymmetries

$$\varepsilon_{LL} = P_A P_B \cdot A_{LL} = \frac{N^{++} - R_2 * N^{+-}}{N^{++} + R_2 * N^{+-}}$$

Does Not Violate Parity

$$\varepsilon_L = P_A \cdot A_L = \frac{N^+ - R_1 * N^-}{N^+ + R_1 * N^-}$$

Violates Parity

$$R_1 = L^+ / L^- \quad R_2 = L^{++} / L^{+-}$$

REQUIRES luminosity ratios to be measured much more accurately than desired asymmetry

TPC-Jet, Single-Spin Asymmetries

Blue Beam

$$\langle \epsilon \rangle = 0.00016$$

$$\delta(\epsilon) = 0.0038$$

$$\chi^2 / dof = 106/95$$

Yellow Beam

$$\langle \epsilon \rangle = -0.005$$

$$\delta(\epsilon) = +0.0038$$

$$\chi^2 / dof = 99/96$$

Parity-Violating Asymmetries are Consistent with Zero

Dylan Thein

2003 Outlook

Data	Statistical Precision on A_{LL}
TPC Jets	+/- 0.057
EMC+TPC Jets	+/- 0.030

RHIC Spin Collaboration Meeting XX

November 18, 2003

RIKEN BNL Research Center

LIST OF REGISTERED PARTICIPANTS

RIKEN PARTICIPANTS

NAME	AFFILIATION AND ADDRESS	E-MAIL ADDRESS
Gerry Bunce	RBRC / BNL Physics, Bldg. 510A Upton, N.Y. 11973-5000	bunce@bnl.gov
Hideto En'yo	RIKEN / RBRC Radiation Laboratory 2-1 Hirosawa, Wako, Saitama Japan 351-0198	enyo@riken.go.jp
Doug Fields	RBRC / UNM Bldg. 510A Upton, N.Y. 11973-5000	fields@umn.edu
Yuji Goto	RBRC / RIKEN Physics, Bldg. 510A Upton, N.Y. 11973-5000	goto@bnl.gov
Osamu Jinnouchi	RBRC / RIKEN Physics, Bldg. 510A Upton, N.Y. 11973-5000	josamu@bnl.gov
Kensuke Okada	RBRC Physics, Bldg. 510A Upton, N.Y. 11973-5000	okada@bnl.gov
Tsugu Tabaru	RBRC / RIKEN Physics, Bldg. 510A Upton, N.Y. 11973-5000	tsugu@riken.jp
Junpei Takano	RIKEN / TITech Radiation Laboratory 2-1 Hirosawa, Wako, Saitama Japan 351-0198	jtakano@rarfaxp.riken.go.jp
Kiyoshi Tanida	RIKEN / RBRC Radiation Laboratory 2-1 Hirosawa, Wako, Saitama Japan 351-0198	tanida@rarfaxp.riken.go.jp
Werner Vogelsang	RBRC / BNL Physics, Bldg. 510A Upton, N.Y. 11973-5000	wvogelsang@bnl.gov
Yasushi Watanabe	RIKEN Radiation Laboratory 2-1 Hirosawa, Wako, Saitama Japan, 351-0198	watanabe@bnl.gov

RHIC Spin Collaboration Meeting XX
November 18, 2003
RIKEN BNL Research Center

LIST OF REGISTERED PARTICIPANTS

NON-RIKEN PARTICIPANTS

NAME	AFFILIATION AND ADDRESS	E-MAIL ADDRESS
Leif Ahrens	BNL Bldg. 911B Upton, N.Y. 11973-5000	ahrens@bnl.gov
Mei Bai	BNL C-A, Bldg. 911B Upton, N.Y. 11973-5000	mbai@bnl.gov
Ken Barish	UC Riverside Department of Physics Riverside, CA 92521	Kenneth.Barish@ucr.edu
Alexander Bazilevsky	BNL Bldg. 510A Upton, N.Y. 11973-5000	shura@bnl.gov
Les Bland	BNL Physics, Bldg. 510A Upton, N.Y. 11973-5000	bland@bnl.gov
Alessandro Bravar	BNL Physics, Bldg. 510 Upton, N.Y. 11973-5000	bravar@bnl.gov
Robert Cadman	BNL / Argonne National Laboratory Bldg. 510 – STAR Upton, N.Y. 11973-5000	rvc@anl.gov
Pete Cameron	BNL Bldg. 817 Upton, N.Y. 11973-5000	cameron@bnl.gov
Haixin Huang	BNL Bldg. 911B Upton, N.Y. 11973-5000	huanghai@bnl.gov
Wolfram Fischer	BNL C-A, Bldg. 911B Upton, N.Y. 11973-5000	wfischer@bnl.gov
David Kawall	Yale University P.O. Box 208120 New Haven, CT. 06520-8120	david.kawall@yale.edu
Ed Kinney	University of Colorado 390 UCB Boulder, CO 80309	Edward.Kinney@colorado.edu
Joanna Kiriyluk	BNL / UCLA Bldg. 510 – STAR Upton, N.Y. 11973-5000	joanna@physics.ucla.edu
Waldo Mackay	BNL C-A, Bldg. 911B Upton, N.Y. 11973-5000	waldo@bnl.gov
Yousef Makdisi	BNL Bldg. 911B Upton, N.Y. 11973-5000	makdisi@bnl.gov
Alexander Nass	BNL C-A, Bldg. 911B Upton, N.Y. 11973-5000	nass@bnl.gov

RHIC Spin Collaboration Meeting XX

November 18, 2003

RIKEN BNL Research Center

LIST OF REGISTERED PARTICIPANTS

NON-RIKEN PARTICIPANTS

NAME	AFFILIATION AND ADDRESS	E-MAIL ADDRESS
Vadim Ptitsyn	BNL Bldg. 911B Upton, N.Y. 11973-5000	vadimp@bnl.gov
Greg Rakness	BNL / Penn State Bldg. 510 – STAR Upton, N.Y. 11973-5000	rakness@rcf2.rhic.bnl.gov
Thomas Roser	BNL Bldg. 911B Upton, N.Y. 11973-5000	roser@bnl.gov
Jim Sowinski	BNL / IUCF Bldg. 510 – STAR Upton, N.Y. 11973-5000	sowinski@iucf.indiana.edu
Rogelio Tomas	BNL C-A, Bldg. 911B Upton, N.Y. 11973-5000	rtomas@bnl.gov
Stephen Trentalange	BNL / UCLA Bldg. 510 – STAR Upton, N.Y. 11973-5000	trent@physics.ucla.edu
Johannes van Zeijts	BNL Bldg. 911C Upton, N.Y. 11973-5000	johannes@bnl.gov
Steve Vigdor	IUCF / LBNL / BNL Bldg. 510 – STAR Upton, N.Y. 11973-5000	vigdor@iucf.indiana.edu
Anatoli Zelenski	BNL Bldg. 930 Upton, N.Y. 11973-5000	zelenski@bnl.gov

RHIC Spin Collaboration Meeting XX

November 18, 2003

RIKEN BNL Research Center

LIST OF REGISTERED PARTICIPANTS

Not in attendance but will be sent proceedings:

NAME	AFFILIATION AND ADDRESS	E-MAIL ADDRESS
Christine Aidala	BNL Physics, Bldg. 510C Upton, N.Y. 11973-5000	caidala@bnl.gov
Igor Alekseev	BNL Bldg. 510 ~ pp2pp Upton, N.Y. 11973-5000	igor.alekseev@itep.ru
Kazuya Aoki	Kyoto University Physics Department Kitashirakawa-Oiwakecho, Sakyo-ku Kyoto, 606-8502 Japan	aoki@nh.scphys.kyoto-u.ac.jp
Bill Christie	BNL Bldg. 510 – STAR Upton, N.Y. 11973-5000	christie@bnl.gov
Abhay Deshpande	RBRC Bldg. 510A Upton, N.Y. 11973-5000	abhay@bnl.gov
Satish Dhawan	Yale University Physics Department P.O. Box 208121 New Haven, CT. 06520-8121	satish.dhawan@yale.edu
Geary Eppley	Rice University MS 315 6100 Main St. Houston, TX. 77005	eppley@physics.rice.edu
Yoshinori Fukao	RBRC / Kyoto University Bldg. 510A Upton, N.Y. 11973-5000	fukao@nh.scphys.kyoto.u.ac.jp
Matthias Grosse Perdekamp	UIUC / RBRC Bldg. 510A Upton, N.Y. 11973-5000	mgp@uiuc.edu
Wlodek Guryn	BNL Bldg. 510C Upton, N.Y. 11973-5000	guryn@bnl.gov
Takuma Horaguchi	RBRC / TITech Bldg. 510A Upton, N.Y. 11973-5000	horaguchi@nucl.phys.titech.ac.jp
George Igo	BNL / UCLA Bldg. 902B – STAR Upton, N.Y. 11973-5000	igo@physics.ucla.edu
Nobuyuki Kamihara	RBRC / TITech Bldg. 510A Upton, N.Y. 11973-5000	kamihara@bnl.gov
Kazu Kurita	Rikkyo University Department of Physics 3-34-1 Nishi-Ikebukuro Toshima, Tokyo 171-8501 Japan	kurita@ne.rikkyo.ac.jp

RHIC Spin Collaboration Meeting XX
November 18, 2003
RIKEN BNL Research Center

LIST OF REGISTERED PARTICIPANTS

Not in attendance but will be sent proceedings:

NAME	AFFILIATION AND ADDRESS	E-MAIL ADDRESS
Naohito Saito	Kyoto University / RBRC / RIKEN Physics Department Kitashirakawa-Oiwakecho, Sakyo-ku Kyoto, 606-8502, Japan	saito@bnl.gov saito@nh.scphys.kyoto-u.ac.jp
Hiroki Sato	Kyoto University / RBRC Physics Department Kitashirakawa-Oiwakecho, Sakyo-ku Kyoto, 606-8502, Japan	satohiro@bnl.gov
Dima Svirida	Institute for Theoretical and Experimental Physics 25 B. Cheremushkinskaya 117259, Moscow, Russia	dmitry.svirida@itep.ru
Hal Spinka	Argonne National Laboratory Bldg. 362 – HEP Argonne, IL. 60439	hms@hep.anl.gov
Atsushi Taketani	RIKEN Radiation Laboratory 2-1 Hirosawa, Wako, Saitama 351-0198 Japan	taketani@bnl.gov
Manabu Togawa	Kyoto University Physics Department Kitashirakawa-Oiwakecho, Sakyo-ku Kyoto, 606-8502, Japan	togawa@nh.scphys.kyoto-u.ac.jp
Hisayuki Torii	RIKEN Radiation Laboratory 2-1 Hirosawa, Wako, Saitama 351-0198 Japan	htorii@bnl.gov
Flemming Videbæk	BNL Bldg. 510D Upton, N.Y. 11973-5000	videbaek@bnl.gov

RIKEN BNL Research Center
RHIC Spin Collaboration Meeting XX
November 18, 2003
Bldg. 1005, Third Floor Conference Room
Brookhaven National Laboratory

*******AGENDA*******

Accelerator topics

9:00 - 9:30 RHIC working point.....Rojelio Tomas and Wolfram Fischer
9:30-10:00 AGS warm-bore helical dipole.....Jun Takano and Waldo MacKay
10:00-10:20 RHIC spin-tune measurement and polarization reversal.....Mei Bai
10:20-10:40 Status of PLL..... Peter Cameron
10:40-11:10 Coffee Break
11:10-11:30 AGS polarized proton commissioning..... Leif Ahrens
11:30-12:00 Run - 4 PP ramp.....Johannes van Zeijts

12:00 - 1:15 Lunch

Polarimetry

1:15 - 1:35 Status of polarized gas jet target.....Alexander Nass
1:35 - 1:55 Polarimeter Status.....Haixin Huang
1:55 - 2:15 CNI polarimetry status.....Osamu Jinnouchi and Sandro Bravar

Future...

2:15 - 2:45 Discussion: Physics opportunities in runs-4, 5
2:45 - 3:15 Discussion: Summary of spin-strategy meeting and plans..S. Vigdor (TBC)
3:15 - 3:45 Coffee Break

Status of Run-3 analyses

3:45 - 4:15 New theoretical studies of pp interactions.....W. Vogelsang
4:15 - 4:45 PHENIX.....Kiyoshi Tanida
4:45 - 5:15 STAR.....Stephen Trentalange

Additional RIKEN BNL Research Center Proceedings:

- Volume 58 – RHIC Spin Collaboration Meeting XX – BNL-
- Volume 57 – High pt Physics at RHIC, December 2-6, 2003 – BNL-
- Volume 56 – RBRC Scientific Review Committee Meeting – BNL-
- Volume 55 – Collective Flow and QGP Properties – BNL -
- Volume 54 – RHIC Spin Collaboration Meetings XVII, XVIII, XIX – BNL-71751-2003
- Volume 53 – Theory Studies for Polarized pp Scattering – BNL-71747-2003
- Volume 52 – RIKEN School on QCD “Topics on the Proton” – BNL-71694-2003
- Volume 51 – RHIC Spin Collaboration Meetings XV, XVI – BNL-71539-2003
- Volume 50 – High Performance Computing with QCDOC and BlueGene – BNL-71147-2003
- Volume 49 – RBRC Scientific Review Committee Meeting – BNL-52679
- Volume 48 – RHIC Spin Collaboration Meeting XIV – BNL-71300-2003
- Volume 47 – RHIC Spin Collaboration Meetings XII, XIII – BNL-71118-2003
- Volume 46 – Large-Scale Computations in Nuclear Physics using the QCDOC – BNL-52678
- Volume 45 – Summer Program: Current and Future Directions at RHIC – BNL-71035
- Volume 44 – RHIC Spin Collaboration Meetings VIII, IX, X, XI – BNL-71117-2003
- Volume 43 – RIKEN Winter School – Quark-Gluon Structure of the Nucleon and QCD – BNL-52672
- Volume 42 – Baryon Dynamics at RHIC – BNL-52669
- Volume 41 – Hadron Structure from Lattice QCD – BNL-52674
- Volume 40 – Theory Studies for RHIC-Spin – BNL-52662
- Volume 39 – RHIC Spin Collaboration Meeting VII – BNL-52659
- Volume 38 – RBRC Scientific Review Committee Meeting – BNL-52649
- Volume 37 – RHIC Spin Collaboration Meeting VI (Part 2) – BNL-52660
- Volume 36 – RHIC Spin Collaboration Meeting VI – BNL-52642
- Volume 35 – RIKEN Winter School – Quarks, Hadrons and Nuclei – QCD Hard Processes and the Nucleon Spin – BNL-52643
- Volume 34 – High Energy QCD: Beyond the Pomeron – BNL-52641
- Volume 33 – Spin Physics at RHIC in Year-1 and Beyond – BNL-52635
- Volume 32 – RHIC Spin Physics V – BNL-52628
- Volume 31 – RHIC Spin Physics III & IV Polarized Partons at High Q^2 Region – BNL-52617
- Volume 30 – RBRC Scientific Review Committee Meeting – BNL-52603
- Volume 29 – Future Transversity Measurements – BNL-52612
- Volume 28 – Equilibrium & Non-Equilibrium Aspects of Hot, Dense QCD – BNL-52613
- Volume 27 – Predictions and Uncertainties for RHIC Spin Physics & Event Generator for RHIC Spin Physics III – Towards Precision Spin Physics at RHIC – BNL-52596
- Volume 26 – Circum-Pan-Pacific RIKEN Symposium on High Energy Spin Physics – BNL-52588
- Volume 25 – RHIC Spin – BNL-52581

Additional RIKEN BNL Research Center Proceedings:

- Volume 24 – Physics Society of Japan Biannual Meeting Symposium on QCD Physics at RIKEN BNL Research Center – BNL-52578
- Volume 23 – Coulomb and Pion-Asymmetry Polarimetry and Hadronic Spin Dependence at RHIC Energies – BNL-52589
- Volume 22 – OSCAR II: Predictions for RHIC – BNL-52591
- Volume 21 – RBRC Scientific Review Committee Meeting – BNL-52568
- Volume 20 – Gauge-Invariant Variables in Gauge Theories – BNL-52590
- Volume 19 – Numerical Algorithms at Non-Zero Chemical Potential – BNL-52573
- Volume 18 – Event Generator for RHIC Spin Physics – BNL-52571
- Volume 17 – Hard Parton Physics in High-Energy Nuclear Collisions – BNL-52574
- Volume 16 – RIKEN Winter School - Structure of Hadrons - Introduction to QCD Hard Processes – BNL-52569
- Volume 15 – QCD Phase Transitions – BNL-52561
- Volume 14 – Quantum Fields In and Out of Equilibrium – BNL-52560
- Volume 13 – Physics of the 1 Teraflop RIKEN-BNL-Columbia QCD Project First Anniversary Celebration – BNL-66299
- Volume 12 – Quarkonium Production in Relativistic Nuclear Collisions – BNL-52559
- Volume 11 – Event Generator for RHIC Spin Physics – BNL-66116
- Volume 10 – Physics of Polarimetry at RHIC – BNL-65926
- Volume 9 – High Density Matter in AGS, SPS and RHIC Collisions – BNL-65762
- Volume 8 – Fermion Frontiers in Vector Lattice Gauge Theories – BNL-65634
- Volume 7 – RHIC Spin Physics – BNL-65615
- Volume 6 – Quarks and Gluons in the Nucleon – BNL-65234
- Volume 5 – Color Superconductivity, Instantons and Parity (Non?)-Conservation at High Baryon Density – BNL-65105
- Volume 4 – Inauguration Ceremony, September 22 and Non -Equilibrium Many Body Dynamics – BNL-64912
- Volume 3 – Hadron Spin-Flip at RHIC Energies – BNL-64724
- Volume 2 – Perturbative QCD as a Probe of Hadron Structure – BNL-64723
- Volume 1 – Open Standards for Cascade Models for RHIC – BNL-64722

For information please contact:

Ms. Pamela Esposito
RIKEN BNL Research Center
Building 510A
Brookhaven National Laboratory
Upton, NY 11973-5000 USA

Phone: (631) 344-3097
Fax: (631) 344-4067
E-Mail: pesposit@bnl.gov

Ms. Tammy Heinz
RIKEN BNL Research Center
Building 510A
Brookhaven National Laboratory
Upton, NY 11973-5000 USA

(631) 344-5864
(631) 344-2562
theinz@bnl.gov

Homepage: <http://www.bnl.gov/riken>

RIKEN BNL RESEARCH CENTER

RHIC Spin Collaboration Meeting XX

November 18, 2003

Li Keran

*Nuclei as heavy as bulls
Through collision
Generate new states of matter.
T.D. Lee*

Copyright©CCASTA

Speakers:

L. Ahrens
P. Cameron
O. Jinnouchi
J. Takano
S. Trentalange
W. Vogelsang

M. Bai
W. Fischer
W. MacKay
K. Tanida
J. van Zeijts

S. Bravar
H. Huang
A. Nass
R. Tomas
S. Vigdor

Organizer: L. Bland