BNL-73470-2005-CP

*Comparison of Off-line IR Bump and Action-Angle
Kick Minimization*

Y. Luo, F. Pilat, V. Ptitsyn, D. Trbojevic, J. Wei

*Presented at the Particle Accelerator Conference (PAC'05)
Knoxville, Tennessee
May 16-20, 2005*

Collider-Accelerator Department

Brookhaven National Laboratory

P.O. Box 5000

Upton, NY 11973-5000

www.bnl.gov

Managed by

Brookhaven Science Associates, LLC

for the United States Department of Energy under

Contract No. DE-AC02-98CH10886

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

FOR UNCLASSIFIED, UNLIMITED STI PRODUCTS

Available electronically at:

OSTI:

<http://www.osti.gov/bridge>

Available for a processing fee to U.S. Department of Energy and its contractors, in paper from:

US. Department of Energy
Office of Scientific and Technical Information
P.O. Box **62**
Oak Ridge, TN **37831**
Phone: **(865) 576-8401**
Facsimile: **(865) 576-5728**
E-mail: reports@adonis.osti.gov

National Technical Information Service (NTIS):

Available for sale to the public from:

US. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA **22131**
Phone: **(800) 553-6847**
Facsimile: **(703) 605-6900**
Online ordering: <http://www.ntis.gov/ordering.htm>

Printed on recycled paper

Comparison of Off-line IR Bump and Action-Angle Kick Minimization*

Y. Luo, F. Pilat, V. Ptitsyn, D. Trbojevic, J. Wei
Brookhaven National Laboratory, Upton, NY 11973, USA

Abstract

The interaction region bump (IR bump) nonlinear correction method has been used for the sextupole and octupole field error on-line corrections in the Relativistic Heavy Ion Collider (RHIC). Some differences were found for the sextupole and octupole corrector strengths between the on-line IR bump correction and the predictions from the action-angle kick minimization. In this article, we compare the corrector strengths from these two methods based on the RHIC Blue ring lattice with the IR nonlinear modeling. The comparison confirms the differences between resulting corrector strengths. And the reason for the differences is found and discussed.

ACTION-ANGLE KICK MINIMIZATION

To *minimize* the action change for each order of IR nonlinear field error, it is equivalent to minimize the following quantities simultaneously [1],

$$\oint_L ds C_z c_n + (-1)^{n+1} \oint_R ds C_z c_n, \quad (1)$$

where L and R mean the left and right sides of the interaction region, z stands for x or y plane, c_n stands for the normal or skew field errors b_n or a_n , n is the field error order. C_z is the weight factor, which is different for different order of errors and for different x or y plane. The integrations or summations are taken across the interaction region. The correction weight factors are given according to the multiples as

$$C_x = \begin{cases} \beta_x^{(n+1)/2} & \text{for } b_n \\ \beta_x^{n/2} \beta_y^{1/2} & \text{for } a_n \end{cases}, \quad (2)$$

$$C_y = \begin{cases} \beta_y^{(n+1)/2} & \text{for odd } b_n \text{ and even } a_n \\ \beta_x^{1/2} \beta_y^{n/2} & \text{for even } b_n \text{ and odd } a_n \end{cases} \quad (3)$$

For each order, there are two quantities to be minimized, one in the horizontal plane, one in the vertical plane. Therefore, at least two nonlinear correctors are needed for each order correction. A natural choice is to place one corrector in either side of the interaction region. In order to facilitate the following discussion, Table 1 lists the weight factors for different field error b , and a .

The action-angle kick minimization assumes that the phase advances in the horizontal and vertical planes across the interaction point are close to π . And it ignores the phase advance in the triplet because of the small β^* . It assumes the beam is round beam, therefore, only the leading resonances in horizontal and vertical planes are corrected.

*Work supported by U.S. DOE under contract No DE-AC02-98CH10886

Table 1: Weight factors for action-angle kick minimization

Order	C_x	C_y	$k_n l / k_{ns} l$	$(-1)^{n+1}$
b_2	$\beta_x^{3/2}$	$\beta_x^{1/2} \beta_y$	$k_2 l$	-1
b_3	β_x^2	β_y^2	$k_3 l$	1
b_4	$\beta_x^{5/2}$	$\beta_x^{1/2} \beta_y^2$	$k_4 l$	-1
b_5	β_x^3	β_y^3	$k_5 l$	1
a_2	$\beta_x \beta_y^{1/2}$	$\beta_y^{3/2}$	$k_{2s} l$	-1
a_3	$\beta_x^{3/2} \beta_y^{1/2}$	$\beta_x^{1/2} \beta_y^{3/2}$	$k_{3s} l$	1
a_4	$\beta_x^2 \beta_y^{1/2}$	$\beta_y^{5/2}$	$k_{4s} l$	-1
a_5	$\beta_x^{5/2} \beta_y^{1/2}$	$\beta_x^{1/2} \beta_y^{3/2}$	$k_{5s} l$	1

IR BUMP CORRECTION

The IR bump correction method [2, 3, 4] is an elegant way for the operational IR nonlinear corrections in a real machine. It creates a local horizontal or vertical orbit bump across the interaction region. The small tune shifts due to the bump are measured with a high resolution phase lock loop (PLL) tune measurement system. Since the relations between the tune shifts and the bump amplitudes are different for different orders of nonlinear errors, the IR bump correction is performed order by order by minimizing the polynomial fitting coefficients of the tune shifts.

For example, to correct the sextupole errors in the IR, we minimize the linear term of the tune shifts from the horizontal IR bump with respect to the bump amplitude. According to Hamiltonian perturbation theory, the tune shifts from the sextupoles with the horizontal beam orbit offsets are given by

$$\begin{cases} \Delta Q_x = \frac{1}{2\pi} \sum (6b_2 x_{co} \beta_x) ds \\ \Delta Q_y = \frac{1}{2\pi} \sum (-6b_2 x_{co} \beta_y) ds \end{cases} \quad (4)$$

Considering the phase advance π between the interaction region, we assume that the local horizontal orbit bump is proportional to the $\beta_x^{1/2}$,

$$\begin{cases} x_{co} \propto \beta_x^{1/2}, \text{ in one side of IR} \\ x_{co} \propto -\beta_x^{1/2}, \text{ in another side of IR} \end{cases} \quad (5)$$

For simplicity, in the following analysis and simulation, we assume that the orbit at one point in the IR bump is proportional to the bump steps. The orbit bump at different bump steps are given by

$$x_{co} = x_{co,1} \times k, \quad (6)$$

$x_{co,1}$ is the orbit bump amplitude increment, $k = -N, -(N-1), \dots, (N-1), N$. N is the maximum bump

Figure 1: The orbits with the horizontal bump in the RHIC Blue ring IR8.

step No.. Then, the total tune shifts due to the x plane IR bump are

$$\begin{cases} \Delta Q_x \propto (\sum_L b_2 ds \beta_x^{3/2} - \sum_R b_2 ds \beta_x^{3/2}) \cdot k \\ \Delta Q_y \propto (\sum_L b_2 ds \beta_x^{1/2} \beta_y - \sum_R b_2 ds \beta_x^{1/2} \beta_y) \cdot k \end{cases} \quad (7)$$

If we use the sextupole correctors in the IR bump to minimize the linear polynomial term of the tune shifts with the bump steps k , we obtain

$$\begin{cases} \sum_L b_2 ds \beta_x^{3/2} - \sum_R b_2 ds \beta_x^{3/2} = 0 \\ \sum_L b_2 ds \beta_x^{1/2} \beta_y - \sum_R b_2 ds \beta_x^{1/2} \beta_y = 0 \end{cases} \quad (8)$$

Eq. (8) gives the same weight factors $\beta_x^{3/2}$ and $\beta_x^{1/2} \beta_y$ as that from the action-angle kick minimization method shown in Table 1.

OFF-LINE COMPARISON

IR bump Simulation

The MADX code [5] is used to simulate the IR bump correction. At each bump step, we set the right dipole kicker strengths to produce the desired orbit amplitude. MADX calculates the tune shifts at each bump step. By fitting the tune shift with the bump steps, the different orders of polynomial coefficients are obtained.

For simplicity, in the following we only simulate the IR bump correction in the 8 o'clock interaction region in the RHIC Blue ring. In the simulation, the maximum orbit amplitude in the IR bump is chosen below 5 mm. Figure 1 shows an example of the orbit with the horizontal IR bump. The orbit leakage due to the IR bump is seen in Figure 1. However, the bump closure is still acceptable. Fig. 2 gives the tune shifts from the simulation. The tune shifts are obtained with respect to that at zero bump amplitude. Up to seven order polynomial fittings are performed, the fitting curve is also shown in Fig. 2. Fig.3 shows the tune shift contributions from the first three polynomial terms.

Figure 2: The tune shifts and the polynomial fitting.

Figure 3: The tune shift contributions from the first three polynomial terms.

Sextupole Correction

For the horizontal IR bump, the linear terms of the tune shifts are contributed from the sextupoles. We use two sextupole correctors bo7-sx3 and bi8-sx3 to minimize these linear term tune shifts in the Blue IR8. The three dipole kickers bo7-th4, bi8-th3 and bi8-th5 bump produce the desired IR bumps.

Table 2 gives the linear orders from the off-line IR bump simulations. The second block gives the residual linear term of the tune shifts from the IR bump in the IR8. The third block gives the *two* correctors' contributions to the linear terms with $k_2 l = 0.001$. Based on Table 2, the correction strengths for bo7-sx3 and bi8-sx3 are calculated to cancel the residual linear terms.

From IR bump simulation, the correction strengths for bo7-sx3 and bi8-sx3 are $-4.54 \times 10^{-3} \text{ m}^{-2}$ and $2.74 \times 10^{-3} \text{ m}^{-2}$, respectively. While from the action-angle kick minimization analytical calculation, based on the nonlinear optics model and Eq.(1), the integrated correction strengths for bo7-sx3 and bi8-sx3 are $-3.99 \times 10^{-3} \text{ m}^{-2}$ and $2.97 \times 10^{-3} \text{ m}^{-2}$, respectively. There is about a 10% difference in the correction strength of bo7-sx3.

Conditions	Plane	Linear term Coefficient
only b_2 errors	x	10.08×10^{-5}
	y	-9.26×10^{-5}
$bo7-sx3$ $k_{2l} = 0.001 \text{ m}^{-2}$	x	5.65×10^{-6}
	y	-1.43×10^{-5}
$bi8-sx3$ $k_{2l} = 0.001 \text{ m}^{-2}$	x	-2.74×10^{-5}
	y	1.01×10^{-5}

Octupole Correction

We use the two octupole correctors $bo7-oct2$ and $bi8-oct2$ to minimize the quadratic terms of the tune shifts from the horizontal IR bump in Blue IR8. The three dipole kickers $bo7-th4$, $bi8-th3$ and $B18-TH5$ bump produce the desired IR bumps. Table 3 gives the residual quadratic terms from the octupole errors from the IR bump simulations. The correctors $bo7-oct2$ and $bi8-oct2$'s contributions to the quadratic terms with $k_{3l} = 0.001$ are given, too.

Table 3: IR bump simulation for octupole correction in Blue IR8.

Conditions	Plane	Quadratic Term coefficient
only b_3 errors	x	-1.68×10^{-7}
	y	6.40×10^{-8}
$bo7-oct2$ $k_{3l} = 0.001 \text{ m}^{-3}$	x	1.58×10^{-7}
	y	-8.74×10^{-8}
$bi8-oct2$ $k_{3l} = 0.001 \text{ m}^{-3}$	x	7.94×10^{-8}
	y	-1.43×10^{-7}

From the IR bump correction simulation, the integrated correction strengths for $bo7-oct2$ and $bi8-oct2$ are 0.121 m^{-3} and -0.029 m^{-3} , respectively. While from the action-angle kick minimization analytical calculation, based on the IR nonlinear modeling, the integrated correction strengths for $bo7-sx3$ and $bi8-sx3$ are 0.0768 m^{-3} and -0.023 m^{-3} , respectively. There are about 30% difference in the correction strengths of $bo7-oct2$.

ANALYSIS

From the off-line IR bump correction and the action angle minimization analytical calculation, we found that there are about 10% difference for the sextupole corrector strengths, and about 30% difference for the octupole corrector strengths. They verified the discrepancies found in the sextupole and octupole correction strengths from the operational IR bump corrections and the off-line action-angle kicker minimization analytical calculations.

Here we check the ratios of the linear and quadratic terms of two individual sextupoles and octupoles, respectively.

For sextupoles, according to Eq. (8), the horizontal linear term should be proportional to $\beta_x^{3/2}$ for two sextupoles if they have the same integrated strength. From IR bump simulations, the ratio of the linear term of the horizontal tune shifts from two sextupoles $bo7-sx3$ and $bi8-sx3$ with the same integrated strength 0.01 m^{-2} is

$$\begin{aligned} \Delta\nu_x|_{bo7-sx3} : \Delta\nu_x|_{bi8-sx3} \\ = 5.29 \times 10^{-5} : 27.79 \times 10^{-5} \\ = 1 : 5.25 \end{aligned} \quad (9)$$

The ratio of the $\beta_x^{3/2}$ of the two sextupoles are:

$$\begin{aligned} \beta_x^{3/2}|_{bo7-sx3} : \beta_x^{3/2}|_{bi8-sx3} \\ = 479.03^{3/2} : 1297.89^{3/2} \\ = 1 : 4.46 \end{aligned} \quad (10)$$

Therefore, from the IR bump simulation, the ratio of the horizontal tune shift terms is not proportional to the ratio of $\beta_x^{3/2}$ for two sextupoles with the same integrated strength. This only reason for the inequality is

$$\begin{aligned} x_{co}|_{bo7-sx3} : x_{co}|_{bi8-sx3} \\ \neq \beta_x^{1/2}|_{bo7-sx3} : \beta_x^{1/2}|_{bi8-sx3}. \end{aligned} \quad (11)$$

This guess is verified by the following orbit bump check at these two sextupole correctors. From the IR bump simulation with MADX,

$$\begin{aligned} x_{co}|_{bo7-sx3} : x_{co}|_{bi8-sx3} \\ = 1 : 1.955, \\ \beta_x^{1/2}|_{bo7-sx3} : \beta_x^{1/2}|_{bi8-sx3} \\ = 1 : 1.646. \end{aligned} \quad (12)$$

If we substitute the orbit ratio of x_{co} instead of the ratio of $\beta_x^{1/2}$ into Eq. (4), we get the horizontal tune shift contribution ratio from the two sextupoles:

$$\begin{aligned} (x_{co}\beta_x)|_{bo7-sx3} : (x_{co}\beta_x)|_{bi8-sx3} \\ = 1 : 5.30, \end{aligned} \quad (13)$$

which is much closer to the linear term tune shift ratio we get from the IR bump simulation.

Similarly, we check the quadratic term tune shift ratios of two individual octupoles $bo7-oct2$ and $bi8-oct2$ with the same integrated strength 0.001 m^{-3} . From the IR bump simulation, we get

$$\begin{aligned} \Delta\nu_x|_{bo7-oct2} : \Delta\nu_x|_{bi8-oct2} \\ = 1.58 \times 10^{-7} : 7.94 \times 10^{-8} \\ = 1.996 : 1 \end{aligned} \quad (14)$$

However,

$$\begin{aligned} \beta_x^2|_{bo7-sx3} : \beta_x^2|_{bi8-sx3} \\ = 1042.11' : 577.37' \\ = 3.261 : 1 \end{aligned} \quad (15)$$

The ratio of the tune shifts are not equal to the ratio of β_x^2 as assumed from Table 1, either. The ratio of orbit amplitudes at the two octupoles is:

$$\begin{aligned} x_{co}|_{bo7-oct2} : x_{co}|_{bi8-oct2} \\ = 1.051 : 1 \end{aligned} \quad (16)$$

Substituting x_{co} ratio instead of $\beta_x^{1/2}$ ratio to calculate the tune shifts due to octupoles, we obtain the horizontal tune shift contribution ratio from the two octupoles:

$$\frac{(x_{co}^2 \beta_x)|_{b_{07-sx3}} : (x_{co}^2 \beta_x)|_{b_{i8-sx3}}}{= 1.995 : 1} \quad (17)$$

which is almost the same as that from the IR bump simulation.

DISCUSSION

From the above calculation, we find the difference of the correction strengths from the IR bump correction and the action-angle kick minimization comes from the fact that the horizontal orbit is not exactly proportional to the $\beta_x^{1/2}$. The source for this difference is that the phase advance over the interaction region is not exactly equal to π and there is a small phase advance in the triplet.

Presenting the phase advance as $\Delta\Psi = \pi + \Delta\psi$ in another IR side, one can get the ratio of orbit positions on the left and right sides of the interaction region as

$$\begin{aligned} \frac{x_{co,L}}{x_{co,R}} &= \frac{\sqrt{\beta_{x,L}} \sin(\Psi_0)}{\sqrt{\beta_{x,R}} \sin(\Psi_0 + \Delta\Psi)} \\ &\simeq -\frac{\sqrt{\beta_{x,L}}}{\sqrt{\beta_{x,R}}} (1 - \cot(\Psi_0) \Delta\psi). \end{aligned} \quad (18)$$

Although $\Delta\psi$ is small, the $\cot\Psi_0$ can reach 15 units for RHIC IR bump. It leads to considerable difference between the ratio of the orbits and the ratio of $\sqrt{\beta_x}$. Then the correction strengths from the IR bump correction are not the same as that from the action-angle kick minimization analytical prediction.

Action-angle kick minimization ignores the small phase change in the IR bump on both sides of the IP, while the IR bump correction method takes into account the phase shifts. Based on the Hamiltonian perturbation theory, sextupoles could introduce $Q_x = p$, $3Q_x = p$, $Q_x \pm 2Q_y = p$ resonances. To fully correct all the resonances, we should minimize all the following resonance strengths [6], that is,

$$\left\{ \begin{array}{ll} \sum_j k_2 l \beta_x^{1/2} \beta_y e^{i\Psi_x} & \longrightarrow 0 \\ \sum_j k_2 l \beta_x^{3/2} e^{i\Psi_x} & \longrightarrow 0 \\ \sum_j k_2 l \beta_x^{3/2} e^{i3\Psi_x} & \longrightarrow 0 \\ \sum_j k_2 l \beta_x^{1/2} \beta_y e^{i(\Psi_x - 2\Psi_y)} & \longrightarrow 0 \\ \sum_j k_2 l \beta_x^{1/2} \beta_y e^{i(\Psi_x + 2\Psi_y)} & \longrightarrow 0 \end{array} \right. \quad (19)$$

Octupoles induce $4Q_x = p$, $4Q_y = p$, $2Q_x = p$, $2Q_y = p$, $2Q_x \pm 2Q_y = p$ resonances. To correct all the resonances, we should minimize all the following resonance strengths,

that is,

$$\left\{ \begin{array}{ll} \sum_j k_3 l \beta_x^2 e^{i4\Psi_x} & \longrightarrow 0 \\ \sum_j k_3 l \beta_y^2 e^{i4\Psi_y} & \longrightarrow 0 \\ \sum_j k_3 l \beta_x \beta_y e^{i2\Psi_x} & \longrightarrow 0 \\ \sum_j k_3 l \beta_x^2 e^{i2\Psi_x} & \longrightarrow 0 \\ \sum_j k_3 l \beta_x \beta_y e^{i2\Psi_y} & \longrightarrow 0 \\ \sum_j k_3 l \beta_y^2 e^{i2\Psi_y} & \longrightarrow 0 \\ \sum_j k_3 l \beta_x \beta_y e^{i(2\Psi_x + 2\Psi_y)} & \longrightarrow 0 \\ \sum_j k_3 l \beta_x \beta_y e^{i(2\Psi_x - 2\Psi_y)} & \longrightarrow 0 \end{array} \right. \quad (20)$$

So the two methods, IR bump correction and the action-angle kick minimization, have different approximations in the betatron phase advance. Action-angle kicker minimization ignores the not exact π phase jump across the IP and the small phase change in the one side triplet. The IR bump correction method uses a local orbit bump to minimize the introduced polynomial terms of tune shifts. However, the tune shifts from IR bump are proportional to $\sin^n \Phi$, instead of $e^{in\Phi}$ from Eqs. (19) - (20).

The action-angle kick minimization is used for IR nonlinear field correction off line up to b_{10} at CERN [7]. The IR bump correction is applicable for the on-line IR nonlinear field correction. It has been verified and used in the RHIC IR nonlinear corrections. Limited by the bump amplitude and the tune measurement resolution, it is generally used for the lower order nonlinear field error corrections.

CONCLUSION

The correction strengths from the off-line IR bump correction simulation are compared to that from the action-angle kick minimization. It verifies that there are some discrepancies in the correction strengths from these two methods. This is caused by the fact that the not exact π phase advance between the two sides of the interaction region, which makes that the bump orbit not exactly proportionally to the $\beta^{1/2}$. Both methods make different approximations in the betatron phase advances. The action-angle kick minimization is used for IR nonlinear field correction off line, while the IR bump correction is applicable for the on-line IR nonlinear field correction.

REFERENCES

- [1] J. Wei, Proceedings of the Workshop on LHC Interaction Region Correction Systems, BNL, May 6-7, 1999.
- [2] J-P. Koutchouk, F. Pilat, V. Ptitsyn, et al, PAC2001.
- [3] F. Pilat, V. Ptitsyn, et al., EPAC2002, Paris, France.
- [4] Y. Luo, F. Pilat, V. Ptitsyn, et al., **ENL.C-AD AP** Note 160, Aug. 2004.
- [5] Madx code, CERN, <http://madhoney.cern.ch/madx/>.
- [6] H. Wiedemann, Particle Accelerator Physics: Nonlinear and Higher-order Beam Dynamics, Springer Press, 1994.
- [7] H. Grote, et al., LHC-project-note 197.