CHARACTERISTICS OF POTENTIAL REPOSITORY WASTES

JULY 1992

Prepared for the
U.S. DEPARTMENT OF ENERGY
OFFICE OF CIVILIAN RADIOACTIVE WASTE MANAGEMENT
Washington, D.C. 20585

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under
Contract No. DE-AC05-84OR21400

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831; prices available from (615) 576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Characteristics of Potential Repository Wastes

July 1992

**Appendix 3A. ORIGEN2 Decay Tables for
Immobilized High-Level Waste**

Appendix 3B. Interim High-Level Waste Forms

Prepared for the
U.S. DEPARTMENT OF ENERGY
OFFICE OF CIVILIAN RADIOACTIVE WASTE MANAGEMENT
Washington, D.C. 20585

Prepared by the
OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
managed by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
under
Contract No. DE-AC05-84OR21400

MASTER

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

CONTENTS — VOLUME 3

LIST OF ACRONYMS	v
APPENDIX 3A. ORIGEN2 DECAY TABLES FOR IMMOBILIZED	
HIGH-LEVEL WASTE	3A-1
Tables 3A.1 through 3A.5: radionuclide contents per canister, all sites	3A-5
Tables of radioactivity per canister as a function of time from 0 to 10 ⁶ years, all sites:	
WEST VALLEY DEMONSTRATION PROJECT	3A-18
SAVANNAH RIVER SITE	3A-46
HANFORD SITE	3A-74
IDAHO NATIONAL ENGINEERING LABORATORY	3A-128
APPENDIX 3B. INTERIM HIGH-LEVEL WASTE FORMS	3B-1
3B.1 INTRODUCTION	3B-3
3B.2 WEST VALLEY DEMONSTRATION PROJECT	3B-3
3B.3 SAVANNAH RIVER SITE	3B-4
3B.4 HANFORD SITE	3B-6
3B.5 IDAHO NATIONAL ENGINEERING LABORATORY	3B-9

LIST OF ACRONYMS

AC	Allis Chalmers
ANF	Advanced Nuclear Fuels Corporation
ANL	Argonne National Laboratory
AP	Activation products
APSR	Axial power shaping rod
ASTM	American Society for Testing and Materials
B-C	Battelle-Columbus
B&W	Babcock and Wilcox
BPRA	Burnable poison rod assembly
BWR	Boiling-water reactor
CC	Complexant concentrate
CDB	Characteristics Data Base
CE	Combustion Engineering
CEA	Control element assembly
CEU	Consolidated Edison uranium
CFR	Code of Federal Regulations
CH	Contact handled
CWMS	Civilian Waste Management System
DHLW	Defense high-level waste
DOE	Department of Energy
DWPF	Defense Waste Processing Facility
ECF	Expended Core Facility
EFPD	Equivalent full-power days
EIA	Energy Information Administration
EIS	Environmental impact statement
EPRI	Electric Power Research Institute
FFTF	Fast Flux Test Facility
FIS	Federal Interim Storage
FP	Fission products
FSV	Fort St. Vrain
GAPSR	Gray axial power shaping rod
GE	General Electric
GTCC	Greater than Class C
GWd	Gigawatt-days
GW(e)	Gigawatts (electric)
HANF	Hanford Site
HEDL	Hanford Engineering Development Laboratory
HEPA	High-efficiency particulate air
HLW	High-level waste
HTGR	High-temperature gas-cooled reactor
HWVP	Hanford Waste Vitrification Plant
ICPP	Idaho Chemical Processing Plant
IDB	Integrated Data Base
INEL	Idaho National Engineering Laboratory
LANL	Los Alamos National Laboratory
LER	Licensee Event Report
LLW	Low-level waste
LTA	Lead test assembly
LWBR	Light-water breeder reactor
LWR	Light-water reactor
MOX	Mixed oxide
MRS	Monitored retrievable storage

LIST OF ACRONYMS (continued)

MSRE	Molten Salt Reactor Experiment
MT	Metric tons
MTIHM	Metric tons of initial heavy metal
MTR	Materials Test Reactor
MW(t)	Megawatts (thermal)
MW(e)	Megawatts (electric)
NCAW	Neutralized current acid waste
NCRW	Neutralized cladding removal waste
NFA	Non-fuel assembly
NFB	Non-fuel bearing
NFS	Nuclear Fuel Services, Inc.
NMMSS	Nuclear Materials Management and Safeguards System
NPR	New Production Reactor
NRC	Nuclear Regulatory Commission
NRF	Naval Reactors Facility
NWTSP	National Waste Terminal Storage Program
O/U	Oxygen/uranium atom ratio
OCRWM	Office of Civilian Radioactive Waste Management
OFA	Optimized fuel assembly
ORA	Orifice rod assembly
ORNL	Oak Ridge National Laboratory
PB1	Peach Bottom Unit 1
PC	Personal computer
PCI	Pellet-clad interaction
PFP	Plutonium finishing plant
PIE	Postirradiation examination
PNL	Pacific Northwest Laboratory
PNS	Primary neutron source
PWR	Pressurized-water reactor
QA	Quality assurance
QC	Quality control
RH	Remotely handled
RNS	Regenerative neutron source
SAS	Statistical Analysis System
SFD	Spent fuel disassembly
SNF	Spent nuclear fuel
SRL	Savannah River Laboratory
SRP	Savannah River Plant
SRS	Savannah River Site
SS	Stainless steel
SST	Single-shell tanks
TMI-2	Three Mile Island 2
TRIGA	Training Research Isotopes – General Atomics
TRU	Transuranic (waste)
TRUW	Transuranic waste
UN	United Nuclear
VEPCO	Virginia Electric Power Co.
WAC	Waste acceptance criteria
WAPS	Waste Acceptance Preliminary Specification
WE or <u>W</u>	Westinghouse
WIPP	Waste Isolation Pilot Plant
WVDP	West Valley Demonstration Project

**APPENDIX 3A. ORIGEN2 DECAY TABLES FOR
IMMOBILIZED HIGH-LEVEL WASTE**

APPENDIX 3A. ORIGEN2 DECAY TABLES FOR IMMOBILIZED HIGH-LEVEL WASTE

This appendix presents the results of decay calculations using the ORIGEN2 code to determine the radiological properties of canisters of immobilized high-level waste as a function of decay time for decay times up to one million years. These calculations were made for the four HLW sites (West Valley Demonstration Project, Savannah River Site, Hanford Site, and Idaho National Engineering Laboratory) using the composition data discussed in the HLW section of this report. Calculated (α , n) neutron production rates are also shown. These were calculated by the ALPHN code (Salmon 1992).

The reference canister compositions for the four sites are shown in Tables 3A.1-3A.4. The reference compositions for SRS and HANF represent the maximum activity canisters anticipated from the site. In the case of HANF, an average or nominal composition for neutralized current acid waste glass (NCAW) is also shown. For WVDP, the average composition from the latest mass balance (Revision 7) was used; a maximum composition was not available. For INEL, no canister composition data have been made available, so an estimate was made based on a 1982 report (IDO-1982). The INEL canister load was based on the assumption that the high-density glass-ceramic form is used. Canister outside dimensions are the same for WVDP, SRS, and HANF, approximately 2 ft in diameter and 10 ft in length, and it is assumed that the INEL canisters will have approximately these same dimensions.

The initial time point for the West Valley canister decay calculations was selected as the end of year 1989 because the reference glass composition data were based on that year.

The three defense sites continue to reprocess fuel and produce fresh HLW. The composition of the waste of maximum activity was estimated for each site, and the starting point for the decay calculations was taken as the year in which immobilization of this waste occurs. Thus, decay time for these wastes means the time elapsed since filling the canister.

The ORIGEN2 and ALPHN output tables are grouped according to site and are presented in the following order: WVDP, SRS (DWPF), HANF, and INEL. The output quantities represent one canister in each case. The order of presentation for each site except INEL is as follows:

- Table 1. Actinides and daughters, grams, by nuclides
- Table 2. Actinides and daughters, grams, by elements
- Table 3. Actinides and daughters, curies, by nuclides
- Table 4. Actinides and daughters, curies, by elements
- Table 5. Actinides and daughters, watts, by nuclides
- Table 6. Actinides and daughters, watts, by elements
- Table 7. Actinides and daughters, alpha curies, by nuclides

- Table 8. Actinides and daughters, alpha curies, by elements
- Table 9. Actinides and daughters, (alpha, n) neutron sources, by nuclides
- Table 10. Actinides and daughters, spontaneous fission neutron sources, by nuclides
- Table 11. Actinides and daughters, photon spectrum, by energy groups
- Table 12. Fission products, grams, by nuclides
- Table 13. Fission products, grams, by elements
- Table 14. Fission products, curies, by nuclides
- Table 15. Fission products, curies, by elements
- Table 16. Fission products, watts, by nuclides
- Table 17. Fission products, watts, by elements
- Table 18. Fission products, photon spectrum, by energy groups
- Table 19. Activation products, grams, by nuclides
- Table 20. Activation products, grams, by elements
- Table 21. Activation products, curies, by nuclides
- Table 22. Activation products, curies, by elements
- Table 23. Activation products, watts, by nuclides
- Table 24. Activation products, watts, by elements
- Table 25. Activation products, photon spectrum, by energy groups

In the case of INEL the order is the same as above except that fission and activation products are shown together in Tables 12 through 18, because no data were available to permit a separate breakdown. Table 9 for INEL is not included; this table is in the process of being revised and will be added when available.

In Tables 1-25, "sum" means the sum of the radionuclides listed in the table, and "total" means the sum of all of the radionuclides in the ORIGEN2 run, including those not listed in the table. Cutoffs were chosen to get tables of reasonable size, while still including all radionuclides contributing a significant fraction of the total radioactivity. The cutoffs used are shown at the bottoms of the tables. A radionuclide is shown if, and only if, its value exceeds the cutoff at any time point within the table. If a radionuclide is shown at some time point in a table, it is shown at all time points in that table. It was not practical to use the same cutoff in every table, so cutoff values vary from table to table. The computerized HLW data base, which was based on the ORIGEN2 outputs, contains values for many radionuclides not shown in Tables 1-25.

The data presented here are also available in magnetic diskette form, as described in Appendix 3C. In the hard copy presented here, decay times of 1, 10, 100, 1,000, 10,000, 100,000, and 1,000,000 years are shown. The diskette data include a number of intermediate decay times, as follows: 1, 2, 5, 10, 15, 20, 30, 50, 100, 200, 300, 350, 500, 1,000, 1,050, 2,000, 5,000, 10,000, 20,000, 50,000,

100,000, 500,000, and 1,000,000 years. The years 350 and 1,050 were included at the request of the project sponsor, because of the possible need for calculations at emplacement age plus 300 or 1,000 years after repository closure.

References

Baxter 1988. R. G. Baxter, *Defense Waste Processing Facility Wasteform and Canister Description*, DP-1606, Rev. 2, December 1988.

Crocker 1989. R. L. Crocker, *West Valley Vitirification Process Mass Balance*, Revision 7, October 10, 1989.

IDO 1982. *Environmental Evaluation of Alternatives for Long-Term Management of Defense High-Level Radioactive Wastes at the ICPP*, IDO-10105, U.S. DOE, Idaho Operations Office, September 1982.

Mitchell and Nelson 1988. D. E. Mitchell and J. L. Nelson, *Hanford Waste Vitirification Plant Preliminary Description of Waste Form and Canister - FY 1988 Update*, WHC-EP-0008 Rev. 1, June 1988.

Salmon 1992. R. Salmon and O. W. Hermann, *ALPHN - A Computer Program for Calculating (α , n) Neutron Production in Canisters of High-Level Waste*, ORNL/TM-12016 (in preparation).

Table 3A.1. West Valley Demonstration Project: estimated radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	0.1104E-02	0.2760E+01	0.9313E-04
Co-60	0.2679E-02	0.3030E+01	0.4666E-01
Ni-59	0.5491E+01	0.4160E+00	0.1650E-04
Ni-63	0.4895E+00	0.3020E+02	0.3039E-02
Se-79	0.1980E+00	0.1380E-01	0.3431E-05
Sr-90	0.1928E+03	0.2630E+05	0.3048E+02
Y-90	0.4833E-01	0.2630E+05	0.1456E+03
Zr-93	0.4257E+03	0.1070E+01	0.1242E-03
Nb-93m	0.2529E-02	0.7150E+00	0.1265E-03
Tc-99	0.2524E+02	0.4280E+00	0.2144E-03
Ru-106	0.1655E-04	0.5540E-01	0.3290E-05
Rh-106	0.1556E-10	0.5540E-01	0.5307E-03
Pd-107	0.8416E+02	0.4330E-01	0.2563E-05
Cd-113m	0.3845E-01	0.8340E+01	0.1402E-01
Sn-121m	0.1160E-02	0.6860E-01	0.1373E-03
Sn-126	0.1441E+02	0.4090E+00	0.5095E-03
Sb-125	0.2769E-01	0.2860E+02	0.8929E-01
Sb-126	0.6852E-06	0.5730E-01	0.1057E-02
Sb-126m	0.5206E-08	0.4090E+00	0.5201E-02
Te-125m	0.3885E-03	0.7000E+01	0.5876E-02
Cs-134	0.1569E-01	0.2030E+02	0.2063E+00
Cs-135	0.5505E+03	0.6340E+00	0.2113E-03
Cs-137	0.3252E+03	0.2830E+05	0.3126E+02
Ba-137m	0.4981E-04	0.2680E+05	0.1051E+03
Ce-144	0.8023E-06	0.2560E-02	0.1696E-05
Pr-144	0.3387E-10	0.2560E-02	0.1879E-04
Pm-146	0.9566E-04	0.4260E-01	0.2146E-03
Pm-147	0.3721E+00	0.3450E+03	0.1236E+00
Sm-151	0.1258E+02	0.3310E+03	0.3876E-01
Eu-152	0.8267E-02	0.1430E+01	0.1080E-01
Eu-154	0.1389E+01	0.3750E+03	0.3350E+01
Eu-155	0.2014E+00	0.9370E+02	0.6806E-01
Tl-207	0.1690E-09	0.3220E-01	0.9444E-04
Tl-208	0.4312E-10	0.1270E-01	0.2985E-03
Pb-209	0.1815E-09	0.8250E-03	0.9475E-06
Pb-211	0.1308E-08	0.3230E-01	0.9666E-04
Pb-212	0.2540E-07	0.3530E-01	0.6712E-04
Bi-211	0.7718E-10	0.3230E-01	0.1287E-02
Bi-212	0.2409E-08	0.3530E-01	0.5995E-03

Table 3A.1 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Bi-213	0.4265E-10	0.8250E-03	0.3464E-05
Po-212	0.1274E-18	0.2260E-01	0.1196E-02
Po-213	0.6231E-19	0.7860E-03	0.3972E-04
Po-215	0.1095E-14	0.3230E-01	0.1440E-02
Po-216	0.1013E-12	0.3530E-01	0.1443E-02
At-217	0.5124E-15	0.8250E-03	0.3516E-04
Rn-219	0.2482E-11	0.3230E-01	0.1339E-02
Rn-220	0.3826E-10	0.3530E-01	0.1339E-02
Fr-221	0.4653E-11	0.8250E-03	0.3180E-04
Fr-223	0.1117E-10	0.4320E-03	0.1120E-05
Ra-223	0.6306E-06	0.3230E-01	0.1149E-02
Ra-224	0.2216E-06	0.3530E-01	0.1210E-02
Ra-225	0.2104E-07	0.8250E-03	0.5778E-06
Ra-228	0.2550E-04	0.5970E-02	0.4595E-06
Ac-225	0.1421E-07	0.8250E-03	0.2878E-04
Ac-227	0.4464E-05	0.3230E-03	0.1562E-06
Ac-228	0.2662E-08	0.5970E-02	0.5153E-04
Th-227	0.1034E-05	0.3180E-01	0.1159E-02
Th-228	0.4306E-04	0.3530E-01	0.1153E-02
Th-229	0.3877E-02	0.8250E-03	0.2521E-04
Th-230	0.1169E-01	0.2360E-03	0.6670E-05
Th-231	0.6657E-09	0.3540E-03	0.1984E-06
Th-232	0.5880E+05	0.6450E-02	0.1559E-03
Th-234	0.1356E-06	0.3140E-02	0.1271E-05
Pa-231	0.1264E+01	0.5970E-01	0.1796E-02
Pa-233	0.4422E-05	0.9180E-01	0.2081E-03
Pa-234m	0.4571E-11	0.3140E-02	0.1550E-04
U-232	0.1270E-02	0.2720E-01	0.8721E-03
U-233	0.3666E+01	0.3550E-01	0.1031E-02
U-234	0.2640E+01	0.1650E-01	0.4746E-03
U-235	0.1637E+03	0.3540E-03	0.9259E-05
U-236	0.1700E+02	0.1100E-02	0.2976E-04
U-238	0.9337E+04	0.3140E-02	0.7954E-04
Np-236	0.2823E+01	0.3720E-01	0.7494E-04
Np-237	0.1302E+03	0.9180E-01	0.2802E-02
Np-239	0.5861E-05	0.1360E+01	0.3283E-02
Pu-236	0.6209E-05	0.3300E-02	0.1147E-03
Pu-238	0.1904E+01	0.3260E+02	0.1079E+01
Pu-239	0.1028E+03	0.6390E+01	0.1967E+00

Table 3A.1 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Pu-240	0.2053E+02	0.4680E+01	0.1455E+00
Pu-241	0.3076E+01	0.3170E+03	0.9815E-02
Pu-242	0.1668E+01	0.6370E-02	0.1879E-03
Am-241	0.6117E+02	0.2100E+03	0.6967E+01
Am-242	0.1435E-05	0.1160E+01	0.1315E-02
Am-242m	0.1204E+00	0.1170E+01	0.4616E-03
Am-243	0.6820E+01	0.1360E+01	0.4366E-01
Cm-242	0.2912E-03	0.9630E+00	0.3544E-01
Cm-243	0.1021E-01	0.5270E+00	0.1931E-01
Cm-244	0.3707E+00	0.3000E+02	0.1048E+01
Cm-245	0.2015E-01	0.3460E-02	0.1147E-03
Cm-246	0.1279E-02	0.3930E-03	0.1285E-04
Total	0.7029E+05	0.1096E+06	0.3260E+03

^aThis table represents the radionuclide content of a canister containing 1,900 kg of HLW glass having the radionuclide composition described in the WVDP Mass Balance Revision 7 (Crocker 1989). Radioactivity shown is as of the end of year 1989.

Table 3A.2. Savannah River Site: radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Cr-51	0.1008E-20	0.9312E-16	0.1996E-19
Co-60	0.1502E+00	0.1699E+03	0.2619E+01
Ni-59	0.3163E+00	0.2397E-01	0.9519E-06
Ni-63	0.4824E-01	0.2975E+01	0.3000E-03
Se-79	0.2439E+01	0.1699E+00	0.4232E-04
Rb-87	0.9961E+01	0.8719E-06	0.7278E-09
Sr-89	0.1470E-08	0.4267E-04	0.1473E-06
Sr-90	0.3426E+03	0.4675E+05	0.5426E+02
Y-90	0.8795E-01	0.4786E+05	0.2653E+03
Y-91	0.3085E-07	0.7568E-03	0.2715E-05
Zr-93	0.4443E+03	0.1117E+01	0.1298E-03
Zr-95	0.4680E-06	0.1005E-01	0.5084E-04
Nb-94	0.5147E-03	0.9646E-04	0.9830E-06
Nb-95	0.5407E-06	0.2115E-01	0.1013E-03
Nb-95m	0.3272E-09	0.1247E-03	0.1730E-06
Tc-99	0.1816E+03	0.3079E+01	0.1545E-02
Ru-103	0.5217E-12	0.1684E-07	0.5827E-10
Ru-106	0.6729E+00	0.2252E+04	0.1339E-00
Rh-103m	0.5028E-15	0.1636E-07	0.3761E-11
Rh-106	0.6346E-06	0.2259E+04	0.2167E+02
Pd-107	0.2863E+02	0.1473E-01	0.8732E-06
Ag-110m	0.2647E-04	0.1258E+00	0.2098E-02
Cd-113	0.1472E+00	0.5009E-13	0.8420E-16
Cd-115m	0.4763E-13	0.1213E-08	0.4518E-11
Sn-121m	0.1336E-02	0.7902E-01	0.1581E-03
Sn-123	0.3101E-04	0.2549E+00	0.7951E-03
Sn-126	0.1556E+02	0.4415E+00	0.5508E-03
Sb-124	0.4071E-11	0.7123E-07	0.9445E-09
Sb-125	0.8226E+00	0.8496E+03	0.2656E+01
Sb-126	0.7365E-06	0.6159E-01	0.1138E-02
Sb-126m	0.5619E-08	0.4415E+00	0.5622E-02
Te-126m	0.1532E-01	0.2760E+03	0.2320E+00

Table 3A.2 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-127	0.4555E-07	0.1202E+00	0.1622E-03
Te-128m	0.1302E-04	0.1228E+00	0.6597E-04
Te-129	0.1457E-18	0.3053E-11	0.1089E-13
Te-129m	0.1576E-15	0.4749E-11	0.8316E-14
Cs-134	0.2606E+00	0.3372E+03	0.3433E+01
Cs-135	0.8633E+02	0.9943E-01	0.3319E-04
Cs-136	0.1068E-43	0.7828E-39	0.1066E-41
Cs-137	0.4989E+03	0.4341E+05	0.4802E+02
Ba-136m	0.3195E-49	0.8607E-38	0.1040E-40
Ba-137m	0.7724E-04	0.4155E+05	0.1632E+03
Ba-140	0.1404E-40	0.1024E-35	0.2853E-38
La-140	0.7734E-42	0.4304E-36	0.7205E-38
Ce-141	0.1260E-14	0.3591E-10	0.5250E-13
Ce-142	0.4005E+03	0.9609E-05	0.0000E+00
Ce-144	0.3093E+01	0.9869E+04	0.6547E+01
Pr-143	0.1780E-38	0.1198E-33	0.2291E-37
Pr-144	0.1306E-03	0.9869E+04	0.7255E+02
Pr-144m	0.6545E-06	0.1187E+03	0.4063E-01
Nd-144	0.4110E+03	0.4860E-09	0.0000E+00
Nd-147	0.1570E-48	0.1261E-43	0.3038E-46
Pm-147	0.2609E+02	0.2419E+05	0.8679E+01
Pm-148	0.4243E-15	0.6975E-10	0.5364E-12
Pm-148m	0.4722E-13	0.1009E-08	0.1277E-10
Sm-147	0.8796E+02	0.2000E-05	0.2738E-07
Sm-148	0.1916E+02	0.5788E-11	0.6901E-13
Sm-149	0.7420E+01	0.1781E-11	0.0000E+00
Sm-151	0.9418E+01	0.2478E+03	0.2906E-01
Eu-152	0.2132E-01	0.3688E+01	0.2790E-01
Eu-154	0.2295E+01	0.6196E+03	0.5543E+01
Eu-155	0.1021E+01	0.4749E+03	0.3455E+00
Eu-156	0.9489E-36	0.5231E-31	0.5392E-33
Tb-160	0.9923E-10	0.1120E-05	0.9110E-08

Table 3A.2 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Tl-208	0.3829E-11	0.1128E-02	0.2645E-04
U-232	0.6256E-03	0.1339E-01	0.4301E-03
U-233	0.1636E-03	0.1584E-05	0.4605E-07
U-234	0.5485E+01	0.3428E-01	0.9875E-03
U-235	0.7278E+02	0.1573E-03	0.4122E-05
U-236	0.1742E+02	0.1128E-02	0.3054E-04
U-238	0.3122E+05	0.1050E-01	0.2663E-03
Np-236	0.1323E-05	0.1744E-07	0.3514E-10
Np-237	0.1263E+02	0.8904E-02	0.2722E-03
Pu-236	0.2297E-03	0.1221E+00	0.4249E-02
Pu-237	0.7401E-15	0.8941E-11	0.3292E-14
Pu-238	0.8667E+02	0.1484E+04	0.4919E+02
Pu-239	0.2076E+03	0.1291E+02	0.3979E+00
Pu-240	0.3809E+02	0.8681E+01	0.2704E+00
Pu-241	0.1620E+02	0.1670E+04	0.5176E-01
Pu-242	0.3206E+01	0.1224E-01	0.3616E-03
Am-241	0.3210E+01	0.1102E+02	0.3661E+00
Am-242	0.1776E-07	0.1436E-01	0.1628E-04
Am-242m	0.1488E-02	0.1447E-01	0.5709E-05
Am-243	0.2902E-01	0.5788E-02	0.1860E-03
Cm-242	0.1057E-04	0.3495E-01	0.1288E-02
Cm-243	0.1078E-03	0.5565E-02	0.2039E-03
Cm-244	0.1329E+01	0.1076E+03	0.3763E+01
Cm-245	0.3910E-04	0.6715E-05	0.2225E-06
Cm-246	0.1739E-05	0.5342E-06	0.1747E-07
Cm-247	0.7116E-08	0.6604E-12	0.2107E-13
Cm-248	0.1614E-09	0.6864E-12	0.8533E-13
Totals	0.3427E+05	0.2344E+06	0.7093E+03

^aQuantities shown are for sludge-precipitate glass and are based on Baxter 1988, assuming sludge aged an average of 5 years and supernate aged an average of 15 years, with a canister load of 3,710 lb of glass (1,682 kg). Radionuclide contents are at time of filling canister.

Table 3A.3. Hanford Site: radionuclide content per HLW canister, NCAW glass, maximum case^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	5.64E-02	1.41E+02	4.765E-03
Ni-59	1.80E+00	1.36E-01	5.402E-06
Co-60	3.79E-03	4.29E+00	6.615E-02
Ni-63	2.54E-01	1.57E+01	6.236E-03
Se-79	5.60E-02	3.90E-03	9.711E-07
Sr-89	2.24E-06	6.52E-02	2.254E-04
Sr-90	3.06E+02	4.18E+04	4.852E+01
Y-90	7.68E-02	4.18E+04	2.317E+02
Y-91	2.96E-05	7.26E-01	2.608E-03
Nb-93m	2.04E-03	5.77E-01	1.023E-04
Zr-93	5.13E+02	1.29E+00	1.499E-04
Zr-95	1.28E-04	2.76E+00	1.398E-02
Nb-95	1.45E-04	5.67E+00	2.720E-02
Tc-99	5.51E+02	9.35E+00	4.689E-03
Ru-103	9.23E-09	2.98E-04	9.971E-07
Rh-103m	8.27E-12	2.69E-04	6.192E-08
Ru-106	1.49E+00	4.99E+03	2.967E-01
Rh-106	1.40E-06	4.99E+03	4.786E+01
Pd-107	7.91E+01	4.07E-02	2.413E-06
Ag-110m	3.35E-04	1.59E+00	2.655E-02
Cd-113m	6.73E-02	1.46E+01	2.458E-02
In-113m	1.51E-09	2.52E-02	5.871E-05
Sn-113	2.51E-06	2.52E-02	4.198E-06
Cd-115m	3.45E-10	8.78E-06	3.275E-08
Sn-119m	1.21E-03	5.42E+00	2.802E-03
Sn-121m	1.79E-03	1.06E-01	2.124E-04
Sn-123	3.52E-04	2.89E+00	9.027E-03
Sn-126	1.62E+01	4.60E-01	5.738E-04
Sb-124	2.00E-09	3.50E-05	4.648E-07
Sb-126	7.75E-07	6.48E-02	1.197E-03
Sb-126m	5.86E-09	4.60E-01	5.858E-03
Sb-125	1.70E+00	1.76E+03	5.503E+00
Te-125m	2.38E-02	4.29E+02	3.606E-01

Table 3A.3 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-127	1.12E-06	2.95E+00	3.984E-03
Te-127m	3.18E-04	3.00E+00	1.614E-03
Te-129	1.79E-14	3.75E-07	1.340E-09
Te-129m	1.91E-11	5.77E-07	1.012E-09
I-129	9.23E-02	1.63E-05	7.541E-09
Cs-134	9.27E-01	1.20E+03	1.221E+01
Cs-135	2.18E+02	2.51E-01	8.378E-05
Cs-137	5.86E+02	5.10E+04	5.642E+01
Ba-137m	8.96E-05	4.82E+04	1.893E+02
Ce-141	3.97E-10	1.13E-05	1.655E-08
Ce-144	9.34E+00	2.98E+04	1.977E+01
Pr-144	3.94E-04	2.98E+04	2.191E+02
Pr-144m	1.97E-06	3.58E+02	1.225E-01
Pm-147	4.28E+01	3.97E+04	1.424E+01
Pm-148m	6.18E-10	1.32E-05	1.674E-07
Sm-151	3.18E+01	8.36E+02	9.803E-02
Eu-152	1.58E-02	2.74E+00	2.073E-02
Gd-153	3.26E-06	1.15E-02	1.039E-05
Eu-154	1.24E+00	3.36E+02	3.006E+00
Eu-155	8.83E-01	4.11E+02	2.990E-01
Tb-160	9.74E-09	1.10E-04	8.961E-07
U-234	7.71E-01	4.82E-03	1.388E-04
U-235	9.11E+01	1.97E-04	5.160E-06
U-236	7.34E+00	4.75E-04	1.287E-05
U-238	1.11E+04	3.72E-03	9.437E-05
Np-237	2.82E+02	1.99E-01	6.083E-03
Pu-238	4.48E-02	7.68E-01	2.546E-02
Pu-239	2.27E+01	1.41E+00	4.346E-02
Pu-240	2.38E+00	5.42E-01	1.688E-02
Pu-241	2.50E-01	2.58E+01	7.999E-04
Pu-242	3.43E-02	1.31E-04	3.869E-06
Am-241	1.68E+02	5.77E+02	1.917E+01
Am-242	5.12E-07	4.14E-01	4.700E-04

Table 3A.3 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Am-243	3.39E-01	6.76E-02	2.173E-03
Cm-242	1.51E-04	4.99E-01	1.839E-02
Cm-244	1.54E-01	1.25E+01	4.374E-01
Total	1.40E+04	2.98E+05	8.687E+02

^aThis table identifies the maximum expected activity of HWVP canisters at the time of vitrification. The maximum is principally based on close-coupling the final accumulated tank of NCAW (21 months from fuel discharge to HWVP). Canister contains 1,650 kg of HLW glass (85% fill). Source: Mitchell and Nelson 1988.

Table 3A.4. Hanford Site: radionuclide content per HLW canister,
NCAW glass, nominal case^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	7.20E-03	1.80E+01	6.083E-04
Ni-59	1.44E+00	1.09E-01	4.330E-06
Co-60	1.33E-03	1.50E+00	2.313E-02
Ni-63	1.96E-01	1.21E+01	4.806E-03
Se-79	4.52E-02	3.15E-03	7.843E-07
Sr-89	1.84E-17	5.35E-13	1.850E-15
Sr-90	2.18E+02	2.98E+04	3.459E+01
Y-90	5.48E-02	2.98E+04	1.652E+02
Y-91	5.63E-15	1.38E-10	4.957E-13
Nb-93m	2.18E-03	6.16E-01	1.092E-04
Zr-93	4.18E+02	1.05E+00	1.220E-04
Zr-95	1.36E-13	2.92E-09	1.479E-11
Nb-95	1.72E-13	6.73E-09	3.229E-11
Tc-99	4.43E+02	7.51E+00	3.767E-03
Ru-103	1.04E-22	3.37E-18	1.126E-20
Rh-103m	9.34E-26	3.04E-18	6.988E-22
Ru-106	1.25E-02	4.18E+01	2.486E-03
Rh-106	1.17E-08	4.18E+01	4.009E-01
Pd-107	5.87E+01	3.02E-02	1.790E-06
Ag-110m	4.67E-07	2.22E-03	3.708E-05
Cd-113m	3.93E-02	8.53E+00	1.436E-02
In-113m	6.04E-15	1.01E-07	2.353E-10
Sn-113	1.01E-11	1.01E-07	1.683E-11
Cd-115m	1.26E-22	3.20E-18	1.192E-20
Sn-119m	1.52E-06	6.80E-03	3.516E-06
Sn-121m	1.31E-03	7.76E-02	1.555E-04
Sn-123	4.44E-09	3.65E-05	1.140E-07
Sn-126	1.29E+01	3.65E-01	4.553E-04
Sb-124	6.57E-19	1.15E-14	1.527E-16
Sb-126	6.10E-07	5.10E-02	9.424E-04
Sb-126m	4.65E-09	3.65E-01	4.648E-03
Sb-125	2.46E-01	2.54E+02	7.942E-01
Te-125m	3.44E-03	6.20E+01	5.212E-02
Te-127	2.48E-12	6.55E-06	8.846E-09
Te-127m	7.06E-10	6.66E-06	3.583E-09
Te-129	1.49E-30	3.14E-23	1.120E-25

Table 3A.4 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-129m	1.60E-27	4.82E-23	8.440E-26
I-129	7.31E-02	1.29E-05	5.968E-09
Cs-134	7.19E-02	9.31E+01	9.476E-01
Cs-135	1.75E+02	2.02E-01	6.742E-05
Cs-137	4.15E+02	3.61E+04	3.994E+01
Ba-137m	6.32E-05	3.40E+04	1.335E+02
Ce-141	1.03E-26	2.93E-22	4.284E-25
Ce-144	2.51E-02	8.00E+01	5.307E-02
Pr-144	1.06E-06	8.00E+01	5.881E-01
Pr-144m	5.29E-09	9.60E-01	3.285E-04
Pm-147	5.62E+00	5.21E+03	1.869E+00
Pm-148m	2.92E-23	6.23E-19	7.889E-21
Sm-151	2.65E+01	6.98E+02	8.185E-02
Eu-152	8.09E-03	1.40E-00	1.059E-02
Gd-153	3.83E-09	1.35E-05	1.220E-08
Eu-154	5.37E-01	1.45E+02	1.297E+00
Eu-155	2.94E-01	1.37E+02	9.965E-02
Tb-160	8.41E-17	9.49E-13	7.730E-15
U-234	7.31E-01	4.57E-03	1.316E-04
U-235	8.83E+01	1.91E-04	5.003E-06
U-236	6.51E+00	4.21E-04	1.141E-05
U-238	1.04E+04	3.51E-03	8.904E-05
Np-237	2.21E+02	1.56E-01	4.769E-03
Pu-238	2.59E-02	4.43E-01	1.468E-02
Pu-239	1.88E+01	1.17E+00	3.606E-02
Pu-240	1.72E+00	3.93E-01	1.224E-02
Pu-241	1.22E-01	1.26E+01	3.907E-04
Pu-242	1.99E-02	7.61E-05	2.248E-06
Am-241	8.27E+01	2.84E+02	9.436E+00
Am-242	2.73E-07	2.21E-01	2.509E-04
Am-243	1.90E-01	3.79E-02	1.218E-03
Cm-242	5.50E-05	1.82E-01	6.707E-03
Cm-244	6.22E-02	5.03E+00	1.760E-01
Total	1.26E+04	1.37E+05	3.892E+02

^aThis table identifies the nominal expected activity of HWVP canisters at the time of vitrification. Canister contains 1,650 kg of HLW glass (85% fill). Source: Mitchell and Nelson 1988.

Table 3A.5. Idaho National Engineering Laboratory: radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Se-79	0.1173E+01	0.8173E-01	0.2035E-04
Rb-87	0.5252E+02	0.4597E-05	0.3843E-08
Sr-90	0.1217E+03	0.1660E+05	0.1927E+02
Y-90	0.3051E-01	0.1660E+05	0.9204E+02
Zr-93	0.1575E+03	0.3959E+00	0.4600E-04
Nb-93m	0.3387E-03	0.9577E-01	0.1697E-04
Tc-99	0.1582E+03	0.2682E+01	0.1346E-02
Ru-106	0.3701E+00	0.1239E+04	0.7365E-01
Rh-106	0.3479E-06	0.1239E+04	0.1188E+02
Pd-107	0.4965E+01	0.2554E-02	0.1514E-06
Sn-126	0.1440E+01	0.4086E-01	0.5097E-04
Sb-126m	0.5201E-09	0.4086E-01	0.5203E-03
Sb-126	0.4887E-06	0.4086E-01	0.7552E-03
Cs-134	0.3256E+01	0.4214E+04	0.4290E+02
Cs-135	0.8316E+02	0.9577E-01	0.3197E-04
Cs-137	0.1908E+03	0.1660E+05	0.1837E+02
Ba-137m	0.2848E-04	0.1532E+05	0.6017E+02
Ce-144	0.3282E+01	0.1047E+05	0.6947E+01
Pr-144	0.1386E-03	0.1047E+05	0.7700E+02
Pm-147	0.1653E+02	0.1532E+05	0.5499E+01
Sm-151	0.8250E+01	0.2171E+03	0.2546E-01
Eu-154	0.8513E+00	0.2299E+03	0.2056E+01
U-233	0.1583E-06	0.1532E-08	0.4456E-10
U-234	0.8785E-04	0.5491E-06	0.1582E-07
U-235	0.1063E+01	0.2299E-05	0.6020E-07
U-236	0.1973E+00	0.1277E-04	0.3459E-06
U-237	0.7507E-13	0.6130E-08	0.1158E-10
U-238	0.3797E-04	0.1277E-10	0.3235E-12
Np-237	0.8693E-01	0.6130E-04	0.1874E-05
Pu-238	0.5221E+01	0.8939E+02	0.2963E+01
Pu-239	0.1437E+02	0.8939E+00	0.2754E-01
Pu-240	0.3642E+01	0.8300E+00	0.2585E-01
Pu-241	0.1983E+01	0.2043E+03	0.6336E-02
Pu-242	0.6018E+00	0.2299E-02	0.6788E-04
Am-241	0.3385E+00	0.1162E+01	0.3861E-01
Am-243	0.5315E-01	0.1060E-01	0.3407E-03

Table 3A.5 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Se-79	0.1173E+01	0.8173E-01	0.2035E-04
Cm-242	0.2510E-03	0.8300E+00	0.3059E-01
Cm-244	0.8201E-02	0.6640E+00	0.2322E-01
Total	0.8315E+03	0.1088E+06	0.3393E+03

^aQuantities are at time of filling canister and are based on 3-year-old calcine immobilized in glass-ceramic with a load of 1,277 kg of calcine per canister (1,825 kg of glass-ceramic per canister). Based on IDO 1982 and Berreth 1986c.

TABLE 1-WVDP. MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TH232	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04
U235	1.637E+02	1.637E+02	1.637E+02	1.640E+02	1.666E+02	1.896E+02	2.652E+02	2.711E+02
U238	9.337E+03	9.337E+03	9.337E+03	9.337E+03	9.337E+03	9.337E+03	9.337E+03	9.337E+03
NP237	1.302E+02	1.303E+02	1.312E+02	1.395E+02	1.806E+02	1.928E+02	1.872E+02	1.399E+02
PU239	1.028E+02	1.028E+02	1.028E+02	1.026E+02	1.005E+02	8.055E+01	6.311E+00	3.479E-11
SUM	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04
TOTAL	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 2-WVDP. MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE	0.000E+00	2.220E-03	2.180E-02	1.910E-01	9.919E-01	1.897E+00	3.857E+00	1.017E+01
PB	2.689E-08	1.613E-05	1.708E-04	2.722E-03	2.802E-02	2.596E-01	1.729E+00	8.624E+00
BI	2.529E-09	3.372E-07	3.410E-06	4.003E-05	9.789E-04	5.499E-02	1.914E+00	4.092E+01
TH	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04	5.880E+04
PA	1.264E+00	1.264E+00	1.264E+00	1.261E+00	1.238E+00	1.025E+00	1.624E-01	1.241E-02
U	9.524E+03	9.524E+03	9.524E+03	9.526E+03	9.531E+03	9.566E+03	9.652E+03	9.659E+03
NP	1.330E+02	1.331E+02	1.340E+02	1.423E+02	1.834E+02	1.954E+02	1.888E+02	1.399E+02
PU	1.300E+02	1.298E+02	1.287E+02	1.258E+02	1.210E+02	8.945E+01	7.724E+00	2.817E-01
AM	6.811E+01	6.816E+01	6.829E+01	6.162E+01	1.916E+01	2.667E+00	5.689E-04	0.000E+00
CM	4.026E-01	3.885E-01	2.825E-01	3.040E-02	1.968E-02	9.209E-03	5.784E-06	0.000E+00
SUM	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04
TOTAL	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04	6.866E+04

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 3-WVDP. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

CURIES

NUCLIDE	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	3.220E-02	2.179E-03	1.649E-02	5.704E-02	5.836E-02	4.831E-02	7.659E-03	5.847E-04
TL208	1.270E-02	1.249E-02	1.145E-02	6.895E-03	3.506E-03	3.451E-03	2.976E-03	2.321E-03
TL209	0.000E+00	1.789E-05	1.853E-05	2.486E-05	8.564E-05	5.071E-04	1.454E-03	2.284E-03
PB209	8.250E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
PB210	0.000E+00	1.549E-09	1.435E-07	7.155E-06	1.266E-04	2.112E-03	1.507E-02	5.337E-03
PB211	3.230E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
PB212	3.531E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
PB214	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.338E-03
BI210	0.000E+00	1.549E-09	1.435E-07	7.156E-06	1.266E-04	2.112E-03	1.507E-02	5.337E-03
BI211	3.230E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
BI212	3.530E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
BI213	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
BI214	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.338E-03
PO210	0.000E+00	6.213E-10	1.435E-07	7.156E-06	1.266E-04	2.112E-03	1.507E-02	5.337E-03
PO211	0.000E+00	6.119E-06	4.630E-05	1.602E-04	1.639E-04	1.356E-04	2.150E-05	1.642E-06
PO212	2.261E-02	2.227E-02	2.043E-02	1.230E-02	6.252E-03	6.153E-03	5.307E-03	4.138E-03
PO213	7.861E-04	8.104E-04	8.393E-04	1.126E-03	3.879E-03	2.297E-02	6.586E-02	1.035E-01
PO214	0.000E+00	1.022E-07	1.023E-06	1.037E-05	1.266E-04	2.112E-03	1.507E-02	5.337E-03
PO215	3.229E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
PO216	3.529E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
PO218	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.339E-03
AT217	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RN219	3.230E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
RN220	3.530E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
RN222	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.339E-03
FR221	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
FR223	4.322E-04	3.014E-05	2.280E-04	7.891E-04	8.076E-04	6.685E-04	1.060E-04	8.091E-06
RA223	3.231E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
RA224	3.531E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
RA225	8.252E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RA226	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.339E-03
RA228	5.970E-03	6.017E-03	6.280E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03
AC225	8.250E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
AC227	3.230E-04	2.184E-03	1.652E-02	5.718E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
AC228	5.971E-03	6.018E-03	6.281E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03
TH227	3.179E-02	2.155E-03	1.631E-02	5.641E-02	5.772E-02	4.778E-02	7.574E-03	5.782E-04

TABLE 3-WVDP. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES (CONTINUED)

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TH228	3.531E-02	3.463E-02	3.185E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
TH229	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
TH230	2.361E-04	2.362E-04	2.376E-04	2.540E-04	4.768E-04	2.651E-03	1.494E-02	5.336E-03
TH231	3.540E-04	3.540E-04	3.541E-04	3.546E-04	3.602E-04	4.100E-04	5.734E-04	5.863E-04
TH232	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03
TH234	3.141E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03
PA231	5.973E-02	5.973E-02	5.971E-02	5.960E-02	5.848E-02	4.841E-02	7.675E-03	5.863E-04
PA233	9.181E-02	9.189E-02	9.250E-02	9.835E-02	1.274E-01	1.359E-01	1.320E-01	9.866E-02
PA234M	3.141E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03
U232	2.719E-02	2.696E-02	2.500E-02	1.243E-02	3.305E-03	3.152E-03	1.833E-03	8.074E-06
U233	3.550E-02	3.550E-02	3.550E-02	3.553E-02	3.585E-02	3.969E-02	7.033E-02	1.051E-01
U234	1.650E-02	1.659E-02	1.739E-02	2.296E-02	2.872E-02	2.810E-02	2.248E-02	4.648E-03
U235	3.540E-04	3.540E-04	3.541E-04	3.546E-04	3.602E-04	4.100E-04	5.734E-04	5.863E-04
U236	1.100E-03	1.100E-03	1.102E-03	1.114E-03	1.235E-03	1.979E-03	2.498E-03	2.521E-03
U237	0.000E+00	7.412E-03	4.806E-03	6.321E-05	7.829E-08	3.758E-08	2.438E-11	0.000E+00
U238	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03
NP236	3.720E-02	3.720E-02	3.720E-02	3.718E-02	3.698E-02	3.503E-02	2.036E-02	8.971E-05
NP237	9.182E-02	9.189E-02	9.250E-02	9.835E-02	1.274E-01	1.359E-01	1.320E-01	9.866E-02
NP238	0.000E+00	5.826E-03	5.592E-03	3.710E-03	6.124E-05	9.201E-23	0.000E+00	0.000E+00
NP239	1.360E+00	1.360E+00	1.359E+00	1.347E+00	1.238E+00	5.317E-01	1.134E-04	0.000E+00
PU236	3.300E-03	3.311E-03	3.344E-03	3.346E-03	3.328E-03	3.152E-03	1.833E-03	8.074E-06
PU238	3.261E+01	3.236E+01	3.020E+01	1.521E+01	3.523E-02	3.614E-20	0.000E+00	0.000E+00
PU239	6.393E+00	6.393E+00	6.391E+00	6.379E+00	6.249E+00	5.009E+00	3.924E-01	2.164E-12
PU240	4.680E+00	4.683E+00	4.701E+00	4.711E+00	4.284E+00	1.650E+00	1.183E-04	0.000E+00
PU241	3.170E+02	3.021E+02	1.959E+02	2.577E+00	3.195E-03	1.534E-03	9.952E-07	0.000E+00
PU242	6.371E-03	6.372E-03	6.375E-03	6.399E-03	6.439E-03	6.340E-03	5.397E-03	1.076E-03
AM241	2.100E+02	2.102E+02	2.107E+02	1.881E+02	4.445E+01	1.558E-03	9.951E-07	0.000E+00
AM242M	1.171E+00	1.165E+00	1.118E+00	7.419E-01	1.225E-02	1.840E-20	0.000E+00	0.000E+00
AM242	1.161E+00	1.159E+00	1.113E+00	7.382E-01	1.219E-02	1.831E-20	0.000E+00	0.000E+00
AM243	1.360E+00	1.360E+00	1.359E+00	1.347E+00	1.238E+00	5.317E-01	1.134E-04	0.000E+00
CM242	9.632E-01	9.599E-01	9.207E-01	6.105E-01	1.008E-02	1.519E-20	0.000E+00	0.000E+00
CM243	5.273E-01	5.146E-01	4.134E-01	4.632E-02	1.444E-11	0.000E+00	0.000E+00	0.000E+00
CM244	3.000E+01	2.888E+01	2.046E+01	6.530E-01	7.155E-16	0.000E+00	0.000E+00	0.000E+00
CM245	3.461E-03	3.461E-03	3.458E-03	3.433E-03	3.190E-03	1.531E-03	9.935E-07	0.000E+00
CM246	3.930E-04	3.930E-04	3.925E-04	3.873E-04	3.395E-04	9.081E-05	1.704E-10	0.000E+00
SUM	6.082E+02	5.918E+02	4.754E+02	2.235E+02	5.856E+01	8.859E+00	1.628E+00	1.291E+00
TOTAL	6.082E+02	5.918E+02	4.754E+02	2.235E+02	5.856E+01	8.859E+00	1.628E+00	1.291E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 4-WVDP. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: ELEMENTS

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	4.490E-02	1.469E-02	2.796E-02	6.396E-02	6.195E-02	5.227E-02	1.209E-02	5.190E-03
PB	6.843E-02	3.777E-02	4.927E-02	7.756E-02	7.250E-02	8.575E-02	1.134E-01	1.235E-01
BI	6.843E-02	3.777E-02	4.927E-02	7.756E-02	7.250E-02	8.575E-02	1.134E-01	1.235E-01
PO	9.098E-02	6.003E-02	6.973E-02	9.000E-02	7.896E-02	9.364E-02	1.324E-01	1.307E-01
AT	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RN	6.760E-02	3.694E-02	4.842E-02	7.640E-02	6.841E-02	6.016E-02	3.104E-02	1.238E-02
FR	1.257E-03	8.585E-04	1.086E-03	1.940E-03	4.773E-03	2.415E-02	6.742E-02	1.058E-01
RA	7.441E-02	4.379E-02	5.555E-02	8.400E-02	7.882E-02	9.009E-02	1.048E-01	1.246E-01
AC	7.119E-03	9.031E-03	2.366E-02	6.478E-02	6.894E-02	7.837E-02	8.144E-02	1.128E-01
TH	7.811E-02	4.779E-02	5.920E-02	8.695E-02	8.187E-02	9.351E-02	1.083E-01	1.283E-01
PA	1.547E-01	1.548E-01	1.554E-01	1.611E-01	1.890E-01	1.875E-01	1.429E-01	1.024E-01
U	8.379E-02	9.107E-02	8.730E-02	7.559E-02	7.261E-02	7.647E-02	1.009E-01	1.160E-01
NP	1.489E+00	1.495E+00	1.494E+00	1.487E+00	1.403E+00	7.027E-01	1.525E-01	9.875E-02
PU	3.607E+02	3.456E+02	2.372E+02	2.889E+01	1.058E+01	6.670E+00	3.998E-01	1.084E-03
AM	2.137E+02	2.139E+02	2.143E+02	1.909E+02	4.571E+01	5.333E-01	1.144E-04	0.000E+00
CM	3.150E+01	3.036E+01	2.180E+01	1.314E+00	1.361E-02	1.622E-03	9.936E-07	0.000E+00
SUM	6.082E+02	5.918E+02	4.754E+02	2.235E+02	5.856E+01	8.859E+00	1.628E+00	1.291E+00
TOTAL	6.082E+02	5.918E+02	4.754E+02	2.235E+02	5.856E+01	8.859E+00	1.628E+00	1.291E+00

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 5-WVDP. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	9.455E-05	6.399E-06	4.842E-05	1.675E-04	1.714E-04	1.419E-04	2.249E-05	1.717E-06
TL208	2.989E-04	2.939E-04	2.696E-04	1.623E-04	8.250E-05	8.120E-05	7.004E-05	5.462E-05
TL209	0.000E+00	2.973E-07	3.079E-07	4.131E-07	1.423E-06	8.426E-06	2.416E-05	3.795E-05
PB209	9.488E-07	9.526E-07	9.864E-07	1.324E-06	4.560E-06	2.700E-05	7.741E-05	1.216E-04
PB210	0.000E+00	3.588E-13	3.324E-11	1.658E-09	2.932E-08	4.892E-07	3.492E-06	1.236E-06
PB211	9.679E-05	6.548E-06	4.955E-05	1.714E-04	1.754E-04	1.452E-04	2.301E-05	1.757E-06
PB212	6.722E-05	6.618E-05	6.070E-05	3.654E-05	1.858E-05	1.829E-05	1.577E-05	1.230E-05
PB214	0.000E+00	3.261E-10	3.264E-09	3.307E-08	4.037E-07	6.735E-06	4.808E-05	1.702E-05
BI210	0.000E+00	3.571E-12	3.309E-10	1.650E-08	2.918E-07	4.869E-06	3.475E-05	1.231E-05
BI211	1.289E-03	8.716E-05	6.596E-04	2.282E-03	2.334E-03	1.932E-03	3.063E-04	2.339E-05
BI212	6.004E-04	5.911E-04	5.422E-04	3.264E-04	1.659E-04	1.633E-04	1.409E-04	1.098E-04
BI213	3.469E-06	3.482E-06	3.606E-06	4.839E-06	1.667E-05	9.870E-05	2.830E-04	4.446E-04
BI214	0.000E+00	1.311E-09	1.312E-08	1.329E-07	1.622E-06	2.707E-05	1.932E-04	6.841E-05
PO210	0.000E+00	1.992E-11	4.601E-09	2.294E-07	4.057E-06	6.769E-05	4.832E-04	1.711E-04
PO211	0.000E+00	2.754E-07	2.084E-06	7.208E-06	7.374E-06	6.104E-06	9.678E-07	7.388E-08
PO212	1.198E-03	1.180E-03	1.082E-03	6.516E-04	3.313E-04	3.261E-04	2.813E-04	2.193E-04
PO213	3.978E-05	4.101E-05	4.247E-05	5.700E-05	1.963E-04	1.162E-03	3.333E-03	5.236E-03
PO214	0.000E+00	4.747E-09	4.751E-08	4.813E-07	5.876E-06	9.804E-05	6.998E-04	2.478E-04
PO215	1.442E-03	9.755E-05	7.382E-04	2.554E-03	2.613E-03	2.163E-03	3.429E-04	2.617E-05
PO216	1.445E-03	1.423E-03	1.305E-03	7.856E-04	3.994E-04	3.931E-04	3.391E-04	2.644E-04
PO218	0.000E+00	3.706E-09	3.710E-08	3.758E-07	4.588E-06	7.655E-05	5.464E-04	1.935E-04
AT217	3.521E-05	3.535E-05	3.660E-05	4.912E-05	1.692E-04	1.002E-03	2.872E-03	4.513E-03
RN219	1.340E-03	9.067E-05	6.861E-04	2.373E-03	2.428E-03	2.010E-03	3.187E-04	2.433E-05
RN220	1.340E-03	1.320E-03	1.210E-03	7.286E-04	3.705E-04	3.646E-04	3.145E-04	2.452E-04
RN222	0.000E+00	3.389E-09	3.392E-08	3.437E-07	4.195E-06	7.000E-05	4.996E-04	1.769E-04
FR221	3.184E-05	3.197E-05	3.311E-05	4.443E-05	1.530E-04	9.061E-04	2.598E-03	4.081E-03
RA223	1.150E-03	7.781E-05	5.888E-04	2.037E-03	2.084E-03	1.725E-03	2.735E-04	2.088E-05
RA224	1.212E-03	1.193E-03	1.094E-03	6.586E-04	3.349E-04	3.296E-04	2.843E-04	2.217E-04
RA225	5.787E-07	5.809E-07	6.015E-07	8.072E-07	2.780E-06	1.646E-05	4.720E-05	7.416E-05
RA226	0.000E+00	2.953E-09	2.956E-08	2.995E-07	3.656E-06	6.099E-05	4.354E-04	1.542E-04
RA228	4.601E-07	4.637E-07	4.840E-07	4.971E-07	4.971E-07	4.971E-07	4.971E-07	4.971E-07
AC225	2.882E-05	2.894E-05	2.996E-05	4.021E-05	1.385E-04	8.201E-04	2.351E-03	3.694E-03
AC227	1.564E-07	1.058E-06	7.999E-06	2.769E-05	2.834E-05	2.346E-05	3.719E-06	2.839E-07
AC228	5.160E-05	5.201E-05	5.428E-05	5.576E-05	5.576E-05	5.576E-05	5.576E-05	5.575E-05
TH227	1.160E-03	7.865E-05	5.952E-04	2.059E-03	2.106E-03	1.744E-03	2.764E-04	2.110E-05
TH228	1.155E-03	1.132E-03	1.042E-03	6.276E-04	3.191E-04	3.141E-04	2.709E-04	2.112E-04

TABLE 5-WVDP. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES (CONTINUED)

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TH229	2.524E-05	2.534E-05	2.624E-05	3.522E-05	1.213E-04	7.182E-04	2.059E-03	3.235E-03
TH230	6.680E-06	6.685E-06	6.723E-06	7.187E-06	1.349E-05	7.501E-05	4.229E-04	1.510E-04
TH231	1.987E-07	1.986E-07	1.987E-07	1.990E-07	2.021E-07	2.301E-07	3.217E-07	3.290E-07
TH232	1.562E-04	1.562E-04	1.562E-04	1.562E-04	1.562E-04	1.562E-04	1.562E-04	1.562E-04
TH234	1.274E-06	1.273E-06	1.273E-06	1.273E-06	1.273E-06	1.273E-06	1.273E-06	1.273E-06
PA231	1.800E-03	1.800E-03	1.799E-03	1.796E-03	1.762E-03	1.459E-03	2.313E-04	1.767E-05
PA233	2.084E-04	2.086E-04	2.099E-04	2.232E-04	2.891E-04	3.086E-04	2.997E-04	2.239E-04
PA234M	1.552E-05	1.552E-05	1.552E-05	1.552E-05	1.552E-05	1.552E-05	1.552E-05	1.552E-05
U232	8.730E-04	8.657E-04	8.027E-04	3.991E-04	1.061E-04	1.012E-04	5.883E-05	2.592E-07
U233	1.032E-03	1.032E-03	1.032E-03	1.033E-03	1.042E-03	1.154E-03	2.044E-03	3.056E-03
U234	4.753E-04	4.780E-04	5.009E-04	6.614E-04	8.273E-04	8.092E-04	6.474E-04	1.339E-04
U235	9.270E-06	9.271E-06	9.272E-06	9.287E-06	9.433E-06	1.074E-05	1.502E-05	1.535E-05
U236	2.981E-05	2.981E-05	2.984E-05	3.019E-05	3.346E-05	5.360E-05	6.768E-05	6.830E-05
U238	7.965E-05	7.965E-05	7.965E-05	7.965E-05	7.965E-05	7.965E-05	7.965E-05	7.965E-05
NP236	7.504E-05	7.504E-05	7.504E-05	7.500E-05	7.459E-05	7.065E-05	4.107E-05	1.810E-07
NP237	2.806E-03	2.808E-03	2.827E-03	3.006E-03	3.892E-03	4.155E-03	4.036E-03	3.015E-03
NP239	3.288E-03	3.287E-03	3.285E-03	3.257E-03	2.993E-03	1.285E-03	2.741E-07	0.000E+00
PU236	1.149E-04	1.152E-04	1.164E-04	1.165E-04	1.158E-04	1.097E-04	6.377E-05	2.810E-07
PU238	1.081E+00	1.072E+00	1.001E+00	5.042E-01	1.168E-03	1.198E-21	0.000E+00	0.000E+00
PU239	1.970E-01	1.970E-01	1.970E-01	1.966E-01	1.926E-01	1.544E-01	1.209E-02	6.668E-14
PU240	1.457E-01	1.458E-01	1.464E-01	1.467E-01	1.334E-01	5.137E-02	3.684E-06	0.000E+00
PU241	9.828E-03	9.366E-03	6.073E-03	7.988E-05	9.906E-08	4.755E-08	3.085E-11	0.000E+00
PU242	1.882E-04	1.882E-04	1.883E-04	1.890E-04	1.901E-04	1.872E-04	1.594E-04	3.178E-05
AM241	6.977E+00	6.982E+00	6.999E+00	6.249E+00	1.476E+00	5.177E-05	3.306E-08	0.000E+00
AM242M	4.624E-04	4.603E-04	4.418E-04	2.931E-04	4.838E-06	7.268E-24	0.000E+00	0.000E+00
AM242	1.317E-03	1.316E-03	1.263E-03	8.380E-04	1.383E-05	2.078E-23	0.000E+00	0.000E+00
AM243	4.372E-02	4.372E-02	4.368E-02	4.331E-02	3.980E-02	1.709E-02	3.645E-06	0.000E+00
CM242	3.549E-02	3.537E-02	3.392E-02	2.250E-02	3.713E-04	5.595E-22	0.000E+00	0.000E+00
CM243	1.934E-02	1.888E-02	1.517E-02	1.699E-03	5.299E-13	0.000E+00	0.000E+00	0.000E+00
CM244	1.050E+00	1.010E+00	7.157E-01	2.284E-02	2.503E-17	0.000E+00	0.000E+00	0.000E+00
CM245	1.148E-04	1.148E-04	1.148E-04	1.139E-04	1.059E-04	5.081E-05	3.297E-08	0.000E+00
CM246	1.287E-05	1.287E-05	1.285E-05	1.268E-05	1.111E-05	2.973E-06	5.579E-12	0.000E+00
SUM	9.585E+00	9.536E+00	9.181E+00	7.215E+00	1.870E+00	2.501E-01	4.037E-02	3.096E-02
TOTAL	9.585E+00	9.536E+00	9.181E+00	7.215E+00	1.870E+00	2.501E-01	4.037E-02	3.096E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 6-WVDP. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	3.935E-04	3.006E-04	3.183E-04	3.302E-04	2.553E-04	2.315E-04	1.167E-04	9.428E-05
PB	1.650E-04	7.368E-05	1.112E-04	2.093E-04	1.989E-04	1.977E-04	1.678E-04	1.539E-04
BI	1.892E-03	6.817E-04	1.205E-03	2.613E-03	2.519E-03	2.226E-03	9.581E-04	6.585E-04
PO	4.124E-03	2.742E-03	3.170E-03	4.056E-03	3.561E-03	4.293E-03	6.026E-03	6.358E-03
AT	3.521E-05	3.535E-05	3.660E-05	4.912E-05	1.692E-04	1.002E-03	2.872E-03	4.513E-03
RN	2.681E-03	1.410E-03	1.897E-03	3.102E-03	2.803E-03	2.445E-03	1.133E-03	4.465E-04
FR	3.297E-05	3.205E-05	3.370E-05	4.648E-05	1.551E-04	9.079E-04	2.598E-03	4.081E-03
RA	2.363E-03	1.272E-03	1.684E-03	2.697E-03	2.426E-03	2.133E-03	1.041E-03	4.714E-04
AC	8.058E-05	8.200E-05	9.224E-05	1.237E-04	2.226E-04	8.993E-04	2.411E-03	3.750E-03
TH	2.505E-03	1.401E-03	1.827E-03	2.886E-03	2.718E-03	3.009E-03	3.187E-03	3.776E-03
PA	2.024E-03	2.024E-03	2.025E-03	2.035E-03	2.067E-03	1.783E-03	5.465E-04	2.572E-04
U	2.499E-03	2.508E-03	2.464E-03	2.212E-03	2.098E-03	2.208E-03	2.913E-03	3.354E-03
NP	6.169E-03	6.199E-03	6.214E-03	6.356E-03	6.960E-03	5.511E-03	4.077E-03	3.015E-03
PU	1.434E+00	1.425E+00	1.351E+00	8.478E-01	3.274E-01	2.060E-01	1.232E-02	3.206E-05
AM	7.022E+00	7.027E+00	7.044E+00	6.294E+00	1.516E+00	1.714E-02	3.678E-06	0.000E+00
CM	1.104E+00	1.064E+00	7.650E-01	4.716E-02	4.883E-04	5.378E-05	3.297E-08	0.000E+00
SUM	9.585E+00	9.536E+00	9.181E+00	7.215E+00	1.870E+00	2.501E-01	4.037E-02	3.096E-02
TOTAL	9.585E+00	9.536E+00	9.181E+00	7.215E+00	1.870E+00	2.501E-01	4.037E-02	3.096E-02

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 7-WVDP. ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI211	3.221E-02	2.179E-03	1.649E-02	5.704E-02	5.836E-02	4.831E-02	7.659E-03	5.847E-04
BI212	1.268E-02	1.249E-02	1.145E-02	6.895E-03	3.506E-03	3.451E-03	2.976E-03	2.321E-03
BI213	1.782E-05	1.789E-05	1.853E-05	2.486E-05	8.564E-05	5.071E-04	1.454E-03	2.284E-03
PO210	0.000E+00	6.213E-10	1.435E-07	7.156E-06	1.266E-04	2.112E-03	1.507E-02	5.337E-03
PO211	0.000E+00	6.119E-06	4.630E-05	1.602E-04	1.639E-04	1.356E-04	2.150E-05	1.642E-06
PO212	2.261E-02	2.227E-02	2.043E-02	1.230E-02	6.252E-03	6.153E-03	5.307E-03	4.138E-03
PO213	7.861E-04	8.104E-04	8.393E-04	1.126E-03	3.879E-03	2.297E-02	6.586E-02	1.035E-01
PO214	0.000E+00	1.022E-07	1.023E-06	1.037E-05	1.266E-04	2.112E-03	1.507E-02	5.337E-03
PO215	3.229E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
PO216	3.529E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
PO218	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.338E-03
AT217	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RN219	3.230E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
RN220	3.530E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
RN222	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.339E-03
FR221	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RA223	3.231E-02	2.185E-03	1.654E-02	5.720E-02	5.852E-02	4.844E-02	7.680E-03	5.863E-04
RA224	3.531E-02	3.476E-02	3.188E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
RA226	0.000E+00	1.023E-07	1.024E-06	1.037E-05	1.266E-04	2.112E-03	1.508E-02	5.339E-03
AC225	8.250E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
AC227	4.458E-06	3.014E-05	2.279E-04	7.891E-04	8.076E-04	6.685E-04	1.060E-04	8.091E-06
TH227	3.179E-02	2.155E-03	1.631E-02	5.641E-02	5.772E-02	4.778E-02	7.574E-03	5.782E-04
TH228	3.531E-02	3.463E-02	3.185E-02	1.919E-02	9.758E-03	9.604E-03	8.284E-03	6.459E-03
TH229	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
TH230	2.361E-04	2.362E-04	2.376E-04	2.540E-04	4.768E-04	2.651E-03	1.494E-02	5.336E-03
TH232	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03	6.451E-03
PA231	5.973E-02	5.973E-02	5.971E-02	5.960E-02	5.848E-02	4.841E-02	7.675E-03	5.863E-04
U232	2.719E-02	2.696E-02	2.500E-02	1.243E-02	3.305E-03	3.152E-03	1.833E-03	8.074E-06
U233	3.550E-02	3.550E-02	3.550E-02	3.553E-02	3.585E-02	3.969E-02	7.033E-02	1.051E-01
U234	1.650E-02	1.659E-02	1.739E-02	2.296E-02	2.872E-02	2.810E-02	2.248E-02	4.648E-03
U235	3.540E-04	3.540E-04	3.541E-04	3.546E-04	3.602E-04	4.100E-04	5.734E-04	5.863E-04
U236	1.100E-03	1.100E-03	1.102E-03	1.114E-03	1.235E-03	1.979E-03	2.498E-03	2.521E-03
U238	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03	3.140E-03
NP237	9.182E-02	9.189E-02	9.250E-02	9.835E-02	1.274E-01	1.359E-01	1.320E-01	9.866E-02
PU236	3.300E-03	3.311E-03	3.344E-03	3.346E-03	3.328E-03	3.152E-03	1.833E-03	8.074E-06
PU238	3.261E+01	3.236E+01	3.020E+01	1.521E+01	3.523E-02	3.614E-20	0.000E+00	0.000E+00

TABLE 7-WVP. ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: NUCLIDES (CONTINUED)

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
PU239	6.393E+00	6.393E+00	6.391E+00	6.379E+00	6.249E+00	5.009E+00	3.924E-01	2.164E-12
PU240	4.680E+00	4.683E+00	4.701E+00	4.711E+00	4.284E+00	1.650E+00	1.183E-04	0.000E+00
PU241	7.767E-03	7.402E-03	4.800E-03	6.313E-05	7.829E-08	3.758E-08	2.438E-11	0.000E+00
PU242	6.371E-03	6.372E-03	6.375E-03	6.399E-03	6.439E-03	6.340E-03	5.397E-03	1.076E-03
AM241	2.100E+02	2.102E+02	2.107E+02	1.881E+02	4.445E+01	1.558E-03	9.951E-07	0.000E+00
AM242M	5.853E-03	5.826E-03	5.592E-03	3.710E-03	6.124E-05	9.200E-23	0.000E+00	0.000E+00
AM243	1.360E+00	1.360E+00	1.359E+00	1.347E+00	1.238E+00	5.317E-01	1.134E-04	0.000E+00
CM242	9.632E-01	9.599E-01	9.207E-01	6.105E-01	1.008E-02	1.519E-20	0.000E+00	0.000E+00
CM243	5.260E-01	5.134E-01	4.124E-01	4.621E-02	1.441E-11	0.000E+00	0.000E+00	0.000E+00
CM244	3.000E+01	2.888E+01	2.046E+01	6.530E-01	7.155E-16	0.000E+00	0.000E+00	0.000E+00
CM245	3.461E-03	3.461E-03	3.458E-03	3.433E-03	3.190E-03	1.531E-03	9.935E-07	0.000E+00
CM246	3.929E-04	3.929E-04	3.923E-04	3.872E-04	3.394E-04	9.079E-05	1.704E-10	0.000E+00
SUM	2.872E+02	2.858E+02	2.756E+02	2.177E+02	5.690E+01	7.891E+00	1.154E+00	8.188E-01
TOTAL	2.872E+02	2.858E+02	2.756E+02	2.177E+02	5.690E+01	7.891E+00	1.154E+00	8.188E-01

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 8-WVDP. ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI	4.491E-02	1.469E-02	2.796E-02	6.396E-02	6.195E-02	5.227E-02	1.209E-02	5.191E-03
PO	9.098E-02	6.003E-02	6.973E-02	9.000E-02	7.896E-02	9.364E-02	1.324E-01	1.307E-01
AT	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RN	6.760E-02	3.694E-02	4.842E-02	7.640E-02	6.841E-02	6.016E-02	3.104E-02	1.238E-02
FR	8.251E-04	8.283E-04	8.578E-04	1.151E-03	3.965E-03	2.348E-02	6.731E-02	1.058E-01
RA	6.762E-02	3.694E-02	4.842E-02	7.640E-02	6.841E-02	6.016E-02	3.104E-02	1.238E-02
AC	8.295E-04	8.585E-04	1.086E-03	1.940E-03	4.773E-03	2.415E-02	6.742E-02	1.058E-01
TH	7.461E-02	4.430E-02	5.571E-02	8.346E-02	7.837E-02	8.996E-02	1.046E-01	1.246E-01
PA	5.973E-02	5.973E-02	5.971E-02	5.960E-02	5.848E-02	4.841E-02	7.675E-03	5.863E-04
U	8.379E-02	8.366E-02	8.250E-02	7.553E-02	7.261E-02	7.647E-02	1.009E-01	1.160E-01
NP	9.182E-02	9.189E-02	9.250E-02	9.835E-02	1.274E-01	1.359E-01	1.320E-01	9.866E-02
PU	4.370E+01	4.345E+01	4.131E+01	2.631E+01	1.058E+01	6.668E+00	3.998E-01	1.084E-03
AM	2.114E+02	2.115E+02	2.120E+02	1.895E+02	4.569E+01	5.333E-01	1.144E-04	0.000E+00
CM	3.150E+01	3.035E+01	2.180E+01	1.314E+00	1.361E-02	1.622E-03	9.936E-07	0.000E+00
SUM	2.872E+02	2.858E+02	2.756E+02	2.177E+02	5.690E+01	7.891E+00	1.154E+00	8.188E-01
TOTAL	2.872E+02	2.858E+02	2.756E+02	2.177E+02	5.690E+01	7.891E+00	1.154E+00	8.188E-01

TABLE 9-WVDP. (ALPHA,N) NEUTRON SOURCES IN DECAY OF WVDP HLW
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

		NEUTRONS/SEC						
	IMMOBLZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KYR	100.0KYR	1.0MYR
PO210	0.000E-01	2.118E-05	4.891E-03	2.439E-01	4.315E+00	7.199E+01	5.137E+02	1.819E+02
PO213	1.209E+02	1.247E+02	1.291E+02	1.732E+02	5.968E+02	3.534E+03	1.013E+04	1.592E+04
PO214	0.000E-01	1.244E-02	1.246E-01	1.263E+00	1.542E+01	2.572E+02	1.835E+03	6.499E+02
PO218	0.000E-01	5.343E-03	5.348E-02	5.416E-01	6.612E+00	1.103E+02	7.876E+02	2.788E+02
AT217	7.591E+01	7.620E+01	7.892E+01	1.059E+02	3.648E+02	2.160E+03	6.192E+03	9.733E+03
RN222	0.000E-01	3.924E-03	3.928E-02	3.978E-01	4.856E+00	8.101E+01	5.784E+02	2.048E+02
FR221	5.281E+01	5.302E+01	5.491E+01	7.367E+01	2.538E+02	1.503E+03	4.308E+03	6.772E+03
RA226	0.000E-01	2.315E-03	2.317E-02	2.346E-01	2.864E+00	4.778E+01	3.412E+02	1.208E+02
AC225	3.747E+01	3.761E+01	3.895E+01	5.227E+01	1.801E+02	1.066E+03	3.057E+03	4.805E+03
TH229	2.009E+01	2.017E+01	2.088E+01	2.802E+01	9.653E+01	5.716E+02	1.639E+03	2.576E+03
TH230	4.838E+00	4.840E+00	4.869E+00	5.205E+00	9.770E+00	5.432E+01	3.061E+02	1.093E+02
U233	8.309E+02	8.309E+02	8.309E+02	8.316E+02	8.391E+02	9.290E+02	1.646E+03	2.460E+03
U234	3.730E+02	3.750E+02	3.931E+02	5.190E+02	6.493E+02	6.352E+02	5.082E+02	1.051E+02
U238	3.981E+01	3.981E+01	3.981E+01	3.981E+01	3.981E+01	3.981E+01	3.981E+01	3.981E+01
NP237	2.052E+03	2.054E+03	2.067E+03	2.198E+03	2.847E+03	3.037E+03	2.950E+03	2.205E+03
PU238	1.259E+06	1.249E+06	1.166E+06	5.873E+05	1.360E+03	1.395E-15	0.000E-01	0.000E-01
PU239	1.951E+05	1.951E+05	1.950E+05	1.947E+05	1.907E+05	1.529E+05	1.197E+04	6.604E-08
PU240	1.434E+05	1.435E+05	1.440E+05	1.443E+05	1.312E+05	5.055E+04	3.624E+00	0.000E-01
PU242	1.586E+02	1.586E+02	1.587E+02	1.593E+02	1.603E+02	1.578E+02	1.344E+02	2.679E+01
AM241	8.011E+06	8.018E+06	8.037E+06	7.175E+06	1.696E+06	5.943E+01	3.796E-02	0.000E-01
CM242	5.347E+04	5.329E+04	5.111E+04	3.389E+04	5.596E+02	8.432E-16	0.000E-01	0.000E-01
CM244	1.403E+06	1.351E+06	9.569E+05	3.054E+04	3.346E-11	0.000E-01	0.000E-01	0.000E-01
CM246	1.405E+01	1.405E+01	1.402E+01	1.384E+01	1.213E+01	3.246E+00	6.091E-06	0.000E-01
TOTAL	1.107E+07	1.101E+07	1.055E+07	8.170E+06	2.026E+06	2.177E+05	4.695E+04	4.619E+04

TABLE 10-WVDP. SPONTANEOUS FISSION NEUTRON SOURCES IN DECAY OF WEST VALLEY HLW
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	NEUTRONS/SEC							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	1.185E+02	1.185E+02	1.185E+02	1.185E+02	1.185E+02	1.185E+02	1.185E+02	1.185E+02
PU238	5.059E+03	5.021E+03	4.686E+03	2.360E+03	5.466E+00	5.607E-18	0.000E+00	0.000E+00
PU240	1.869E+04	1.870E+04	1.878E+04	1.882E+04	1.711E+04	6.590E+03	4.726E-01	0.000E+00
PU242	2.812E+03	2.812E+03	2.813E+03	2.824E+03	2.842E+03	2.798E+03	2.382E+03	4.749E+02
CM242	6.274E+03	6.253E+03	5.997E+03	3.977E+03	6.565E+01	9.892E-17	0.000E+00	0.000E+00
CM244	4.122E+06	3.968E+06	2.811E+06	8.972E+04	9.831E-11	0.000E+00	0.000E+00	0.000E+00
CM246	1.138E+04	1.138E+04	1.137E+04	1.122E+04	9.831E+03	2.630E+03	4.935E-03	0.000E+00
SUM	4.167E+06	4.012E+06	2.855E+06	1.291E+05	3.001E+04	1.215E+04	2.501E+03	5.937E+02
TOTAL	4.167E+06	4.012E+06	2.855E+06	1.291E+05	3.001E+04	1.215E+04	2.501E+03	5.937E+02

TABLE 11-WVDP. PHOTON SPECTRUM OF ACTINIDES AND DAUGHTERS IN DECAY OF WEST VALLEY HLW
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

PHOTONS/SEC								
EMEAN	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	2.257E+12	2.249E+12	2.188E+12	1.776E+12	4.694E+11	5.888E+10	1.667E+10	1.425E+10
2.500E-02	1.987E+11	1.987E+11	1.992E+11	1.781E+11	4.328E+10	1.394E+09	1.007E+09	8.588E+08
3.750E-02	2.134E+10	2.125E+10	2.117E+10	1.888E+10	7.084E+09	2.072E+09	1.292E+09	1.885E+09
5.750E-02	2.892E+12	2.894E+12	2.901E+12	2.591E+12	6.133E+11	1.069E+09	4.742E+08	4.515E+08
8.500E-02	4.914E+10	4.831E+10	4.789E+10	4.532E+10	3.986E+10	2.033E+10	5.522E+09	5.210E+09
1.250E-01	3.607E+10	3.590E+10	3.467E+10	2.928E+10	2.357E+10	1.059E+10	1.022E+09	8.950E+08
2.250E-01	2.517E+10	2.432E+10	2.336E+10	1.950E+10	1.735E+10	8.656E+09	1.627E+09	1.467E+09
3.750E-01	3.834E+09	3.401E+09	3.611E+09	4.236E+09	4.495E+09	3.861E+09	3.184E+09	2.897E+09
5.750E-01	6.060E+08	5.886E+08	5.526E+08	3.770E+08	2.069E+08	2.415E+08	5.015E+08	3.069E+08
8.500E-01	4.656E+08	4.919E+08	4.953E+08	4.431E+08	3.261E+08	3.189E+08	3.211E+08	2.563E+08
1.250E+00	2.951E+07	7.305E+07	7.043E+07	5.128E+07	2.123E+07	4.786E+07	2.152E+08	9.590E+07
1.750E+00	6.451E+07	6.453E+07	6.140E+07	4.545E+07	3.680E+07	7.078E+07	2.327E+08	1.584E+08
2.250E+00	8.248E+05	7.947E+05	5.702E+05	6.304E+04	4.063E+05	6.651E+06	4.746E+07	1.680E+07
2.750E+00	4.464E+08	4.388E+08	4.024E+08	2.421E+08	1.231E+08	1.212E+08	1.053E+08	8.176E+07
3.500E+00	4.297E+05	4.139E+05	2.952E+05	1.537E+04	5.240E+03	2.327E+04	1.554E+05	5.497E+04
5.000E+00	1.836E+05	1.768E+05	1.261E+05	6.352E+03	1.641E+03	6.573E+02	1.438E+02	3.558E+01
7.000E+00	2.116E+04	2.037E+04	1.452E+04	7.111E+02	1.836E+02	7.495E+01	1.645E+01	4.018E+00
9.500E+00	2.429E+03	2.339E+03	1.666E+03	8.034E+01	2.075E+01	8.573E+00	1.886E+00	4.571E-01
TOTAL	5.485E+12	5.477E+12	5.420E+12	4.663E+12	1.219E+12	1.077E+11	3.222E+10	2.883E+10

TABLE 12-WVDP. MASS OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR 90	1.928E+02	1.883E+02	1.520E+02	1.784E+01	8.872E-09	0.000E+00	0.000E+00	0.000E+00
ZR 90	0.000E+00	4.536E+00	4.085E+01	1.750E+02	1.928E+02	1.928E+02	1.928E+02	1.928E+02
ZR 93	4.257E+02	4.257E+02	4.257E+02	4.257E+02	4.255E+02	4.238E+02	4.068E+02	2.706E+02
NB 93	0.000E+00	1.399E-04	1.503E-03	1.823E-02	1.918E-01	1.923E+00	1.886E+01	1.551E+02
TC 99	2.524E+01	2.524E+01	2.524E+01	2.523E+01	2.516E+01	2.443E+01	1.823E+01	9.747E-01
RU 99	0.000E+00	8.213E-05	8.213E-04	8.212E-03	8.200E-02	8.081E-01	7.011E+00	2.427E+01
PD107	8.416E+01	8.416E+01	8.416E+01	8.416E+01	8.415E+01	8.407E+01	8.327E+01	7.564E+01
AG107	0.000E+00	8.980E-06	8.980E-05	8.980E-04	8.980E-03	8.975E-02	8.932E-01	8.518E+00
SN126	1.441E+01	1.441E+01	1.441E+01	1.440E+01	1.431E+01	1.345E+01	7.205E+00	1.408E-02
TE126	0.000E+00	9.987E-05	9.987E-04	9.984E-03	9.953E-02	9.649E-01	7.205E+00	1.440E+01
CS135	5.505E+02	5.505E+02	5.505E+02	5.505E+02	5.503E+02	5.488E+02	5.342E+02	4.073E+02
BA135	0.000E+00	1.659E-04	1.659E-03	1.659E-02	1.659E-01	1.657E+00	1.634E+01	1.432E+02
CS137	3.252E+02	3.178E+02	2.581E+02	3.226E+01	3.003E-08	0.000E+00	0.000E+00	0.000E+00
BA137	0.000E+00	7.428E+00	6.709E+01	2.929E+02	3.252E+02	3.252E+02	3.252E+02	3.252E+02
SM151	1.258E+01	1.248E+01	1.165E+01	5.823E+00	5.685E-03	4.464E-33	0.000E+00	0.000E+00
EU151	0.000E+00	9.652E-02	9.326E-01	6.757E+00	1.257E+01	1.258E+01	1.258E+01	1.258E+01
SUM	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03
TOTAL	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 13-WVDP. MASS OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	1.928E+02	1.883E+02	1.520E+02	1.784E+01	8.872E-09	0.000E+00	0.000E+00	0.000E+00
ZR	4.257E+02	4.302E+02	4.665E+02	6.007E+02	6.184E+02	6.166E+02	5.997E+02	4.635E+02
NB	2.529E-03	2.722E-03	4.458E-03	2.182E-02	1.954E-01	1.927E+00	1.886E+01	1.551E+02
TC	2.524E+01	2.524E+01	2.524E+01	2.523E+01	2.516E+01	2.443E+01	1.823E+01	9.747E-01
RU	1.655E-05	9.045E-05	8.213E-04	8.212E-03	8.200E-02	8.081E-01	7.011E+00	2.427E+01
PD	8.416E+01	8.416E+01	8.416E+01	8.416E+01	8.415E+01	8.407E+01	8.327E+01	7.564E+01
AG	0.000E+00	8.980E-06	8.980E-05	8.980E-04	8.980E-03	8.975E-02	8.932E-01	8.518E+00
SN	1.441E+01	1.441E+01	1.441E+01	1.440E+01	1.431E+01	1.345E+01	7.205E+00	1.408E-02
TE	3.885E-04	6.619E-03	2.681E-02	3.806E-02	1.276E-01	9.930E-01	7.233E+00	1.442E+01
CS	8.757E+02	8.683E+02	8.086E+02	5.827E+02	5.503E+02	5.488E+02	5.342E+02	4.073E+02
BA	4.981E-05	7.432E+00	6.711E+01	2.930E+02	3.254E+02	3.269E+02	3.416E+02	4.685E+02
PM	3.722E-01	2.858E-01	2.652E-02	3.225E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	1.258E+01	1.257E+01	1.200E+01	6.202E+00	3.838E-01	3.781E-01	3.781E-01	3.781E-01
EU	1.599E+00	1.561E+00	1.608E+00	6.757E+00	1.257E+01	1.258E+01	1.258E+01	1.258E+01
GD	0.000E+00	1.339E-01	9.211E-01	1.592E+00	1.593E+00	1.593E+00	1.593E+00	1.593E+00
SUM	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03
TOTAL	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03	1.633E+03

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 14-WVDP. RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR 90	2.631E+04	2.569E+04	2.074E+04	2.434E+03	1.211E-06	0.000E+00	0.000E+00	0.000E+00
Y 90	2.631E+04	2.570E+04	2.074E+04	2.435E+03	1.211E-06	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.070E+00	1.070E+00	1.070E+00	1.070E+00	1.070E+00	1.065E+00	1.023E+00	6.802E-01
NB 93M	7.151E-01	7.301E-01	8.355E-01	1.015E+00	1.016E+00	1.012E+00	9.715E-01	6.462E-01
TC 99	4.281E-01	4.281E-01	4.281E-01	4.279E-01	4.267E-01	4.144E-01	3.092E-01	1.653E-02
PD107	4.330E-02	4.330E-02	4.330E-02	4.330E-02	4.330E-02	4.326E-02	4.284E-02	3.892E-02
CD113M	8.341E+00	7.954E+00	5.187E+00	7.209E-02	1.939E-20	0.000E+00	0.000E+00	0.000E+00
SB125	2.860E+01	2.227E+01	2.342E+00	3.874E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	7.000E+00	5.434E+00	5.714E-01	9.453E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	4.090E-01	4.090E-01	4.090E-01	4.087E-01	4.062E-01	3.816E-01	2.045E-01	3.996E-04
SB126	5.731E-02	5.726E-02	5.726E-02	5.722E-02	5.686E-02	5.342E-02	2.863E-02	5.595E-05
SB126M	4.091E-01	4.090E-01	4.090E-01	4.087E-01	4.062E-01	3.816E-01	2.045E-01	3.996E-04
CS134	2.031E+01	1.451E+01	7.043E-01	5.086E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	6.341E-01	6.341E-01	6.341E-01	6.341E-01	6.339E-01	6.322E-01	6.153E-01	4.691E-01
CS137	2.830E+04	2.765E+04	2.246E+04	2.807E+03	2.613E-06	0.000E+00	0.000E+00	0.000E+00
BA137M	2.680E+04	2.616E+04	2.125E+04	2.656E+03	2.472E-06	0.000E+00	0.000E+00	0.000E+00
PM147	3.451E+02	2.650E+02	2.457E+01	1.157E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM151	3.311E+02	3.285E+02	3.065E+02	1.533E+02	1.496E-01	0.000E+00	0.000E+00	0.000E+00
EU152	1.430E+00	1.359E+00	8.591E-01	8.751E-03	1.051E-22	0.000E+00	0.000E+00	0.000E+00
EU154	3.751E+02	3.460E+02	1.675E+02	1.185E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
EU155	9.371E+01	8.148E+01	2.316E+01	7.973E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	1.089E+05	1.063E+05	8.572E+04	1.049E+04	4.208E+00	3.984E+00	3.399E+00	1.852E+00
TOTAL	1.089E+05	1.063E+05	8.572E+04	1.049E+04	4.208E+00	3.984E+00	3.399E+00	1.852E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 15-WVDP. RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	2.631E+04	2.569E+04	2.074E+04	2.434E+03	1.211E-06	0.000E+00	0.000E+00	0.000E+00
Y	2.631E+04	2.570E+04	2.074E+04	2.435E+03	1.211E-06	0.000E+00	0.000E+00	0.000E+00
ZR	1.070E+00	1.070E+00	1.070E+00	1.070E+00	1.070E+00	1.065E+00	1.023E+00	6.802E-01
NB	7.151E-01	7.301E-01	8.355E-01	1.015E+00	1.016E+00	1.012E+00	9.715E-01	6.462E-01
TC	4.281E-01	4.281E-01	4.281E-01	4.279E-01	4.267E-01	4.144E-01	3.092E-01	1.653E-02
PD	4.330E-02	4.330E-02	4.330E-02	4.330E-02	4.330E-02	4.326E-02	4.284E-02	3.892E-02
SN	4.776E-01	4.767E-01	4.687E-01	4.258E-01	4.062E-01	3.816E-01	2.045E-01	3.996E-04
SB	2.907E+01	2.274E+01	2.808E+00	4.659E-01	4.630E-01	4.350E-01	2.331E-01	4.556E-04
CS	2.832E+04	2.767E+04	2.246E+04	2.808E+03	6.339E-01	6.322E-01	6.153E-01	4.691E-01
BA	2.680E+04	2.616E+04	2.125E+04	2.656E+03	2.472E-06	0.000E+00	0.000E+00	0.000E+00
PM	3.451E+02	2.650E+02	2.459E+01	1.442E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	3.311E+02	3.285E+02	3.065E+02	1.533E+02	1.496E-01	9.699E-09	9.697E-09	9.686E-09
EU	4.702E+02	4.289E+02	1.916E+02	1.274E-01	1.051E-22	0.000E+00	0.000E+00	0.000E+00
SUM	1.089E+05	1.063E+05	8.572E+04	1.049E+04	4.208E+00	3.984E+00	3.399E+00	1.852E+00
TOTAL	1.089E+05	1.063E+05	8.572E+04	1.049E+04	4.208E+00	3.984E+00	3.399E+00	1.852E+00

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 16-WVDP. THERMAL POWER OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR 90	3.054E+01	2.982E+01	2.407E+01	2.826E+00	1.405E-09	0.000E+00	0.000E+00	0.000E+00
Y 90	1.458E+02	1.424E+02	1.150E+02	1.350E+01	6.712E-09	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.243E-04	1.243E-04	1.243E-04	1.243E-04	1.243E-04	1.238E-04	1.188E-04	7.903E-05
NB 93M	1.267E-04	1.294E-04	1.480E-04	1.798E-04	1.800E-04	1.793E-04	1.721E-04	1.145E-04
TC 99	2.147E-04	2.147E-04	2.147E-04	2.146E-04	2.140E-04	2.078E-04	1.550E-04	8.290E-06
PD107	2.567E-06	2.567E-06	2.567E-06	2.567E-06	2.567E-06	2.564E-06	2.540E-06	2.307E-06
CD113M	1.404E-02	1.339E-02	8.732E-03	1.214E-04	3.264E-23	0.000E+00	0.000E+00	0.000E+00
SB125	8.942E-02	6.962E-02	7.322E-03	1.211E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	5.884E-03	4.568E-03	4.803E-04	7.946E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	5.101E-04	5.101E-04	5.101E-04	5.097E-04	5.066E-04	4.759E-04	2.551E-04	4.984E-07
SB126	1.059E-03	1.058E-03	1.058E-03	1.057E-03	1.051E-03	9.871E-04	5.290E-04	1.034E-06
SB126M	5.208E-03	5.208E-03	5.207E-03	5.204E-03	5.172E-03	4.859E-03	2.604E-03	5.088E-06
CS134	2.067E-01	1.477E-01	7.168E-03	5.176E-16	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	2.116E-04	2.116E-04	2.116E-04	2.116E-04	2.116E-04	2.110E-04	2.053E-04	1.566E-04
CS137	3.130E+01	3.059E+01	2.484E+01	3.105E+00	2.890E-09	0.000E+00	0.000E+00	0.000E+00
BA137M	1.052E+02	1.027E+02	8.343E+01	1.043E+01	9.706E-09	0.000E+00	0.000E+00	0.000E+00
PM147	1.238E-01	9.504E-02	8.814E-03	4.149E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM151	3.882E-02	3.852E-02	3.594E-02	1.797E-02	1.754E-05	1.378E-35	0.000E+00	0.000E+00
EU152	1.082E-02	1.028E-02	6.498E-03	6.619E-05	7.952E-25	0.000E+00	0.000E+00	0.000E+00
EU154	3.355E+00	3.095E+00	1.499E+00	1.060E-03	3.333E-35	0.000E+00	0.000E+00	0.000E+00
EU155	6.816E-02	5.927E-02	1.685E-02	5.799E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	3.168E+02	3.091E+02	2.489E+02	2.988E+01	7.479E-03	7.046E-03	4.042E-03	3.673E-04
TOTAL	3.168E+02	3.091E+02	2.489E+02	2.988E+01	7.479E-03	7.046E-03	4.042E-03	3.673E-04

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 17-WVDP. THERMAL POWER OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	3.054E+01	2.982E+01	2.407E+01	2.826E+00	1.405E-09	0.000E+00	0.000E+00	0.000E+00
Y	1.458E+02	1.424E+02	1.150E+02	1.350E+01	6.712E-09	0.000E+00	0.000E+00	0.000E+00
ZR	1.243E-04	1.243E-04	1.243E-04	1.243E-04	1.243E-04	1.238E-04	1.188E-04	7.903E-05
NB	1.267E-04	1.294E-04	1.480E-04	1.798E-04	1.800E-04	1.793E-04	1.721E-04	1.145E-04
TC	2.147E-04	2.147E-04	2.147E-04	2.146E-04	2.140E-04	2.078E-04	1.550E-04	8.290E-06
PD	2.567E-06	2.567E-06	2.567E-06	2.567E-06	2.567E-06	2.564E-06	2.540E-06	2.307E-06
SN	6.476E-04	6.457E-04	6.297E-04	5.441E-04	5.066E-04	4.759E-04	2.551E-04	4.984E-07
SB	9.569E-02	7.589E-02	1.359E-02	6.261E-03	6.222E-03	5.846E-03	3.133E-03	6.122E-06
CS	3.151E+01	3.073E+01	2.485E+01	3.106E+00	2.116E-04	2.110E-04	2.053E-04	1.566E-04
BA	1.052E+02	1.027E+02	8.343E+01	1.043E+01	9.706E-09	0.000E+00	0.000E+00	0.000E+00
PM	1.240E-01	9.523E-02	8.875E-03	7.221E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	3.882E-02	3.852E-02	3.594E-02	1.797E-02	1.754E-05	1.345E-10	1.345E-10	1.343E-10
EU	3.434E+00	3.165E+00	1.522E+00	1.126E-03	7.952E-25	0.000E+00	0.000E+00	0.000E+00
SUM	3.168E+02	3.091E+02	2.489E+02	2.988E+01	7.479E-03	7.046E-03	4.042E-03	3.673E-04
TOTAL	3.168E+02	3.091E+02	2.489E+02	2.988E+01	7.479E-03	7.046E-03	4.042E-03	3.673E-04

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 18-WVDP. PHOTON SPECTRUM OF FISSION PRODUCTS IN DECAY OF WEST VALLEY HLW
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

EMEAN	PHOTONS/SEC							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	7.705E+14	7.524E+14	6.066E+14	7.177E+13	1.521E+10	1.473E+10	1.145E+10	5.222E+09
2.500E-02	1.597E+14	1.559E+14	1.253E+14	1.482E+13	7.428E+09	6.985E+09	3.820E+09	9.836E+07
3.750E-02	1.720E+14	1.677E+14	1.342E+14	1.601E+13	1.002E+09	9.465E+08	5.457E+08	4.465E+07
5.750E-02	1.480E+14	1.445E+14	1.163E+14	1.370E+13	2.800E+09	2.636E+09	1.449E+09	4.162E+07
8.500E-02	8.875E+13	8.656E+13	6.935E+13	8.152E+12	7.758E+09	7.290E+09	3.918E+09	1.780E+07
1.250E-01	6.258E+13	6.075E+13	4.709E+13	5.240E+12	4.136E+08	3.891E+08	2.115E+08	2.536E+06
2.250E-01	7.523E+13	7.341E+13	5.889E+13	6.871E+12	7.322E+08	6.880E+08	3.692E+08	8.884E+05
3.750E-01	3.237E+13	3.154E+13	2.519E+13	2.961E+12	1.647E+10	1.548E+10	8.294E+09	1.621E+07
5.750E-01	1.040E+15	1.015E+15	8.233E+14	1.028E+14	3.657E+10	3.436E+10	1.841E+10	3.598E+07
8.500E-01	1.242E+13	1.160E+13	7.033E+12	4.793E+11	1.831E+09	1.720E+09	9.217E+08	1.801E+06
1.250E+00	9.066E+12	8.447E+12	4.611E+12	1.587E+11	4.443E+08	4.175E+08	2.237E+08	4.372E+05
1.750E+00	3.507E+11	3.307E+11	2.016E+11	1.223E+10	1.895E+04	1.780E+04	9.539E+03	1.864E+01
2.250E+00	1.823E+07	1.597E+07	1.143E+07	1.342E+06	6.676E-04	1.493E-07	1.493E-07	1.493E-07
2.750E+00	4.413E+05	2.218E+05	4.546E+02	7.479E-08	7.479E-08	7.479E-08	7.479E-08	7.479E-08
3.500E+00	5.759E+04	2.895E+04	5.941E+01	5.508E-08	5.508E-08	5.508E-08	5.508E-08	5.508E-08
5.000E+00	0.000E+00	3.807E-09	1.523E-08	1.640E-08	1.640E-08	1.640E-08	1.640E-08	1.640E-08
7.000E+00	0.000E+00	2.470E-10	9.882E-10	1.064E-09	1.064E-09	1.064E-09	1.064E-09	1.064E-09
9.500E+00	0.000E+00	1.562E-11	6.249E-11	6.728E-11	6.728E-11	6.728E-11	6.728E-11	6.728E-11
TOTAL	2.572E+15	2.508E+15	2.018E+15	2.429E+14	9.067E+10	8.564E+10	4.962E+10	5.482E+09

TABLE 19-WVDP. MASS OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES

(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO 59	0.000E+00	4.758E-05	4.757E-04	4.756E-03	4.737E-02	4.557E-01	3.182E+00	5.490E+00
NI 59	5.491E+00	5.491E+00	5.491E+00	5.486E+00	5.444E+00	5.035E+00	2.309E+00	9.479E-04
NI 63	4.895E-01	4.858E-01	4.540E-01	2.304E-01	2.616E-04	9.313E-34	0.000E+00	0.000E+00
CU 63	0.000E+00	3.674E-03	3.552E-02	2.591E-01	4.892E-01	4.895E-01	4.895E-01	4.895E-01
SE 79	1.980E-01	1.980E-01	1.980E-01	1.978E-01	1.959E-01	1.780E-01	6.812E-02	4.600E-06
BR 79	0.000E+00	2.113E-06	2.113E-05	2.112E-04	2.101E-03	2.004E-02	1.299E-01	1.980E-01
SUM	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00
TOTAL	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 20-WVDP. MASS OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	GRAMS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
MN	0.000E+00	2.584E-04	1.027E-03	1.104E-03	1.104E-03	1.104E-03	1.104E-03	1.104E-03
FE	1.104E-03	8.456E-04	7.676E-05	2.916E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	2.679E-03	2.396E-03	1.195E-03	4.756E-03	4.737E-02	4.557E-01	3.182E+00	5.490E+00
NI	5.981E+00	5.977E+00	5.946E+00	5.719E+00	5.447E+00	5.038E+00	2.311E+00	3.627E-03
CU	0.000E+00	3.674E-03	3.552E-02	2.591E-01	4.892E-01	4.895E-01	4.895E-01	4.895E-01
SE	1.980E-01	1.980E-01	1.980E-01	1.978E-01	1.959E-01	1.780E-01	6.812E-02	4.600E-06
BR	0.000E+00	2.113E-06	2.113E-05	2.112E-04	2.101E-03	2.004E-02	1.299E-01	1.980E-01
SUM	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00
TOTAL	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00	6.182E+00

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 21-WVDP. RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	2.761E+00	2.115E+00	1.920E-01	7.293E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	3.030E+00	2.657E+00	8.133E-01	5.868E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	4.161E-01	4.161E-01	4.160E-01	4.157E-01	4.125E-01	3.815E-01	1.749E-01	7.183E-05
NI 63	3.021E+01	2.998E+01	2.801E+01	1.422E+01	1.614E-02	0.000E+00	0.000E+00	0.000E+00
SE 79	1.380E-02	1.380E-02	1.380E-02	1.378E-02	1.365E-02	1.240E-02	4.747E-03	3.206E-07
SUM	3.643E+01	3.518E+01	2.945E+01	1.465E+01	4.423E-01	3.939E-01	1.797E-01	7.215E-05
TOTAL	3.643E+01	3.518E+01	2.945E+01	1.465E+01	4.423E-01	3.939E-01	1.797E-01	7.215E-05

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 22-WVDP. RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	CURIES							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	2.761E+00	2.115E+00	1.920E-01	7.293E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	3.030E+00	2.657E+00	8.133E-01	5.868E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	3.062E+01	3.039E+01	2.843E+01	1.463E+01	4.286E-01	3.815E-01	1.749E-01	7.183E-05
SE	1.380E-02	1.380E-02	1.380E-02	1.378E-02	1.365E-02	1.240E-02	4.747E-03	3.206E-07
SUM	3.643E+01	3.518E+01	2.945E+01	1.465E+01	4.423E-01	3.939E-01	1.797E-01	7.215E-05
TOTAL	3.643E+01	3.518E+01	2.945E+01	1.465E+01	4.423E-01	3.939E-01	1.797E-01	7.215E-05

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 23-WVDP. THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: NUCLIDES
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

NUCLIDE	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	9.329E-05	7.146E-05	6.487E-06	2.464E-16	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	4.672E-02	4.096E-02	1.254E-02	9.047E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	1.652E-05	1.652E-05	1.652E-05	1.651E-05	1.638E-05	1.515E-05	6.948E-06	2.853E-09
NI 63	3.044E-03	3.021E-03	2.823E-03	1.433E-03	1.627E-06	5.791E-36	0.000E+00	0.000E+00
SE 79	3.435E-06	3.435E-06	3.435E-06	3.432E-06	3.399E-06	3.088E-06	1.182E-06	7.981E-11
SUM	4.988E-02	4.407E-02	1.539E-02	1.453E-03	2.141E-05	1.824E-05	8.129E-06	2.932E-09
TOTAL	4.988E-02	4.407E-02	1.539E-02	1.453E-03	2.141E-05	1.824E-05	8.129E-06	2.932E-09

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 24-WVDP. THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW: ELEMENTS
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

ELEMENT	WATTS							
	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	9.329E-05	7.146E-05	6.487E-06	2.464E-16	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	4.672E-02	4.096E-02	1.254E-02	9.047E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	3.060E-03	3.037E-03	2.839E-03	1.449E-03	1.801E-05	1.515E-05	6.948E-06	2.853E-09
SE	3.435E-06	3.435E-06	3.435E-06	3.432E-06	3.399E-06	3.088E-06	1.182E-06	7.981E-11
SUM	4.988E-02	4.407E-02	1.539E-02	1.453E-03	2.141E-05	1.824E-05	8.129E-06	2.932E-09
TOTAL	4.988E-02	4.407E-02	1.539E-02	1.453E-03	2.141E-05	1.824E-05	8.129E-06	2.932E-09

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 25-WVDP. PHOTON SPECTRUM OF ACTIVATION PRODUCTS IN DECAY OF WEST VALLEY HLW
(BASED ON ONE CANISTER, END OF 1989 DATA, 1900 KG GLASS)

PHOTONS/SEC								
EMEAN	JAN 1990	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	1.213E+10	1.135E+10	7.339E+09	2.923E+09	1.714E+07	1.257E+07	4.811E+06	3.249E+02
2.500E-02	1.405E+09	1.276E+09	6.268E+08	1.793E+08	2.246E+06	1.858E+06	7.112E+05	4.803E+01
3.750E-02	6.636E+08	5.907E+08	2.291E+08	3.695E+07	1.061E+06	9.267E+05	3.547E+05	2.395E+01
5.750E-02	6.699E+08	5.884E+08	1.855E+08	4.502E+06	8.626E+05	7.799E+05	2.985E+05	2.016E+01
8.500E-02	2.602E+08	2.282E+08	6.996E+07	1.632E+05	1.611E+05	1.463E+05	5.602E+04	3.783E+00
1.250E-01	9.989E+07	8.758E+07	2.682E+07	1.452E+04	1.419E+04	1.289E+04	4.934E+03	3.332E-01
2.250E-01	3.284E+07	2.879E+07	8.814E+06	6.750E+01	3.863E+00	3.509E+00	1.343E+00	9.072E-05
3.750E-01	9.214E+06	8.078E+06	2.473E+06	1.784E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.750E-01	5.291E+05	4.639E+05	1.420E+05	1.024E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
8.500E-01	8.373E+06	7.341E+06	2.247E+06	1.621E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.250E+00	2.242E+11	1.966E+11	6.017E+10	4.341E+05	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.750E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.250E+00	1.188E+06	1.042E+06	3.189E+05	2.301E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.750E+00	3.677E+03	3.223E+03	9.867E+02	7.119E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00
3.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
7.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
9.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TOTAL	2.395E+11	2.107E+11	6.866E+10	3.144E+09	2.148E+07	1.629E+07	6.237E+06	4.212E+02

TABLE 1-SRS (DWPF). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
PB206	0.0	1.884E-17	7.273E-13	1.358E-08	9.105E-05	6.167E-02	6.859E+00	7.565E+01
TH232	0.0	5.070E-07	5.075E-06	5.128E-05	5.646E-04	9.392E-03	1.526E-01	1.600E+00
U234	5.485E+00	6.155E+00	1.196E+01	5.201E+01	9.045E+01	8.824E+01	6.875E+01	6.909E+00
U235	7.278E+01	7.279E+01	7.284E+01	7.337E+01	7.858E+01	1.239E+02	2.655E+02	2.767E+02
U236	1.742E+01	1.742E+01	1.746E+01	1.782E+01	2.131E+01	4.273E+01	5.600E+01	5.453E+01
U238	3.122E+04	3.122E+04	3.122E+04	3.122E+04	3.122E+04	3.122E+04	3.122E+04	3.122E+04
PU238	8.667E+01	8.599E+01	8.009E+01	3.934E+01	3.216E-02	2.609E-23	0.0	0.0
PU239	2.076E+02	2.076E+02	2.075E+02	2.070E+02	2.017E+02	1.557E+02	1.165E+01	7.439E-11
PU240	3.809E+01	3.814E+01	3.847E+01	3.896E+01	3.544E+01	1.365E+01	9.772E-04	9.805E-15
SUM	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.169E+04
TOTAL	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.169E+04

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 2-SRS (DWPf). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE	0.0	1.254E-02	1.210E-01	8.739E-01	1.902E+00	3.154E+00	6.977E+00	1.730E+01
PB	0.0	8.827E-07	4.575E-05	4.610E-04	8.654E-04	6.300E-02	6.874E+00	7.587E+01
BI	0.0	2.883E-10	1.079E-09	1.065E-09	3.816E-07	4.493E-04	1.387E-01	6.177E+00
RA	0.0	2.658E-08	1.084E-07	2.980E-06	7.768E-04	3.739E-02	2.934E-01	5.880E-02
TH	0.0	2.233E-05	2.689E-04	8.914E-03	2.221E-01	2.367E+00	1.443E+01	4.559E+00
U	3.132E+04	3.132E+04	3.132E+04	3.136E+04	3.141E+04	3.147E+04	3.161E+04	3.156E+04
NP	1.263E+01	1.264E+01	1.273E+01	1.499E+01	2.776E+01	3.162E+01	3.071E+01	2.294E+01
PU	3.518E+02	3.504E+02	3.393E+02	2.886E+02	2.404E+02	1.725E+02	1.433E+01	5.345E-01
AM	3.241E+00	3.996E+00	9.325E+00	1.690E+01	4.043E+00	1.135E-02	2.422E-06	3.168E-12
CM	1.329E+00	1.279E+00	9.065E-01	2.898E-02	3.758E-05	1.771E-05	1.844E-08	6.828E-09
SUM	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.169E+04
TOTAL	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.168E+04	3.169E+04

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 3-SRS (DWPF). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.0	5.229E-11	4.778E-09	2.327E-07	3.418E-06	4.131E-05	4.467E-04	5.966E-04
TL209	0.0	3.272E-12	3.645E-11	7.398E-10	6.245E-08	6.888E-06	1.531E-04	3.730E-04
PB209	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
PB210	0.0	6.973E-13	8.359E-10	1.436E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
PB211	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
PB214	0.0	6.956E-11	9.353E-09	2.902E-06	7.682E-04	3.697E-02	2.902E-01	5.815E-02
BI210	0.0	6.973E-13	8.359E-10	1.436E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
BI211	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
BI213	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
BI214	0.0	6.956E-11	9.353E-09	2.902E-06	7.682E-04	3.697E-02	2.902E-01	5.815E-02
PO210	0.0	2.250E-13	8.359E-10	1.436E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
PO213	0.0	1.482E-10	1.651E-09	3.351E-08	2.829E-06	3.120E-04	6.933E-03	1.689E-02
PO214	0.0	6.955E-11	9.351E-09	2.902E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
PO215	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
PO218	0.0	6.958E-11	9.355E-09	2.903E-06	7.684E-04	3.698E-02	2.902E-01	5.816E-02
AT217	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RN219	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
RN222	0.0	6.958E-11	9.355E-09	2.903E-06	7.684E-04	3.698E-02	2.902E-01	5.816E-02
FR221	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RA223	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
RA225	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RA226	0.0	6.958E-11	9.355E-09	2.903E-06	7.684E-04	3.698E-02	2.902E-01	5.816E-02
AC225	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
AC227	0.0	5.244E-11	4.789E-09	2.333E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
TH227	0.0	5.172E-11	4.725E-09	2.302E-07	3.381E-06	4.085E-05	4.418E-04	5.900E-04
TH229	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
TH230	0.0	3.275E-07	4.931E-06	1.788E-04	4.473E-03	4.758E-02	2.877E-01	5.812E-02
TH231	0.0	1.574E-04	1.575E-04	1.587E-04	1.699E-04	2.679E-04	5.740E-04	5.983E-04
TH232	0.0	5.563E-14	5.569E-13	5.627E-12	6.194E-11	1.030E-09	1.674E-08	1.756E-07
TH234	0.0	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02
PA231	0.0	3.329E-09	3.335E-08	3.340E-07	3.427E-06	4.142E-05	4.479E-04	5.983E-04
PA233	0.0	8.911E-03	8.979E-03	1.057E-02	1.958E-02	2.230E-02	2.166E-02	1.618E-02
PA234M	0.0	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02
PA234	0.0	1.365E-05	1.365E-05	1.365E-05	1.365E-05	1.365E-05	1.365E-05	1.365E-05
U233	1.584E-06	1.623E-06	1.983E-06	5.812E-06	6.913E-05	9.209E-04	7.765E-03	1.717E-02
U234	3.429E-02	3.848E-02	7.475E-02	3.251E-01	5.654E-01	5.516E-01	4.297E-01	4.319E-02

TABLE 3-SRS (DWPf). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES (CONTINUED)

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U235	1.574E-04	1.574E-04	1.575E-04	1.587E-04	1.699E-04	2.679E-04	5.740E-04	5.983E-04
U236	1.127E-03	1.128E-03	1.130E-03	1.154E-03	1.379E-03	2.766E-03	3.625E-03	3.530E-03
U237	0.0	3.903E-02	2.531E-02	3.324E-04	1.519E-10	7.291E-11	4.731E-14	6.258E-46
U238	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02
NP237	8.907E-03	8.911E-03	8.979E-03	1.057E-02	1.958E-02	2.230E-02	2.166E-02	1.618E-02
NP239	0.0	5.787E-03	5.782E-03	5.733E-03	5.269E-03	2.263E-03	4.829E-07	6.318E-13
PU236	1.221E-01	9.575E-02	1.074E-02	1.572E-09	1.560E-09	1.477E-09	8.588E-10	3.784E-12
PU238	1.484E+03	1.473E+03	1.372E+03	6.737E+02	5.507E-01	4.468E-22	0.0	0.0
PU239	1.291E+01	1.291E+01	1.291E+01	1.287E+01	1.254E+01	9.680E+00	7.245E-01	4.626E-12
PU240	8.683E+00	8.693E+00	8.768E+00	8.880E+00	8.078E+00	3.111E+00	2.228E-04	2.235E-15
PU241	1.670E+03	1.591E+03	1.032E+03	1.355E+01	6.200E-06	2.976E-06	1.931E-09	2.554E-41
PU242	1.225E-02	1.225E-02	1.225E-02	1.224E-02	1.222E-02	1.203E-02	1.024E-02	2.042E-03
AM241	1.102E+01	1.362E+01	3.191E+01	5.793E+01	1.379E+01	1.042E-05	1.931E-09	2.691E-41
AM242M	1.447E-02	1.440E-02	1.382E-02	9.169E-03	1.514E-04	2.275E-22	0.0	0.0
AM242	1.436E-02	1.433E-02	1.375E-02	9.124E-03	1.506E-04	2.264E-22	0.0	0.0
AM243	5.787E-03	5.787E-03	5.782E-03	5.733E-03	5.269E-03	2.263E-03	4.829E-07	6.318E-13
CM242	3.496E-02	1.675E-02	1.138E-02	7.545E-03	1.246E-04	1.877E-22	0.0	0.0
CM244	1.076E+02	1.035E+02	7.336E+01	2.341E+00	2.575E-15	5.234E-21	4.781E-20	2.239E-19
SUM	3.294E+03	3.203E+03	2.531E+03	7.697E+02	3.564E+01	1.382E+01	4.212E+00	8.560E-01
TOTAL	3.294E+03	3.203E+03	2.531E+03	7.697E+02	3.564E+01	1.382E+01	4.212E+00	8.560E-01

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 4-SRS (DWPFF). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: ELEMENTS
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	1.128E-03	1.518E-03	5.664E-03	2.598E-03	3.930E-06	4.820E-05	5.998E-04	9.696E-04
PB	0.0	4.225E-03	1.576E-02	7.234E-03	1.544E-03	7.430E-02	5.878E-01	1.342E-01
BI	0.0	4.225E-03	1.576E-02	7.234E-03	1.544E-03	7.430E-02	5.878E-01	1.342E-01
PO	0.0	6.932E-03	2.586E-02	1.187E-02	2.313E-03	1.113E-01	8.778E-01	1.919E-01
AT	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RN	0.0	4.225E-03	1.576E-02	7.232E-03	7.730E-04	3.702E-02	2.907E-01	5.876E-02
FR	0.0	1.522E-10	1.754E-09	3.747E-08	2.938E-06	3.194E-04	7.093E-03	1.728E-02
RA	0.0	4.225E-03	1.576E-02	7.232E-03	7.759E-04	3.734E-02	2.978E-01	7.603E-02
AC	0.0	2.039E-10	6.477E-09	2.676E-07	6.319E-06	3.603E-04	7.535E-03	1.787E-02
TH	0.0	1.488E-02	2.642E-02	1.807E-02	1.515E-02	5.871E-02	3.063E-01	8.707E-02
PA	0.0	1.942E-02	1.949E-02	2.109E-02	3.009E-02	3.285E-02	3.262E-02	2.729E-02
U	5.947E-02	1.036E-01	1.281E-01	3.443E-01	5.775E-01	5.661E-01	4.522E-01	7.499E-02
NP	8.907E-03	1.477E-02	1.483E-02	1.635E-02	2.485E-02	2.456E-02	2.166E-02	1.618E-02
PU	3.176E+03	3.086E+03	2.425E+03	7.090E+02	2.118E+01	1.280E+01	7.350E-01	2.042E-03
AM	1.106E+01	1.365E+01	3.195E+01	5.796E+01	1.380E+01	2.273E-03	4.848E-07	6.318E-13
CM	1.076E+02	1.035E+02	7.337E+01	2.349E+00	1.312E-04	3.094E-06	1.929E-09	7.207E-13
SUM	3.294E+03	3.203E+03	2.531E+03	7.697E+02	3.564E+01	1.382E+01	4.212E+00	8.560E-01
TOTAL	3.294E+03	3.203E+03	2.531E+03	7.697E+02	3.564E+01	1.382E+01	4.212E+00	8.560E-01

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 5-SRS (DWPf). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.0	1.536E-13	1.403E-11	6.834E-10	1.004E-08	1.213E-07	1.312E-06	1.752E-06
TL209	0.0	5.436E-14	6.056E-13	1.229E-11	1.038E-09	1.144E-07	2.543E-06	6.197E-06
PB209	0.0	1.742E-13	1.941E-12	3.938E-11	3.325E-09	3.667E-07	8.149E-06	1.986E-05
PB210	0.0	1.615E-16	1.936E-13	3.327E-10	1.779E-07	8.564E-06	6.721E-05	1.347E-05
PB211	0.0	1.571E-13	1.436E-11	6.993E-10	1.027E-08	1.241E-07	1.342E-06	1.793E-06
PB214	0.0	2.218E-13	2.983E-11	9.256E-09	2.450E-06	1.179E-04	9.254E-04	1.854E-04
BI210	0.0	1.608E-15	1.927E-12	3.312E-09	1.771E-06	8.524E-05	6.690E-04	1.341E-04
BI211	0.0	2.092E-12	1.911E-10	9.309E-09	1.367E-07	1.652E-06	1.787E-05	2.386E-05
BI213	0.0	6.368E-13	7.094E-12	1.440E-10	1.215E-08	1.341E-06	2.979E-05	7.259E-05
BI214	0.0	8.915E-13	1.199E-10	3.720E-08	9.845E-06	4.739E-04	3.719E-03	7.452E-04
PO210	0.0	7.211E-15	2.680E-11	4.604E-08	2.462E-05	1.185E-03	9.300E-03	1.864E-03
PO212	0.0	1.434E-04	5.352E-04	2.455E-04	4.240E-08	8.515E-11	5.976E-10	5.961E-09
PO213	0.0	7.499E-12	8.355E-11	1.696E-09	1.431E-07	1.579E-05	3.509E-04	8.549E-04
PO214	0.0	3.229E-12	4.342E-10	1.347E-07	3.566E-05	1.716E-03	1.347E-02	2.699E-03
PO215	0.0	2.341E-12	2.139E-10	1.042E-08	1.530E-07	1.849E-06	2.000E-05	2.671E-05
PO216	0.0	1.729E-04	6.453E-04	2.959E-04	5.112E-08	1.027E-10	7.205E-10	7.187E-09
PO218	0.0	2.521E-12	3.390E-10	1.052E-07	2.784E-05	1.340E-03	1.052E-02	2.108E-03
AT217	0.0	6.464E-12	7.201E-11	1.462E-09	1.234E-07	1.361E-05	3.024E-04	7.368E-04
RN219	0.0	2.176E-12	1.988E-10	9.684E-09	1.422E-07	1.719E-06	1.859E-05	2.483E-05
RN220	0.0	1.604E-04	5.985E-04	2.745E-04	4.741E-08	9.521E-11	6.682E-10	6.665E-09
RN222	0.0	2.305E-12	3.100E-10	9.619E-08	2.546E-05	1.225E-03	9.617E-03	1.927E-03
FR221	0.0	5.846E-12	6.513E-11	1.322E-09	1.116E-07	1.231E-05	2.735E-04	6.664E-04
RA223	0.0	1.867E-12	1.706E-10	8.310E-09	1.221E-07	1.475E-06	1.595E-05	2.130E-05
RA224	0.0	1.450E-04	5.410E-04	2.481E-04	4.286E-08	8.607E-11	6.041E-10	6.025E-09
RA225	0.0	1.062E-13	1.183E-12	2.402E-11	2.027E-09	2.236E-07	4.969E-06	1.211E-05
RA226	0.0	2.009E-12	2.701E-10	8.382E-08	2.219E-05	1.068E-03	8.380E-03	1.679E-03
AC225	0.0	5.291E-12	5.895E-11	1.196E-09	1.010E-07	1.114E-05	2.475E-04	6.032E-04
AC227	0.0	2.539E-14	2.319E-12	1.130E-10	1.660E-09	2.006E-08	2.169E-07	2.897E-07
TH227	0.0	1.887E-12	1.725E-10	8.400E-09	1.234E-07	1.491E-06	1.612E-05	2.153E-05
TH228	0.0	1.382E-04	5.152E-04	2.364E-04	4.084E-08	8.201E-11	5.756E-10	5.741E-09
TH229	0.0	4.634E-12	5.163E-11	1.048E-09	8.845E-08	9.755E-06	2.168E-04	5.282E-04
TH230	0.0	9.269E-09	1.395E-07	5.059E-06	1.266E-04	1.347E-03	8.140E-03	1.645E-03
TH231	0.0	8.832E-08	8.838E-08	8.902E-08	9.534E-08	1.503E-07	3.221E-07	3.357E-07
TH232	0.0	1.347E-15	1.348E-14	1.362E-13	1.500E-12	2.494E-11	4.053E-10	4.250E-09
TH234	0.0	4.257E-06	4.257E-06	4.257E-06	4.257E-06	4.257E-06	4.257E-06	4.257E-06
PA231	0.0	1.003E-10	1.005E-09	1.006E-08	1.033E-07	1.248E-06	1.350E-05	1.803E-05

TABLE 5-SRS (DWPf). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES (CONTINUED)

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
PA233	0.0	2.022E-05	2.038E-05	2.400E-05	4.443E-05	5.061E-05	4.916E-05	3.673E-05
PA234M	0.0	5.189E-05	5.189E-05	5.189E-05	5.189E-05	5.189E-05	5.189E-05	5.189E-05
PA234	0.0	1.961E-07	1.961E-07	1.961E-07	1.961E-07	1.961E-07	1.961E-07	1.960E-07
U232	4.301E-04	4.593E-04	5.232E-04	2.259E-04	3.903E-08	4.743E-11	2.757E-11	1.215E-13
U233	4.605E-08	4.719E-08	5.765E-08	1.689E-07	2.009E-06	2.677E-05	2.257E-04	4.991E-04
U234	9.875E-04	1.108E-03	2.153E-03	9.365E-03	1.628E-02	1.589E-02	1.238E-02	1.244E-03
U235	4.122E-06	4.122E-06	4.125E-06	4.155E-06	4.450E-06	7.015E-06	1.503E-05	1.567E-05
U236	3.054E-05	3.055E-05	3.061E-05	3.125E-05	3.737E-05	7.492E-05	9.819E-05	9.561E-05
U238	2.663E-04	2.663E-04	2.663E-04	2.663E-04	2.663E-04	2.663E-04	2.663E-04	2.663E-04
NP237	2.722E-04	2.723E-04	2.744E-04	3.232E-04	5.983E-04	6.815E-04	6.619E-04	4.945E-04
NP239	0.0	1.399E-05	1.398E-05	1.386E-05	1.274E-05	5.469E-06	1.167E-09	1.527E-15
PU236	4.249E-03	3.332E-03	3.736E-04	5.469E-11	5.428E-11	5.141E-11	2.989E-11	1.317E-13
PU238	4.919E+01	4.881E+01	4.546E+01	2.233E+01	1.825E-02	1.481E-23	0.0	0.0
PU239	3.979E-01	3.978E-01	3.977E-01	3.967E-01	3.866E-01	2.983E-01	2.233E-02	1.426E-13
PU240	2.704E-01	2.707E-01	2.730E-01	2.765E-01	2.515E-01	9.686E-02	6.936E-06	6.959E-17
PU241	5.176E-02	4.933E-02	3.198E-02	4.201E-04	1.922E-10	9.226E-11	5.986E-14	7.919E-46
PU242	3.616E-04	3.616E-04	3.616E-04	3.616E-04	3.610E-04	3.552E-04	3.023E-04	6.029E-05
AM241	3.661E-01	4.523E-01	1.060E+00	1.924E+00	4.581E-01	3.460E-07	6.414E-11	8.940E-43
AM243	1.860E-04	1.860E-04	1.859E-04	1.843E-04	1.694E-04	7.273E-05	1.552E-08	2.031E-14
CM242	1.288E-03	6.172E-04	4.193E-04	2.780E-04	4.589E-06	6.918E-24	0.0	0.0
CM244	3.763E+00	3.621E+00	2.566E+00	8.189E-02	9.008E-17	1.831E-22	1.672E-21	7.833E-21
SUM	5.405E+01	5.360E+01	4.979E+01	2.502E+01	1.133E+00	4.213E-01	1.027E-01	1.941E-02
TOTAL	5.405E+01	5.360E+01	4.979E+01	2.502E+01	1.133E+00	4.213E-01	1.027E-01	1.941E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 6-SRS (DWPF). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

WATTS								
ELEMENT	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	2.654E-05	3.572E-05	1.333E-04	6.113E-05	2.163E-08	2.358E-07	3.855E-06	7.950E-06
PB	0.0	8.044E-06	3.001E-05	1.377E-05	2.644E-06	1.270E-04	1.002E-03	2.206E-04
BI	0.0	7.185E-05	2.681E-04	1.230E-04	1.179E-05	5.621E-04	4.435E-03	9.757E-04
PO	0.0	3.164E-04	1.181E-03	5.417E-04	8.851E-05	4.259E-03	3.366E-02	7.552E-03
AT	0.0	6.464E-12	7.201E-11	1.462E-09	1.234E-07	1.361E-05	3.024E-04	7.368E-04
RN	0.0	1.604E-04	5.985E-04	2.746E-04	2.565E-05	1.227E-03	9.636E-03	1.952E-03
FR	0.0	5.848E-12	6.530E-11	1.330E-09	1.117E-07	1.231E-05	2.735E-04	6.664E-04
RA	0.0	1.450E-04	5.410E-04	2.482E-04	2.235E-05	1.070E-03	8.401E-03	1.713E-03
AC	0.0	5.316E-12	6.127E-11	1.309E-09	1.027E-07	1.116E-05	2.478E-04	6.035E-04
TH	0.0	1.425E-04	5.196E-04	2.458E-04	1.312E-04	1.362E-03	8.378E-03	2.199E-03
PA	0.0	7.231E-05	7.247E-05	7.610E-05	9.662E-05	1.039E-04	1.147E-04	1.068E-04
U	1.719E-03	1.942E-03	3.025E-03	9.893E-03	1.659E-02	1.626E-02	1.298E-02	2.121E-03
NP	2.722E-04	2.867E-04	2.887E-04	3.373E-04	6.111E-04	6.870E-04	6.619E-04	4.945E-04
PU	4.992E+01	4.953E+01	4.616E+01	2.300E+01	6.567E-01	3.955E-01	2.264E-02	6.029E-05
AM	3.663E-01	4.525E-01	1.060E+00	1.925E+00	4.583E-01	7.308E-05	1.559E-08	2.031E-14
CM	3.764E+00	3.622E+00	2.567E+00	8.219E-02	4.810E-06	1.026E-07	6.406E-11	3.125E-14
SUM	5.405E+01	5.360E+01	4.979E+01	2.502E+01	1.133E+00	4.213E-01	1.027E-01	1.941E-02
TOTAL	5.405E+01	5.360E+01	4.979E+01	2.502E+01	1.133E+00	4.213E-01	1.027E-01	1.941E-02

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 7-SRS (DWPf). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI211	0.0	5.229E-11	4.778E-09	2.327E-07	3.418E-06	4.131E-05	4.467E-04	5.966E-04
BI213	0.0	3.272E-12	3.645E-11	7.398E-10	6.245E-08	6.888E-06	1.531E-04	3.730E-04
BI214	0.0	1.461E-14	1.964E-12	6.095E-10	1.613E-07	7.765E-06	6.094E-05	1.221E-05
PO210	0.0	2.250E-13	8.359E-10	1.436E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
PO213	0.0	1.482E-10	1.651E-09	3.351E-08	2.829E-06	3.120E-04	6.933E-03	1.689E-02
PO214	0.0	6.955E-11	9.351E-09	2.902E-06	7.680E-04	3.697E-02	2.901E-01	5.814E-02
PO215	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
PO216	0.0	4.225E-03	1.576E-02	7.229E-03	1.249E-06	2.508E-09	1.760E-08	1.756E-07
PO218	0.0	6.956E-11	9.353E-09	2.902E-06	7.682E-04	3.697E-02	2.902E-01	5.815E-02
AT217	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RN219	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
RN220	0.0	4.225E-03	1.576E-02	7.229E-03	1.249E-06	2.508E-09	1.760E-08	1.756E-07
RN222	0.0	6.958E-11	9.355E-09	2.903E-06	7.684E-04	3.698E-02	2.902E-01	5.816E-02
FR221	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RA223	0.0	5.244E-11	4.791E-09	2.334E-07	3.428E-06	4.143E-05	4.480E-04	5.983E-04
RA224	0.0	4.225E-03	1.576E-02	7.229E-03	1.249E-06	2.508E-09	1.760E-08	1.756E-07
RA226	0.0	6.958E-11	9.355E-09	2.903E-06	7.684E-04	3.698E-02	2.902E-01	5.816E-02
AC225	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
AC227	0.0	7.237E-13	6.609E-11	3.220E-09	4.731E-08	5.717E-07	6.182E-06	8.256E-06
TH227	0.0	5.172E-11	4.725E-09	2.302E-07	3.381E-06	4.085E-05	4.418E-04	5.900E-04
TH228	0.0	4.225E-03	1.575E-02	7.229E-03	1.249E-06	2.508E-09	1.760E-08	1.756E-07
TH229	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
TH230	0.0	3.275E-07	4.931E-06	1.788E-04	4.473E-03	4.758E-02	2.877E-01	5.812E-02
PA231	0.0	3.329E-09	3.335E-08	3.340E-07	3.427E-06	4.142E-05	4.479E-04	5.983E-04
U232	1.340E-02	1.431E-02	1.630E-02	7.037E-03	1.216E-06	1.477E-09	8.588E-10	3.784E-12
U233	1.584E-06	1.623E-06	1.983E-06	5.812E-06	6.913E-05	9.209E-04	7.765E-03	1.717E-02
U234	3.429E-02	3.848E-02	7.475E-02	3.251E-01	5.654E-01	5.516E-01	4.297E-01	4.319E-02
U235	1.574E-04	1.574E-04	1.575E-04	1.587E-04	1.699E-04	2.679E-04	5.740E-04	5.983E-04
U236	1.127E-03	1.128E-03	1.130E-03	1.154E-03	1.379E-03	2.766E-03	3.625E-03	3.530E-03
U238	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02	1.050E-02
NP237	8.907E-03	8.911E-03	8.979E-03	1.057E-02	1.958E-02	2.230E-02	2.166E-02	1.618E-02
PU236	1.221E-01	9.575E-02	1.074E-02	1.572E-09	1.560E-09	1.477E-09	8.588E-10	3.784E-12
PU238	1.484E+03	1.473E+03	1.372E+03	6.737E+02	5.507E-01	4.468E-22	0.0	0.0
PU239	1.291E+01	1.291E+01	1.291E+01	1.287E+01	1.254E+01	9.680E+00	7.245E-01	4.626E-12
PU240	8.683E+00	8.693E+00	8.768E+00	8.880E+00	8.078E+00	3.111E+00	2.228E-04	2.235E-15

TABLE 7-SRS (DWPF). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
PU241	4.091E-02	3.898E-02	2.528E-02	3.320E-04	1.519E-10	7.291E-11	4.731E-14	6.258E-46
PU242	1.225E-02	1.225E-02	1.225E-02	1.224E-02	1.222E-02	1.203E-02	1.024E-02	2.042E-03
AM241	1.102E+01	1.362E+01	3.191E+01	5.793E+01	1.379E+01	1.042E-05	1.931E-09	2.691E-41
AM243	5.787E-03	5.787E-03	5.782E-03	5.733E-03	5.269E-03	2.263E-03	4.829E-07	6.318E-13
CM242	3.496E-02	1.675E-02	1.138E-02	7.545E-03	1.246E-04	1.877E-22	0.0	0.0
CM244	1.076E+02	1.035E+02	7.336E+01	2.341E+00	2.575E-15	5.234E-21	4.781E-20	2.239E-19
SUM	1.625E+03	1.612E+03	1.499E+03	7.561E+02	3.559E+01	1.363E+01	2.986E+00	5.320E-01
TOTAL	1.625E+03	1.612E+03	1.499E+03	7.561E+02	3.559E+01	1.363E+01	2.986E+00	5.320E-01

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 8-SRS (DWPF). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI	0.0	1.518E-03	5.664E-03	2.598E-03	4.091E-06	5.596E-05	6.607E-04	9.819E-04
PO	0.0	6.932E-03	2.586E-02	1.187E-02	2.313E-03	1.113E-01	8.778E-01	1.919E-01
AT	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RN	0.0	4.225E-03	1.576E-02	7.232E-03	7.730E-04	3.702E-02	2.907E-01	5.876E-02
FR	0.0	1.515E-10	1.688E-09	3.425E-08	2.891E-06	3.189E-04	7.087E-03	1.727E-02
RA	0.0	4.225E-03	1.576E-02	7.232E-03	7.730E-04	3.702E-02	2.907E-01	5.876E-02
AC	0.0	1.522E-10	1.754E-09	3.747E-08	2.938E-06	3.194E-04	7.093E-03	1.728E-02
TH	0.0	4.225E-03	1.576E-02	7.408E-03	4.481E-03	4.794E-02	2.952E-01	7.597E-02
PA	0.0	3.329E-09	3.335E-08	3.340E-07	3.427E-06	4.142E-05	4.479E-04	5.983E-04
U	5.947E-02	6.457E-02	1.028E-01	3.440E-01	5.775E-01	5.661E-01	4.522E-01	7.499E-02
NP	8.907E-03	8.911E-03	8.979E-03	1.057E-02	1.958E-02	2.230E-02	2.166E-02	1.618E-02
PU	1.506E+03	1.494E+03	1.393E+03	6.954E+02	2.118E+01	1.280E+01	7.350E-01	2.042E-03
AM	1.103E+01	1.362E+01	3.192E+01	5.794E+01	1.380E+01	2.273E-03	4.848E-07	6.318E-13
CM	1.076E+02	1.035E+02	7.337E+01	2.349E+00	1.312E-04	3.094E-06	1.929E-09	7.133E-13
SUM	1.625E+03	1.612E+03	1.499E+03	7.561E+02	3.559E+01	1.363E+01	2.986E+00	5.320E-01
TOTAL	1.625E+03	1.612E+03	1.499E+03	7.561E+02	3.559E+01	1.363E+01	2.986E+00	5.320E-01

TABLE 9-SRS (DWPFF). (ALPHA,N) NEUTRON SOURCES IN DECAY OF SRP HLW

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

		NEUTRONS/SEC						
	IMMOBLZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KYR	100.0KYR	1.0MYR
PO210	0.000E-01	7.738E-09	2.875E-05	4.939E-02	2.641E+01	1.271E+03	9.977E+03	2.000E+03
PO213	0.000E-01	2.363E-05	2.633E-04	5.344E-03	4.511E-01	4.976E+01	1.106E+03	2.693E+03
PO214	0.000E-01	8.737E-06	1.175E-03	3.646E-01	9.648E+01	4.644E+03	3.644E+04	7.304E+03
PO218	0.000E-01	3.693E-06	4.965E-04	1.541E-01	4.078E+01	1.963E+03	1.541E+04	3.087E+03
AT217	0.000E-01	1.432E-05	1.596E-04	3.237E-03	2.733E-01	3.014E+01	6.699E+02	1.632E+03
RN222	0.000E-01	2.697E-06	3.627E-04	1.125E-01	2.979E+01	1.434E+03	1.125E+04	2.255E+03
FR221	0.000E-01	9.900E-06	1.103E-04	2.238E-03	1.889E-01	2.084E+01	4.631E+02	1.129E+03
RA226	0.000E-01	1.580E-06	2.125E-04	6.593E-02	1.745E+01	8.399E+02	6.591E+03	1.321E+03
AC225	0.000E-01	6.976E-06	7.772E-05	1.577E-03	1.331E-01	1.468E+01	3.263E+02	7.952E+02
TH229	0.000E-01	3.705E-06	4.128E-05	8.376E-04	7.070E-02	7.799E+00	1.733E+02	4.223E+02
TH230	0.000E-01	6.730E-03	1.013E-01	3.674E+00	9.192E+01	9.778E+02	5.912E+03	1.194E+03
U233	3.723E-02	3.815E-02	4.661E-02	1.366E-01	1.625E+00	2.164E+01	1.825E+02	4.036E+02
U234	7.782E+02	8.733E+02	1.696E+03	7.378E+03	1.283E+04	1.252E+04	9.751E+03	9.801E+02
U238	1.331E+02	1.331E+02	1.331E+02	1.331E+02	1.331E+02	1.331E+02	1.331E+02	1.331E+02
NP237	1.998E+02	1.999E+02	2.014E+02	2.371E+02	4.393E+02	5.003E+02	4.859E+02	3.630E+02
PU238	5.791E+07	5.748E+07	5.354E+07	2.629E+07	2.149E+04	1.744E-17	0.000E-01	0.000E-01
PU239	3.969E+05	3.969E+05	3.969E+05	3.957E+05	3.855E+05	2.976E+05	2.227E+04	1.422E-07
PU240	2.680E+05	2.683E+05	2.706E+05	2.741E+05	2.493E+05	9.602E+04	6.876E+00	6.898E-11
PU242	3.064E+02	3.064E+02	3.064E+02	3.061E+02	3.056E+02	3.009E+02	2.561E+02	5.107E+01
AM241	4.248E+05	5.250E+05	1.230E+06	2.233E+06	5.316E+05	4.017E-01	7.444E-05	0.000E-01
CM242	1.975E+03	9.464E+02	6.430E+02	4.263E+02	7.040E+00	1.061E-17	0.000E-01	0.000E-01
CM244	5.105E+06	4.911E+06	3.481E+06	1.111E+05	1.222E-10	2.483E-16	2.268E-15	1.062E-14
CM246	1.928E-02	1.928E-02	1.926E-02	1.900E-02	1.665E-02	4.454E-03	8.363E-09	0.000E-01
TOTAL	6.411E+07	6.359E+07	5.892E+07	2.931E+07	1.202E+06	4.183E+05	1.214E+05	2.576E+04

TABLE 10-SRS (DWPF). SPONTANEOUS FISSION NEUTRON SOURCES IN DECAY OF SRS HLW
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	NEUTRONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	3.961E+02	3.961E+02	3.961E+02	3.961E+02	3.961E+02	3.961E+02	3.961E+02	3.961E+02
PU238	2.303E+05	2.285E+05	2.128E+05	1.045E+05	8.545E+01	6.932E-20	0.0	0.0
PU240	3.468E+04	3.472E+04	3.502E+04	3.547E+04	3.227E+04	1.243E+04	8.898E-01	8.928E-12
PU242	5.404E+03	5.404E+03	5.404E+03	5.404E+03	5.395E+03	5.309E+03	4.518E+03	9.010E+02
CM244	1.478E+07	1.422E+07	1.008E+07	3.217E+05	3.538E-10	7.192E-16	6.568E-15	3.077E-14
SUM	1.505E+07	1.449E+07	1.033E+07	4.675E+05	3.817E+04	1.814E+04	4.916E+03	1.298E+03
TOTAL	1.505E+07	1.449E+07	1.033E+07	4.675E+05	3.817E+04	1.814E+04	4.916E+03	1.298E+03

TABLE 11-SRS (DWPF). PHOTON SPECTRUM OF ACTINIDES AND DAUGHTERS IN DECAY OF SRS HLW
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

PHOTONS/SEC								
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	9.408E+12	9.340E+12	8.727E+12	4.479E+12	1.918E+11	4.398E+10	1.947E+10	5.517E+09
2.500E-02	1.036E+10	1.289E+10	3.003E+10	5.433E+10	1.304E+10	3.553E+08	1.809E+09	5.062E+08
3.750E-02	2.689E+10	2.690E+10	2.630E+10	1.582E+10	1.142E+09	1.998E+08	1.215E+09	5.410E+08
5.750E-02	1.519E+11	1.882E+11	4.398E+11	7.975E+11	1.901E+11	4.644E+08	2.114E+09	5.235E+08
8.500E-02	5.138E+09	5.637E+09	5.557E+09	3.245E+09	7.170E+08	9.707E+08	3.916E+09	1.534E+09
1.250E-01	2.689E+08	8.936E+08	8.909E+08	7.647E+08	3.651E+08	2.549E+08	7.081E+08	2.785E+08
2.250E-01	5.065E+08	1.031E+09	1.122E+09	4.800E+08	1.324E+08	6.804E+08	4.825E+09	1.181E+09
3.750E-01	3.644E+07	2.122E+08	2.325E+08	2.416E+08	3.522E+08	8.989E+08	4.584E+09	1.304E+09
5.750E-01	4.869E+07	7.142E+07	2.437E+08	1.264E+08	2.557E+07	7.523E+08	5.864E+09	1.192E+09
8.500E-01	5.581E+07	8.186E+07	1.530E+08	6.999E+07	8.464E+06	1.843E+08	1.423E+09	2.908E+08
1.250E+00	1.157E+07	1.565E+07	1.713E+07	6.404E+06	1.222E+07	4.703E+08	3.674E+09	7.403E+08
1.750E+00	5.180E+06	1.065E+07	2.367E+07	9.645E+06	8.243E+06	3.781E+08	2.968E+09	6.063E+08
2.250E+00	2.997E+06	2.887E+06	2.067E+06	1.210E+05	2.428E+06	1.164E+08	9.134E+08	1.830E+08
2.750E+00	4.133E+07	5.496E+07	2.000E+08	9.124E+07	6.370E+04	2.027E+06	1.589E+07	3.186E+06
3.500E+00	1.560E+06	1.502E+06	1.075E+06	5.630E+04	1.313E+04	3.827E+05	2.985E+06	5.981E+05
5.000E+00	6.661E+05	6.416E+05	4.587E+05	2.347E+04	2.216E+03	1.048E+03	2.848E+02	7.590E+01
7.000E+00	7.667E+04	7.385E+04	5.276E+04	2.631E+03	2.517E+02	1.194E+02	3.250E+01	8.691E+00
9.500E+00	8.799E+03	8.475E+03	6.053E+03	2.975E+02	2.873E+01	1.365E+01	3.722E+00	9.966E-01
TOTAL	9.604E+12	9.576E+12	9.232E+12	5.352E+12	3.977E+11	4.971E+10	5.351E+10	1.440E+10

TABLE 12-SRS (DWPF). MASS OF FISSION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
RB 87	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00
SR 90	3.426E+02	3.345E+02	2.700E+02	3.170E+01	1.576E-08	0.0	0.0	0.0
ZR 90	0.0	8.062E+00	7.259E+01	3.110E+02	3.427E+02	3.427E+02	3.427E+02	3.427E+02
ZR 93	4.443E+02	4.443E+02	4.443E+02	4.443E+02	4.441E+02	4.423E+02	4.246E+02	2.824E+02
NB 93	0.0	1.486E-05	5.147E-04	1.640E-02	1.975E-01	2.005E+00	1.968E+01	1.619E+02
TC 99	1.816E+02	1.816E+02	1.816E+02	1.815E+02	1.810E+02	1.758E+02	1.312E+02	7.013E+00
RU 99	0.0	5.909E-04	5.909E-03	5.908E-02	5.900E-01	5.814E+00	5.044E+01	1.746E+02
PD107	2.863E+01	2.863E+01	2.863E+01	2.863E+01	2.863E+01	2.860E+01	2.833E+01	2.573E+01
AG107	0.0	3.055E-06	3.055E-05	3.055E-04	3.055E-03	3.053E-02	3.039E-01	2.898E+00
SN126	1.556E+01	1.556E+01	1.556E+01	1.555E+01	1.545E+01	1.452E+01	7.780E+00	1.520E-02
TE126	0.0	1.078E-04	1.078E-03	1.078E-02	1.075E-01	1.042E+00	7.780E+00	1.554E+01
CS135	8.633E+01	8.633E+01	8.633E+01	8.633E+01	8.630E+01	8.607E+01	8.377E+01	6.387E+01
BA135	0.0	2.602E-05	2.602E-04	2.602E-03	2.601E-02	2.598E-01	2.563E+00	2.246E+01
CS137	4.989E+02	4.875E+02	3.960E+02	4.949E+01	4.607E-08	0.0	0.0	0.0
BA137	0.0	1.140E+01	1.029E+02	4.494E+02	4.989E+02	4.989E+02	4.989E+02	4.989E+02
CE142	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02
CE144	3.093E+00	1.269E+00	4.194E-04	6.469E-39	0.0	0.0	0.0	0.0
ND144	4.110E+02	4.128E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02
PM147	2.609E+01	2.003E+01	1.858E+00	8.758E-11	0.0	0.0	0.0	0.0
SM147	8.796E+01	9.402E+01	1.122E+02	1.140E+02	1.140E+02	1.140E+02	1.140E+02	1.140E+02
SM148	1.916E+01	1.916E+01	1.916E+01	1.916E+01	1.916E+01	1.916E+01	1.916E+01	1.916E+01
SM149	7.420E+00	7.420E+00	7.420E+00	7.420E+00	7.420E+00	7.420E+00	7.420E+00	7.420E+00
SM151	9.418E+00	9.346E+00	8.720E+00	4.360E+00	4.256E-03	3.342E-33	0.0	0.0
EU151	0.0	7.226E-02	6.982E-01	5.058E+00	9.414E+00	9.418E+00	9.418E+00	9.418E+00
SUM	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03
TOTAL	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 13-SRS (DWPf). MASS OF FISSION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	2.439E+00	2.439E+00	2.439E+00	2.436E+00	2.413E+00	2.192E+00	8.391E-01	5.679E-05
BR	0.0	2.602E-05	2.602E-04	2.601E-03	2.589E-02	2.468E-01	1.600E+00	2.439E+00
RB	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00	9.961E+00
SR	3.426E+02	3.345E+02	2.700E+02	3.170E+01	1.628E-07	1.470E-06	1.470E-05	1.470E-04
ZR	4.443E+02	4.524E+02	5.169E+02	7.553E+02	7.868E+02	7.850E+02	7.673E+02	6.251E+02
NB	5.152E-04	7.160E-04	2.527E-03	2.064E-02	2.017E-01	2.009E+00	1.968E+01	1.619E+02
TC	1.816E+02	1.816E+02	1.816E+02	1.815E+02	1.810E+02	1.758E+02	1.312E+02	7.013E+00
RU	6.729E-01	3.389E-01	6.604E-03	5.908E-02	5.900E-01	5.814E+00	5.044E+01	1.746E+02
PD	2.863E+01	2.896E+01	2.930E+01	2.930E+01	2.930E+01	2.927E+01	2.900E+01	2.641E+01
AG	2.647E-05	1.267E-05	3.055E-05	3.055E-04	3.055E-03	3.053E-02	3.039E-01	2.898E+00
SN	1.556E+01	1.556E+01	1.556E+01	1.555E+01	1.545E+01	1.452E+01	7.780E+00	1.520E-02
SB	8.226E-01	6.405E-01	6.756E-02	1.034E-03	1.368E-03	1.368E-03	1.367E-03	1.367E-03
TE	1.533E-02	1.975E-01	7.716E-01	8.487E-01	9.454E-01	1.880E+00	8.617E+00	1.638E+01
CS	5.855E+02	5.740E+02	4.823E+02	1.358E+02	8.630E+01	8.607E+01	8.377E+01	6.387E+01
BA	7.724E-05	1.147E+01	1.032E+02	4.497E+02	4.992E+02	4.994E+02	5.017E+02	5.216E+02
CE	4.036E+02	4.018E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02	4.005E+02
ND	4.110E+02	4.128E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02	4.141E+02
PM	2.609E+01	2.003E+01	1.858E+00	8.758E-11	0.0	0.0	0.0	0.0
SM	1.240E+02	1.299E+02	1.475E+02	1.450E+02	1.406E+02	1.406E+02	1.406E+02	1.406E+02
EU	3.337E+00	3.098E+00	1.988E+00	5.059E+00	9.414E+00	9.418E+00	9.418E+00	9.418E+00
GD	0.0	3.112E-01	2.041E+00	3.321E+00	3.322E+00	3.322E+00	3.322E+00	3.322E+00
SUM	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03
TOTAL	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03	2.580E+03

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 14-SRS (DWPF). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	1.700E-01	1.700E-01	1.700E-01	1.698E-01	1.682E-01	1.528E-01	5.848E-02	3.958E-06
SR 90	4.675E+04	4.565E+04	3.685E+04	4.326E+03	2.151E-06	0.0	0.0	0.0
Y 90	4.787E+04	4.566E+04	3.686E+04	4.327E+03	2.152E-06	0.0	0.0	0.0
ZR 93	1.117E+00	1.117E+00	1.117E+00	1.117E+00	1.116E+00	1.112E+00	1.067E+00	7.099E-01
NB 93M	0.0	5.272E-02	4.236E-01	1.054E+00	1.061E+00	1.056E+00	1.014E+00	6.744E-01
NB 94	9.647E-05	9.647E-05	9.644E-05	9.614E-05	9.323E-05	6.857E-05	3.173E-06	1.429E-19
TC 99	3.080E+00	3.080E+00	3.080E+00	3.079E+00	3.070E+00	2.981E+00	2.224E+00	1.189E-01
RU106	2.252E+03	1.132E+03	2.324E+00	3.232E-27	0.0	0.0	0.0	0.0
RH106	2.260E+03	1.132E+03	2.324E+00	3.232E-27	0.0	0.0	0.0	0.0
PD107	1.473E-02	1.473E-02	1.473E-02	1.473E-02	1.473E-02	1.472E-02	1.457E-02	1.324E-02
SB125	8.497E+02	6.616E+02	6.958E+01	1.151E-08	0.0	0.0	0.0	0.0
TE125M	2.760E+02	1.623E+02	1.698E+01	2.809E-09	0.0	0.0	0.0	0.0
SN126	4.416E-01	4.416E-01	4.416E-01	4.413E-01	4.386E-01	4.121E-01	2.208E-01	4.315E-04
SB126	6.160E-02	6.183E-02	6.182E-02	6.179E-02	6.140E-02	5.769E-02	3.092E-02	6.041E-05
SB126M	4.415E-01	4.416E-01	4.416E-01	4.413E-01	4.386E-01	4.121E-01	2.208E-01	4.315E-04
CS134	3.373E+02	2.410E+02	1.170E+01	8.453E-13	0.0	0.0	0.0	0.0
CS135	9.944E-02	9.944E-02	9.944E-02	9.944E-02	9.941E-02	9.914E-02	9.649E-02	7.357E-02
CS137	4.342E+04	4.242E+04	3.446E+04	4.307E+03	4.009E-06	0.0	0.0	0.0
BA137M	4.156E+04	4.013E+04	3.260E+04	4.074E+03	3.792E-06	0.0	0.0	0.0
CE144	9.871E+03	4.051E+03	1.339E+00	2.064E-35	0.0	0.0	0.0	0.0
PR144	9.871E+03	4.051E+03	1.339E+00	2.065E-35	0.0	0.0	0.0	0.0
PR144M	1.187E+02	4.861E+01	1.606E-02	2.477E-37	0.0	0.0	0.0	0.0
PM147	2.420E+04	1.858E+04	1.723E+03	8.122E-08	0.0	0.0	0.0	0.0
SM151	2.479E+02	2.460E+02	2.295E+02	1.147E+02	1.120E-01	8.796E-32	0.0	0.0
EU152	3.688E+00	3.505E+00	2.216E+00	2.257E-02	2.713E-22	0.0	0.0	0.0
EU154	6.197E+02	5.718E+02	2.768E+02	1.959E-01	6.157E-33	0.0	0.0	0.0
EU155	4.750E+02	4.131E+02	1.174E+02	4.042E-04	0.0	0.0	0.0	0.0
SUM	2.310E+05	2.052E+05	1.432E+05	1.716E+04	6.580E+00	6.298E+00	4.948E+00	1.591E+00
TOTAL	2.310E+05	2.052E+05	1.432E+05	1.716E+04	6.580E+00	6.298E+00	4.948E+00	1.591E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 15-SRS (DWPf). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	1.700E-01	1.700E-01	1.700E-01	1.698E-01	1.682E-01	1.528E-01	5.848E-02	3.958E-06
SR	4.675E+04	4.565E+04	3.685E+04	4.326E+03	2.151E-06	0.0	0.0	0.0
Y	4.787E+04	4.566E+04	3.686E+04	4.327E+03	2.152E-06	0.0	0.0	0.0
ZR	1.127E+00	1.117E+00	1.117E+00	1.117E+00	1.116E+00	1.112E+00	1.067E+00	7.099E-01
NB	2.137E-02	5.326E-02	4.237E-01	1.055E+00	1.061E+00	1.056E+00	1.014E+00	6.744E-01
TC	3.080E+00	3.080E+00	3.080E+00	3.079E+00	3.070E+00	2.981E+00	2.224E+00	1.189E-01
RU	2.252E+03	1.132E+03	2.324E+00	3.232E-27	0.0	0.0	0.0	0.0
RH	2.260E+03	1.132E+03	2.324E+00	3.232E-27	0.0	0.0	0.0	0.0
PD	1.473E-02	1.473E-02	1.473E-02	1.473E-02	1.473E-02	1.472E-02	1.457E-02	1.324E-02
SN	7.756E-01	5.555E-01	5.104E-01	4.611E-01	4.386E-01	4.121E-01	2.208E-01	4.315E-04
SB	8.502E+02	6.621E+02	7.008E+01	5.031E-01	5.000E-01	4.697E-01	2.517E-01	4.919E-04
TE	2.763E+02	1.623E+02	1.698E+01	2.809E-09	0.0	0.0	0.0	0.0
CS	4.375E+04	4.266E+04	3.447E+04	4.307E+03	9.942E-02	9.914E-02	9.649E-02	7.357E-02
BA	4.156E+04	4.013E+04	3.260E+04	4.074E+03	3.792E-06	0.0	0.0	0.0
CE	9.871E+03	4.051E+03	1.339E+00	9.614E-06	9.614E-06	9.614E-06	9.614E-06	9.614E-06
PR	9.990E+03	4.100E+03	1.355E+00	2.089E-35	0.0	0.0	0.0	0.0
PM	2.420E+04	1.858E+04	1.723E+03	8.122E-08	0.0	0.0	0.0	0.0
SM	2.479E+02	2.460E+02	2.295E+02	1.147E+02	1.120E-01	2.593E-06	2.593E-06	2.593E-06
EU	1.098E+03	9.883E+02	3.964E+02	2.188E-01	2.713E-22	0.0	0.0	0.0
SUM	2.310E+05	2.052E+05	1.432E+05	1.716E+04	6.580E+00	6.298E+00	4.948E+00	1.591E+00
TOTAL	2.310E+05	2.052E+05	1.432E+05	1.716E+04	6.580E+00	6.298E+00	4.948E+00	1.591E+00

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 16-SRS (DWPf). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	4.232E-05	4.232E-05	4.231E-05	4.227E-05	4.187E-05	3.803E-05	1.456E-05	9.854E-10
SR 90	5.426E+01	5.298E+01	4.277E+01	5.021E+00	2.497E-09	0.0	0.0	0.0
Y 90	2.653E+02	2.531E+02	2.043E+02	2.398E+01	1.193E-08	0.0	0.0	0.0
ZR 93	1.298E-04	1.298E-04	1.298E-04	1.297E-04	1.297E-04	1.292E-04	1.240E-04	8.248E-05
NB 93M	0.0	9.340E-06	7.506E-05	1.868E-04	1.879E-04	1.871E-04	1.797E-04	1.195E-04
NB 94	9.830E-07	9.830E-07	9.827E-07	9.797E-07	9.500E-07	6.987E-07	3.233E-08	1.456E-21
TC 99	1.545E-03	1.545E-03	1.544E-03	1.544E-03	1.540E-03	1.495E-03	1.115E-03	5.964E-05
RU106	1.339E-01	6.733E-02	1.382E-04	1.922E-31	0.0	0.0	0.0	0.0
RH106	2.167E+01	1.086E+01	2.229E-02	3.100E-29	0.0	0.0	0.0	0.0
PD107	8.732E-07	8.732E-07	8.732E-07	8.732E-07	8.731E-07	8.723E-07	8.640E-07	7.848E-07
SB125	2.656E+00	2.068E+00	2.175E-01	3.599E-11	0.0	0.0	0.0	0.0
TE125M	2.320E-01	1.364E-01	1.427E-02	2.361E-12	0.0	0.0	0.0	0.0
SN126	5.508E-04	5.508E-04	5.508E-04	5.504E-04	5.470E-04	5.139E-04	2.754E-04	5.382E-07
SB126	1.138E-03	1.142E-03	1.142E-03	1.142E-03	1.134E-03	1.066E-03	5.712E-04	1.116E-06
SB126M	5.622E-03	5.623E-03	5.623E-03	5.619E-03	5.584E-03	5.247E-03	2.812E-03	5.494E-06
CS134	3.433E+00	2.453E+00	1.191E-01	8.604E-15	0.0	0.0	0.0	0.0
CS135	3.319E-05	3.319E-05	3.319E-05	3.319E-05	3.318E-05	3.309E-05	3.220E-05	2.455E-05
CS137	4.802E+01	4.692E+01	3.811E+01	4.764E+00	4.434E-09	0.0	0.0	0.0
BA137M	1.632E+02	1.576E+02	1.280E+02	1.600E+01	1.489E-08	0.0	0.0	0.0
CE144	6.547E+00	2.687E+00	8.879E-04	1.369E-38	0.0	0.0	0.0	0.0
PR144	7.255E+01	2.978E+01	9.839E-03	1.518E-37	0.0	0.0	0.0	0.0
PR144M	4.063E-02	1.663E-02	5.496E-06	8.476E-41	0.0	0.0	0.0	0.0
PM147	8.679E+00	6.664E+00	6.180E-01	2.913E-11	0.0	0.0	0.0	0.0
SM147	2.738E-08	2.927E-08	3.493E-08	3.551E-08	3.551E-08	3.551E-08	3.551E-08	3.551E-08
SM151	2.906E-02	2.884E-02	2.691E-02	1.345E-02	1.313E-05	1.031E-35	0.0	0.0
EU152	2.790E-02	2.651E-02	1.676E-02	1.707E-04	2.052E-24	0.0	0.0	0.0
EU154	5.543E+00	5.114E+00	2.476E+00	1.752E-03	5.507E-35	0.0	0.0	0.0
EU155	3.455E-01	3.004E-01	8.540E-02	2.940E-07	0.0	0.0	0.0	0.0
SUM	6.527E+02	5.708E+02	4.168E+02	4.979E+01	9.213E-03	8.710E-03	5.125E-03	2.941E-04
TOTAL	6.527E+02	5.708E+02	4.168E+02	4.979E+01	9.213E-03	8.710E-03	5.125E-03	2.941E-04

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 17-SRS (DWPF). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	4.232E-05	4.232E-05	4.231E-05	4.227E-05	4.187E-05	3.803E-05	1.456E-05	9.854E-10
SR	5.426E+01	5.298E+01	4.277E+01	5.021E+00	2.497E-09	0.0	0.0	0.0
Y	2.653E+02	2.531E+02	2.043E+02	2.398E+01	1.193E-08	0.0	0.0	0.0
ZR	1.807E-04	1.307E-04	1.298E-04	1.297E-04	1.297E-04	1.292E-04	1.240E-04	8.248E-05
NB	1.026E-04	1.244E-05	7.604E-05	1.878E-04	1.888E-04	1.878E-04	1.797E-04	1.195E-04
TC	1.545E-03	1.545E-03	1.544E-03	1.544E-03	1.540E-03	1.495E-03	1.115E-03	5.964E-05
RU	1.339E-01	6.733E-02	1.382E-04	1.922E-31	0.0	0.0	0.0	0.0
RH	2.167E+01	1.086E+01	2.229E-02	3.100E-29	0.0	0.0	0.0	0.0
PD	8.732E-07	8.732E-07	8.732E-07	8.732E-07	8.731E-07	8.723E-07	8.640E-07	7.848E-07
SN	1.505E-03	8.191E-04	6.886E-04	5.900E-04	5.470E-04	5.139E-04	2.754E-04	5.382E-07
SB	2.663E+00	2.075E+00	2.243E-01	6.761E-03	6.719E-03	6.312E-03	3.383E-03	6.610E-06
TE	2.323E-01	1.364E-01	1.427E-02	2.361E-12	0.0	0.0	0.0	0.0
CS	5.145E+01	4.938E+01	3.823E+01	4.764E+00	3.318E-05	3.309E-05	3.220E-05	2.455E-05
BA	1.632E+02	1.576E+02	1.280E+02	1.600E+01	1.489E-08	0.0	0.0	0.0
CE	6.547E+00	2.687E+00	8.879E-04	1.369E-38	0.0	0.0	0.0	0.0
PR	7.260E+01	2.979E+01	9.844E-03	1.518E-37	0.0	0.0	0.0	0.0
PM	8.679E+00	6.664E+00	6.180E-01	2.913E-11	0.0	0.0	0.0	0.0
SM	2.906E-02	2.884E-02	2.691E-02	1.345E-02	1.317E-05	3.551E-08	3.551E-08	3.551E-08
EU	5.917E+00	5.441E+00	2.578E+00	1.923E-03	2.052E-24	0.0	0.0	0.0
SUM	6.527E+02	5.708E+02	4.168E+02	4.979E+01	9.213E-03	8.710E-03	5.125E-03	2.941E-04
TOTAL	6.527E+02	5.708E+02	4.168E+02	4.979E+01	9.213E-03	8.710E-03	5.125E-03	2.941E-04

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 18-SRS (DWPF). PHOTON SPECTRUM OF FISSION PRODUCTS IN DECAY OF SRS HLW
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

EMEAN	IMMOBILZN	PHOTONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
		18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND						
1.000E-02	1.751E+15	1.487E+15	1.059E+15	1.249E+14	2.015E+10	1.950E+10	1.479E+10	5.178E+09
2.500E-02	3.863E+14	3.236E+14	2.209E+14	2.585E+13	8.700E+09	8.200E+09	4.586E+09	6.491E+07
3.750E-02	3.907E+14	3.249E+14	2.248E+14	2.661E+13	1.467E+09	1.393E+09	8.511E+08	2.840E+07
5.750E-02	3.417E+14	2.876E+14	2.041E+14	2.397E+13	3.448E+09	3.256E+09	1.857E+09	2.954E+07
8.500E-02	2.181E+14	1.791E+14	1.227E+14	1.430E+13	8.535E+09	8.026E+09	4.342E+09	1.729E+07
1.250E-01	1.876E+14	1.392E+14	8.314E+13	9.215E+12	5.032E+08	4.750E+08	2.687E+08	3.375E+06
2.250E-01	1.835E+14	1.508E+14	1.041E+14	1.213E+13	8.045E+08	7.564E+08	4.087E+08	1.426E+06
3.750E-01	9.244E+13	7.417E+13	4.554E+13	5.238E+12	1.779E+10	1.671E+10	8.955E+09	1.750E+07
5.750E-01	1.691E+15	1.603E+15	1.268E+15	1.580E+14	3.949E+10	3.710E+10	1.988E+10	3.885E+07
8.500E-01	3.687E+13	2.932E+13	1.248E+13	8.501E+11	1.983E+09	1.862E+09	9.954E+08	1.945E+06
1.250E+00	2.116E+13	1.699E+13	7.824E+12	2.816E+11	4.798E+08	4.508E+08	2.416E+08	4.721E+05
1.750E+00	1.358E+12	9.089E+11	3.465E+11	2.172E+10	2.047E+04	1.922E+04	1.030E+04	2.013E+01
2.250E+00	2.901E+12	1.202E+12	5.235E+08	2.385E+06	1.401E-03	2.154E-04	2.154E-04	2.154E-04
2.750E+00	2.046E+10	1.003E+10	1.882E+07	1.079E-04	1.079E-04	1.079E-04	1.079E-04	1.079E-04
3.500E+00	2.349E+09	1.177E+09	2.416E+06	7.947E-05	7.947E-05	7.947E-05	7.947E-05	7.947E-05
5.000E+00	2.251E-05	2.278E-05	2.358E-05	2.366E-05	2.366E-05	2.366E-05	2.366E-05	2.366E-05
7.000E+00	1.461E-06	1.478E-06	1.530E-06	1.535E-06	1.535E-06	1.535E-06	1.535E-06	1.535E-06
9.500E+00	9.237E-08	9.346E-08	9.675E-08	9.708E-08	9.708E-08	9.708E-08	9.708E-08	9.708E-08
TOTAL	5.305E+15	4.618E+15	3.353E+15	4.013E+14	1.033E+11	9.773E+10	5.718E+10	5.382E+09

TABLE 19-SRS (DWPF). MASS OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	GRAMS							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO 59	0.0	2.741E-06	2.740E-05	2.739E-04	2.729E-03	2.625E-02	1.833E-01	3.162E-01
CO 60	1.502E-01	1.317E-01	4.031E-02	2.903E-07	0.0	0.0	0.0	0.0
NI 59	3.163E-01	3.163E-01	3.163E-01	3.160E-01	3.136E-01	2.900E-01	1.330E-01	5.459E-05
NI 60	0.0	1.851E-02	1.099E-01	1.502E-01	1.502E-01	1.502E-01	1.502E-01	1.502E-01
NI 63	4.824E-02	4.788E-02	4.474E-02	2.271E-02	2.578E-05	9.179E-35	0.0	0.0
CU 63	0.0	3.621E-04	3.501E-03	2.553E-02	4.821E-02	4.824E-02	4.824E-02	4.824E-02
SUM	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01
TOTAL	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 20-SRS (DWPF). MASS OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS

(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO	1.502E-01	1.317E-01	4.034E-02	2.742E-04	2.729E-03	2.625E-02	1.833E-01	3.162E-01
NI	3.645E-01	3.827E-01	4.709E-01	4.889E-01	4.638E-01	4.402E-01	2.832E-01	1.503E-01
CU	0.0	3.621E-04	3.501E-03	2.553E-02	4.821E-02	4.824E-02	4.824E-02	4.824E-02
SUM	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01
TOTAL	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01	5.147E-01

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 21-SRS (DWPF). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO 60	1.699E+02	1.490E+02	4.560E+01	3.283E-04	0.0	0.0	0.0	0.0
NI 59	2.397E-02	2.397E-02	2.396E-02	2.395E-02	2.376E-02	2.198E-02	1.008E-02	4.137E-06
NI 63	2.977E+00	2.954E+00	2.761E+00	1.401E+00	1.591E-03	5.664E-33	0.0	0.0
SUM	1.729E+02	1.519E+02	4.838E+01	1.426E+00	2.535E-02	2.198E-02	1.008E-02	4.137E-06
TOTAL	1.729E+02	1.519E+02	4.838E+01	1.426E+00	2.535E-02	2.198E-02	1.008E-02	4.137E-06

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 22-SRS (DWPF). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO	1.699E+02	1.490E+02	4.560E+01	3.283E-04	0.0	0.0	0.0	0.0
NI	3.001E+00	2.978E+00	2.785E+00	1.425E+00	2.535E-02	2.198E-02	1.008E-02	4.137E-06
SUM	1.729E+02	1.519E+02	4.838E+01	1.426E+00	2.535E-02	2.198E-02	1.008E-02	4.137E-06
TOTAL	1.729E+02	1.519E+02	4.838E+01	1.426E+00	2.535E-02	2.198E-02	1.008E-02	4.137E-06

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 23-SRS (DWPF). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: NUCLIDES
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO 60	2.619E+00	2.297E+00	7.030E-01	5.062E-06	0.0	0.0	0.0	0.0
NI 59	9.519E-07	9.518E-07	9.518E-07	9.510E-07	9.436E-07	8.728E-07	4.002E-07	1.643E-10
NI 63	3.000E-04	2.977E-04	2.782E-04	1.412E-04	1.603E-07	5.707E-37	0.0	0.0
SUM	2.620E+00	2.297E+00	7.033E-01	1.472E-04	1.104E-06	8.728E-07	4.002E-07	1.643E-10
TOTAL	2.620E+00	2.297E+00	7.033E-01	1.472E-04	1.104E-06	8.728E-07	4.002E-07	1.643E-10

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 24-SRS (DWPF). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW: ELEMENTS
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
CO	2.619E+00	2.297E+00	7.030E-01	5.062E-06	0.0	0.0	0.0	0.0
NI	3.009E-04	2.987E-04	2.791E-04	1.422E-04	1.104E-06	8.728E-07	4.002E-07	1.643E-10
SUM	2.620E+00	2.297E+00	7.033E-01	1.472E-04	1.104E-06	8.728E-07	4.002E-07	1.643E-10
TOTAL	2.620E+00	2.297E+00	7.033E-01	1.472E-04	1.104E-06	8.728E-07	4.002E-07	1.643E-10

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 25-SRS (DWPF). PHOTON SPECTRUM OF ACTIVATION PRODUCTS IN DECAY OF SRS HLW
(BASED ON ONE CANISTER, SLUDGE + PRECIPITATE GLASS, 1682 KG GLASS)

		PHOTONS/SEC						
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	3.337E+11	2.926E+11	8.996E+10	2.873E+08	3.254E+05	1.159E-24	0.0	0.0
2.500E-02	5.760E+10	5.051E+10	1.548E+10	1.758E+07	1.983E+04	7.060E-26	0.0	0.0
3.750E-02	3.287E+10	2.882E+10	8.828E+09	3.604E+06	4.019E+03	1.431E-26	0.0	0.0
5.750E-02	3.708E+10	3.251E+10	9.952E+09	4.298E+05	4.067E+02	1.448E-27	0.0	0.0
8.500E-02	1.458E+10	1.278E+10	3.913E+09	2.817E+04	0.0	0.0	0.0	0.0
1.250E-01	5.599E+09	4.909E+09	1.503E+09	1.082E+04	0.0	0.0	0.0	0.0
2.250E-01	1.841E+09	1.614E+09	4.942E+08	3.558E+03	0.0	0.0	0.0	0.0
3.750E-01	5.166E+08	4.529E+08	1.386E+08	9.983E+02	0.0	0.0	0.0	0.0
5.750E-01	2.966E+07	2.601E+07	7.961E+06	5.732E+01	0.0	0.0	0.0	0.0
8.500E-01	4.694E+08	4.116E+08	1.260E+08	9.072E+02	0.0	0.0	0.0	0.0
1.250E+00	1.257E+13	1.102E+13	3.373E+12	2.429E+07	0.0	0.0	0.0	0.0
1.750E+00	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
2.250E+00	6.661E+07	5.840E+07	1.788E+07	1.287E+02	0.0	0.0	0.0	0.0
2.750E+00	2.061E+05	1.807E+05	5.532E+04	3.983E-01	0.0	0.0	0.0	0.0
3.500E+00	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
5.000E+00	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
7.000E+00	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
9.500E+00	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
TOTAL	1.305E+13	1.144E+13	3.504E+12	3.332E+08	3.497E+05	1.245E-24	0.0	0.0

TABLE 1-HANF (MAX). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE 4	0.000E+00	4.591E-03	4.544E-02	4.185E-01	2.253E+00	2.950E+00	3.698E+00	1.558E+01
BI209	0.000E+00	5.003E-15	5.773E-12	5.697E-09	6.375E-06	6.577E-03	1.963E+00	8.722E+01
U233	0.000E+00	8.992E-05	9.221E-04	9.405E-03	1.161E-01	1.362E+00	1.133E+01	2.503E+01
U235	9.111E+01	9.111E+01	9.112E+01	9.118E+01	9.174E+01	9.672E+01	1.124E+02	1.136E+02
U238	1.106E+04	1.106E+04	1.106E+04	1.106E+04	1.106E+04	1.106E+04	1.106E+04	1.106E+04
NP237	2.822E+02	2.825E+02	2.849E+02	3.067E+02	4.144E+02	4.464E+02	4.335E+02	3.239E+02
PU239	2.268E+01	2.268E+01	2.267E+01	2.261E+01	2.206E+01	1.717E+01	1.299E+00	7.163E-12
AM241	1.681E+02	1.678E+02	1.655E+02	1.434E+02	3.387E+01	1.826E-05	0.000E+00	0.000E+00
SUM	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04
TOTAL	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 2-HANF (MAX). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE	0.000E+00	4.591E-03	4.544E-02	4.185E-01	2.253E+00	2.950E+00	3.698E+00	1.558E+01
PB	0.000E+00	9.035E-15	9.376E-12	5.952E-09	2.014E-06	6.566E-04	7.164E-02	1.595E+00
BI	0.000E+00	5.006E-15	5.774E-12	5.698E-09	6.375E-06	6.577E-03	1.963E+00	8.722E+01
TH	0.000E+00	2.524E-06	2.385E-05	2.421E-04	2.691E-03	4.571E-02	6.424E-01	1.617E+00
U	1.116E+04	1.116E+04	1.116E+04	1.116E+04	1.116E+04	1.117E+04	1.120E+04	1.121E+04
NP	2.822E+02	2.825E+02	2.849E+02	3.067E+02	4.144E+02	4.464E+02	4.335E+02	3.239E+02
PU	2.538E+01	2.538E+01	2.533E+01	2.517E+01	2.437E+01	1.808E+01	1.328E+00	5.718E-03
AM	1.684E+02	1.682E+02	1.659E+02	1.437E+02	3.418E+01	1.325E-01	2.827E-05	0.000E+00
CM	1.547E-01	1.487E-01	1.054E-01	3.363E-03	3.690E-18	0.000E+00	0.000E+00	0.000E+00
SUM	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04
TOTAL	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04	1.164E+04

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 3-HANF (MAX). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.000E+00	6.546E-11	5.980E-09	2.905E-07	4.129E-06	3.872E-05	2.070E-04	2.450E-04
TL209	0.000E+00	8.881E-13	9.272E-11	9.161E-09	1.049E-06	9.960E-05	2.164E-03	5.265E-03
PB209	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
PB210	0.000E+00	9.513E-14	9.005E-11	5.199E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
PB211	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
PB214	0.000E+00	9.398E-12	9.398E-10	9.377E-08	8.518E-06	3.399E-04	2.929E-03	3.838E-03
BI210	0.000E+00	9.513E-14	9.006E-11	5.199E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
BI211	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
BI213	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
BI214	0.000E+00	9.398E-12	9.398E-10	9.377E-08	8.518E-06	3.399E-04	2.929E-03	3.838E-03
PO210	0.000E+00	3.084E-14	9.006E-11	5.199E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
PO213	0.000E+00	4.023E-11	4.200E-09	4.150E-07	4.753E-05	4.512E-03	9.801E-02	2.385E-01
PO214	0.000E+00	9.396E-12	9.396E-10	9.375E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
PO215	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
PO218	0.000E+00	9.400E-12	9.400E-10	9.379E-08	8.519E-06	3.400E-04	2.929E-03	3.838E-03
AT217	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RN219	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
RN222	0.000E+00	9.400E-12	9.400E-10	9.379E-08	8.519E-06	3.400E-04	2.929E-03	3.838E-03
FR221	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RA223	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
RA225	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RA226	0.000E+00	9.400E-12	9.400E-10	9.379E-08	8.519E-06	3.400E-04	2.929E-03	3.838E-03
AC225	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
AC227	0.000E+00	6.565E-11	5.994E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
TH227	0.000E+00	6.474E-11	5.914E-09	2.873E-07	4.083E-06	3.829E-05	2.047E-04	2.423E-04
TH229	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
TH230	0.000E+00	4.340E-08	4.349E-07	4.414E-06	4.535E-05	4.368E-04	2.905E-03	3.837E-03
TH231	0.000E+00	1.970E-04	1.970E-04	1.972E-04	1.984E-04	2.092E-04	2.432E-04	2.457E-04
TH234	0.000E+00	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03
PA231	0.000E+00	4.168E-09	4.173E-08	4.166E-07	4.139E-06	3.882E-05	2.075E-04	2.457E-04
PA233	0.000E+00	1.992E-01	2.009E-01	2.163E-01	2.922E-01	3.148E-01	3.057E-01	2.284E-01
PA234M	0.000E+00	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03
U233	0.000E+00	8.708E-07	8.929E-06	9.108E-05	1.124E-03	1.319E-02	1.097E-01	2.424E-01
U234	4.821E-03	4.823E-03	4.842E-03	4.971E-03	5.093E-03	5.059E-03	4.757E-03	3.801E-03
U235	1.970E-04	1.970E-04	1.970E-04	1.972E-04	1.984E-04	2.092E-04	2.432E-04	2.457E-04
U236	4.751E-04	4.751E-04	4.752E-04	4.767E-04	4.912E-04	5.801E-04	6.342E-04	6.176E-04

TABLE 3-HANF (MAX). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES (CONTINUED)

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03
NP237	1.990E-01	1.992E-01	2.009E-01	2.163E-01	2.922E-01	3.148E-01	3.057E-01	2.284E-01
NP239	0.000E+00	6.760E-02	6.754E-02	6.697E-02	6.155E-02	2.643E-02	5.637E-06	0.000E+00
PU238	7.681E-01	7.640E-01	7.121E-01	3.498E-01	2.858E-04	0.000E+00	0.000E+00	0.000E+00
PU239	1.410E+00	1.410E+00	1.410E+00	1.406E+00	1.372E+00	1.068E+00	8.080E-02	4.454E-13
PU240	5.421E-01	5.433E-01	5.525E-01	5.700E-01	5.188E-01	1.998E-01	1.433E-05	0.000E+00
PU241	2.580E+01	2.459E+01	1.594E+01	2.094E-01	3.202E-20	0.000E+00	0.000E+00	0.000E+00
PU242	1.310E-04	1.310E-04	1.310E-04	1.310E-04	1.308E-04	1.287E-04	1.095E-04	2.184E-05
AM241	5.772E+02	5.763E+02	5.683E+02	4.924E+02	1.163E+02	6.271E-05	0.000E+00	0.000E+00
AM242	4.141E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
AM243	6.761E-02	6.760E-02	6.754E-02	6.697E-02	6.155E-02	2.643E-02	5.637E-06	0.000E+00
CM242	4.991E-01	1.061E-01	9.161E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	1.250E+01	1.204E+01	8.528E+00	2.722E-01	2.987E-16	0.000E+00	0.000E+00	0.000E+00
SUM	6.194E+02	6.163E+02	5.960E+02	4.958E+02	1.189E+02	2.022E+00	1.652E+00	2.706E+00
TOTAL	6.194E+02	6.163E+02	5.960E+02	4.958E+02	1.189E+02	2.022E+00	1.652E+00	2.706E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 4-HANF (MAX). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.000E+00	6.635E-11	6.073E-09	2.997E-07	5.178E-06	1.383E-04	2.371E-03	5.510E-03
PB	0.000E+00	1.163E-10	1.132E-08	8.612E-07	6.975E-05	5.330E-03	1.062E-01	2.517E-01
BI	0.000E+00	1.163E-10	1.132E-08	8.612E-07	6.975E-05	5.330E-03	1.062E-01	2.517E-01
PO	0.000E+00	1.249E-10	1.218E-08	9.466E-07	7.723E-05	5.570E-03	1.070E-01	2.503E-01
AT	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RN	0.000E+00	7.505E-11	6.937E-09	3.851E-07	1.266E-05	3.788E-04	3.137E-03	4.084E-03
FR	0.000E+00	4.202E-11	4.375E-09	4.282E-07	4.864E-05	4.612E-03	1.002E-01	2.438E-01
RA	0.000E+00	1.162E-10	1.123E-08	8.092E-07	6.124E-05	4.990E-03	1.033E-01	2.478E-01
AC	0.000E+00	1.068E-10	1.029E-08	7.154E-07	5.272E-05	4.650E-03	1.004E-01	2.440E-01
TH	0.000E+00	3.918E-03	3.918E-03	3.923E-03	4.017E-03	9.016E-03	1.072E-01	2.518E-01
PA	0.000E+00	2.029E-01	2.046E-01	2.200E-01	2.959E-01	3.185E-01	3.097E-01	2.324E-01
U	9.213E-03	9.820E-03	9.634E-03	9.462E-03	1.063E-02	2.275E-02	1.191E-01	2.508E-01
NP	1.990E-01	2.668E-01	2.684E-01	2.833E-01	3.538E-01	3.412E-01	3.057E-01	2.284E-01
PU	2.852E+01	2.731E+01	1.862E+01	2.536E+00	1.891E+00	1.268E+00	8.093E-02	2.184E-05
AM	5.776E+02	5.763E+02	5.684E+02	4.925E+02	1.163E+02	2.649E-02	5.637E-06	0.000E+00
CM	1.300E+01	1.214E+01	8.528E+00	2.722E-01	2.987E-16	0.000E+00	0.000E+00	0.000E+00
SUM	6.194E+02	6.163E+02	5.960E+02	4.958E+02	1.189E+02	2.022E+00	1.652E+00	2.706E+00
TOTAL	6.194E+02	6.163E+02	5.960E+02	4.958E+02	1.189E+02	2.022E+00	1.652E+00	2.706E+00

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 5-HANF (MAX). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.000E+00	1.922E-13	1.756E-11	8.531E-10	1.212E-08	1.137E-07	6.078E-07	7.195E-07
TL209	0.000E+00	1.476E-14	1.541E-12	1.522E-10	1.743E-08	1.655E-06	3.595E-05	8.748E-05
PB209	0.000E+00	4.728E-14	4.936E-12	4.877E-10	5.587E-08	5.303E-06	1.152E-04	2.803E-04
PB210	0.000E+00	2.204E-17	2.086E-14	1.204E-11	1.973E-09	7.873E-08	6.783E-07	8.888E-07
PB211	0.000E+00	1.967E-13	1.797E-11	8.729E-10	1.241E-08	1.163E-07	6.219E-07	7.363E-07
PB214	0.000E+00	2.997E-14	2.997E-12	2.991E-10	2.716E-08	1.084E-06	9.340E-06	1.224E-05
BI210	0.000E+00	2.194E-16	2.077E-13	1.199E-10	1.964E-08	7.837E-07	6.752E-06	8.847E-06
BI211	0.000E+00	2.618E-12	2.392E-10	1.162E-08	1.651E-07	1.549E-06	8.278E-06	9.801E-06
BI213	0.000E+00	1.728E-13	1.805E-11	1.783E-09	2.042E-07	1.938E-05	4.211E-04	1.025E-03
BI214	0.000E+00	1.204E-13	1.204E-11	1.202E-09	1.092E-07	4.356E-06	3.753E-05	4.918E-05
PO210	0.000E+00	9.885E-16	2.887E-12	1.667E-09	2.730E-07	1.089E-05	9.386E-05	1.230E-04
PO213	0.000E+00	2.036E-12	2.125E-10	2.100E-08	2.405E-06	2.283E-04	4.960E-03	1.207E-02
PO214	0.000E+00	4.363E-13	4.363E-11	4.353E-09	3.954E-07	1.578E-05	1.360E-04	1.782E-04
PO215	0.000E+00	2.930E-12	2.677E-10	1.300E-08	1.848E-07	1.733E-06	9.265E-06	1.097E-05
PO218	0.000E+00	3.406E-13	3.406E-11	3.399E-09	3.087E-07	1.232E-05	1.061E-04	1.391E-04
AT217	0.000E+00	1.754E-12	1.832E-10	1.810E-08	2.073E-06	1.968E-04	4.275E-03	1.040E-02
RN219	0.000E+00	2.724E-12	2.488E-10	1.209E-08	1.718E-07	1.611E-06	8.612E-06	1.020E-05
RN222	0.000E+00	3.115E-13	3.115E-11	3.108E-09	2.823E-07	1.127E-05	9.706E-05	1.272E-04
FR221	0.000E+00	1.587E-12	1.657E-10	1.637E-08	1.875E-06	1.780E-04	3.866E-03	9.408E-03
RA223	0.000E+00	2.337E-12	2.135E-10	1.037E-08	1.474E-07	1.383E-06	7.390E-06	8.749E-06
RA225	0.000E+00	2.883E-14	3.010E-12	2.974E-10	3.407E-08	3.234E-06	7.025E-05	1.709E-04
RA226	0.000E+00	2.714E-13	2.714E-11	2.708E-09	2.460E-07	9.817E-06	8.458E-05	1.108E-04
AC225	0.000E+00	1.436E-12	1.500E-10	1.482E-08	1.697E-06	1.611E-04	3.499E-03	8.515E-03
TH227	0.000E+00	2.363E-12	2.158E-10	1.049E-08	1.490E-07	1.398E-06	7.470E-06	8.844E-06
TH229	0.000E+00	1.258E-12	1.313E-10	1.298E-08	1.486E-06	1.411E-04	3.065E-03	7.457E-03
TH230	0.000E+00	1.228E-09	1.231E-08	1.249E-07	1.283E-06	1.236E-05	8.220E-05	1.086E-04
TH234	0.000E+00	1.508E-06	1.508E-06	1.508E-06	1.508E-06	1.508E-06	1.508E-06	1.508E-06
PA231	0.000E+00	1.256E-10	1.257E-09	1.255E-08	1.247E-07	1.170E-06	6.252E-06	7.403E-06
PA233	0.000E+00	4.522E-04	4.559E-04	4.910E-04	6.632E-04	7.145E-04	6.939E-04	5.185E-04
PA234M	0.000E+00	1.839E-05	1.839E-05	1.839E-05	1.839E-05	1.839E-05	1.839E-05	1.838E-05
U233	0.000E+00	2.531E-08	2.596E-07	2.648E-06	3.268E-05	3.833E-04	3.190E-03	7.046E-03
U234	1.388E-04	1.389E-04	1.395E-04	1.432E-04	1.467E-04	1.457E-04	1.370E-04	1.095E-04
U235	5.160E-06	5.160E-06	5.160E-06	5.163E-06	5.196E-06	5.477E-06	6.368E-06	6.435E-06
U236	1.287E-05	1.287E-05	1.287E-05	1.291E-05	1.331E-05	1.572E-05	1.718E-05	1.673E-05
U238	9.437E-05	9.437E-05	9.437E-05	9.437E-05	9.437E-05	9.437E-05	9.437E-05	9.435E-05
NP237	6.083E-03	6.089E-03	6.140E-03	6.611E-03	8.931E-03	9.621E-03	9.344E-03	6.981E-03

TABLE 5-HANF (MAX). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES (CONTINUED)

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
NP239	0.000E+00	1.634E-04	1.633E-04	1.619E-04	1.488E-04	6.389E-05	1.363E-08	0.000E+00
PU238	2.546E-02	2.532E-02	2.360E-02	1.159E-02	9.473E-06	1.258E-36	0.000E+00	0.000E+00
PU239	4.346E-02	4.346E-02	4.345E-02	4.334E-02	4.228E-02	3.291E-02	2.490E-03	1.373E-14
PU240	1.688E-02	1.692E-02	1.720E-02	1.775E-02	1.615E-02	6.221E-03	4.461E-07	0.000E+00
PU241	7.999E-04	7.623E-04	4.943E-04	6.493E-06	9.928E-25	0.000E+00	0.000E+00	0.000E+00
PU242	3.869E-06	3.869E-06	3.869E-06	3.868E-06	3.862E-06	3.800E-06	3.235E-06	6.450E-07
AM241	1.917E+01	1.914E+01	1.888E+01	1.636E+01	3.863E+00	2.083E-06	0.000E+00	0.000E+00
AM242	4.700E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
AM243	2.173E-03	2.173E-03	2.171E-03	2.153E-03	1.978E-03	8.496E-04	1.812E-07	0.000E+00
CM242	1.839E-02	3.909E-03	3.376E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	4.374E-01	4.210E-01	2.983E-01	9.520E-03	1.045E-17	0.000E+00	0.000E+00	0.000E+00
SUM	1.972E+01	1.966E+01	1.927E+01	1.645E+01	3.933E+00	5.208E-02	3.701E-02	6.512E-02
TOTAL	1.972E+01	1.966E+01	1.927E+01	1.645E+01	3.933E+00	5.208E-02	3.701E-02	6.512E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 6-HANF (MAX). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.000E+00	2.070E-13	1.910E-11	1.005E-09	2.956E-08	1.769E-06	3.656E-05	8.820E-05
PB	0.000E+00	2.740E-13	2.592E-11	1.672E-09	9.741E-08	6.582E-06	1.258E-04	2.942E-04
BI	0.000E+00	2.912E-12	2.695E-10	1.472E-08	4.982E-07	2.607E-05	4.737E-04	1.093E-03
PO	0.000E+00	5.752E-12	5.616E-10	4.346E-08	3.568E-06	2.690E-04	5.305E-03	1.252E-02
AT	0.000E+00	1.754E-12	1.832E-10	1.810E-08	2.073E-06	1.968E-04	4.275E-03	1.040E-02
RN	0.000E+00	3.035E-12	2.800E-10	1.520E-08	4.541E-07	1.288E-05	1.057E-04	1.374E-04
FR	0.000E+00	1.589E-12	1.659E-10	1.638E-08	1.875E-06	1.780E-04	3.866E-03	9.408E-03
RA	0.000E+00	2.638E-12	2.437E-10	1.338E-08	4.275E-07	1.443E-05	1.622E-04	2.905E-04
AC	0.000E+00	1.468E-12	1.529E-10	1.496E-08	1.699E-06	1.611E-04	3.499E-03	8.515E-03
TH	0.000E+00	1.620E-06	1.632E-06	1.767E-06	4.538E-06	1.565E-04	3.156E-03	7.576E-03
PA	0.000E+00	4.706E-04	4.744E-04	5.094E-04	6.818E-04	7.341E-04	7.186E-04	5.443E-04
U	2.512E-04	2.525E-04	2.529E-04	2.583E-04	2.923E-04	6.446E-04	3.445E-03	7.273E-03
NP	6.083E-03	6.252E-03	6.303E-03	6.773E-03	9.080E-03	9.685E-03	9.344E-03	6.981E-03
PU	8.660E-02	8.646E-02	8.475E-02	7.269E-02	5.845E-02	3.914E-02	2.494E-03	6.450E-07
AM	1.918E+01	1.915E+01	1.888E+01	1.636E+01	3.865E+00	8.517E-04	1.812E-07	0.000E+00
CM	4.558E-01	4.249E-01	2.983E-01	9.520E-03	1.045E-17	0.000E+00	0.000E+00	0.000E+00
SUM	1.972E+01	1.966E+01	1.927E+01	1.645E+01	3.933E+00	5.208E-02	3.701E-02	6.512E-02
TOTAL	1.972E+01	1.966E+01	1.927E+01	1.645E+01	3.933E+00	5.208E-02	3.701E-02	6.512E-02

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 7-HANF (MAX). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI211	0.000E+00	6.546E-11	5.980E-09	2.905E-07	4.129E-06	3.872E-05	2.070E-04	2.450E-04
BI213	0.000E+00	8.881E-13	9.272E-11	9.161E-09	1.049E-06	9.960E-05	2.164E-03	5.265E-03
PO210	0.000E+00	3.084E-14	9.006E-11	5.199E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
PO213	0.000E+00	4.023E-11	4.200E-09	4.150E-07	4.753E-05	4.512E-03	9.801E-02	2.385E-01
PO214	0.000E+00	9.396E-12	9.396E-10	9.375E-08	8.516E-06	3.399E-04	2.928E-03	3.837E-03
PO215	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
PO218	0.000E+00	9.398E-12	9.398E-10	9.377E-08	8.518E-06	3.399E-04	2.929E-03	3.838E-03
AT217	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RN219	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
RN222	0.000E+00	9.400E-12	9.400E-10	9.379E-08	8.519E-06	3.400E-04	2.929E-03	3.838E-03
FR221	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RA223	0.000E+00	6.565E-11	5.997E-09	2.913E-07	4.140E-06	3.883E-05	2.075E-04	2.457E-04
RA226	0.000E+00	9.400E-12	9.400E-10	9.379E-08	8.519E-06	3.400E-04	2.929E-03	3.838E-03
AC225	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
TH227	0.000E+00	6.474E-11	5.914E-09	2.873E-07	4.083E-06	3.829E-05	2.047E-04	2.423E-04
TH229	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
TH230	0.000E+00	4.340E-08	4.349E-07	4.414E-06	4.535E-05	4.368E-04	2.905E-03	3.837E-03
PA231	0.000E+00	4.168E-09	4.173E-08	4.166E-07	4.139E-06	3.882E-05	2.075E-04	2.457E-04
U233	0.000E+00	8.708E-07	8.929E-06	9.108E-05	1.124E-03	1.319E-02	1.097E-01	2.424E-01
U234	4.821E-03	4.823E-03	4.842E-03	4.971E-03	5.093E-03	5.059E-03	4.757E-03	3.801E-03
U235	1.970E-04	1.970E-04	1.970E-04	1.972E-04	1.984E-04	2.092E-04	2.432E-04	2.457E-04
U236	4.751E-04	4.751E-04	4.752E-04	4.767E-04	4.912E-04	5.801E-04	6.342E-04	6.176E-04
U238	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03	3.720E-03
NP237	1.990E-01	1.992E-01	2.009E-01	2.163E-01	2.922E-01	3.148E-01	3.057E-01	2.284E-01
PU238	7.681E-01	7.640E-01	7.121E-01	3.498E-01	2.858E-04	0.000E+00	0.000E+00	0.000E+00
PU239	1.410E+00	1.410E+00	1.410E+00	1.406E+00	1.372E+00	1.068E+00	8.080E-02	4.454E-13
PU240	5.421E-01	5.433E-01	5.525E-01	5.700E-01	5.188E-01	1.998E-01	1.433E-05	0.000E+00
PU242	1.310E-04	1.310E-04	1.310E-04	1.310E-04	1.308E-04	1.287E-04	1.095E-04	2.184E-05
AM241	5.772E+02	5.763E+02	5.683E+02	4.924E+02	1.163E+02	6.271E-05	0.000E+00	0.000E+00
AM243	6.761E-02	6.760E-02	6.754E-02	6.697E-02	6.155E-02	2.643E-02	5.637E-06	0.000E+00
CM242	4.991E-01	1.061E-01	9.161E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	1.250E+01	1.204E+01	8.528E+00	2.722E-01	2.987E-16	0.000E+00	0.000E+00	0.000E+00
SUM	5.932E+02	5.914E+02	5.798E+02	4.953E+02	1.185E+02	1.657E+00	1.025E+00	1.723E+00
TOTAL	5.932E+02	5.914E+02	5.798E+02	4.953E+02	1.185E+02	1.657E+00	1.025E+00	1.723E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 8-HANF (MAX). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI	0.000E+00	6.635E-11	6.073E-09	2.997E-07	5.180E-06	1.384E-04	2.371E-03	5.511E-03
PO	0.000E+00	1.249E-10	1.218E-08	9.466E-07	7.723E-05	5.570E-03	1.070E-01	2.503E-01
AT	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RN	0.000E+00	7.505E-11	6.937E-09	3.851E-07	1.266E-05	3.788E-04	3.137E-03	4.084E-03
FR	0.000E+00	4.111E-11	4.293E-09	4.241E-07	4.858E-05	4.611E-03	1.002E-01	2.438E-01
RA	0.000E+00	7.505E-11	6.937E-09	3.851E-07	1.266E-05	3.788E-04	3.137E-03	4.084E-03
AC	0.000E+00	4.202E-11	4.375E-09	4.282E-07	4.864E-05	4.612E-03	1.002E-01	2.438E-01
TH	0.000E+00	4.351E-08	4.451E-07	5.126E-06	9.801E-05	5.086E-03	1.033E-01	2.478E-01
PA	0.000E+00	4.168E-09	4.173E-08	4.166E-07	4.139E-06	3.882E-05	2.075E-04	2.457E-04
U	9.213E-03	9.216E-03	9.243E-03	9.457E-03	1.063E-02	2.275E-02	1.191E-01	2.508E-01
NP	1.990E-01	1.992E-01	2.009E-01	2.163E-01	2.922E-01	3.148E-01	3.057E-01	2.284E-01
PU	2.721E+00	2.718E+00	2.675E+00	2.326E+00	1.891E+00	1.268E+00	8.093E-02	2.184E-05
AM	5.772E+02	5.763E+02	5.684E+02	4.925E+02	1.163E+02	2.649E-02	5.637E-06	0.000E+00
CM	1.300E+01	1.214E+01	8.528E+00	2.722E-01	2.987E-16	0.000E+00	0.000E+00	0.000E+00
SUM	5.932E+02	5.914E+02	5.798E+02	4.953E+02	1.185E+02	1.657E+00	1.025E+00	1.723E+00
TOTAL	5.932E+02	5.914E+02	5.798E+02	4.953E+02	1.185E+02	1.657E+00	1.025E+00	1.723E+00

TABLE 9-HANF (MAX). (ALPHA,N) NEUTRON SOURCES IN DECAY OF HANFORD HLW
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

		NEUTRONS/SEC						
	IMMOBLZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KYR	100.0KYR	1.0MYR
PO210	0.000E-01	1.146E-09	3.346E-06	1.932E-03	3.164E-01	1.263E+01	1.088E+02	1.426E+02
PO213	0.000E-01	6.897E-06	7.201E-04	7.115E-02	8.149E+00	7.736E+02	1.680E+04	4.089E+04
PO214	0.000E-01	1.272E-06	1.272E-04	1.269E-02	1.153E+00	4.601E+01	3.963E+02	5.194E+02
PO218	0.000E-01	5.369E-07	5.369E-05	5.357E-03	4.866E-01	1.942E+01	1.673E+02	2.193E+02
AT217	0.000E-01	4.180E-06	4.365E-04	4.312E-02	4.939E+00	4.688E+02	1.019E+04	2.479E+04
RN222	0.000E-01	3.934E-07	3.934E-05	3.925E-03	3.565E-01	1.423E+01	1.226E+02	1.606E+02
FR221	0.000E-01	2.885E-06	3.013E-04	2.977E-02	3.410E+00	3.236E+02	7.033E+03	1.711E+04
RA226	0.000E-01	2.314E-07	2.314E-05	2.309E-03	2.097E-01	8.370E+00	7.211E+01	9.449E+01
AC225	0.000E-01	2.040E-06	2.130E-04	2.104E-02	2.410E+00	2.288E+02	4.972E+03	1.210E+04
TH229	0.000E-01	1.089E-06	1.137E-04	1.124E-02	1.287E+00	1.222E+02	2.655E+03	6.460E+03
TH230	0.000E-01	9.675E-04	9.695E-03	9.840E-02	1.011E+00	9.738E+00	6.476E+01	8.554E+01
U233	0.000E-01	2.218E-02	2.275E-01	2.320E+00	2.863E+01	3.360E+02	2.795E+03	6.175E+03
U234	1.186E+02	1.186E+02	1.191E+02	1.223E+02	1.253E+02	1.245E+02	1.170E+02	9.351E+01
U235	3.307E+00	3.307E+00	3.307E+00	3.310E+00	3.330E+00	3.511E+00	4.082E+00	4.124E+00
U238	5.130E+01	5.130E+01	5.130E+01	5.130E+01	5.130E+01	5.130E+01	5.130E+01	5.130E+01
NP237	4.840E+03	4.845E+03	4.886E+03	5.261E+03	7.107E+03	7.656E+03	7.435E+03	5.555E+03
PU238	3.236E+04	3.218E+04	3.000E+04	1.473E+04	1.204E+01	0.000E-01	0.000E-01	0.000E-01
PU239	4.688E+04	4.688E+04	4.688E+04	4.674E+04	4.561E+04	3.551E+04	2.686E+03	1.481E-08
PU240	1.809E+04	1.813E+04	1.844E+04	1.902E+04	1.732E+04	6.669E+03	4.783E-01	0.000E-01
PU242	3.549E+00	3.549E+00	3.549E+00	3.549E+00	3.544E+00	3.487E+00	2.967E+00	5.918E-01
AM241	2.402E+07	2.398E+07	2.365E+07	2.049E+07	4.840E+06	2.610E+00	0.000E-01	0.000E-01
CM242	3.033E+04	6.448E+03	5.567E-03	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01
CM244	6.389E+05	6.154E+05	4.359E+05	1.391E+04	1.527E-11	0.000E-01	0.000E-01	0.000E-01
CM246	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01
TOTAL	2.479E+07	2.471E+07	2.419E+07	2.059E+07	4.910E+06	5.238E+04	5.567E+04	1.144E+05

TABLE 10-HANF (MAX). SPONTANEOUS FISSION NEUTRON SOURCES IN DECAY OF HANFORD HLW GLASS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	NEUTRONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	1.404E+02	1.404E+02	1.404E+02	1.404E+02	1.404E+02	1.404E+02	1.403E+02	1.403E+02
PU238	1.192E+02	1.185E+02	1.105E+02	5.427E+01	4.435E-02	5.888E-33	0.000E+00	0.000E+00
PU240	2.165E+03	2.170E+03	2.207E+03	2.277E+03	2.072E+03	7.980E+02	5.723E-02	0.000E+00
PU242	5.782E+01	5.782E+01	5.782E+01	5.781E+01	5.772E+01	5.679E+01	4.834E+01	9.639E+00
AM241	2.086E+02	2.082E+02	2.054E+02	1.779E+02	4.202E+01	2.266E-05	0.000E+00	0.000E+00
CM242	3.251E+03	6.911E+02	5.968E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	1.718E+06	1.654E+06	1.172E+06	3.739E+04	4.104E-11	0.000E+00	0.000E+00	0.000E+00
SUM	1.724E+06	1.657E+06	1.174E+06	4.010E+04	2.314E+03	9.960E+02	1.889E+02	1.501E+02
TOTAL	1.724E+06	1.657E+06	1.174E+06	4.010E+04	2.314E+03	9.960E+02	1.889E+02	1.501E+02

TABLE 11-HANF (MAX). PHOTON SPECTRUM OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

PHOTONS/SEC								
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	4.818E+12	4.803E+12	4.718E+12	4.050E+12	9.753E+11	2.247E+10	2.498E+10	3.135E+10
2.500E-02	5.412E+11	5.402E+11	5.328E+11	4.618E+11	1.102E+11	1.466E+09	1.730E+09	1.837E+09
3.750E-02	4.466E+10	4.449E+10	4.384E+10	3.793E+10	9.124E+09	2.077E+08	1.787E+09	4.250E+09
5.750E-02	7.942E+12	7.930E+12	7.820E+12	6.776E+12	1.600E+12	1.984E+08	4.737E+08	8.884E+08
8.500E-02	8.906E+09	1.092E+10	1.087E+10	1.046E+10	8.373E+09	6.923E+09	8.863E+09	1.162E+10
1.250E-01	6.361E+09	6.875E+09	6.808E+09	6.192E+09	3.113E+09	1.583E+09	1.555E+09	2.003E+09
2.250E-01	4.092E+08	1.288E+09	1.282E+09	1.245E+09	1.061E+09	6.482E+08	1.353E+09	2.790E+09
3.750E-01	2.776E+08	3.296E+09	3.316E+09	3.508E+09	4.447E+09	4.745E+09	5.791E+09	6.425E+09
5.750E-01	1.480E+08	1.499E+08	1.478E+08	1.284E+08	3.255E+07	1.298E+07	1.417E+08	2.752E+08
8.500E-01	4.594E+07	4.700E+07	4.568E+07	3.828E+07	1.003E+07	3.581E+06	2.943E+07	5.342E+07
1.250E+00	1.240E+06	2.064E+06	1.728E+06	9.358E+05	9.957E+05	5.763E+06	5.073E+07	8.059E+07
1.750E+00	6.056E+05	7.170E+05	5.530E+05	1.668E+05	2.610E+05	6.891E+06	1.014E+08	2.130E+08
2.250E+00	3.493E+05	3.360E+05	2.411E+05	1.707E+04	2.949E+04	1.070E+06	9.219E+06	1.208E+07
2.750E+00	2.016E+05	1.939E+05	1.389E+05	9.047E+03	1.856E+03	1.877E+04	1.604E+05	2.105E+05
3.500E+00	1.807E+05	1.738E+05	1.243E+05	7.409E+03	1.162E+03	3.638E+03	3.015E+04	3.950E+04
5.000E+00	7.682E+04	7.387E+04	5.268E+04	2.752E+03	3.602E+02	5.982E+01	1.324E+01	1.240E+01
7.000E+00	8.804E+03	8.463E+03	6.020E+03	2.690E+02	2.994E+01	6.726E+00	1.425E+00	1.261E+00
9.500E+00	1.008E+03	9.686E+02	6.879E+02	2.768E+01	2.668E+00	7.631E-01	1.573E-01	1.338E-01
TOTAL	1.336E+13	1.334E+13	1.314E+13	1.135E+13	2.712E+12	3.827E+10	4.686E+10	6.180E+10

TABLE 12-HANF (MAX). MASS OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR 90	3.064E+02	2.992E+02	2.415E+02	2.835E+01	1.410E-08	0.000E+00	0.000E+00	0.000E+00
ZR 90	0.000E+00	7.208E+00	6.491E+01	2.781E+02	3.064E+02	3.064E+02	3.064E+02	3.064E+02
ZR 93	5.133E+02	5.133E+02	5.133E+02	5.132E+02	5.130E+02	5.109E+02	4.905E+02	3.263E+02
NB 93	0.000E+00	1.186E-04	1.410E-03	2.097E-02	2.302E-01	2.318E+00	2.273E+01	1.870E+02
TC 99	5.514E+02	5.514E+02	5.513E+02	5.512E+02	5.496E+02	5.337E+02	3.982E+02	2.129E+01
RU 99	0.000E+00	1.794E-03	1.794E-02	1.794E-01	1.791E+00	1.765E+01	1.532E+02	5.301E+02
PD107	7.911E+01	7.911E+01	7.911E+01	7.911E+01	7.910E+01	7.903E+01	7.827E+01	7.110E+01
AG107	0.000E+00	8.441E-06	8.441E-05	8.441E-04	8.441E-03	8.437E-02	8.396E-01	8.007E+00
SN126	1.621E+01	1.621E+01	1.621E+01	1.620E+01	1.610E+01	1.512E+01	8.105E+00	1.584E-02
TE126	0.000E+00	1.124E-04	1.123E-03	1.123E-02	1.120E-01	1.085E+00	8.105E+00	1.619E+01
CS135	2.179E+02	2.179E+02	2.179E+02	2.179E+02	2.179E+02	2.173E+02	2.115E+02	1.612E+02
BA135	0.000E+00	6.568E-05	6.568E-04	6.568E-03	6.567E-02	6.558E-01	6.470E+00	5.671E+01
CS137	5.861E+02	5.728E+02	4.652E+02	5.815E+01	5.412E-08	0.000E+00	0.000E+00	0.000E+00
BA137	0.000E+00	1.339E+01	1.209E+02	5.280E+02	5.861E+02	5.861E+02	5.861E+02	5.861E+02
CE144	9.339E+00	3.833E+00	1.266E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
ND144	0.000E+00	5.507E+00	9.338E+00	9.339E+00	9.339E+00	9.339E+00	9.339E+00	9.339E+00
PM147	4.281E+01	3.287E+01	3.049E+00	1.435E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM147	0.000E+00	9.941E+00	3.977E+01	4.281E+01	4.281E+01	4.281E+01	4.281E+01	4.281E+01
SM151	3.177E+01	3.153E+01	2.941E+01	1.471E+01	1.435E-02	1.127E-32	0.000E+00	0.000E+00
EU151	0.000E+00	2.437E-01	2.355E+00	1.706E+01	3.175E+01	3.177E+01	3.177E+01	3.177E+01
SUM	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03
TOTAL	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 13-HANF (MAX). MASS OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	3.064E+02	2.992E+02	2.415E+02	2.835E+01	1.410E-08	0.000E+00	0.000E+00	0.000E+00
ZR	5.133E+02	5.205E+02	5.782E+02	7.913E+02	8.195E+02	8.174E+02	7.970E+02	6.327E+02
NB	2.186E-03	2.277E-03	4.366E-03	2.530E-02	2.345E-01	2.322E+00	2.274E+01	1.870E+02
TC	5.514E+02	5.514E+02	5.513E+02	5.512E+02	5.496E+02	5.337E+02	3.982E+02	2.129E+01
RU	1.491E+00	7.514E-01	1.948E-02	1.794E-01	1.791E+00	1.765E+01	1.532E+02	5.301E+02
PD	7.911E+01	7.985E+01	8.060E+01	8.060E+01	8.059E+01	8.052E+01	7.976E+01	7.260E+01
AG	3.346E-04	1.299E-04	8.443E-05	8.441E-04	8.441E-03	8.437E-02	8.396E-01	8.007E+00
SN	1.621E+01	1.621E+01	1.621E+01	1.620E+01	1.610E+01	1.513E+01	8.107E+00	1.705E-02
SB	1.704E+00	1.327E+00	1.401E-01	1.697E-03	2.145E-03	2.145E-03	2.144E-03	2.144E-03
TE	2.413E-02	4.012E-01	1.589E+00	1.739E+00	1.840E+00	2.813E+00	9.832E+00	1.792E+01
CS	8.050E+02	7.914E+02	6.832E+02	2.761E+02	2.179E+02	2.173E+02	2.115E+02	1.612E+02
BA	8.959E-05	1.365E+01	1.218E+02	5.289E+02	5.871E+02	5.877E+02	5.935E+02	6.438E+02
CE	9.339E+00	3.833E+00	1.266E-03	2.965E-13	2.996E-12	2.999E-11	2.999E-10	2.999E-09
ND	0.000E+00	5.507E+00	9.338E+00	9.339E+00	9.339E+00	9.339E+00	9.340E+00	9.340E+00
PM	4.281E+01	3.287E+01	3.049E+00	1.435E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	3.177E+01	4.147E+01	6.918E+01	5.753E+01	4.284E+01	4.283E+01	4.283E+01	4.283E+01
EU	2.144E+00	2.175E+00	3.139E+00	1.706E+01	3.175E+01	3.177E+01	3.177E+01	3.177E+01
GD	3.260E-06	2.118E-01	1.355E+00	2.132E+00	2.132E+00	2.132E+00	2.132E+00	2.132E+00
SUM	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03
TOTAL	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03	2.361E+03

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 14-HANF (MAX). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

CURIES

NUCLIDE	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	3.901E-03	3.901E-03	3.900E-03	3.896E-03	3.859E-03	3.506E-03	1.342E-03	9.061E-08
SR 90	4.181E+04	4.082E+04	3.295E+04	3.868E+03	1.924E-06	0.000E+00	0.000E+00	0.000E+00
Y 90	4.181E+04	4.083E+04	3.296E+04	3.869E+03	1.924E-06	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.290E+00	1.290E+00	1.290E+00	1.290E+00	1.290E+00	1.284E+00	1.233E+00	8.201E-01
NB 93M	5.771E-01	6.093E-01	8.361E-01	1.222E+00	1.225E+00	1.220E+00	1.171E+00	7.791E-01
NB 95	5.671E+00	1.209E-01	4.004E-17	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TC 99	9.351E+00	9.351E+00	9.351E+00	9.348E+00	9.321E+00	9.052E+00	6.754E+00	3.611E-01
RU106	4.991E+03	2.509E+03	5.149E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH106	4.991E+03	2.509E+03	5.149E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD107	4.071E-02	4.071E-02	4.071E-02	4.071E-02	4.070E-02	4.066E-02	4.027E-02	3.659E-02
CD113M	1.460E+01	1.392E+01	9.080E+00	1.262E-01	3.394E-20	0.000E+00	0.000E+00	0.000E+00
SN119M	5.421E+00	1.929E+00	1.765E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SB125	1.760E+03	1.371E+03	1.441E+02	2.385E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	4.291E+02	3.344E+02	3.517E+01	5.819E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	4.601E-01	4.601E-01	4.601E-01	4.598E-01	4.569E-01	4.293E-01	2.301E-01	4.495E-04
SB126	6.481E-02	6.441E-02	6.441E-02	6.437E-02	6.397E-02	6.010E-02	3.221E-02	6.293E-05
SB126M	4.601E-01	4.601E-01	4.601E-01	4.598E-01	4.569E-01	4.293E-01	2.301E-01	4.495E-04
TE127M	3.000E+00	2.941E-01	2.457E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS134	1.200E+03	8.575E+02	4.162E+01	3.004E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	2.510E-01	2.510E-01	2.510E-01	2.510E-01	2.510E-01	2.503E-01	2.436E-01	1.857E-01
CS137	5.101E+04	4.984E+04	4.048E+04	5.060E+03	4.710E-06	0.000E+00	0.000E+00	0.000E+00
BA137M	4.821E+04	4.715E+04	3.830E+04	4.787E+03	4.455E-06	0.000E+00	0.000E+00	0.000E+00
CE144	2.980E+04	1.223E+04	4.039E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144	2.980E+04	1.223E+04	4.039E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144M	3.580E+02	1.468E+02	4.847E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM147	3.971E+04	3.049E+04	2.828E+03	1.331E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM151	8.361E+02	8.297E+02	7.741E+02	3.871E+02	3.778E-01	0.000E+00	0.000E+00	0.000E+00
EU152	2.741E+00	2.604E+00	1.646E+00	1.677E-02	2.015E-22	0.000E+00	0.000E+00	0.000E+00
EU154	3.360E+02	3.100E+02	1.501E+02	1.062E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
EU155	4.111E+02	3.574E+02	1.016E+02	3.497E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	2.975E+05	2.429E+05	1.488E+05	1.799E+04	1.349E+01	1.277E+01	9.936E+00	2.184E+00
TOTAL	2.975E+05	2.429E+05	1.488E+05	1.799E+04	1.349E+01	1.277E+01	9.936E+00	2.184E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 15-HANF (MAX). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	3.901E-03	3.901E-03	3.900E-03	3.896E-03	3.859E-03	3.506E-03	1.342E-03	9.061E-08
SR	4.181E+04	4.082E+04	3.295E+04	3.868E+03	1.924E-06	0.000E+00	0.000E+00	0.000E+00
Y	4.181E+04	4.083E+04	3.296E+04	3.869E+03	1.924E-06	0.000E+00	0.000E+00	0.000E+00
ZR	4.051E+00	1.343E+00	1.290E+00	1.290E+00	1.290E+00	1.284E+00	1.233E+00	8.201E-01
NB	6.248E+00	7.307E-01	8.361E-01	1.222E+00	1.225E+00	1.220E+00	1.171E+00	7.791E-01
TC	9.351E+00	9.351E+00	9.351E+00	9.348E+00	9.321E+00	9.052E+00	6.754E+00	3.611E-01
RU	4.991E+03	2.509E+03	5.149E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH	4.991E+03	2.509E+03	5.149E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD	4.071E-02	4.071E-02	4.071E-02	4.071E-02	4.070E-02	4.066E-02	4.027E-02	3.659E-02
SN	8.877E+00	2.901E+00	5.525E-01	4.862E-01	4.569E-01	4.293E-01	2.301E-01	4.495E-04
SB	1.761E+03	1.371E+03	1.447E+02	5.241E-01	5.209E-01	4.894E-01	2.623E-01	5.125E-04
TE	4.350E+02	3.350E+02	3.517E+01	5.819E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS	5.221E+04	5.070E+04	4.053E+04	5.060E+03	2.510E-01	2.503E-01	2.436E-01	1.857E-01
BA	4.821E+04	4.715E+04	3.830E+04	4.787E+03	4.455E-06	0.000E+00	0.000E+00	0.000E+00
CE	2.980E+04	1.223E+04	4.039E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR	3.016E+04	1.238E+04	4.088E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM	3.971E+04	3.049E+04	2.828E+03	1.331E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	8.361E+02	8.297E+02	7.741E+02	3.871E+02	3.778E-01	9.734E-07	9.734E-07	9.734E-07
EU	7.498E+02	6.701E+02	2.533E+02	1.233E-01	2.015E-22	0.000E+00	0.000E+00	0.000E+00
SUM	2.975E+05	2.429E+05	1.488E+05	1.799E+04	1.349E+01	1.277E+01	9.936E+00	2.184E+00
TOTAL	2.975E+05	2.429E+05	1.488E+05	1.799E+04	1.349E+01	1.277E+01	9.936E+00	2.184E+00

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 16-HANF (MAX). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

WATTS								
NUCLIDE	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	9.711E-07	9.711E-07	9.710E-07	9.701E-07	9.608E-07	8.728E-07	3.341E-07	2.256E-11
SR 90	4.852E+01	4.738E+01	3.824E+01	4.490E+00	2.233E-09	0.000E+00	0.000E+00	0.000E+00
Y 90	2.317E+02	2.263E+02	1.827E+02	2.145E+01	1.066E-08	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.499E-04	1.499E-04	1.499E-04	1.499E-04	1.498E-04	1.492E-04	1.433E-04	9.529E-05
NB 93M	1.023E-04	1.080E-04	1.481E-04	2.165E-04	2.171E-04	2.162E-04	2.075E-04	1.380E-04
ZR 95	1.398E-02	2.673E-04	9.135E-20	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NB 95	2.720E-02	5.800E-04	1.920E-19	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TC 99	4.689E-03	4.689E-03	4.689E-03	4.688E-03	4.674E-03	4.539E-03	3.387E-03	1.811E-04
RU106	2.967E-01	1.492E-01	3.062E-04	4.078E-31	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH106	4.786E+01	2.407E+01	4.939E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD107	2.413E-06	2.413E-06	2.413E-06	2.413E-06	2.413E-06	2.410E-06	2.387E-06	2.169E-06
AG110M	2.655E-02	9.641E-03	1.057E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CD113M	2.458E-02	2.344E-02	1.528E-02	2.124E-04	5.714E-23	0.000E+00	0.000E+00	0.000E+00
SN123	9.027E-03	1.272E-03	2.772E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SB125	5.503E+00	4.285E+00	4.506E-01	7.456E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	3.606E-01	2.811E-01	2.956E-02	4.891E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	5.738E-04	5.738E-04	5.738E-04	5.734E-04	5.698E-04	5.354E-04	2.869E-04	5.606E-07
SB126	1.197E-03	1.190E-03	1.190E-03	1.189E-03	1.182E-03	1.110E-03	5.951E-04	1.163E-06
SB126M	5.858E-03	5.858E-03	5.858E-03	5.854E-03	5.818E-03	5.466E-03	2.929E-03	5.724E-06
I129	7.541E-09	7.541E-09	7.541E-09	7.541E-09	7.541E-09	7.538E-09	7.508E-09	7.215E-09
CS134	1.221E+01	8.727E+00	4.236E-01	3.058E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	8.378E-05	8.378E-05	8.378E-05	8.378E-05	8.375E-05	8.353E-05	8.129E-05	6.198E-05
CS137	5.642E+01	5.513E+01	4.478E+01	5.597E+00	5.209E-09	0.000E+00	0.000E+00	0.000E+00
BA137M	1.893E+02	1.851E+02	1.504E+02	1.880E+01	1.749E-08	0.000E+00	0.000E+00	0.000E+00
CE144	1.977E+01	8.113E+00	2.679E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144	2.191E+02	8.991E+01	2.969E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144M	1.225E-01	5.022E-02	1.658E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM147	1.424E+01	1.094E+01	1.014E+00	4.774E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM147	0.000E+00	3.095E-09	1.238E-08	1.333E-08	1.333E-08	1.333E-08	1.333E-08	1.333E-08
SM151	9.803E-02	9.728E-02	9.077E-02	4.538E-02	4.430E-05	3.479E-35	0.000E+00	0.000E+00
EU152	2.073E-02	1.970E-02	1.245E-02	1.268E-04	1.524E-24	0.000E+00	0.000E+00	0.000E+00
EU154	3.006E+00	2.773E+00	1.343E+00	9.499E-04	2.986E-35	0.000E+00	0.000E+00	0.000E+00
EU155	2.990E-01	2.600E-01	7.389E-02	2.544E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	8.489E+02	6.637E+02	4.196E+02	5.039E+01	1.274E-02	1.210E-02	7.633E-03	4.860E-04
TOTAL	8.489E+02	6.637E+02	4.196E+02	5.039E+01	1.274E-02	1.210E-02	7.633E-03	4.860E-04

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 17-HANF (MAX). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	4.852E+01	4.738E+01	3.824E+01	4.490E+00	2.233E-09	0.000E+00	0.000E+00	0.000E+00
Y	2.317E+02	2.263E+02	1.827E+02	2.145E+01	1.066E-08	0.000E+00	0.000E+00	0.000E+00
ZR	1.413E-02	4.172E-04	1.499E-04	1.499E-04	1.498E-04	1.492E-04	1.433E-04	9.529E-05
NB	2.730E-02	6.886E-04	1.481E-04	2.165E-04	2.171E-04	2.162E-04	2.075E-04	1.380E-04
TC	4.689E-03	4.689E-03	4.689E-03	4.688E-03	4.674E-03	4.539E-03	3.387E-03	1.811E-04
RU	2.967E-01	1.492E-01	3.062E-04	4.078E-31	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH	4.786E+01	2.407E+01	4.939E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD	2.413E-06	2.413E-06	2.413E-06	2.413E-06	2.413E-06	2.410E-06	2.387E-06	2.169E-06
SN	1.262E-02	3.052E-03	7.587E-04	6.265E-04	5.698E-04	5.354E-04	2.869E-04	5.606E-07
SB	5.510E+00	4.292E+00	4.577E-01	7.043E-03	6.999E-03	6.576E-03	3.524E-03	6.887E-06
TE	3.662E-01	2.816E-01	2.956E-02	4.891E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS	6.863E+01	6.386E+01	4.520E+01	5.597E+00	8.376E-05	8.353E-05	8.129E-05	6.198E-05
BA	1.893E+02	1.851E+02	1.504E+02	1.880E+01	1.749E-08	0.000E+00	0.000E+00	0.000E+00
CE	1.977E+01	8.113E+00	2.679E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR	2.192E+02	8.996E+01	2.971E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM	1.424E+01	1.094E+01	1.014E+00	4.774E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	9.803E-02	9.728E-02	9.077E-02	4.538E-02	4.431E-05	1.333E-08	1.333E-08	1.333E-08
EU	3.326E+00	3.053E+00	1.429E+00	1.077E-03	1.524E-24	0.000E+00	0.000E+00	0.000E+00
SUM	8.489E+02	6.637E+02	4.196E+02	5.039E+01	1.274E-02	1.210E-02	7.633E-03	4.860E-04
TOTAL	8.489E+02	6.637E+02	4.196E+02	5.039E+01	1.274E-02	1.210E-02	7.633E-03	4.860E-04

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 18-HANF (MAX). PHOTON SPECTRUM OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

		PHOTONS/SEC						
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	2.309E+15	1.687E+15	9.859E+14	1.164E+14	3.239E+10	3.135E+10	2.392E+10	6.472E+09
2.500E-02	5.229E+14	3.821E+14	2.065E+14	2.396E+13	1.083E+10	1.026E+10	6.070E+09	1.552E+08
3.750E-02	5.651E+14	4.030E+14	2.269E+14	2.706E+13	2.517E+09	2.412E+09	1.610E+09	7.728E+07
5.750E-02	4.549E+14	3.264E+14	1.877E+14	2.208E+13	4.668E+09	4.438E+09	2.721E+09	8.082E+07
8.500E-02	3.046E+14	2.095E+14	1.123E+14	1.311E+13	9.287E+09	8.745E+09	4.811E+09	3.540E+07
1.250E-01	3.157E+14	1.862E+14	7.427E+13	8.409E+12	6.621E+08	6.287E+08	3.801E+08	9.253E+06
2.250E-01	2.540E+14	1.753E+14	9.410E+13	1.098E+13	8.710E+08	8.199E+08	4.496E+08	2.777E+06
3.750E-01	1.396E+14	9.581E+13	4.195E+13	4.707E+12	1.853E+10	1.741E+10	9.330E+09	1.823E+07
5.750E-01	2.064E+15	1.939E+15	1.485E+15	1.849E+14	4.114E+10	3.865E+10	2.071E+10	4.048E+07
8.500E-01	6.904E+13	4.814E+13	1.046E+13	7.595E+11	2.059E+09	1.935E+09	1.037E+09	2.026E+06
1.250E+00	2.589E+13	1.675E+13	5.162E+12	2.505E+11	4.999E+08	4.696E+08	2.517E+08	4.918E+05
1.750E+00	2.370E+12	1.273E+12	2.535E+11	1.939E+10	2.132E+04	2.002E+04	1.073E+04	2.097E+01
2.250E+00	8.661E+12	3.580E+12	1.435E+09	2.133E+06	1.078E-03	1.718E-05	1.718E-05	1.718E-05
2.750E+00	4.721E+10	2.304E+10	4.197E+07	8.606E-06	8.606E-06	8.606E-06	8.606E-06	8.606E-06
3.500E+00	5.187E+09	2.608E+09	5.352E+06	6.337E-06	6.337E-06	6.337E-06	6.337E-06	6.337E-06
5.000E+00	0.000E+00	4.381E-07	1.752E-06	1.887E-06	1.887E-06	1.887E-06	1.887E-06	1.887E-06
7.000E+00	0.000E+00	2.843E-08	1.137E-07	1.224E-07	1.224E-07	1.224E-07	1.224E-07	1.224E-07
9.500E+00	0.000E+00	1.798E-09	7.190E-09	7.742E-09	7.742E-09	7.742E-09	7.742E-09	7.742E-09
TOTAL	7.036E+15	5.474E+15	3.430E+15	4.126E+14	1.235E+11	1.171E+11	7.129E+10	6.894E+09

TABLE 19-HANF (MAX). MASS OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
MN 55	0.000E+00	1.319E-02	5.246E-02	5.638E-02	5.638E-02	5.638E-02	5.638E-02	5.638E-02
FE 55	5.638E-02	4.319E-02	3.921E-03	1.489E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 59	0.000E+00	1.555E-05	1.555E-04	1.555E-03	1.549E-02	1.490E-01	1.040E+00	1.795E+00
CO 60	3.793E-03	3.326E-03	1.018E-03	7.352E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	1.795E+00	1.795E+00	1.795E+00	1.794E+00	1.780E+00	1.646E+00	7.547E-01	3.099E-04
NI 60	0.000E+00	4.675E-04	2.775E-03	3.793E-03	3.793E-03	3.793E-03	3.793E-03	3.793E-03
NI 63	2.545E-01	2.526E-01	2.360E-01	1.198E-01	1.360E-04	4.842E-34	0.000E+00	0.000E+00
CU 63	0.000E+00	1.910E-03	1.847E-02	1.347E-01	2.543E-01	2.545E-01	2.545E-01	2.545E-01
SUM	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00
TOTAL	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 20-HANF (MAX). MASS OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

		GRAMS						
ELEMENT	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
MN	0.000E+00	1.319E-02	5.246E-02	5.638E-02	5.638E-02	5.638E-02	5.638E-02	5.638E-02
FE	5.638E-02	4.319E-02	3.921E-03	1.489E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	3.793E-03	3.341E-03	1.174E-03	1.555E-03	1.549E-02	1.490E-01	1.040E+00	1.795E+00
NI	2.050E+00	2.048E+00	2.034E+00	1.917E+00	1.784E+00	1.650E+00	7.585E-01	4.103E-03
CU	0.000E+00	1.910E-03	1.847E-02	1.347E-01	2.543E-01	2.545E-01	2.545E-01	2.545E-01
SUM	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00
TOTAL	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00	2.110E+00

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 21-HANF (MAX). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	1.410E+02	1.080E+02	9.805E+00	3.724E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	4.291E+00	3.762E+00	1.152E+00	8.316E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	1.360E-01	1.360E-01	1.360E-01	1.359E-01	1.348E-01	1.247E-01	5.719E-02	2.348E-05
NI 63	1.570E+01	1.558E+01	1.456E+01	7.392E+00	8.393E-03	0.000E+00	0.000E+00	0.000E+00
IN113M	0.000E+00	2.796E-03	7.066E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN113	2.520E-02	2.794E-03	7.061E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	1.612E+02	1.275E+02	2.566E+01	7.528E+00	1.432E-01	1.247E-01	5.719E-02	2.348E-05
TOTAL	1.612E+02	1.275E+02	2.566E+01	7.528E+00	1.432E-01	1.247E-01	5.719E-02	2.348E-05

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 22-HANF (MAX). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	1.410E+02	1.080E+02	9.805E+00	3.724E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	4.291E+00	3.762E+00	1.152E+00	8.316E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	1.584E+01	1.572E+01	1.470E+01	7.528E+00	1.432E-01	1.247E-01	5.719E-02	2.348E-05
IN	0.000E+00	2.796E-03	7.066E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN	2.520E-02	2.794E-03	7.061E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	1.612E+02	1.275E+02	2.566E+01	7.528E+00	1.432E-01	1.247E-01	5.719E-02	2.348E-05
TOTAL	1.612E+02	1.275E+02	2.566E+01	7.528E+00	1.432E-01	1.247E-01	5.719E-02	2.348E-05

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 23-HANF (MAX). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	4.765E-03	3.650E-03	3.313E-04	1.258E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	6.615E-02	5.800E-02	1.775E-02	1.282E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	5.402E-06	5.402E-06	5.402E-06	5.397E-06	5.355E-06	4.954E-06	2.271E-06	9.326E-10
NI 63	1.582E-03	1.570E-03	1.467E-03	7.449E-04	8.457E-07	3.011E-36	0.000E+00	0.000E+00
IN113M	0.000E+00	6.513E-06	1.646E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN113	4.198E-06	4.654E-07	1.176E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	7.251E-02	6.323E-02	1.956E-02	7.504E-04	6.201E-06	4.954E-06	2.271E-06	9.326E-10
TOTAL	7.251E-02	6.323E-02	1.956E-02	7.504E-04	6.201E-06	4.954E-06	2.271E-06	9.326E-10

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 24-HANF (MAX). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	4.765E-03	3.650E-03	3.313E-04	1.258E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	6.615E-02	5.800E-02	1.775E-02	1.282E-07	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	1.588E-03	1.576E-03	1.473E-03	7.503E-04	6.201E-06	4.954E-06	2.271E-06	9.326E-10
IN	0.000E+00	6.513E-06	1.646E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN	4.198E-06	4.654E-07	1.176E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	7.251E-02	6.323E-02	1.956E-02	7.504E-04	6.201E-06	4.954E-06	2.271E-06	9.326E-10
TOTAL	7.251E-02	6.323E-02	1.956E-02	7.504E-04	6.201E-06	4.954E-06	2.271E-06	9.326E-10

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 25-HANF (MAX). PHOTON SPECTRUM OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS
[BASED ON ONE CANISTER, MAXIMUM CASE (1989), 1650 KG GLASS]

EMEAN	IMMOBILZN	PHOTONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	1.164E+10	1.056E+10	5.236E+09	1.512E+09	1.717E+06	6.111E-24	0.000E+00	0.000E+00
2.500E-02	2.319E+09	1.543E+09	5.717E+08	9.215E+07	1.046E+05	3.724E-25	0.000E+00	0.000E+00
3.750E-02	8.697E+08	7.671E+08	2.596E+08	1.868E+07	2.120E+04	7.548E-26	0.000E+00	0.000E+00
5.750E-02	9.404E+08	8.249E+08	2.550E+08	1.891E+06	2.145E+03	7.636E-27	0.000E+00	0.000E+00
8.500E-02	3.682E+08	3.228E+08	9.882E+07	7.136E+02	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.250E-01	1.414E+08	1.240E+08	3.795E+07	2.741E+02	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.250E-01	6.887E+07	4.325E+07	1.248E+07	9.013E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
3.750E-01	1.305E+07	1.144E+07	3.501E+06	2.528E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.750E-01	7.491E+05	6.568E+05	2.010E+05	1.452E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
8.500E-01	1.186E+07	1.039E+07	3.182E+06	2.298E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.250E+00	3.174E+11	2.783E+11	8.519E+10	6.152E+05	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.750E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.250E+00	1.682E+06	1.475E+06	4.515E+05	3.261E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.750E+00	5.206E+03	4.564E+03	1.397E+03	1.009E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00
3.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
7.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
9.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TOTAL	3.338E+11	2.925E+11	9.167E+10	1.625E+09	1.845E+06	6.567E-24	0.000E+00	0.000E+00

TABLE 1-HANF (NOM.). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI209	0.000E+00	3.922E-15	4.521E-12	4.430E-09	4.718E-06	4.520E-03	1.329E+00	5.897E+01
U233	0.000E+00	7.048E-05	7.216E-04	7.256E-03	8.330E-02	9.263E-01	7.665E+00	1.692E+01
U235	8.834E+01	8.834E+01	8.834E+01	8.839E+01	8.886E+01	9.298E+01	1.060E+02	1.069E+02
U238	1.044E+04	1.044E+04	1.044E+04	1.044E+04	1.044E+04	1.044E+04	1.044E+04	1.044E+04
NP237	2.213E+02	2.214E+02	2.225E+02	2.333E+02	2.863E+02	3.018E+02	2.931E+02	2.190E+02
PU239	1.882E+01	1.882E+01	1.881E+01	1.876E+01	1.830E+01	1.420E+01	1.071E+00	5.904E-12
AM241	8.273E+01	8.260E+01	8.146E+01	7.058E+01	1.667E+01	8.988E-06	0.000E+00	0.000E+00
SUM	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04
TOTAL	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 2-HANF (NOM.). MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE	0.000E+00	2.256E-03	2.235E-02	2.063E-01	1.114E+00	1.495E+00	2.045E+00	1.014E+01
PB	0.000E+00	8.732E-15	9.063E-12	5.739E-09	1.908E-06	6.128E-04	6.668E-02	1.495E+00
BI	0.000E+00	3.924E-15	4.522E-12	4.430E-09	4.719E-06	4.520E-03	1.329E+00	5.897E+01
TH	0.000E+00	2.379E-06	2.246E-05	2.265E-04	2.448E-03	3.698E-02	4.757E-01	1.190E+00
U	1.053E+04	1.053E+04	1.053E+04	1.053E+04	1.053E+04	1.054E+04	1.056E+04	1.057E+04
NP	2.213E+02	2.214E+02	2.225E+02	2.333E+02	2.863E+02	3.018E+02	2.931E+02	2.190E+02
PU	2.071E+01	2.071E+01	2.067E+01	2.056E+01	1.993E+01	1.484E+01	1.088E+00	3.322E-03
AM	8.292E+01	8.279E+01	8.165E+01	7.077E+01	1.684E+01	7.431E-02	1.585E-05	0.000E+00
CM	6.221E-02	5.983E-02	4.239E-02	1.353E-03	1.483E-18	0.000E+00	0.000E+00	0.000E+00
SUM	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04
TOTAL	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04	1.086E+04

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 3-HANF (NOM.). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.000E+00	6.347E-11	5.798E-09	2.816E-07	4.001E-06	3.737E-05	1.958E-04	2.306E-04
TL209	0.000E+00	6.961E-13	7.261E-11	7.104E-09	7.672E-07	6.806E-05	1.464E-03	3.560E-03
PB209	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
PB210	0.000E+00	9.020E-14	8.534E-11	4.909E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
PB211	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
PB214	0.000E+00	8.910E-12	8.906E-10	8.848E-08	7.939E-06	3.156E-04	2.723E-03	3.614E-03
BI210	0.000E+00	9.020E-14	8.535E-11	4.909E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
BI211	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
BI213	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
BI214	0.000E+00	8.910E-12	8.906E-10	8.848E-08	7.939E-06	3.156E-04	2.723E-03	3.614E-03
PO210	0.000E+00	2.924E-14	8.535E-11	4.909E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
PO213	0.000E+00	3.153E-11	3.289E-09	3.218E-07	3.475E-05	3.083E-03	6.630E-02	1.612E-01
PO214	0.000E+00	8.908E-12	8.904E-10	8.846E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
PO215	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
PO218	0.000E+00	8.912E-12	8.907E-10	8.849E-08	7.940E-06	3.156E-04	2.723E-03	3.615E-03
AT217	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RN219	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
RN222	0.000E+00	8.912E-12	8.907E-10	8.849E-08	7.940E-06	3.156E-04	2.723E-03	3.615E-03
FR221	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RA223	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
RA225	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RA226	0.000E+00	8.912E-12	8.907E-10	8.849E-08	7.940E-06	3.156E-04	2.723E-03	3.615E-03
AC225	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
AC227	0.000E+00	6.365E-11	5.811E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
TH227	0.000E+00	6.277E-11	5.734E-09	2.785E-07	3.957E-06	3.695E-05	1.936E-04	2.280E-04
TH229	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
TH230	0.000E+00	4.115E-08	4.120E-07	4.157E-06	4.219E-05	4.055E-04	2.701E-03	3.614E-03
TH231	0.000E+00	1.910E-04	1.910E-04	1.911E-04	1.922E-04	2.011E-04	2.291E-04	2.312E-04
TH234	0.000E+00	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.510E-03
PA231	0.000E+00	4.041E-09	4.046E-08	4.039E-07	4.011E-06	3.746E-05	1.963E-04	2.312E-04
PA233	0.000E+00	1.561E-01	1.569E-01	1.645E-01	2.019E-01	2.128E-01	2.067E-01	1.544E-01
PA234M	0.000E+00	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.510E-03
U233	0.000E+00	6.825E-07	6.988E-06	7.027E-05	8.067E-04	8.970E-03	7.423E-02	1.639E-01
U234	4.571E-03	4.572E-03	4.583E-03	4.657E-03	4.727E-03	4.696E-03	4.429E-03	3.582E-03
U235	1.910E-04	1.910E-04	1.910E-04	1.911E-04	1.922E-04	2.011E-04	2.291E-04	2.312E-04
U236	4.211E-04	4.211E-04	4.212E-04	4.222E-04	4.325E-04	4.952E-04	5.331E-04	5.191E-04

TABLE 3-HANF (NOM.). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES (CONTINUED)

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.510E-03
NP237	1.560E-01	1.561E-01	1.569E-01	1.645E-01	2.019E-01	2.128E-01	2.067E-01	1.544E-01
NP239	0.000E+00	3.790E-02	3.787E-02	3.755E-02	3.451E-02	1.482E-02	3.161E-06	0.000E+00
PU238	4.431E-01	4.403E-01	4.103E-01	2.015E-01	1.647E-04	0.000E+00	0.000E+00	0.000E+00
PU239	1.170E+00	1.170E+00	1.170E+00	1.167E+00	1.138E+00	8.833E-01	6.660E-02	3.671E-13
PU240	3.931E-01	3.935E-01	3.971E-01	4.024E-01	3.661E-01	1.410E-01	1.011E-05	0.000E+00
PU241	1.260E+01	1.201E+01	7.787E+00	1.023E-01	1.564E-20	0.000E+00	0.000E+00	0.000E+00
PU242	7.611E-05	7.611E-05	7.611E-05	7.609E-05	7.597E-05	7.476E-05	6.362E-05	1.269E-05
AM241	2.840E+02	2.836E+02	2.797E+02	2.423E+02	5.722E+01	3.086E-05	0.000E+00	0.000E+00
AM242	2.210E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
AM243	3.791E-02	3.790E-02	3.787E-02	3.755E-02	3.451E-02	1.482E-02	3.161E-06	0.000E+00
CM242	1.820E-01	3.874E-02	3.346E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	5.031E+00	4.842E+00	3.431E+00	1.095E-01	1.200E-16	0.000E+00	0.000E+00	0.000E+00
SUM	3.043E+02	3.029E+02	2.933E+02	2.447E+02	5.922E+01	1.534E+00	1.141E+00	1.844E+00
TOTAL	3.043E+02	3.029E+02	2.933E+02	2.447E+02	5.922E+01	1.534E+00	1.141E+00	1.844E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 4-HANF (NOM.). RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.000E+00	6.416E-11	5.870E-09	2.887E-07	4.768E-06	1.054E-04	1.659E-03	3.790E-03
PB	0.000E+00	1.049E-10	1.015E-08	7.489E-07	5.541E-05	3.819E-03	7.340E-02	1.723E-01
BI	0.000E+00	1.049E-10	1.015E-08	7.489E-07	5.541E-05	3.819E-03	7.340E-02	1.723E-01
PO	0.000E+00	1.132E-10	1.099E-08	8.310E-07	6.259E-05	4.067E-03	7.466E-02	1.723E-01
AT	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RN	0.000E+00	7.256E-11	6.705E-09	3.709E-07	1.195E-05	3.531E-04	2.920E-03	3.846E-03
FR	0.000E+00	3.311E-11	3.442E-09	3.328E-07	3.558E-05	3.151E-03	6.776E-02	1.648E-01
RA	0.000E+00	1.048E-10	1.007E-08	6.998E-07	4.747E-05	3.504E-03	7.068E-02	1.686E-01
AC	0.000E+00	9.587E-11	9.173E-09	6.113E-07	3.953E-05	3.188E-03	6.796E-02	1.650E-01
TH	0.000E+00	3.702E-03	3.702E-03	3.706E-03	3.784E-03	7.305E-03	7.439E-02	1.724E-01
PA	0.000E+00	1.596E-01	1.605E-01	1.680E-01	2.054E-01	2.164E-01	2.104E-01	1.582E-01
U	8.693E-03	8.990E-03	8.904E-03	8.854E-03	9.669E-03	1.787E-02	8.293E-02	1.717E-01
NP	1.560E-01	1.940E-01	1.948E-01	2.021E-01	2.364E-01	2.276E-01	2.067E-01	1.544E-01
PU	1.461E+01	1.401E+01	9.764E+00	1.873E+00	1.504E+00	1.024E+00	6.667E-02	1.269E-05
AM	2.843E+02	2.836E+02	2.797E+02	2.424E+02	5.726E+01	1.485E-02	3.161E-06	0.000E+00
CM	5.213E+00	4.881E+00	3.431E+00	1.095E-01	1.200E-16	0.000E+00	0.000E+00	0.000E+00
SUM	3.043E+02	3.029E+02	2.933E+02	2.447E+02	5.922E+01	1.534E+00	1.141E+00	1.844E+00
TOTAL	3.043E+02	3.029E+02	2.933E+02	2.447E+02	5.922E+01	1.534E+00	1.141E+00	1.844E+00

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 5-HANF (NOM.). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.000E+00	1.864E-13	1.703E-11	8.270E-10	1.175E-08	1.097E-07	5.749E-07	6.771E-07
TL209	0.000E+00	1.157E-14	1.206E-12	1.180E-10	1.275E-08	1.131E-06	2.432E-05	5.914E-05
PB209	0.000E+00	3.706E-14	3.865E-12	3.782E-10	4.085E-08	3.623E-06	7.792E-05	1.895E-04
PB210	0.000E+00	2.089E-17	1.977E-14	1.137E-11	1.839E-09	7.309E-08	6.306E-07	8.371E-07
PB211	0.000E+00	1.907E-13	1.742E-11	8.463E-10	1.202E-08	1.123E-07	5.882E-07	6.928E-07
PB214	0.000E+00	2.842E-14	2.840E-12	2.822E-10	2.532E-08	1.006E-06	8.683E-06	1.153E-05
BI210	0.000E+00	2.080E-16	1.968E-13	1.132E-10	1.830E-08	7.275E-07	6.277E-06	8.332E-06
BI211	0.000E+00	2.539E-12	2.319E-10	1.127E-08	1.600E-07	1.495E-06	7.830E-06	9.222E-06
BI213	0.000E+00	1.355E-13	1.413E-11	1.383E-09	1.493E-07	1.325E-05	2.849E-04	6.928E-04
BI214	0.000E+00	1.142E-13	1.141E-11	1.134E-09	1.017E-07	4.044E-06	3.489E-05	4.632E-05
PO210	0.000E+00	9.372E-16	2.736E-12	1.574E-09	2.544E-07	1.011E-05	8.726E-05	1.158E-04
PO213	0.000E+00	1.596E-12	1.664E-10	1.628E-08	1.759E-06	1.560E-04	3.355E-03	8.159E-03
PO214	0.000E+00	4.136E-13	4.134E-11	4.107E-09	3.685E-07	1.465E-05	1.264E-04	1.678E-04
PO215	0.000E+00	2.841E-12	2.595E-10	1.261E-08	1.791E-07	1.673E-06	8.764E-06	1.032E-05
PO218	0.000E+00	3.229E-13	3.228E-11	3.207E-09	2.877E-07	1.144E-05	9.868E-05	1.310E-04
AT217	0.000E+00	1.375E-12	1.434E-10	1.403E-08	1.516E-06	1.345E-04	2.892E-03	7.032E-03
RN219	0.000E+00	2.641E-12	2.412E-10	1.172E-08	1.665E-07	1.555E-06	8.146E-06	9.594E-06
RN222	0.000E+00	2.953E-13	2.952E-11	2.932E-09	2.631E-07	1.046E-05	9.024E-05	1.198E-04
FR221	0.000E+00	1.244E-12	1.297E-10	1.269E-08	1.371E-06	1.216E-04	2.615E-03	6.360E-03
RA223	0.000E+00	2.266E-12	2.070E-10	1.006E-08	1.429E-07	1.334E-06	6.990E-06	8.233E-06
RA225	0.000E+00	2.260E-14	2.357E-12	2.306E-10	2.491E-08	2.210E-06	4.752E-05	1.156E-04
RA226	0.000E+00	2.573E-13	2.572E-11	2.555E-09	2.293E-07	9.114E-06	7.863E-05	1.044E-04
AC225	0.000E+00	1.126E-12	1.174E-10	1.149E-08	1.241E-06	1.101E-04	2.367E-03	5.757E-03
TH227	0.000E+00	2.291E-12	2.093E-10	1.017E-08	1.444E-07	1.349E-06	7.066E-06	8.322E-06
TH229	0.000E+00	9.859E-13	1.028E-10	1.006E-08	1.087E-06	9.639E-05	2.073E-03	5.041E-03
TH230	0.000E+00	1.164E-09	1.166E-08	1.176E-07	1.194E-06	1.148E-05	7.642E-05	1.023E-04
TH234	0.000E+00	1.423E-06	1.423E-06	1.423E-06	1.423E-06	1.423E-06	1.423E-06	1.423E-06
PA231	0.000E+00	1.218E-10	1.219E-09	1.217E-08	1.208E-07	1.129E-06	5.914E-06	6.966E-06
PA233	0.000E+00	3.543E-04	3.562E-04	3.734E-04	4.582E-04	4.830E-04	4.691E-04	3.505E-04
PA234M	0.000E+00	1.735E-05	1.735E-05	1.735E-05	1.735E-05	1.735E-05	1.735E-05	1.735E-05
U233	0.000E+00	1.984E-08	2.031E-07	2.043E-06	2.345E-05	2.607E-04	2.158E-03	4.763E-03
U234	1.316E-04	1.317E-04	1.320E-04	1.341E-04	1.361E-04	1.353E-04	1.276E-04	1.032E-04
U235	5.003E-06	5.003E-06	5.003E-06	5.006E-06	5.032E-06	5.266E-06	6.001E-06	6.055E-06
U236	1.141E-05	1.141E-05	1.141E-05	1.144E-05	1.172E-05	1.341E-05	1.444E-05	1.406E-05
U238	8.904E-05	8.904E-05	8.904E-05	8.904E-05	8.904E-05	8.904E-05	8.904E-05	8.903E-05
NP237	4.769E-03	4.771E-03	4.797E-03	5.029E-03	6.170E-03	6.504E-03	6.317E-03	4.720E-03

TABLE 5-HANF (NOM.). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES (CONTINUED)

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
NP239	0.000E+00	9.162E-05	9.154E-05	9.077E-05	8.341E-05	3.582E-05	7.641E-09	0.000E+00
PU238	1.468E-02	1.459E-02	1.360E-02	6.678E-03	5.457E-06	7.246E-37	0.000E+00	0.000E+00
PU239	3.606E-02	3.606E-02	3.605E-02	3.596E-02	3.507E-02	2.722E-02	2.052E-03	1.131E-14
PU240	1.224E-02	1.225E-02	1.236E-02	1.253E-02	1.140E-02	4.390E-03	3.147E-07	0.000E+00
PU241	3.907E-04	3.723E-04	2.414E-04	3.171E-06	4.849E-25	0.000E+00	0.000E+00	0.000E+00
PU242	2.248E-06	2.248E-06	2.248E-06	2.247E-06	2.244E-06	2.208E-06	1.879E-06	3.747E-07
AM241	9.436E+00	9.421E+00	9.291E+00	8.050E+00	1.901E+00	1.025E-06	0.000E+00	0.000E+00
AM242	2.509E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
AM243	1.218E-03	1.218E-03	1.217E-03	1.207E-03	1.109E-03	4.764E-04	1.016E-07	0.000E+00
CM242	6.707E-03	1.427E-03	1.233E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	1.760E-01	1.694E-01	1.200E-01	3.830E-03	4.198E-18	0.000E+00	0.000E+00	0.000E+00
SUM	9.688E+00	9.662E+00	9.480E+00	8.116E+00	1.956E+00	4.036E-02	2.564E-02	4.433E-02
TOTAL	9.688E+00	9.662E+00	9.480E+00	8.116E+00	1.956E+00	4.036E-02	2.564E-02	4.433E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 6-HANF (NOM.). THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.000E+00	1.979E-13	1.823E-11	9.451E-10	2.450E-08	1.241E-06	2.489E-05	5.982E-05
PB	0.000E+00	2.562E-13	2.415E-11	1.518E-09	8.003E-08	4.815E-06	8.782E-05	2.026E-04
BI	0.000E+00	2.789E-12	2.576E-10	1.389E-08	4.294E-07	1.951E-05	3.339E-04	7.567E-04
PO	0.000E+00	5.182E-12	5.031E-10	3.781E-08	2.849E-06	1.939E-04	3.676E-03	8.584E-03
AT	0.000E+00	1.375E-12	1.434E-10	1.403E-08	1.516E-06	1.345E-04	2.892E-03	7.032E-03
RN	0.000E+00	2.936E-12	2.708E-10	1.465E-08	4.296E-07	1.201E-05	9.838E-05	1.294E-04
FR	0.000E+00	1.246E-12	1.299E-10	1.270E-08	1.371E-06	1.216E-04	2.615E-03	6.360E-03
RA	0.000E+00	2.546E-12	2.351E-10	1.284E-08	3.970E-07	1.266E-05	1.331E-04	2.282E-04
AC	0.000E+00	1.157E-12	1.202E-10	1.163E-08	1.243E-06	1.101E-04	2.367E-03	5.757E-03
TH	0.000E+00	1.532E-06	1.543E-06	1.668E-06	3.956E-06	1.108E-04	2.158E-03	5.154E-03
PA	0.000E+00	3.718E-04	3.736E-04	3.909E-04	4.757E-04	5.015E-04	4.925E-04	3.749E-04
U	2.371E-04	2.377E-04	2.380E-04	2.417E-04	2.654E-04	5.037E-04	2.395E-03	4.976E-03
NP	4.769E-03	4.863E-03	4.888E-03	5.119E-03	6.253E-03	6.540E-03	6.317E-03	4.720E-03
PU	6.338E-02	6.328E-02	6.226E-02	5.518E-02	4.648E-02	3.161E-02	2.055E-03	3.747E-07
AM	9.437E+00	9.422E+00	9.292E+00	8.051E+00	1.902E+00	4.774E-04	1.016E-07	0.000E+00
CM	1.827E-01	1.708E-01	1.200E-01	3.830E-03	4.198E-18	0.000E+00	0.000E+00	0.000E+00
SUM	9.688E+00	9.662E+00	9.480E+00	8.116E+00	1.956E+00	4.036E-02	2.564E-02	4.433E-02
TOTAL	9.688E+00	9.662E+00	9.480E+00	8.116E+00	1.956E+00	4.036E-02	2.564E-02	4.433E-02

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 7-HANF (NOM.). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI211	0.000E+00	6.347E-11	5.798E-09	2.816E-07	4.001E-06	3.737E-05	1.958E-04	2.306E-04
BI213	0.000E+00	6.961E-13	7.261E-11	7.104E-09	7.672E-07	6.806E-05	1.464E-03	3.560E-03
PO210	0.000E+00	2.924E-14	8.535E-11	4.909E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
PO213	0.000E+00	3.153E-11	3.289E-09	3.218E-07	3.475E-05	3.083E-03	6.630E-02	1.612E-01
PO214	0.000E+00	8.908E-12	8.904E-10	8.846E-08	7.937E-06	3.155E-04	2.722E-03	3.614E-03
PO215	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
PO218	0.000E+00	8.910E-12	8.906E-10	8.848E-08	7.939E-06	3.156E-04	2.723E-03	3.614E-03
AT217	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RN219	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
RN222	0.000E+00	8.912E-12	8.907E-10	8.849E-08	7.940E-06	3.156E-04	2.723E-03	3.615E-03
FR221	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RA223	0.000E+00	6.365E-11	5.814E-09	2.824E-07	4.012E-06	3.747E-05	1.963E-04	2.312E-04
RA226	0.000E+00	8.912E-12	8.907E-10	8.849E-08	7.940E-06	3.156E-04	2.723E-03	3.615E-03
AC225	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
TH227	0.000E+00	6.277E-11	5.734E-09	2.785E-07	3.957E-06	3.695E-05	1.936E-04	2.280E-04
TH229	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
TH230	0.000E+00	4.115E-08	4.120E-07	4.157E-06	4.219E-05	4.055E-04	2.701E-03	3.614E-03
PA231	0.000E+00	4.041E-09	4.046E-08	4.039E-07	4.011E-06	3.746E-05	1.963E-04	2.312E-04
U233	0.000E+00	6.825E-07	6.988E-06	7.027E-05	8.067E-04	8.970E-03	7.423E-02	1.639E-01
U234	4.571E-03	4.572E-03	4.583E-03	4.657E-03	4.727E-03	4.696E-03	4.429E-03	3.582E-03
U235	1.910E-04	1.910E-04	1.910E-04	1.911E-04	1.922E-04	2.011E-04	2.291E-04	2.312E-04
U236	4.211E-04	4.211E-04	4.212E-04	4.222E-04	4.325E-04	4.952E-04	5.331E-04	5.191E-04
U238	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.511E-03	3.510E-03
NP237	1.560E-01	1.561E-01	1.569E-01	1.645E-01	2.019E-01	2.128E-01	2.067E-01	1.544E-01
PU238	4.431E-01	4.403E-01	4.103E-01	2.015E-01	1.647E-04	0.000E+00	0.000E+00	0.000E+00
PU239	1.170E+00	1.170E+00	1.170E+00	1.167E+00	1.138E+00	8.833E-01	6.660E-02	3.671E-13
PU240	3.931E-01	3.935E-01	3.971E-01	4.024E-01	3.661E-01	1.410E-01	1.011E-05	0.000E+00
PU242	7.611E-05	7.611E-05	7.611E-05	7.609E-05	7.597E-05	7.476E-05	6.362E-05	1.269E-05
AM241	2.840E+02	2.836E+02	2.797E+02	2.423E+02	5.722E+01	3.086E-05	0.000E+00	0.000E+00
AM243	3.791E-02	3.790E-02	3.787E-02	3.755E-02	3.451E-02	1.482E-02	3.161E-06	0.000E+00
CM242	1.820E-01	3.874E-02	3.346E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	5.031E+00	4.842E+00	3.431E+00	1.095E-01	1.200E-16	0.000E+00	0.000E+00	0.000E+00
SUM	2.915E+02	2.907E+02	2.853E+02	2.444E+02	5.898E+01	1.288E+00	7.126E-01	1.173E+00
TOTAL	2.915E+02	2.907E+02	2.853E+02	2.444E+02	5.898E+01	1.288E+00	7.126E-01	1.173E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 8-HANF (NOM.). ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI	0.000E+00	6.417E-11	5.871E-09	2.888E-07	4.770E-06	1.055E-04	1.660E-03	3.791E-03
PO	0.000E+00	1.132E-10	1.099E-08	8.310E-07	6.259E-05	4.067E-03	7.466E-02	1.723E-01
AT	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RN	0.000E+00	7.256E-11	6.705E-09	3.709E-07	1.195E-05	3.531E-04	2.920E-03	3.846E-03
FR	0.000E+00	3.223E-11	3.361E-09	3.289E-07	3.552E-05	3.151E-03	6.776E-02	1.648E-01
RA	0.000E+00	7.256E-11	6.705E-09	3.709E-07	1.195E-05	3.531E-04	2.920E-03	3.846E-03
AC	0.000E+00	3.311E-11	3.442E-09	3.328E-07	3.558E-05	3.151E-03	6.776E-02	1.648E-01
TH	0.000E+00	4.124E-08	4.211E-07	4.764E-06	8.167E-05	3.593E-03	7.065E-02	1.686E-01
PA	0.000E+00	4.041E-09	4.046E-08	4.039E-07	4.011E-06	3.746E-05	1.963E-04	2.312E-04
U	8.693E-03	8.695E-03	8.713E-03	8.852E-03	9.669E-03	1.787E-02	8.293E-02	1.717E-01
NP	1.560E-01	1.561E-01	1.569E-01	1.645E-01	2.019E-01	2.128E-01	2.067E-01	1.544E-01
PU	2.007E+00	2.004E+00	1.977E+00	1.771E+00	1.504E+00	1.024E+00	6.667E-02	1.269E-05
AM	2.841E+02	2.836E+02	2.797E+02	2.424E+02	5.726E+01	1.485E-02	3.161E-06	0.000E+00
CM	5.213E+00	4.881E+00	3.431E+00	1.095E-01	1.200E-16	0.000E+00	0.000E+00	0.000E+00
SUM	2.915E+02	2.907E+02	2.853E+02	2.444E+02	5.898E+01	1.288E+00	7.126E-01	1.173E+00
TOTAL	2.915E+02	2.907E+02	2.853E+02	2.444E+02	5.898E+01	1.288E+00	7.126E-01	1.173E+00

TABLE 9-HANF (NOM.). (ALPHA,N) NEUTRON SOURCES IN DECAY OF HANFORD HLW
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

		NEUTRONS/SEC						
	IMMOBLZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KYR	100.0KYR	1.0MYR
PO210	0.000E-01	1.086E-09	3.171E-06	1.824E-03	2.949E-01	1.172E+01	1.011E+02	1.343E+02
PO213	0.000E-01	5.405E-06	5.639E-04	5.517E-02	5.958E+00	5.285E+02	1.137E+04	2.764E+04
PO214	0.000E-01	1.206E-06	1.205E-04	1.197E-02	1.074E+00	4.270E+01	3.684E+02	4.892E+02
PO218	0.000E-01	5.090E-07	5.088E-05	5.055E-03	4.535E-01	1.803E+01	1.556E+02	2.065E+02
AT217	0.000E-01	3.277E-06	3.417E-04	3.344E-02	3.611E+00	3.203E+02	6.889E+03	1.675E+04
RN222	0.000E-01	3.729E-07	3.727E-05	3.703E-03	3.322E-01	1.321E+01	1.139E+02	1.513E+02
FR221	0.000E-01	2.262E-06	2.359E-04	2.308E-02	2.493E+00	2.211E+02	4.756E+03	1.157E+04
RA226	0.000E-01	2.193E-07	2.192E-05	2.178E-03	1.954E-01	7.766E+00	6.701E+01	8.896E+01
AC225	0.000E-01	1.599E-06	1.667E-04	1.632E-02	1.762E+00	1.563E+02	3.362E+03	8.176E+03
TH229	0.000E-01	8.537E-07	8.903E-05	8.712E-03	9.409E-01	8.346E+01	1.795E+03	4.365E+03
TH230	0.000E-01	9.167E-04	9.178E-03	9.261E-02	9.399E-01	9.033E+00	6.017E+01	8.051E+01
U233	0.000E-01	1.738E-02	1.779E-01	1.789E+00	2.054E+01	2.284E+02	1.890E+03	4.173E+03
U234	1.124E+02	1.124E+02	1.127E+02	1.145E+02	1.162E+02	1.155E+02	1.089E+02	8.807E+01
U238	4.834E+01	4.834E+01	4.834E+01	4.834E+01	4.834E+01	4.834E+01	4.834E+01	4.833E+01
NP237	3.792E+03	3.795E+03	3.814E+03	3.999E+03	4.908E+03	5.173E+03	5.025E+03	3.753E+03
PU238	1.866E+04	1.855E+04	1.728E+04	8.487E+03	6.937E+00	0.000E-01	0.000E-01	0.000E-01
PU239	3.890E+04	3.890E+04	3.890E+04	3.880E+04	3.783E+04	2.936E+04	2.214E+03	1.220E-08
PU240	1.312E+04	1.313E+04	1.325E+04	1.343E+04	1.222E+04	4.706E+03	3.374E-01	0.000E-01
PU242	2.062E+00	2.062E+00	2.062E+00	2.061E+00	2.058E+00	2.025E+00	1.723E+00	3.438E-01
AM241	1.182E+07	1.180E+07	1.164E+07	1.008E+07	2.381E+06	1.284E+00	0.000E-01	0.000E-01
CM242	1.106E+04	2.354E+03	2.033E-03	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01
CM244	2.571E+05	2.475E+05	1.754E+05	5.596E+03	6.133E-12	0.000E-01	0.000E-01	0.000E-01
CM246	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01	0.000E-01
TOTAL	1.216E+07	1.213E+07	1.189E+07	1.015E+07	2.436E+06	4.105E+04	3.832E+04	7.771E+04

TABLE 10-HANF (NOM.). SPONTANEOUS FISSION NEUTRON SOURCES IN DECAY OF HANFORD HLW GLASS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	NEUTRONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U238	1.324E+02	1.324E+02	1.324E+02	1.324E+02	1.324E+02	1.324E+02	1.324E+02	1.324E+02
PU238	6.874E+01	6.832E+01	6.366E+01	3.127E+01	2.555E-02	3.392E-33	0.000E+00	0.000E+00
PU240	1.570E+03	1.572E+03	1.586E+03	1.607E+03	1.462E+03	5.631E+02	4.037E-02	0.000E+00
PU242	3.359E+01	3.359E+01	3.359E+01	3.358E+01	3.353E+01	3.299E+01	2.808E+01	5.599E+00
AM241	1.026E+02	1.025E+02	1.011E+02	8.756E+01	2.068E+01	1.115E-05	0.000E+00	0.000E+00
CM242	1.186E+03	2.523E+02	2.179E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CM244	6.912E+05	6.652E+05	4.714E+05	1.504E+04	1.649E-11	0.000E+00	0.000E+00	0.000E+00
SUM	6.943E+05	6.674E+05	4.733E+05	1.694E+04	1.650E+03	7.292E+02	1.607E+02	1.381E+02
TOTAL	6.943E+05	6.674E+05	4.733E+05	1.694E+04	1.650E+03	7.292E+02	1.607E+02	1.381E+02

TABLE 11-HANF (NOM.). PHOTON SPECTRUM OF ACTINIDES AND DAUGHTERS IN DECAY OF HANFORD HLW GLASS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

		PHOTONS/SEC						
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	2.369E+12	2.363E+12	2.323E+12	1.998E+12	4.850E+11	1.548E+10	1.699E+10	2.128E+10
2.500E-02	2.666E+11	2.661E+11	2.625E+11	2.276E+11	5.450E+10	9.945E+08	1.178E+09	1.253E+09
3.750E-02	2.199E+10	2.191E+10	2.160E+10	1.869E+10	4.515E+09	1.360E+08	1.214E+09	2.880E+09
5.750E-02	3.909E+12	3.903E+12	3.849E+12	3.335E+12	7.876E+11	1.390E+08	3.315E+08	6.136E+08
8.500E-02	4.949E+09	6.622E+09	6.601E+09	6.398E+09	5.359E+09	4.603E+09	6.008E+09	7.872E+09
1.250E-01	3.233E+09	3.670E+09	3.637E+09	3.333E+09	1.811E+09	1.018E+09	1.055E+09	1.358E+09
2.250E-01	2.289E+08	7.435E+08	7.405E+08	7.221E+08	6.271E+08	4.064E+08	9.319E+08	1.908E+09
3.750E-01	1.380E+08	2.487E+09	2.497E+09	2.592E+09	3.053E+09	3.208E+09	3.928E+09	4.360E+09
5.750E-01	7.296E+07	7.473E+07	7.372E+07	6.418E+07	1.707E+07	1.104E+07	1.113E+08	2.071E+08
8.500E-01	2.238E+07	2.347E+07	2.288E+07	1.939E+07	5.518E+06	3.191E+06	2.394E+07	4.152E+07
1.250E+00	5.033E+05	1.306E+06	1.171E+06	8.526E+05	9.328E+05	5.219E+06	4.394E+07	6.762E+07
1.750E+00	2.458E+05	3.633E+05	2.974E+05	1.424E+05	2.358E+05	5.594E+06	7.622E+07	1.545E+08
2.250E+00	1.416E+05	1.363E+05	9.811E+04	8.029E+03	2.645E+04	9.936E+05	8.571E+06	1.138E+07
2.750E+00	8.168E+04	7.860E+04	5.646E+04	4.161E+03	1.200E+03	1.740E+04	1.491E+05	1.982E+05
3.500E+00	7.314E+04	7.037E+04	5.044E+04	3.381E+03	6.822E+02	3.350E+03	2.803E+04	3.719E+04
5.000E+00	3.105E+04	2.987E+04	2.134E+04	1.240E+03	2.075E+02	4.407E+01	1.101E+01	1.063E+01
7.000E+00	3.553E+03	3.417E+03	2.434E+03	1.191E+02	1.815E+01	4.949E+00	1.199E+00	1.110E+00
9.500E+00	4.064E+02	3.907E+02	2.778E+02	1.210E+01	1.703E+00	5.610E-01	1.332E-01	1.200E-01
TOTAL	6.574E+12	6.568E+12	6.469E+12	5.592E+12	1.342E+12	2.601E+10	3.190E+10	4.201E+10

TABLE 12-HANF (NOM.). MASS OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR 90	2.184E+02	2.133E+02	1.721E+02	2.021E+01	1.005E-08	0.000E+00	0.000E+00	0.000E+00
ZR 90	0.000E+00	5.138E+00	4.627E+01	1.982E+02	2.185E+02	2.185E+02	2.185E+02	2.185E+02
ZR 93	4.178E+02	4.178E+02	4.178E+02	4.178E+02	4.176E+02	4.159E+02	3.993E+02	2.656E+02
NB 93	0.000E+00	1.222E-04	1.354E-03	1.759E-02	1.879E-01	1.887E+00	1.850E+01	1.522E+02
TC 99	4.429E+02	4.429E+02	4.428E+02	4.427E+02	4.414E+02	4.287E+02	3.198E+02	1.710E+01
RU 99	0.000E+00	1.441E-03	1.441E-02	1.441E-01	1.439E+00	1.418E+01	1.230E+02	4.258E+02
PD107	5.870E+01	5.870E+01	5.870E+01	5.870E+01	5.870E+01	5.864E+01	5.808E+01	5.276E+01
AG107	0.000E+00	6.264E-06	6.264E-05	6.264E-04	6.263E-03	6.260E-02	6.230E-01	5.941E+00
SN126	1.286E+01	1.286E+01	1.286E+01	1.285E+01	1.277E+01	1.200E+01	6.431E+00	1.257E-02
TE126	0.000E+00	8.914E-05	8.914E-04	8.911E-03	8.884E-02	8.613E-01	6.431E+00	1.285E+01
CS135	1.754E+02	1.754E+02	1.754E+02	1.754E+02	1.753E+02	1.749E+02	1.702E+02	1.298E+02
BA135	0.000E+00	5.286E-05	5.286E-04	5.286E-03	5.285E-02	5.278E-01	5.207E+00	4.564E+01
CS137	4.149E+02	4.054E+02	3.293E+02	4.116E+01	3.831E-08	0.000E+00	0.000E+00	0.000E+00
BA137	0.000E+00	9.477E+00	8.560E+01	3.737E+02	4.149E+02	4.149E+02	4.149E+02	4.149E+02
PM147	5.619E+00	4.314E+00	4.001E-01	1.884E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM147	0.000E+00	1.305E+00	5.219E+00	5.619E+00	5.619E+00	5.619E+00	5.619E+00	5.619E+00
SM151	2.652E+01	2.632E+01	2.456E+01	1.228E+01	1.199E-02	9.413E-33	0.000E+00	0.000E+00
EU151	0.000E+00	2.035E-01	1.966E+00	1.425E+01	2.651E+01	2.652E+01	2.652E+01	2.652E+01
SUM	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03
TOTAL	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 13-HANF (NOM.). MASS OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	2.184E+02	2.133E+02	1.721E+02	2.021E+01	1.005E-08	0.000E+00	0.000E+00	0.000E+00
ZR	4.178E+02	4.229E+02	4.641E+02	6.160E+02	6.361E+02	6.344E+02	6.177E+02	4.840E+02
NB	2.179E-03	2.368E-03	4.072E-03	2.111E-02	1.914E-01	1.891E+00	1.851E+01	1.522E+02
TC	4.429E+02	4.429E+02	4.428E+02	4.427E+02	4.414E+02	4.287E+02	3.198E+02	1.710E+01
RU	1.249E-02	7.721E-03	1.442E-02	1.441E-01	1.439E+00	1.418E+01	1.230E+02	4.258E+02
PD	5.870E+01	5.871E+01	5.871E+01	5.871E+01	5.871E+01	5.865E+01	5.809E+01	5.277E+01
AG	4.672E-07	6.433E-06	6.264E-05	6.264E-04	6.263E-03	6.260E-02	6.230E-01	5.941E+00
SN	1.286E+01	1.286E+01	1.286E+01	1.285E+01	1.277E+01	1.200E+01	6.431E+00	1.257E-02
SB	2.459E-01	1.915E-01	2.031E-02	9.849E-04	1.313E-03	1.313E-03	1.312E-03	1.312E-03
TE	3.441E-03	5.798E-02	2.301E-01	2.583E-01	3.382E-01	1.111E+00	6.680E+00	1.310E+01
CS	5.904E+02	5.809E+02	5.047E+02	2.165E+02	1.753E+02	1.749E+02	1.702E+02	1.298E+02
BA	6.319E-05	9.497E+00	8.567E+01	3.738E+02	4.150E+02	4.155E+02	4.202E+02	4.606E+02
PM	5.619E+00	4.314E+00	4.001E-01	1.884E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	2.652E+01	2.763E+01	2.978E+01	1.790E+01	5.637E+00	5.625E+00	5.625E+00	5.625E+00
EU	8.396E-01	9.627E-01	2.284E+00	1.425E+01	2.651E+01	2.652E+01	2.652E+01	2.652E+01
GD	3.827E-09	8.011E-02	5.198E-01	8.336E-01	8.338E-01	8.338E-01	8.338E-01	8.338E-01
SUM	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03
TOTAL	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03	1.774E+03

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 14-HANF (NOM.). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	3.150E-03	3.150E-03	3.150E-03	3.147E-03	3.117E-03	2.832E-03	1.084E-03	7.319E-08
SR 90	2.980E+04	2.910E+04	2.349E+04	2.758E+03	1.371E-06	0.000E+00	0.000E+00	0.000E+00
Y 90	2.980E+04	2.911E+04	2.350E+04	2.759E+03	1.372E-06	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.050E+00	1.050E+00	1.050E+00	1.050E+00	1.050E+00	1.045E+00	1.004E+00	6.676E-01
NB 93M	6.161E-01	6.351E-01	7.685E-01	9.953E-01	9.972E-01	9.932E-01	9.535E-01	6.342E-01
TC 99	7.511E+00	7.511E+00	7.511E+00	7.508E+00	7.487E+00	7.270E+00	5.425E+00	2.900E-01
RU106	4.181E+01	2.102E+01	4.314E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH106	4.180E+01	2.102E+01	4.314E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD107	3.020E-02	3.020E-02	3.020E-02	3.020E-02	3.020E-02	3.017E-02	2.988E-02	2.715E-02
CD113M	8.531E+00	8.135E+00	5.305E+00	7.373E-02	1.983E-20	0.000E+00	0.000E+00	0.000E+00
SB125	2.540E+02	1.978E+02	2.080E+01	3.442E-09	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	6.201E+01	4.826E+01	5.075E+00	8.398E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	3.651E-01	3.651E-01	3.650E-01	3.648E-01	3.625E-01	3.406E-01	1.825E-01	3.567E-04
SB126	5.101E-02	5.111E-02	5.110E-02	5.107E-02	5.075E-02	4.769E-02	2.556E-02	4.994E-05
SB126M	3.650E-01	3.651E-01	3.650E-01	3.648E-01	3.625E-01	3.406E-01	1.825E-01	3.567E-04
CS134	9.311E+01	6.653E+01	3.229E+00	2.340E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	2.020E-01	2.020E-01	2.020E-01	2.020E-01	2.020E-01	2.014E-01	1.960E-01	1.495E-01
CS137	3.611E+04	3.528E+04	2.866E+04	3.582E+03	3.334E-06	0.000E+00	0.000E+00	0.000E+00
BA137M	3.400E+04	3.338E+04	2.711E+04	3.388E+03	3.154E-06	0.000E+00	0.000E+00	0.000E+00
CE144	8.001E+01	3.284E+01	1.084E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144	8.001E+01	3.284E+01	1.084E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM147	5.211E+03	4.001E+03	3.711E+02	1.747E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM151	6.981E+02	6.927E+02	6.464E+02	3.232E+02	3.155E-01	0.000E+00	0.000E+00	0.000E+00
EU152	1.400E+00	1.331E+00	8.412E-01	8.567E-03	1.029E-22	0.000E+00	0.000E+00	0.000E+00
EU154	1.450E+02	1.338E+02	6.477E+01	4.583E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00
EU155	1.370E+02	1.191E+02	3.387E+01	1.166E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	1.366E+05	1.323E+05	1.039E+05	1.282E+04	1.086E+01	1.027E+01	7.999E+00	1.769E+00
TOTAL	1.366E+05	1.323E+05	1.039E+05	1.282E+04	1.086E+01	1.027E+01	7.999E+00	1.769E+00

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 15-HANF (NOM.). RADIOACTIVITY OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	CURIES							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	3.150E-03	3.150E-03	3.150E-03	3.147E-03	3.117E-03	2.832E-03	1.084E-03	7.319E-08
SR	2.980E+04	2.910E+04	2.349E+04	2.758E+03	1.371E-06	0.000E+00	0.000E+00	0.000E+00
Y	2.980E+04	2.911E+04	2.350E+04	2.759E+03	1.372E-06	0.000E+00	0.000E+00	0.000E+00
ZR	1.050E+00	1.050E+00	1.050E+00	1.050E+00	1.050E+00	1.045E+00	1.004E+00	6.676E-01
NB	6.161E-01	6.351E-01	7.685E-01	9.953E-01	9.972E-01	9.932E-01	9.535E-01	6.342E-01
TC	7.511E+00	7.511E+00	7.511E+00	7.508E+00	7.487E+00	7.270E+00	5.425E+00	2.900E-01
RU	4.181E+01	2.102E+01	4.314E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
RH	4.180E+01	2.102E+01	4.314E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD	3.020E-02	3.020E-02	3.020E-02	3.020E-02	3.020E-02	3.017E-02	2.988E-02	2.715E-02
SN	4.495E-01	4.440E-01	4.326E-01	3.842E-01	3.625E-01	3.406E-01	1.825E-01	3.567E-04
SB	2.545E+02	1.982E+02	2.122E+01	4.159E-01	4.133E-01	3.883E-01	2.081E-01	4.066E-04
TE	6.201E+01	4.826E+01	5.075E+00	8.398E-10	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS	3.620E+04	3.535E+04	2.866E+04	3.582E+03	2.020E-01	2.014E-01	1.960E-01	1.495E-01
BA	3.400E+04	3.338E+04	2.711E+04	3.388E+03	3.154E-06	0.000E+00	0.000E+00	0.000E+00
CE	8.001E+01	3.284E+01	1.084E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR	8.097E+01	3.323E+01	1.097E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM	5.211E+03	4.001E+03	3.711E+02	1.747E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	6.981E+02	6.927E+02	6.464E+02	3.232E+02	3.155E-01	1.277E-07	1.277E-07	1.277E-07
EU	2.834E+02	2.543E+02	9.948E+01	5.451E-02	1.029E-22	0.000E+00	0.000E+00	0.000E+00
SUM	1.366E+05	1.323E+05	1.039E+05	1.282E+04	1.086E+01	1.027E+01	7.999E+00	1.769E+00
TOTAL	1.366E+05	1.323E+05	1.039E+05	1.282E+04	1.086E+01	1.027E+01	7.999E+00	1.769E+00

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 16-HANF (NOM.). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	7.843E-07	7.843E-07	7.843E-07	7.835E-07	7.760E-07	7.050E-07	2.698E-07	1.822E-11
SR 90	3.459E+01	3.378E+01	2.726E+01	3.201E+00	1.592E-09	0.000E+00	0.000E+00	0.000E+00
Y 90	1.652E+02	1.613E+02	1.302E+02	1.529E+01	7.603E-09	0.000E+00	0.000E+00	0.000E+00
ZR 93	1.220E-04	1.220E-04	1.220E-04	1.220E-04	1.220E-04	1.215E-04	1.166E-04	7.756E-05
NB 93M	1.092E-04	1.125E-04	1.362E-04	1.763E-04	1.767E-04	1.760E-04	1.689E-04	1.124E-04
TC 99	3.767E-03	3.767E-03	3.766E-03	3.765E-03	3.754E-03	3.646E-03	2.720E-03	1.454E-04
RH106	4.009E-01	2.016E-01	4.137E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD107	1.790E-06	1.790E-06	1.790E-06	1.790E-06	1.790E-06	1.789E-06	1.771E-06	1.609E-06
CD113M	1.436E-02	1.369E-02	8.930E-03	1.241E-04	3.338E-23	0.000E+00	0.000E+00	0.000E+00
SB125	7.942E-01	6.184E-01	6.503E-02	1.076E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TE125M	5.212E-02	4.056E-02	4.266E-03	7.059E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SN126	4.553E-04	4.553E-04	4.553E-04	4.550E-04	4.521E-04	4.248E-04	2.277E-04	4.449E-07
SB126	9.424E-04	9.443E-04	9.442E-04	9.436E-04	9.378E-04	8.811E-04	4.722E-04	9.227E-07
SB126M	4.648E-03	4.648E-03	4.648E-03	4.645E-03	4.616E-03	4.337E-03	2.324E-03	4.542E-06
I129	5.968E-09	5.968E-09	5.968E-09	5.968E-09	5.968E-09	5.965E-09	5.942E-09	5.710E-09
CS134	9.476E-01	6.771E-01	3.286E-02	2.382E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS135	6.742E-05	6.742E-05	6.742E-05	6.742E-05	6.740E-05	6.722E-05	6.542E-05	4.988E-05
CS137	3.994E+01	3.902E+01	3.170E+01	3.962E+00	3.688E-09	0.000E+00	0.000E+00	0.000E+00
BA137M	1.335E+02	1.310E+02	1.064E+02	1.330E+01	1.238E-08	0.000E+00	0.000E+00	0.000E+00
CE144	5.307E-02	2.178E-02	7.193E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR144	5.881E-01	2.414E-01	7.971E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM147	1.869E+00	1.435E+00	1.331E-01	6.267E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM151	8.185E-02	8.122E-02	7.578E-02	3.789E-02	3.699E-05	2.905E-35	0.000E+00	0.000E+00
EU152	1.059E-02	1.007E-02	6.362E-03	6.480E-05	7.785E-25	0.000E+00	0.000E+00	0.000E+00
EU154	1.297E+00	1.197E+00	5.794E-01	4.099E-04	1.289E-35	0.000E+00	0.000E+00	0.000E+00
EU155	9.965E-02	8.666E-02	2.463E-02	8.479E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SUM	3.794E+02	3.698E+02	2.966E+02	3.580E+01	1.017E-02	9.656E-03	6.097E-03	3.928E-04
TOTAL	3.794E+02	3.698E+02	2.966E+02	3.580E+01	1.017E-02	9.656E-03	6.097E-03	3.928E-04

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 17-HANF (NOM.). THERMAL POWER OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SR	3.459E+01	3.378E+01	2.726E+01	3.201E+00	1.592E-09	0.000E+00	0.000E+00	0.000E+00
Y	1.652E+02	1.613E+02	1.302E+02	1.529E+01	7.603E-09	0.000E+00	0.000E+00	0.000E+00
ZR	1.220E-04	1.220E-04	1.220E-04	1.220E-04	1.220E-04	1.215E-04	1.166E-04	7.756E-05
NB	1.092E-04	1.125E-04	1.362E-04	1.763E-04	1.767E-04	1.760E-04	1.689E-04	1.124E-04
TC	3.767E-03	3.767E-03	3.766E-03	3.765E-03	3.754E-03	3.646E-03	2.720E-03	1.454E-04
RH	4.009E-01	2.016E-01	4.137E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PD	1.790E-06	1.790E-06	1.790E-06	1.790E-06	1.790E-06	1.789E-06	1.771E-06	1.609E-06
SN	6.144E-04	6.099E-04	5.906E-04	4.938E-04	4.521E-04	4.248E-04	2.277E-04	4.449E-07
SB	7.998E-01	6.240E-01	7.062E-02	5.589E-03	5.554E-03	5.218E-03	2.796E-03	5.464E-06
TE	5.212E-02	4.056E-02	4.266E-03	7.059E-13	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CS	4.088E+01	3.970E+01	3.173E+01	3.962E+00	6.741E-05	6.722E-05	6.542E-05	4.988E-05
BA	1.335E+02	1.310E+02	1.064E+02	1.330E+01	1.238E-08	0.000E+00	0.000E+00	0.000E+00
CE	5.307E-02	2.178E-02	7.193E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PR	5.884E-01	2.415E-01	7.975E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
PM	1.869E+00	1.435E+00	1.331E-01	6.267E-12	0.000E+00	0.000E+00	0.000E+00	0.000E+00
SM	8.185E-02	8.122E-02	7.578E-02	3.789E-02	3.699E-05	1.749E-09	1.749E-09	1.749E-09
EU	1.407E+00	1.293E+00	6.104E-01	4.748E-04	7.785E-25	0.000E+00	0.000E+00	0.000E+00
SUM	3.794E+02	3.698E+02	2.966E+02	3.580E+01	1.017E-02	9.656E-03	6.097E-03	3.928E-04
TOTAL	3.794E+02	3.698E+02	2.966E+02	3.580E+01	1.017E-02	9.656E-03	6.097E-03	3.928E-04

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 18-HANF (NOM.). PHOTON SPECTRUM OF FISSION PRODUCTS IN DECAY OF HANFORD HLW GLASS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

		PHOTONS/SEC						
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	8.969E+14	8.725E+14	6.992E+14	8.289E+13	2.604E+10	2.521E+10	1.927E+10	5.259E+09
2.500E-02	1.908E+14	1.844E+14	1.446E+14	1.707E+13	8.623E+09	8.164E+09	4.837E+09	1.247E+08
3.750E-02	2.058E+14	2.003E+14	1.598E+14	1.922E+13	2.012E+09	1.928E+09	1.288E+09	6.212E+07
5.750E-02	1.706E+14	1.660E+14	1.333E+14	1.573E+13	3.720E+09	3.537E+09	2.171E+09	6.491E+07
8.500E-02	1.026E+14	9.968E+13	7.946E+13	9.341E+12	7.375E+09	6.944E+09	3.822E+09	2.836E+07
1.250E-01	6.820E+13	6.601E+13	5.197E+13	5.991E+12	5.274E+08	5.008E+08	3.030E+08	7.429E+06
2.250E-01	8.580E+13	8.341E+13	6.672E+13	7.828E+12	6.916E+08	6.510E+08	3.571E+08	2.223E+06
3.750E-01	3.980E+13	3.807E+13	2.877E+13	3.357E+12	1.470E+10	1.381E+10	7.403E+09	1.447E+07
5.750E-01	1.324E+15	1.297E+15	1.048E+15	1.309E+14	3.264E+10	3.067E+10	1.644E+10	3.212E+07
8.500E-01	1.149E+13	1.025E+13	5.844E+12	5.410E+11	1.634E+09	1.535E+09	8.227E+08	1.608E+06
1.250E+00	5.026E+12	4.669E+12	2.789E+12	1.780E+11	3.966E+08	3.726E+08	1.997E+08	3.902E+05
1.750E+00	2.440E+11	2.286E+11	1.553E+11	1.380E+10	1.692E+04	1.589E+04	8.514E+03	1.664E+01
2.250E+00	2.481E+10	1.040E+10	1.835E+07	1.520E+06	7.583E-04	2.254E-06	2.254E-06	2.254E-06
2.750E+00	3.526E+08	1.754E+08	3.458E+05	1.129E-06	1.129E-06	1.129E-06	1.129E-06	1.129E-06
3.500E+00	4.345E+07	2.185E+07	4.484E+04	8.317E-07	8.317E-07	8.317E-07	8.317E-07	8.317E-07
5.000E+00	0.000E+00	5.749E-08	2.300E-07	2.476E-07	2.476E-07	2.476E-07	2.476E-07	2.476E-07
7.000E+00	0.000E+00	3.730E-09	1.492E-08	1.607E-08	1.607E-08	1.607E-08	1.607E-08	1.607E-08
9.500E+00	0.000E+00	2.359E-10	9.436E-10	1.016E-09	1.016E-09	1.016E-09	1.016E-09	1.016E-09
TOTAL	3.101E+15	3.023E+15	2.421E+15	2.930E+14	9.837E+10	9.332E+10	5.691E+10	5.598E+09

TABLE 19-HANF (NOM.). MASS OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
MN 55	0.000E+00	1.684E-03	6.697E-03	7.198E-03	7.198E-03	7.198E-03	7.198E-03	7.198E-03
FE 55	7.198E-03	5.514E-03	5.005E-04	1.902E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 59	0.000E+00	1.247E-05	1.246E-04	1.246E-03	1.241E-02	1.194E-01	8.338E-01	1.438E+00
NI 59	1.439E+00	1.439E+00	1.439E+00	1.437E+00	1.426E+00	1.319E+00	6.049E-01	2.484E-04
NI 63	1.961E-01	1.946E-01	1.819E-01	9.232E-02	1.048E-04	3.731E-34	0.000E+00	0.000E+00
CU 63	0.000E+00	1.472E-03	1.423E-02	1.038E-01	1.960E-01	1.961E-01	1.961E-01	1.961E-01
SUM	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00
TOTAL	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 20-HANF (NOM.). MASS OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
MN	0.000E+00	1.684E-03	6.697E-03	7.198E-03	7.198E-03	7.198E-03	7.198E-03	7.198E-03
FE	7.198E-03	5.514E-03	5.005E-04	1.902E-14	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	1.326E-03	1.175E-03	4.806E-04	1.246E-03	1.241E-02	1.194E-01	8.338E-01	1.438E+00
NI	1.635E+00	1.633E+00	1.621E+00	1.531E+00	1.428E+00	1.321E+00	6.062E-01	1.575E-03
CU	0.000E+00	1.472E-03	1.423E-02	1.038E-01	1.960E-01	1.961E-01	1.961E-01	1.961E-01
SUM	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00
TOTAL	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00	1.643E+00

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 21-HANF (NOM.). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	1.800E+01	1.379E+01	1.252E+00	4.756E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	1.500E+00	1.315E+00	4.026E-01	2.929E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	1.090E-01	1.090E-01	1.090E-01	1.089E-01	1.081E-01	9.997E-02	4.583E-02	1.882E-05
NI 63	1.210E+01	1.201E+01	1.122E+01	5.697E+00	6.468E-03	0.000E+00	0.000E+00	0.000E+00
SUM	3.171E+01	2.722E+01	1.299E+01	5.806E+00	1.145E-01	9.997E-02	4.583E-02	1.882E-05
TOTAL	3.171E+01	2.722E+01	1.299E+01	5.806E+00	1.145E-01	9.997E-02	4.583E-02	1.882E-05

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 22-HANF (NOM.). RADIOACTIVITY OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	1.800E+01	1.379E+01	1.252E+00	4.756E-11	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	1.500E+00	1.315E+00	4.026E-01	2.929E-06	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	1.221E+01	1.212E+01	1.133E+01	5.806E+00	1.145E-01	9.997E-02	4.583E-02	1.882E-05
SUM	3.171E+01	2.722E+01	1.299E+01	5.806E+00	1.145E-01	9.997E-02	4.583E-02	1.882E-05
TOTAL	3.171E+01	2.722E+01	1.299E+01	5.806E+00	1.145E-01	9.997E-02	4.583E-02	1.882E-05

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 23-HANF (NOM.). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: NUCLIDES
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE 55	6.083E-04	4.659E-04	4.229E-05	1.607E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO 60	2.313E-02	2.028E-02	6.208E-03	4.516E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI 59	4.330E-06	4.330E-06	4.329E-06	4.326E-06	4.292E-06	3.970E-06	1.820E-06	7.475E-10
NI 63	1.219E-03	1.210E-03	1.131E-03	5.741E-04	6.518E-07	2.320E-36	0.000E+00	0.000E+00
SUM	2.496E-02	2.196E-02	7.385E-03	5.785E-04	4.944E-06	3.970E-06	1.820E-06	7.475E-10
TOTAL	2.496E-02	2.196E-02	7.385E-03	5.785E-04	4.944E-06	3.970E-06	1.820E-06	7.475E-10

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 24-HANF (NOM.). THERMAL POWER OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS: ELEMENTS
[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
FE	6.083E-04	4.659E-04	4.229E-05	1.607E-15	0.000E+00	0.000E+00	0.000E+00	0.000E+00
CO	2.313E-02	2.028E-02	6.208E-03	4.516E-08	0.000E+00	0.000E+00	0.000E+00	0.000E+00
NI	1.224E-03	1.215E-03	1.135E-03	5.784E-04	4.944E-06	3.970E-06	1.820E-06	7.475E-10
SUM	2.496E-02	2.196E-02	7.385E-03	5.785E-04	4.944E-06	3.970E-06	1.820E-06	7.475E-10
TOTAL	2.496E-02	2.196E-02	7.385E-03	5.785E-04	4.944E-06	3.970E-06	1.820E-06	7.475E-10

ELEMENTS CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 25-HANF (NOM.). PHOTON SPECTRUM OF ACTIVATION PRODUCTS IN DECAY OF HANFORD HLW GLASS

[BASED ON ONE CANISTER, NOMINAL CASE (1989), 1650 KG GLASS]

		PHOTONS/SEC						
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	5.417E+09	5.035E+09	3.085E+09	1.165E+09	1.323E+06	4.710E-24	0.000E+00	0.000E+00
2.500E-02	6.592E+08	5.954E+08	2.763E+08	7.102E+07	8.063E+04	2.870E-25	0.000E+00	0.000E+00
3.750E-02	3.208E+08	2.848E+08	1.062E+08	1.439E+07	1.634E+04	5.817E-26	0.000E+00	0.000E+00
5.750E-02	3.305E+08	2.901E+08	9.074E+07	1.457E+06	1.653E+03	5.885E-27	0.000E+00	0.000E+00
8.500E-02	1.287E+08	1.129E+08	3.455E+07	2.513E+02	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.250E-01	4.944E+07	4.335E+07	1.327E+07	9.653E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.250E-01	1.626E+07	1.426E+07	4.364E+06	3.174E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
3.750E-01	4.562E+06	3.999E+06	1.224E+06	8.905E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.750E-01	2.619E+05	2.296E+05	7.030E+04	5.113E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00
8.500E-01	4.145E+06	3.634E+06	1.113E+06	8.093E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.250E+00	1.110E+11	9.731E+10	2.979E+10	2.167E+05	0.000E+00	0.000E+00	0.000E+00	0.000E+00
1.750E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.250E+00	5.882E+05	5.157E+05	1.579E+05	1.148E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
2.750E+00	1.820E+03	1.596E+03	4.885E+02	3.553E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00
3.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
5.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
7.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
9.500E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
TOTAL	1.179E+11	1.037E+11	3.340E+10	1.252E+09	1.422E+06	5.061E-24	0.000E+00	0.000E+00

TABLE 1-INEL. MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	GRAMS							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE 4	0.0	7.225E-04	7.045E-03	5.416E-02	1.317E-01	2.302E-01	4.872E-01	1.071E+00
PB206	0.0	2.388E-20	8.592E-15	5.718E-10	5.029E-06	3.484E-03	3.874E-01	4.147E+00
BI209	0.0	2.797E-17	4.717E-15	3.030E-12	1.407E-08	3.124E-05	1.031E-02	4.614E-01
TH230	0.0	5.659E-08	5.506E-06	4.413E-04	1.243E-02	1.332E-01	8.036E-01	1.360E-01
TH232	0.0	5.747E-09	5.797E-08	6.293E-07	1.109E-05	4.580E-04	1.002E-02	1.076E-01
U233	1.583E-07	1.861E-07	4.565E-07	6.921E-06	3.818E-04	6.926E-03	5.979E-02	1.325E-01
U234	8.785E-05	4.048E-02	3.900E-01	2.803E+00	5.119E+00	4.992E+00	3.868E+00	3.016E-01
U235	1.063E+00	1.063E+00	1.067E+00	1.104E+00	1.464E+00	4.604E+00	1.445E+01	1.523E+01
U236	1.973E-01	1.977E-01	2.011E-01	2.351E-01	5.584E-01	2.543E+00	3.776E+00	3.677E+00
U238	3.797E-05	3.903E-05	4.857E-05	1.440E-04	1.097E-03	1.055E-02	9.711E-02	4.932E-01
NP237	8.693E-02	8.754E-02	9.866E-02	3.685E-01	1.897E+00	2.363E+00	2.295E+00	1.715E+00
PU238	5.221E+00	5.180E+00	4.825E+00	2.370E+00	1.936E-03	2.571E-34	0.0	0.0
PU239	1.437E+01	1.437E+01	1.437E+01	1.433E+01	1.397E+01	1.080E+01	8.105E-01	4.468E-12
PU240	3.642E+00	3.642E+00	3.641E+00	3.611E+00	3.283E+00	1.264E+00	9.084E-05	0.0
PU241	1.983E+00	1.890E+00	1.225E+00	1.610E-02	2.463E-21	0.0	0.0	0.0
PU242	6.018E-01	6.018E-01	6.018E-01	6.017E-01	6.007E-01	5.911E-01	5.031E-01	1.003E-01
AM241	3.385E-01	4.311E-01	1.084E+00	2.019E+00	4.807E-01	2.592E-07	0.0	0.0
AM243	5.315E-02	5.315E-02	5.310E-02	5.265E-02	4.839E-02	2.078E-02	4.432E-06	8.638E-43
SUM	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.759E+01
TOTAL	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.759E+01

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 2-INEL. MASS OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	GRAMS							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
HE	0.0	7.225E-04	7.045E-03	5.416E-02	1.317E-01	2.302E-01	4.872E-01	1.071E+00
PB	0.0	1.073E-16	2.582E-13	1.472E-09	5.599E-06	3.513E-03	3.881E-01	4.159E+00
BI	0.0	2.800E-17	4.816E-15	3.553E-12	1.442E-08	3.126E-05	1.031E-02	4.614E-01
RA	0.0	1.673E-13	1.633E-10	1.367E-07	4.327E-05	2.114E-03	1.655E-02	2.779E-03
TH	0.0	6.235E-08	5.564E-06	4.420E-04	1.244E-02	1.338E-01	8.161E-01	2.497E-01
PA	0.0	4.003E-09	1.367E-08	1.173E-07	1.277E-06	2.608E-05	4.977E-04	6.970E-04
U	1.260E+00	1.302E+00	1.658E+00	4.142E+00	7.143E+00	1.216E+01	2.225E+01	1.983E+01
NP	8.693E-02	8.754E-02	9.866E-02	3.685E-01	1.897E+00	2.363E+00	2.295E+00	1.715E+00
PU	2.582E+01	2.568E+01	2.466E+01	2.093E+01	1.785E+01	1.266E+01	1.314E+00	1.003E-01
AM	3.916E-01	4.842E-01	1.137E+00	2.072E+00	5.291E-01	2.078E-02	4.432E-06	8.638E-43
CM	8.452E-03	7.946E-03	5.593E-03	1.785E-04	1.956E-19	0.0	0.0	0.0
SUM	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.759E+01
TOTAL	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.757E+01	2.759E+01

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 3-INEL. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.0	7.638E-13	6.986E-11	3.440E-09	5.713E-08	1.225E-06	2.345E-05	3.284E-05
TL209	0.0	3.401E-15	6.066E-14	5.427E-12	2.765E-09	4.922E-07	1.141E-05	2.787E-05
PB209	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
PB210	0.0	1.261E-15	1.182E-11	6.450E-08	4.278E-05	2.090E-03	1.637E-02	2.748E-03
PB211	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
PB214	0.0	1.655E-13	1.615E-10	1.352E-07	4.279E-05	2.090E-03	1.637E-02	2.748E-03
BI210	0.0	1.261E-15	1.182E-11	6.450E-08	4.278E-05	2.090E-03	1.637E-02	2.748E-03
BI211	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
BI213	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
BI214	0.0	1.655E-13	1.615E-10	1.352E-07	4.279E-05	2.090E-03	1.637E-02	2.748E-03
PO210	0.0	3.446E-16	1.182E-11	6.450E-08	4.278E-05	2.090E-03	1.637E-02	2.748E-03
PO213	0.0	1.541E-13	2.748E-12	2.458E-10	1.253E-07	2.229E-05	5.169E-04	1.262E-03
PO214	0.0	1.654E-13	1.614E-10	1.352E-07	4.278E-05	2.090E-03	1.637E-02	2.748E-03
PO215	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
PO218	0.0	1.655E-13	1.615E-10	1.352E-07	4.280E-05	2.091E-03	1.637E-02	2.749E-03
AT217	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RN219	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
RN222	0.0	1.655E-13	1.615E-10	1.352E-07	4.280E-05	2.091E-03	1.637E-02	2.749E-03
FR221	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
FR223	0.0	1.057E-14	9.662E-13	4.760E-11	7.906E-10	1.696E-08	3.245E-07	4.545E-07
RA223	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
RA225	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RA226	0.0	1.655E-13	1.615E-10	1.352E-07	4.280E-05	2.091E-03	1.637E-02	2.749E-03
AC225	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
AC227	0.0	7.660E-13	7.002E-11	3.449E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
TH227	0.0	7.554E-13	6.909E-11	3.402E-09	5.650E-08	1.212E-06	2.319E-05	3.248E-05
TH229	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
TH230	0.0	1.143E-09	1.112E-07	8.913E-06	2.510E-04	2.691E-03	1.623E-02	2.746E-03
TH231	0.0	2.300E-06	2.307E-06	2.387E-06	3.166E-06	9.956E-06	3.124E-05	3.293E-05
PA231	0.0	4.864E-11	4.878E-10	4.952E-09	5.728E-08	1.229E-06	2.351E-05	3.293E-05
PA233	0.0	6.173E-05	6.957E-05	2.599E-04	1.338E-03	1.666E-03	1.618E-03	1.209E-03
U233	1.533E-09	1.802E-09	4.421E-09	6.703E-08	3.697E-06	6.708E-05	5.790E-04	1.283E-03
U234	5.492E-07	2.530E-04	2.438E-03	1.752E-02	3.200E-02	3.120E-02	2.418E-02	1.885E-03
U235	2.299E-06	2.300E-06	2.307E-06	2.387E-06	3.166E-06	9.956E-06	3.124E-05	3.293E-05
U236	1.277E-05	1.279E-05	1.302E-05	1.522E-05	3.614E-05	1.646E-04	2.444E-04	2.380E-04

TABLE 3-INEL. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES (CONTINUED)

[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
U237	6.130E-09	4.778E-03	3.098E-03	4.069E-05	6.227E-24	0.0	0.0	0.0
NP237	6.130E-05	6.173E-05	6.957E-05	2.599E-04	1.338E-03	1.666E-03	1.618E-03	1.209E-03
NP239	0.0	1.060E-02	1.059E-02	1.050E-02	9.649E-03	4.144E-03	8.839E-07	1.723E-43
PU238	8.942E+01	8.872E+01	8.263E+01	4.058E+01	3.316E-02	4.403E-33	0.0	0.0
PU239	8.936E-01	8.936E-01	8.934E-01	8.911E-01	8.685E-01	6.716E-01	5.040E-02	2.779E-13
PU240	8.302E-01	8.302E-01	8.299E-01	8.232E-01	7.483E-01	2.882E-01	2.071E-05	0.0
PU241	2.044E+02	1.948E+02	1.263E+02	1.659E+00	2.538E-19	0.0	0.0	0.0
PU242	2.299E-03	2.299E-03	2.299E-03	2.298E-03	2.295E-03	2.258E-03	1.922E-03	3.832E-04
AM241	1.162E+00	1.480E+00	3.723E+00	6.932E+00	1.651E+00	8.900E-07	0.0	0.0
AM243	1.060E-02	1.060E-02	1.059E-02	1.050E-02	9.649E-03	4.144E-03	8.838E-07	1.723E-43
CM242	8.302E-01	1.760E-01	1.519E-07	0.0	0.0	0.0	0.0	0.0
CM244	6.638E-01	6.388E-01	4.527E-01	1.445E-02	1.584E-17	0.0	0.0	0.0
SUM	2.982E+02	2.875E+02	2.148E+02	5.095E+01	3.358E+00	1.027E+00	2.486E-01	4.437E-02
TOTAL	2.982E+02	2.875E+02	2.148E+02	5.095E+01	3.358E+00	1.027E+00	2.486E-01	4.437E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 4-INEL. RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.0	7.672E-13	6.992E-11	3.446E-09	5.990E-08	1.718E-06	3.486E-05	6.071E-05
PB	0.0	1.090E-12	2.461E-10	2.034E-07	8.575E-05	4.204E-03	3.329E-02	6.819E-03
BI	0.0	1.090E-12	2.461E-10	2.034E-07	8.575E-05	4.204E-03	3.329E-02	6.819E-03
PO	0.0	1.253E-12	4.077E-10	3.386E-07	1.285E-04	6.294E-03	4.965E-02	9.539E-03
AT	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RN	0.0	9.315E-13	2.315E-10	1.387E-07	4.285E-05	2.092E-03	1.640E-02	2.782E-03
FR	0.0	1.680E-13	3.775E-12	2.989E-10	1.288E-07	2.280E-05	5.286E-04	1.291E-03
RA	0.0	1.089E-12	2.344E-10	1.389E-07	4.298E-05	2.115E-03	1.693E-02	4.072E-03
AC	0.0	9.235E-13	7.283E-11	3.701E-09	1.853E-07	2.401E-05	5.518E-04	1.323E-03
TH	0.0	2.301E-06	2.419E-06	1.130E-05	2.543E-04	2.725E-03	1.681E-02	4.102E-03
PA	0.0	6.173E-05	6.958E-05	2.599E-04	1.338E-03	1.667E-03	1.642E-03	1.242E-03
U	1.563E-05	5.046E-03	5.551E-03	1.758E-02	3.204E-02	3.145E-02	2.503E-02	3.439E-03
NP	6.130E-05	1.066E-02	1.066E-02	1.076E-02	1.099E-02	5.810E-03	1.619E-03	1.209E-03
PU	2.955E+02	2.852E+02	2.106E+02	4.396E+01	1.652E+00	9.621E-01	5.235E-02	3.832E-04
AM	1.173E+00	1.491E+00	3.733E+00	6.943E+00	1.660E+00	4.145E-03	8.838E-07	1.723E-43
CM	1.494E+00	8.148E-01	4.527E-01	1.445E-02	1.584E-17	0.0	0.0	0.0
SUM	2.982E+02	2.875E+02	2.148E+02	5.095E+01	3.358E+00	1.027E+00	2.486E-01	4.437E-02
TOTAL	2.982E+02	2.875E+02	2.148E+02	5.095E+01	3.358E+00	1.027E+00	2.486E-01	4.437E-02

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 5-INEL. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL207	0.0	2.243E-15	2.051E-13	1.010E-11	1.678E-10	3.599E-09	6.886E-08	9.644E-08
TL209	0.0	5.651E-17	1.008E-15	9.018E-14	4.594E-11	8.177E-09	1.896E-07	4.630E-07
PB209	0.0	1.811E-16	3.230E-15	2.889E-13	1.472E-10	2.620E-08	6.075E-07	1.484E-06
PB210	0.0	2.922E-19	2.739E-15	1.494E-11	9.910E-09	4.841E-07	3.792E-06	6.365E-07
PB211	0.0	2.295E-15	2.099E-13	1.034E-11	1.717E-10	3.682E-09	7.047E-08	9.868E-08
PB214	0.0	5.276E-16	5.149E-13	4.311E-10	1.365E-07	6.666E-06	5.221E-05	8.764E-06
BI210	0.0	2.908E-18	2.726E-14	1.487E-10	9.864E-08	4.819E-06	3.774E-05	6.335E-06
BI211	0.0	3.055E-14	2.794E-12	1.376E-10	2.285E-09	4.902E-08	9.380E-07	1.314E-06
BI213	0.0	6.620E-16	1.181E-14	1.056E-12	5.382E-10	9.579E-08	2.221E-06	5.424E-06
BI214	0.0	2.120E-15	2.069E-12	1.732E-09	5.484E-07	2.679E-05	2.098E-04	3.522E-05
PO210	0.0	1.105E-17	3.790E-13	2.068E-09	1.371E-06	6.699E-05	5.247E-04	8.808E-05
PO213	0.0	7.796E-15	1.391E-13	1.244E-11	6.338E-09	1.128E-06	2.616E-05	6.388E-05
PO214	0.0	7.681E-15	7.495E-12	6.275E-09	1.986E-06	9.704E-05	7.600E-04	1.276E-04
PO215	0.0	3.419E-14	3.127E-12	1.540E-10	2.558E-09	5.486E-08	1.050E-06	1.470E-06
PO218	0.0	5.997E-15	5.852E-12	4.899E-09	1.551E-06	7.576E-05	5.933E-04	9.960E-05
AT217	0.0	6.720E-15	1.198E-13	1.072E-11	5.463E-09	9.723E-07	2.254E-05	5.506E-05
RN219	0.0	3.178E-14	2.907E-12	1.431E-10	2.377E-09	5.099E-08	9.758E-07	1.367E-06
RN222	0.0	5.484E-15	5.351E-12	4.480E-09	1.418E-06	6.928E-05	5.426E-04	9.108E-05
FR221	0.0	6.077E-15	1.084E-13	9.698E-12	4.941E-09	8.794E-07	2.039E-05	4.980E-05
RA223	0.0	2.727E-14	2.494E-12	1.228E-10	2.040E-09	4.376E-08	8.374E-07	1.173E-06
RA225	0.0	1.104E-16	1.969E-15	1.762E-13	8.977E-11	1.598E-08	3.705E-07	9.048E-07
RA226	0.0	4.778E-15	4.663E-12	3.904E-09	1.236E-06	6.037E-05	4.728E-04	7.936E-05
AC225	0.0	5.501E-15	9.811E-14	8.777E-12	4.472E-09	7.959E-07	1.845E-05	4.507E-05
AC227	0.0	3.709E-16	3.390E-14	1.670E-12	2.774E-11	5.951E-10	1.139E-08	1.595E-08
TH227	0.0	2.757E-14	2.521E-12	1.242E-10	2.062E-09	4.423E-08	8.464E-07	1.185E-06
TH229	0.0	4.817E-15	8.592E-14	7.687E-12	3.916E-09	6.971E-07	1.616E-05	3.947E-05
TH230	0.0	3.234E-11	3.147E-09	2.522E-07	7.103E-06	7.614E-05	4.593E-04	7.771E-05
TH231	0.0	1.290E-09	1.295E-09	1.339E-09	1.777E-09	5.586E-09	1.753E-08	1.848E-08
PA231	0.0	1.465E-12	1.470E-11	1.492E-10	1.726E-09	3.702E-08	7.085E-07	9.923E-07
PA233	0.0	1.401E-07	1.579E-07	5.899E-07	3.036E-06	3.782E-06	3.673E-06	2.744E-06
U233	4.456E-11	5.238E-11	1.285E-10	1.948E-09	1.075E-07	1.950E-06	1.683E-05	3.729E-05
U234	1.582E-08	7.288E-06	7.022E-05	5.047E-04	9.216E-04	8.988E-04	6.964E-04	5.430E-05
U235	6.020E-08	6.022E-08	6.043E-08	6.250E-08	8.292E-08	2.607E-07	8.181E-07	8.624E-07
U236	3.459E-07	3.466E-07	3.526E-07	4.123E-07	9.790E-07	4.459E-06	6.621E-06	6.447E-06
NP237	1.874E-06	1.887E-06	2.126E-06	7.943E-06	4.088E-05	5.092E-05	4.946E-05	3.695E-05

TABLE 5-INEL. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES (CONTINUED)
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	WATTS							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
NP239	0.0	2.562E-05	2.560E-05	2.538E-05	2.333E-05	1.002E-05	2.137E-09	4.164E-46
PU238	2.963E+00	2.940E+00	2.738E+00	1.345E+00	1.099E-03	1.459E-34	0.0	0.0
PU239	2.754E-02	2.754E-02	2.753E-02	2.746E-02	2.677E-02	2.070E-02	1.553E-03	8.563E-15
PU240	2.585E-02	2.585E-02	2.584E-02	2.563E-02	2.330E-02	8.973E-03	6.448E-07	0.0
PU241	6.336E-03	6.038E-03	3.915E-03	5.143E-05	7.869E-24	0.0	0.0	0.0
PU242	6.788E-05	6.788E-05	6.788E-05	6.787E-05	6.776E-05	6.668E-05	5.675E-05	1.132E-05
AM241	3.861E-02	4.917E-02	1.237E-01	2.303E-01	5.483E-02	2.956E-08	0.0	0.0
AM243	3.407E-04	3.407E-04	3.404E-04	3.375E-04	3.102E-04	1.332E-04	2.841E-08	5.537E-45
CM242	3.059E-02	6.484E-03	5.597E-09	0.0	0.0	0.0	0.0	0.0
CM244	2.322E-02	2.235E-02	1.583E-02	5.053E-04	5.539E-19	0.0	0.0	0.0
SUM	3.116E+00	3.078E+00	2.936E+00	1.630E+00	1.074E-01	3.133E-02	6.153E-03	1.034E-03
TOTAL	3.116E+00	3.078E+00	2.936E+00	1.630E+00	1.074E-01	3.133E-02	6.153E-03	1.034E-03

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 6-INEL. THERMAL POWER OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
TL	0.0	2.300E-15	2.062E-13	1.019E-11	2.137E-10	1.178E-08	2.585E-07	5.596E-07
PB	0.0	3.004E-15	7.308E-13	4.567E-10	1.467E-07	7.180E-06	5.668E-05	1.098E-05
BI	0.0	3.334E-14	4.902E-12	2.020E-09	6.498E-07	3.175E-05	2.507E-04	4.829E-05
PO	0.0	5.578E-14	1.700E-11	1.341E-08	4.917E-06	2.410E-04	1.905E-03	3.806E-04
AT	0.0	6.720E-15	1.198E-13	1.072E-11	5.463E-09	9.723E-07	2.254E-05	5.506E-05
RN	0.0	3.727E-14	8.258E-12	4.623E-09	1.420E-06	6.933E-05	5.435E-04	9.245E-05
FR	0.0	6.105E-15	1.109E-13	9.821E-12	4.943E-09	8.794E-07	2.039E-05	4.980E-05
RA	0.0	3.216E-14	7.159E-12	4.027E-09	1.238E-06	6.043E-05	4.740E-04	8.144E-05
AC	0.0	5.872E-15	1.320E-13	1.045E-11	4.500E-09	7.965E-07	1.847E-05	4.509E-05
TH	0.0	1.323E-09	4.444E-09	2.537E-07	7.110E-06	7.689E-05	4.763E-04	1.184E-04
PA	0.0	1.401E-07	1.579E-07	5.900E-07	3.038E-06	3.819E-06	4.382E-06	3.737E-06
U	4.220E-07	1.674E-05	7.649E-05	5.053E-04	9.228E-04	9.054E-04	7.206E-04	9.891E-05
NP	1.874E-06	2.751E-05	2.772E-05	3.333E-05	6.421E-05	6.094E-05	4.946E-05	3.695E-05
PU	3.023E+00	3.000E+00	2.796E+00	1.398E+00	5.123E-02	2.974E-02	1.611E-03	1.132E-05
AM	3.895E-02	4.951E-02	1.240E-01	2.306E-01	5.514E-02	1.332E-04	2.841E-08	5.537E-45
CM	5.381E-02	2.883E-02	1.583E-02	5.053E-04	5.539E-19	0.0	0.0	0.0
SUM	3.116E+00	3.078E+00	2.936E+00	1.630E+00	1.074E-01	3.133E-02	6.153E-03	1.034E-03
TOTAL	3.116E+00	3.078E+00	2.936E+00	1.630E+00	1.074E-01	3.133E-02	6.153E-03	1.034E-03

ELEMENTS CONTRIBUTING <0.0001% ARE OMITTED.

TABLE 7-INEL. ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI211	0.0	7.638E-13	6.986E-11	3.440E-09	5.713E-08	1.225E-06	2.345E-05	3.284E-05
BI213	0.0	3.401E-15	6.066E-14	5.427E-12	2.765E-09	4.922E-07	1.141E-05	2.787E-05
BI214	0.0	3.475E-17	3.391E-14	2.839E-11	8.986E-09	4.390E-07	3.438E-06	5.771E-07
PO210	0.0	3.446E-16	1.182E-11	6.450E-08	4.278E-05	2.090E-03	1.637E-02	2.748E-03
PO213	0.0	1.541E-13	2.748E-12	2.458E-10	1.253E-07	2.229E-05	5.169E-04	1.262E-03
PO214	0.0	1.654E-13	1.614E-10	1.352E-07	4.278E-05	2.090E-03	1.637E-02	2.748E-03
PO215	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
PO218	0.0	1.655E-13	1.615E-10	1.352E-07	4.279E-05	2.090E-03	1.637E-02	2.748E-03
AT217	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RN219	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
RN222	0.0	1.655E-13	1.615E-10	1.352E-07	4.280E-05	2.091E-03	1.637E-02	2.749E-03
FR221	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RA223	0.0	7.660E-13	7.005E-11	3.450E-09	5.729E-08	1.229E-06	2.352E-05	3.293E-05
RA226	0.0	1.655E-13	1.615E-10	1.352E-07	4.280E-05	2.091E-03	1.637E-02	2.749E-03
AC225	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
AC227	0.0	1.057E-14	9.662E-13	4.760E-11	7.906E-10	1.696E-08	3.245E-07	4.545E-07
TH227	0.0	7.554E-13	6.909E-11	3.402E-09	5.650E-08	1.212E-06	2.319E-05	3.248E-05
TH229	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
TH230	0.0	1.143E-09	1.112E-07	8.913E-06	2.510E-04	2.691E-03	1.623E-02	2.746E-03
PA231	0.0	4.864E-11	4.878E-10	4.952E-09	5.728E-08	1.229E-06	2.351E-05	3.293E-05
U233	1.533E-09	1.802E-09	4.421E-09	6.703E-08	3.697E-06	6.708E-05	5.790E-04	1.283E-03
U234	5.492E-07	2.530E-04	2.438E-03	1.752E-02	3.200E-02	3.120E-02	2.418E-02	1.885E-03
U235	2.299E-06	2.300E-06	2.307E-06	2.387E-06	3.166E-06	9.956E-06	3.124E-05	3.293E-05
U236	1.277E-05	1.279E-05	1.302E-05	1.522E-05	3.614E-05	1.646E-04	2.444E-04	2.380E-04
NP237	6.130E-05	6.173E-05	6.957E-05	2.599E-04	1.338E-03	1.666E-03	1.618E-03	1.209E-03
PU238	8.942E+01	8.872E+01	8.263E+01	4.058E+01	3.316E-02	4.403E-33	0.0	0.0
PU239	8.936E-01	8.936E-01	8.934E-01	8.911E-01	8.685E-01	6.716E-01	5.040E-02	2.779E-13
PU240	8.302E-01	8.302E-01	8.299E-01	8.232E-01	7.483E-01	2.882E-01	2.071E-05	0.0
PU241	5.007E-03	4.772E-03	3.094E-03	4.064E-05	6.219E-24	0.0	0.0	0.0
PU242	2.299E-03	2.299E-03	2.299E-03	2.298E-03	2.295E-03	2.258E-03	1.922E-03	3.832E-04
AM241	1.162E+00	1.480E+00	3.723E+00	6.932E+00	1.651E+00	8.900E-07	0.0	0.0
AM243	1.060E-02	1.060E-02	1.059E-02	1.050E-02	9.649E-03	4.144E-03	8.838E-07	1.723E-43
CM242	8.302E-01	1.760E-01	1.519E-07	0.0	0.0	0.0	0.0	0.0
CM244	6.638E-01	6.388E-01	4.527E-01	1.445E-02	1.584E-17	0.0	0.0	0.0
SUM	9.381E+01	9.275E+01	8.854E+01	4.928E+01	3.346E+00	1.013E+00	1.799E-01	2.817E-02
TOTAL	9.381E+01	9.275E+01	8.854E+01	4.928E+01	3.346E+00	1.013E+00	1.799E-01	2.817E-02

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 8-INEL. ALPHA RADIOACTIVITY OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
BI	0.0	7.673E-13	6.995E-11	3.474E-09	6.888E-08	2.157E-06	3.830E-05	6.129E-05
PO	0.0	1.253E-12	4.077E-10	3.385E-07	1.285E-04	6.294E-03	4.965E-02	9.539E-03
AT	0.0	1.575E-13	2.808E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RN	0.0	9.315E-13	2.315E-10	1.387E-07	4.285E-05	2.092E-03	1.640E-02	2.782E-03
FR	0.0	1.575E-13	2.809E-12	2.513E-10	1.280E-07	2.279E-05	5.283E-04	1.290E-03
RA	0.0	9.315E-13	2.315E-10	1.387E-07	4.285E-05	2.092E-03	1.640E-02	2.782E-03
AC	0.0	1.680E-13	3.775E-12	2.989E-10	1.288E-07	2.280E-05	5.286E-04	1.291E-03
TH	0.0	1.144E-09	1.113E-07	8.916E-06	2.512E-04	2.715E-03	1.678E-02	4.069E-03
PA	0.0	4.864E-11	4.878E-10	4.952E-09	5.728E-08	1.229E-06	2.351E-05	3.293E-05
U	1.562E-05	2.681E-04	2.453E-03	1.754E-02	3.204E-02	3.145E-02	2.503E-02	3.439E-03
NP	6.130E-05	6.173E-05	6.957E-05	2.599E-04	1.338E-03	1.666E-03	1.618E-03	1.209E-03
PU	9.115E+01	9.045E+01	8.436E+01	4.230E+01	1.652E+00	9.621E-01	5.235E-02	3.832E-04
AM	1.173E+00	1.491E+00	3.733E+00	6.943E+00	1.660E+00	4.145E-03	8.838E-07	1.723E-43
CM	1.494E+00	8.148E-01	4.527E-01	1.445E-02	1.584E-17	0.0	0.0	0.0
SUM	9.381E+01	9.275E+01	8.854E+01	4.928E+01	3.346E+00	1.013E+00	1.799E-01	2.817E-02
TOTAL	9.381E+01	9.275E+01	8.854E+01	4.928E+01	3.346E+00	1.013E+00	1.799E-01	2.817E-02

TABLE 10-INEL. SPONTANEOUS FISSION NEUTRON SOURCES IN DECAY OF INEL HLW
 [BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	NEUTRONS/SEC						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
PU238	1.387E+04	1.376E+04	1.282E+04	6.297E+03	5.145E+00	6.832E-31	0.0	0.0
PU240	3.316E+03	3.316E+03	3.315E+03	3.288E+03	2.989E+03	1.151E+03	8.271E-02	0.0
PU242	1.014E+03	1.014E+03	1.014E+03	1.014E+03	1.013E+03	9.964E+02	8.481E+02	1.691E+02
CM242	5.408E+03	1.146E+03	9.895E-04	0.0	0.0	0.0	0.0	0.0
CM244	9.120E+04	8.777E+04	6.220E+04	1.985E+03	2.176E-12	0.0	0.0	0.0
SUM	1.148E+05	1.070E+05	7.935E+04	1.259E+04	4.008E+03	2.148E+03	8.482E+02	1.692E+02
TOTAL	1.148E+05	1.070E+05	7.935E+04	1.259E+04	4.008E+03	2.148E+03	8.482E+02	1.692E+02

TABLE 11-INEL. PHOTON SPECTRUM OF ACTINIDES AND DAUGHTERS IN DECAY OF INEL HLW
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

PHOTONS/SEC								
EMEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	5.469E+11	5.418E+11	5.216E+11	2.995E+11	2.033E+10	3.557E+09	1.139E+09	3.080E+08
2.500E-02	1.088E+09	1.396E+09	3.492E+09	6.493E+09	1.555E+09	2.134E+07	1.019E+08	2.552E+07
3.750E-02	1.631E+09	1.638E+09	1.703E+09	1.240E+09	1.554E+08	2.107E+07	6.931E+07	3.252E+07
5.750E-02	1.601E+10	2.046E+10	5.129E+10	9.542E+10	2.274E+10	2.897E+07	1.167E+08	2.372E+07
8.500E-02	5.276E+08	6.258E+08	6.151E+08	4.895E+08	3.088E+08	1.711E+08	2.295E+08	9.276E+07
1.250E-01	1.790E+07	2.629E+08	2.623E+08	2.534E+08	1.960E+08	8.712E+07	4.099E+07	1.625E+07
2.250E-01	2.457E+07	1.959E+08	1.804E+08	1.430E+08	1.209E+08	8.653E+07	2.725E+08	6.025E+07
3.750E-01	2.651E+06	1.867E+07	1.894E+07	2.154E+07	3.453E+07	6.131E+07	2.655E+08	7.362E+07
5.750E-01	4.668E+05	5.383E+05	1.094E+06	1.917E+06	1.422E+06	4.227E+07	3.307E+08	5.648E+07
8.500E-01	1.870E+06	1.863E+06	1.879E+06	1.272E+06	3.485E+05	1.023E+07	8.009E+07	1.365E+07
1.250E+00	1.383E+05	1.314E+05	1.083E+05	3.740E+04	5.451E+05	2.645E+07	2.072E+08	3.493E+07
1.750E+00	4.373E+04	4.095E+04	3.133E+04	8.233E+03	4.388E+05	2.136E+07	1.675E+08	2.898E+07
2.250E+00	2.511E+04	2.350E+04	1.793E+04	4.273E+03	1.358E+05	6.580E+06	5.153E+07	8.651E+06
2.750E+00	1.441E+04	1.348E+04	1.026E+04	2.167E+03	2.958E+03	1.148E+05	8.964E+05	1.506E+05
3.500E+00	1.284E+04	1.200E+04	9.110E+03	1.870E+03	9.867E+02	2.178E+04	1.685E+05	2.828E+04
5.000E+00	5.419E+03	5.061E+03	3.827E+03	7.604E+02	2.319E+02	1.229E+02	4.819E+01	9.596E+00
7.000E+00	6.155E+02	5.744E+02	4.325E+02	8.270E+01	2.637E+01	1.405E+01	5.532E+00	1.102E+00
9.500E+00	7.015E+01	6.541E+01	4.913E+01	9.190E+00	3.013E+00	1.610E+00	6.356E-01	1.267E-01
TOTAL	5.662E+11	5.664E+11	5.792E+11	4.035E+11	4.545E+10	4.142E+09	3.074E+09	7.755E+08

TABLE 12-INEL. MASS OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	1.173E+00	1.173E+00	1.173E+00	1.172E+00	1.161E+00	1.054E+00	4.035E-01	2.728E-05
BR 79	0.0	1.252E-05	1.252E-04	1.251E-03	1.245E-02	1.187E-01	7.695E-01	1.173E+00
RB 87	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01
SR 90	1.217E+02	1.188E+02	9.592E+01	1.126E+01	5.601E-09	0.0	0.0	0.0
ZR 90	0.0	2.863E+00	2.578E+01	1.105E+02	1.217E+02	1.217E+02	1.217E+02	1.217E+02
ZR 93	1.575E+02	1.575E+02	1.575E+02	1.575E+02	1.574E+02	1.568E+02	1.505E+02	1.001E+02
NB 93	0.0	2.210E-05	3.177E-04	6.150E-03	7.035E-02	7.110E-01	6.976E+00	5.738E+01
TC 99	1.582E+02	1.582E+02	1.582E+02	1.581E+02	1.577E+02	1.531E+02	1.143E+02	6.109E+00
RU 99	0.0	5.148E-04	5.148E-03	5.147E-02	5.140E-01	5.065E+00	4.394E+01	1.521E+02
PD107	4.965E+00	4.965E+00	4.965E+00	4.965E+00	4.964E+00	4.960E+00	4.912E+00	4.463E+00
SN126	1.440E+00	1.440E+00	1.440E+00	1.439E+00	1.430E+00	1.344E+00	7.200E-01	1.407E-03
TE126	0.0	1.040E-05	1.002E-04	9.981E-04	9.946E-03	9.643E-02	7.200E-01	1.439E+00
CS134	3.256E+00	2.326E+00	1.129E-01	8.545E-15	0.0	0.0	0.0	0.0
BA134	0.0	9.296E-01	3.143E+00	3.256E+00	3.256E+00	3.256E+00	3.256E+00	3.256E+00
CS135	8.316E+01	8.316E+01	8.316E+01	8.316E+01	8.313E+01	8.291E+01	8.069E+01	6.152E+01
BA135	0.0	2.506E-05	2.506E-04	2.506E-03	2.506E-02	2.502E-01	2.469E+00	2.164E+01
CS137	1.908E+02	1.864E+02	1.514E+02	1.893E+01	1.762E-08	0.0	0.0	0.0
BA137	0.0	4.358E+00	3.936E+01	1.719E+02	1.908E+02	1.908E+02	1.908E+02	1.908E+02
CE144	3.282E+00	1.347E+00	4.450E-04	6.861E-39	0.0	0.0	0.0	0.0
ND144	0.0	1.935E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00
PM147	1.653E+01	1.269E+01	1.177E+00	5.541E-11	0.0	0.0	0.0	0.0
SM147	0.0	3.838E+00	1.535E+01	1.653E+01	1.653E+01	1.653E+01	1.653E+01	1.653E+01
SM151	8.250E+00	8.187E+00	7.638E+00	3.819E+00	3.728E-03	2.928E-33	0.0	0.0
EU151	0.0	6.330E-02	6.116E-01	4.431E+00	8.246E+00	8.250E+00	8.250E+00	8.250E+00
EU154	8.513E-01	7.854E-01	3.802E-01	2.690E-04	8.457E-36	0.0	0.0	0.0
GD154	0.0	6.592E-02	4.711E-01	8.510E-01	8.513E-01	8.513E-01	8.513E-01	8.513E-01
SUM	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02
TOTAL	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02

NUCLIDES CONTRIBUTING <0.1000% ARE OMITTED.

TABLE 13-INEL. MASS OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	IMMOBILZN	GRAMS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	1.173E+00	1.173E+00	1.173E+00	1.172E+00	1.161E+00	1.054E+00	4.035E-01	2.728E-05
BR	0.0	1.252E-05	1.252E-04	1.251E-03	1.245E-02	1.187E-01	7.695E-01	1.173E+00
RB	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01	5.252E+01
SR	1.217E+02	1.188E+02	9.592E+01	1.126E+01	7.808E-07	7.752E-06	7.752E-05	7.752E-04
ZR	1.575E+02	1.604E+02	1.833E+02	2.680E+02	2.792E+02	2.785E+02	2.723E+02	2.218E+02
NB	3.387E-04	4.101E-04	1.052E-03	7.474E-03	7.168E-02	7.123E-01	6.977E+00	5.738E+01
TC	1.582E+02	1.582E+02	1.582E+02	1.581E+02	1.577E+02	1.531E+02	1.143E+02	6.109E+00
RU	3.701E-01	1.866E-01	5.530E-03	5.147E-02	5.140E-01	5.065E+00	4.394E+01	1.521E+02
PD	4.965E+00	5.149E+00	5.335E+00	5.335E+00	5.335E+00	5.330E+00	5.282E+00	4.833E+00
AG	0.0	5.298E-07	5.298E-06	5.298E-05	5.298E-04	5.295E-03	5.270E-02	5.025E-01
SN	1.440E+00	1.440E+00	1.440E+00	1.439E+00	1.430E+00	1.344E+00	7.200E-01	1.407E-03
TE	0.0	1.040E-05	1.002E-04	9.981E-04	9.946E-03	9.643E-02	7.200E-01	1.439E+00
CS	2.772E+02	2.719E+02	2.347E+02	1.021E+02	8.313E+01	8.291E+01	8.069E+01	6.152E+01
BA	2.848E-05	5.288E+00	4.251E+01	1.751E+02	1.941E+02	1.943E+02	1.965E+02	2.157E+02
CE	3.282E+00	1.347E+00	4.450E-04	1.042E-13	1.053E-12	1.054E-11	1.054E-10	1.054E-09
ND	0.0	1.935E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00	3.282E+00
PM	1.653E+01	1.269E+01	1.177E+00	5.541E-11	0.0	0.0	0.0	0.0
SM	8.250E+00	1.202E+01	2.299E+01	2.035E+01	1.653E+01	1.653E+01	1.653E+01	1.653E+01
EU	8.513E-01	8.487E-01	9.918E-01	4.431E+00	8.246E+00	8.250E+00	8.250E+00	8.250E+00
GD	0.0	6.592E-02	4.711E-01	8.510E-01	8.513E-01	8.513E-01	8.513E-01	8.513E-01
SUM	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02
TOTAL	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02	8.040E+02

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 14-INEL. RADIOACTIVITY OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

NUCLIDE	IMMOBILZN	CURIES						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	8.175E-02	8.175E-02	8.174E-02	8.166E-02	8.088E-02	7.347E-02	2.812E-02	1.901E-06
SR 90	1.661E+04	1.622E+04	1.309E+04	1.537E+03	7.643E-07	0.0	0.0	0.0
Y 90	1.661E+04	1.622E+04	1.309E+04	1.537E+03	7.645E-07	0.0	0.0	0.0
ZR 93	3.959E-01	3.959E-01	3.959E-01	3.959E-01	3.957E-01	3.941E-01	3.784E-01	2.517E-01
NB 93M	9.577E-02	1.097E-01	2.077E-01	3.744E-01	3.759E-01	3.744E-01	3.594E-01	2.391E-01
TC 99	2.683E+00	2.683E+00	2.683E+00	2.682E+00	2.674E+00	2.597E+00	1.938E+00	1.036E-01
RU106	1.239E+03	6.228E+02	1.278E+00	1.723E-27	0.0	0.0	0.0	0.0
RH106	1.239E+03	6.228E+02	1.278E+00	1.723E-27	0.0	0.0	0.0	0.0
PD107	2.555E-03	2.555E-03	2.555E-03	2.555E-03	2.554E-03	2.552E-03	2.528E-03	2.296E-03
SN126	4.087E-02	4.087E-02	4.087E-02	4.084E-02	4.059E-02	3.813E-02	2.044E-02	3.993E-05
SB126	4.087E-02	5.722E-03	5.722E-03	5.718E-03	5.682E-03	5.339E-03	2.861E-03	5.591E-06
SB126M	4.087E-02	4.087E-02	4.087E-02	4.084E-02	4.059E-02	3.813E-02	2.044E-02	3.993E-05
CS134	4.215E+03	3.011E+03	1.462E+02	1.106E-11	0.0	0.0	0.0	0.0
CS135	9.579E-02	9.579E-02	9.579E-02	9.579E-02	9.576E-02	9.550E-02	9.295E-02	7.087E-02
CS137	1.660E+04	1.622E+04	1.318E+04	1.647E+03	1.533E-06	0.0	0.0	0.0
BA137M	1.533E+04	1.535E+04	1.247E+04	1.558E+03	1.450E-06	0.0	0.0	0.0
CE144	1.047E+04	4.298E+03	1.420E+00	2.190E-35	0.0	0.0	0.0	0.0
PR144	1.048E+04	4.299E+03	1.420E+00	2.190E-35	0.0	0.0	0.0	0.0
PR144M	0.0	5.158E+01	1.704E-02	2.628E-37	0.0	0.0	0.0	0.0
PM147	1.533E+04	1.177E+04	1.092E+03	5.139E-08	0.0	0.0	0.0	0.0
SM151	2.171E+02	2.155E+02	2.010E+02	1.005E+02	9.811E-02	7.705E-32	0.0	0.0
EU154	2.299E+02	2.121E+02	1.027E+02	7.265E-02	2.284E-33	0.0	0.0	0.0
SUM	1.086E+05	8.912E+04	5.338E+04	6.383E+03	3.810E+00	3.619E+00	2.843E+00	6.676E-01
TOTAL	1.086E+05	8.912E+04	5.338E+04	6.383E+03	3.810E+00	3.619E+00	2.843E+00	6.676E-01

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 15-INEL. RADIOACTIVITY OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	CURIES							
	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	8.175E-02	8.175E-02	8.174E-02	8.166E-02	8.088E-02	7.347E-02	2.812E-02	1.901E-06
SR	1.661E+04	1.622E+04	1.309E+04	1.537E+03	7.643E-07	0.0	0.0	0.0
Y	1.661E+04	1.622E+04	1.309E+04	1.537E+03	7.645E-07	0.0	0.0	0.0
ZR	3.959E-01	3.959E-01	3.959E-01	3.959E-01	3.957E-01	3.941E-01	3.784E-01	2.517E-01
NB	9.577E-02	1.097E-01	2.077E-01	3.744E-01	3.759E-01	3.744E-01	3.594E-01	2.391E-01
TC	2.683E+00	2.683E+00	2.683E+00	2.682E+00	2.674E+00	2.597E+00	1.938E+00	1.036E-01
RU	1.239E+03	6.228E+02	1.278E+00	1.723E-27	0.0	0.0	0.0	0.0
RH	1.239E+03	6.228E+02	1.278E+00	1.723E-27	0.0	0.0	0.0	0.0
PD	2.555E-03	2.555E-03	2.555E-03	2.555E-03	2.554E-03	2.552E-03	2.528E-03	2.296E-03
SN	4.087E-02	4.087E-02	4.087E-02	4.084E-02	4.059E-02	3.813E-02	2.044E-02	3.993E-05
SB	8.174E-02	4.659E-02	4.659E-02	4.656E-02	4.627E-02	4.347E-02	2.330E-02	4.552E-05
CS	2.082E+04	1.924E+04	1.332E+04	1.647E+03	9.576E-02	9.550E-02	9.295E-02	7.087E-02
BA	1.533E+04	1.535E+04	1.247E+04	1.558E+03	1.450E-06	0.0	0.0	0.0
CE	1.047E+04	4.298E+03	1.420E+00	2.190E-35	0.0	0.0	0.0	0.0
PR	1.048E+04	4.350E+03	1.437E+00	2.216E-35	0.0	0.0	0.0	0.0
PM	1.533E+04	1.177E+04	1.092E+03	5.139E-08	0.0	0.0	0.0	0.0
SM	2.171E+02	2.155E+02	2.010E+02	1.005E+02	9.811E-02	3.758E-07	3.758E-07	3.758E-07
EU	2.299E+02	2.121E+02	1.027E+02	7.265E-02	2.284E-33	0.0	0.0	0.0
SUM	1.086E+05	8.912E+04	5.338E+04	6.383E+03	3.810E+00	3.619E+00	2.843E+00	6.676E-01
TOTAL	1.086E+05	8.912E+04	5.338E+04	6.383E+03	3.810E+00	3.619E+00	2.843E+00	6.676E-01

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 16-INEL. THERMAL POWER OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: NUCLIDES
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

		WATTS						
NUCLIDE	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE 79	2.035E-05	2.035E-05	2.035E-05	2.033E-05	2.014E-05	1.829E-05	7.002E-06	4.733E-10
RB 87	3.843E-09	3.843E-09	3.843E-09	3.843E-09	3.843E-09	3.843E-09	3.843E-09	3.842E-09
SR 90	1.927E+01	1.882E+01	1.519E+01	1.784E+00	8.870E-10	0.0	0.0	0.0
Y 90	9.204E+01	8.990E+01	7.256E+01	8.519E+00	4.237E-09	0.0	0.0	0.0
ZR 93	4.600E-05	4.600E-05	4.600E-05	4.599E-05	4.598E-05	4.579E-05	4.396E-05	2.924E-05
NB 93M	1.697E-05	1.944E-05	3.680E-05	6.633E-05	6.661E-05	6.634E-05	6.369E-05	4.236E-05
TC 99	1.346E-03	1.346E-03	1.345E-03	1.345E-03	1.341E-03	1.302E-03	9.718E-04	5.196E-05
RU106	7.365E-02	3.703E-02	7.599E-05	1.024E-31	0.0	0.0	0.0	0.0
RH106	1.188E+01	5.974E+00	1.226E-02	1.652E-29	0.0	0.0	0.0	0.0
PD107	1.514E-07	1.514E-07	1.514E-07	1.514E-07	1.514E-07	1.513E-07	1.498E-07	1.361E-07
SN126	5.097E-05	5.097E-05	5.097E-05	5.094E-05	5.062E-05	4.756E-05	2.549E-05	4.980E-08
SB126	7.552E-04	1.057E-04	1.057E-04	1.056E-04	1.050E-04	9.864E-05	5.286E-05	1.033E-07
SB126M	5.203E-04	5.204E-04	5.204E-04	5.200E-04	5.168E-04	4.855E-04	2.602E-04	5.085E-07
CS134	4.290E+01	3.065E+01	1.488E+00	1.126E-13	0.0	0.0	0.0	0.0
CS135	3.197E-05	3.197E-05	3.197E-05	3.197E-05	3.196E-05	3.187E-05	3.102E-05	2.365E-05
CS137	1.837E+01	1.795E+01	1.458E+01	1.822E+00	1.696E-09	0.0	0.0	0.0
BA137M	6.017E+01	6.026E+01	4.895E+01	6.118E+00	5.694E-09	0.0	0.0	0.0
CE144	6.947E+00	2.851E+00	9.419E-04	1.452E-38	0.0	0.0	0.0	0.0
PR144	7.700E+01	3.160E+01	1.044E-02	1.609E-37	0.0	0.0	0.0	0.0
PR144M	0.0	1.765E-02	5.830E-06	8.990E-41	0.0	0.0	0.0	0.0
PM147	5.499E+00	4.222E+00	3.916E-01	1.843E-11	0.0	0.0	0.0	0.0
SM147	0.0	1.195E-09	4.780E-09	5.146E-09	5.146E-09	5.146E-09	5.146E-09	5.146E-09
SM151	2.546E-02	2.526E-02	2.357E-02	1.178E-02	1.150E-05	9.034E-36	0.0	0.0
EU154	2.056E+00	1.897E+00	9.184E-01	6.498E-04	2.043E-35	0.0	0.0	0.0
SUM	3.362E+02	2.642E+02	1.541E+02	1.826E+01	2.190E-03	2.097E-03	1.456E-03	1.480E-04
TOTAL	3.362E+02	2.642E+02	1.541E+02	1.826E+01	2.190E-03	2.097E-03	1.456E-03	1.480E-04

NUCLIDES CONTRIBUTING <0.0010% ARE OMITTED.

TABLE 17-INEL. THERMAL POWER OF FISSION AND ACTIVATION PRODUCTS IN DECAY OF INEL HLW: ELEMENTS
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

ELEMENT	IMMOBILZN	WATTS						
		1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
SE	2.035E-05	2.035E-05	2.035E-05	2.033E-05	2.014E-05	1.829E-05	7.002E-06	4.733E-10
SR	1.927E+01	1.882E+01	1.519E+01	1.784E+00	8.870E-10	0.0	0.0	0.0
Y	9.204E+01	8.990E+01	7.256E+01	8.519E+00	4.237E-09	0.0	0.0	0.0
ZR	4.600E-05	4.600E-05	4.600E-05	4.599E-05	4.598E-05	4.579E-05	4.396E-05	2.924E-05
NB	1.697E-05	1.944E-05	3.680E-05	6.633E-05	6.661E-05	6.634E-05	6.369E-05	4.236E-05
TC	1.346E-03	1.346E-03	1.345E-03	1.345E-03	1.341E-03	1.302E-03	9.718E-04	5.196E-05
RU	7.365E-02	3.703E-02	7.599E-05	1.024E-31	0.0	0.0	0.0	0.0
RH	1.188E+01	5.974E+00	1.226E-02	1.652E-29	0.0	0.0	0.0	0.0
PD	1.514E-07	1.514E-07	1.514E-07	1.514E-07	1.514E-07	1.513E-07	1.498E-07	1.361E-07
SN	5.097E-05	5.097E-05	5.097E-05	5.094E-05	5.062E-05	4.756E-05	2.549E-05	4.980E-08
SB	1.276E-03	6.261E-04	6.261E-04	6.257E-04	6.218E-04	5.842E-04	3.131E-04	6.118E-07
CS	6.126E+01	4.860E+01	1.606E+01	1.822E+00	3.196E-05	3.187E-05	3.102E-05	2.365E-05
BA	6.017E+01	6.026E+01	4.895E+01	6.118E+00	5.694E-09	0.0	0.0	0.0
CE	6.947E+00	2.851E+00	9.419E-04	1.452E-38	0.0	0.0	0.0	0.0
PR	7.700E+01	3.161E+01	1.044E-02	1.610E-37	0.0	0.0	0.0	0.0
PM	5.499E+00	4.222E+00	3.916E-01	1.843E-11	0.0	0.0	0.0	0.0
SM	2.546E-02	2.526E-02	2.357E-02	1.178E-02	1.151E-05	5.146E-09	5.146E-09	5.146E-09
EU	2.056E+00	1.897E+00	9.184E-01	6.498E-04	2.043E-35	0.0	0.0	0.0
SUM	3.362E+02	2.642E+02	1.541E+02	1.826E+01	2.190E-03	2.097E-03	1.456E-03	1.480E-04
TOTAL	3.362E+02	2.642E+02	1.541E+02	1.826E+01	2.190E-03	2.097E-03	1.456E-03	1.480E-04

ELEMENTS CONTRIBUTING <0.0100% ARE OMITTED.

TABLE 18-INEL. PHOTON SPECTRUM OF FISSION AND ACTIVATION PRODUCT IN DECAY OF INEL HLW
[BASED ON ONE CANISTER OF INEL HLW IN GLASS-CERAMIC; 1825 KG GLASS-CERAMIC (1277 KG CALCINE)]

		PHOTONS/SEC						
E MEAN	IMMOBILZN	1.0YR	10.0YR	100.0YR	1000.0YR	10.0KY	100.0KY	1.0MY
18 GROUP PHOTON RELEASE RATES, PHOTONS/SECOND								
1.000E-02	8.508E+14	6.360E+14	3.809E+14	4.483E+13	8.146E+09	7.929E+09	6.367E+09	1.986E+09
2.500E-02	1.767E+14	1.315E+14	7.869E+13	9.262E+12	1.524E+09	1.454E+09	9.442E+08	4.594E+07
3.750E-02	1.955E+14	1.421E+14	8.207E+13	9.770E+12	5.346E+08	5.158E+08	3.676E+08	2.346E+07
5.750E-02	1.671E+14	1.229E+14	7.304E+13	8.577E+12	7.499E+08	7.196E+08	4.841E+08	2.395E+07
8.500E-02	1.096E+14	7.709E+13	4.341E+13	5.109E+12	9.462E+08	8.945E+08	5.152E+08	8.809E+06
1.250E-01	1.142E+14	6.924E+13	2.948E+13	3.290E+12	1.006E+08	9.646E+07	6.410E+07	2.633E+06
2.250E-01	9.143E+13	6.494E+13	3.707E+13	4.323E+12	8.729E+07	8.246E+07	4.713E+07	6.398E+05
3.750E-01	4.195E+13	2.892E+13	1.586E+13	1.858E+12	1.646E+09	1.546E+09	8.288E+08	1.619E+06
5.750E-01	8.266E+14	7.523E+14	4.904E+14	6.034E+13	3.655E+09	3.434E+09	1.840E+09	3.596E+06
8.500E-01	1.498E+14	1.073E+14	9.133E+12	3.014E+11	1.829E+08	1.719E+08	9.210E+07	1.800E+05
1.250E+00	1.907E+13	1.359E+13	3.156E+12	9.982E+10	4.440E+07	4.172E+07	2.236E+07	4.369E+04
1.750E+00	8.144E+11	4.702E+11	1.257E+11	7.718E+09	1.897E+03	1.779E+03	9.532E+02	1.863E+00
2.250E+00	3.016E+12	1.244E+12	4.764E+08	8.471E+05	4.280E-04	6.631E-06	6.631E-06	6.631E-06
2.750E+00	1.250E+10	6.038E+09	1.052E+07	3.322E-06	3.322E-06	3.322E-06	3.322E-06	3.322E-06
3.500E+00	1.288E+09	6.474E+08	1.329E+06	2.447E-06	2.447E-06	2.447E-06	2.447E-06	2.447E-06
5.000E+00	0.0	1.691E-07	6.766E-07	7.284E-07	7.284E-07	7.284E-07	7.284E-07	7.284E-07
7.000E+00	0.0	1.097E-08	4.390E-08	4.727E-08	4.727E-08	4.727E-08	4.727E-08	4.726E-08
9.500E+00	0.0	6.940E-10	2.776E-09	2.989E-09	2.989E-09	2.989E-09	2.989E-09	2.989E-09
TOTAL	2.747E+15	2.148E+15	1.243E+15	1.478E+14	1.762E+10	1.689E+10	1.157E+10	2.097E+09

APPENDIX 3B. INTERIM HIGH-LEVEL WASTE FORMS

22

APPENDIX 3B. INTERIM HIGH-LEVEL WASTE FORMS

3B.1 INTRODUCTION

This appendix presents detailed data on the amounts and compositions of the interim high-level wastes (HLW) in storage at each of the sites, both current and projected, and on the conversion of these wastes to immobilized forms for repository disposal. It thus serves to provide the necessary backup data for the calculation of the immobilized waste quantities and compositions presented in Sect. 3.

As stated in Sect. 3, all of the HLW produced thus far is currently (1991) in interim storage at WVDP, SRS, HANF, and INEL. The storage forms vary from site to site and include liquids, sludges, salt cake, slurry, loaded zeolite resins, calcine, and encapsulated strontium fluoride and cesium chloride. These interim forms of HLW have been produced as the result of a number of treatment and separation operations (e.g., neutralization, precipitation, decantation, evaporation, and calcination) carried out on the effluent streams from fuel reprocessing, and their volumes and compositions have changed as a result of these operations as well as from the natural processes of radioactive decay. Further changes in form, density, and composition will take place when the interim forms are converted to borosilicate glass or other solid form for shipment to a repository for final disposal, and the volumes of the final solid waste forms will be much less, in most cases, than those of the interim forms now in storage at the sites.

Quantitative relationships on the conversion to immobilized forms are scarce but are presented where available so that the reader can follow the rationale used in the calculation of immobilized waste quantities, compositions, and radioactivity per canister. The discussion is arranged by site location and is in the same order as in Sect. 3: WVDP, SRS, HANF, and INEL.

3B.2 WEST VALLEY DEMONSTRATION PROJECT

The HLW stored at West Valley is the result of commercial operations during the period 1966–1972 and consists of both alkaline and acidic wastes. The alkaline waste was generated by the reprocessing of commercial power reactor fuels and some Hanford N-Reactor fuels. The acidic waste, which is much smaller in volume, was generated by reprocessing a small amount of commercial fuel containing thorium. The total quantity of fuel reprocessed from 1966 to 1972 was about 662 metric tons; the sources of this fuel, which represented about 4,000,000 MWd of irradiation exposure, are listed in Table 3B.1.

3B.2.1 HLW Inventories at WVDP

HLW inventory in storage at WVDP consists of 2,270 m³ of alkaline waste and 45 m³ of acid waste, both generated during fuel reprocessing operations at the NFS plant; these operations ended in 1972. The alkaline waste was derived from the PUREX reprocessing of low-burnup commercial and Hanford N-Reactor spent fuels. These wastes, originally acidic, were treated with excess sodium hydroxide to produce an alkaline sludge and a clear supernatant alkaline liquid. The acidic waste currently in storage was generated from the THOREX reprocessing of thorium-uranium fuel from the Indian Point 1 Reactor.

The alkaline and acid wastes are stored in underground tanks, carbon steel for the alkaline PUREX waste and stainless steel for the acid THOREX waste. There are two 2,800 m³ carbon steel tanks, 8D-1 and 8D-2. The alkaline waste is stored in tank 8D-2, with 8D-1 serving as a backup for this service. There are also two 57 m³ stainless steel tanks, 8D-3 and 8D-4. The acid waste is stored in 8D-4, with 8D-3 as backup. All four tanks are in underground concrete storage vaults. The acidic waste in Tank 8D-4 is considered to be a single-phase solution (Hannum 1983, Barnes et al. 1986).

Tables 3B.2 and 3B.3 summarize the total volumes and radioactivity of WVDP wastes of all types, both current and projected to the year 2020. Chemical compositions of the interim wastes are shown in Tables 3B.4, 5, and 6. Table 3B.7 gives the estimated radionuclide composition of the PUREX supernatant liquid, the PUREX solids, the THOREX waste, and of the total WVDP waste as of the end of 1989 (IDB 1990, Maestas 1990). No further additions to these wastes are anticipated at present.

Because zeolite ion-exchange pretreatment of the alkaline waste in Tank 8D-2 (described in the next subsection) has already started, the waste inventories at WVDP at the end of 1989 included about 31 m³ of cesium-loaded zeolite ion-exchange resins (Maestas 1990).

3B.2.2 Treatment and Conversion to Glass

Figure 3B.1 shows a schematic flowsheet of the existing and future HLW treatment and vitrification facilities at WVDP. Vitrification of the waste now in storage is scheduled to begin in 1993 and to be completed in 1995.

Zeolite ion exchange was selected as the most suitable technology for pretreatment of the WVDP waste. The high decontamination factor for cesium achievable and the simplicity of the process were contributing factors in this choice.

The initial waste treatment steps consist of decanting the supernatant liquid from Tank 8D-2 and passing it through zeolite ion exchange columns. As the zeolite ion exchange medium becomes loaded with cesium, it is discharged to the bottom of Tank 8D-1. The hydroxide sludge remaining in Tank 8D-2 is washed to remove interstitial liquids and sodium sulfate. This process requires three to four cycles of washing, settling, and decanting. The wash solutions are treated by zeolite ion exchange. The decontaminated supernatant liquid and wash effluents from the ion exchange system will be solidified into concrete and disposed of as low-level waste.

After the supernatant decontamination and the sludge wash cycles, the tank farm configuration will be cesium-loaded zeolite in Tank 8D-1, washed sludge in Tank 8D-2, and the THOREX wastes in Tank 8D-4. These wastes will then be combined and homogenized in Tank 8D-2, and the homogeneous mixture will be batched to the waste concentrator in the vitrification facility for solidification into borosilicate glass (Barnes et al. 1986). The borosilicate glass product will contain about 23% PUREX and THOREX wastes and about 10% cesium-loaded zeolite.

Some of the pretreatment activities have already been started. The supernatant treatment system began processing the high-level supernatant liquid waste in May 1988. By the end of FY 1989, about 200,000 gal (757 m³) of liquid HLW had been processed through the ion exchange system. Other components of the integrated radwaste treatment system have also started operations, including the low-level waste (LLW) cement solidification system. Nonradioactive test runs of the vitrification plant melter have been completed (WVDP 1990).

3B.2.3 Radioactivity and Thermal Power Per Canister

In the previous (1987) edition of this report, the radionuclide compositions of the HLW glass were supplied by WVDP for the year 1990 in terms of average, minimum, and maximum curies per canister (Eisenstatt 1986). For the present edition of this report, new data were obtained from WVDP based on the most recent revision (Revision 7) of the vitrification plant mass balance (Crocker 1989). Table 3B.8 was calculated from these data and shows curies and grams of each radionuclide for an average activity canister loaded to an 85% fill level, with radioactivity calculated as of the end of year 1989. The WVDP Revision 7 mass balance does not give radionuclide contents for a maximum activity canister. The new data show about 2.6% less radioactivity per average canister than the 1987 data (109,600 vs 112,700 Ci, both calculated as of the end of year 1989).

Using the Revision 7 mass balance average radionuclide content per canister supplied by WVDP for the end of year 1989, ORIGEN2 decay calculations were made to determine radioactivity and thermal power per

canister as a function of decay time from 0 to 10⁶ years, the contribution of each isotope being shown individually. These results are given in Appendix 3A, and are the basis for the summary table showing total radioactivity per canister as a function of time given in Sect. 3 of this report. The Revision 7 mass balance did not include any estimate of the range of variation from batch to batch, but based on previous estimates it appears that a maximum activity canister might contain about 10% more radioactivity than the average.

The Revision 7 mass balance shows a total glass production of 484,000 kg. At 100% fill level, a canister would contain 2,229 kg of glass, based on a density of 2.7 g/mL. At the reference-case fill level of 85%, each canister will contain about 1,900 kg of glass. The number of canisters to be produced, assuming an average fill of 85%, is therefore approximately 255. In this report, an allowance was added to account for operational variations, increasing the estimated number of canisters to 275. The volume of glass per canister calculated by WVDP is about 12% higher than that estimated by SRS and HANF; this is due to the fact that the canister design used at WVDP is different from the one used at SRS and HANF. IDB submittals from WVDP have estimated 300 as the upper limit of the number of canisters that would be produced. WVDP is providing facilities for the on-site storage of up to 312 filled canisters until a repository or MRS becomes available.

3B.2.4 Volume Conversion Factor to Glass

As of 1986, the total volume of waste before vitrification was about 2,200 m³, and the total volume of glass to be produced (assuming a density of 2.7 g/mL) is about 180 m³; the ratio of interim waste volume to glass volume is about 12.2 to 1. The volume of LLW produced, in final solidified form, is estimated to be about 2,750 m³ (Rykken 1986d, IDB 1990).

3B.2.5 Assessment of Data

The volumes and radionuclide compositions of the wastes and the vitrification flowsheet at WVDP are well established. The number of canisters of HLW glass and the average activity per canister have been determined within reasonable tolerances.

3B.3 SAVANNAH RIVER SITE

At the Savannah River Site (SRS), the acidic HLW from the reprocessing of defense reactor fuel is made alkaline by the addition of caustic and is stored in large underground tanks. The total volume of HLW produced since plant operations started in 1954 is about 300,000 m³ (Doherty 1986). The volume reduction that occurs as a result of evaporation has reduced the total volume in

storage as of the end of CY 1989 to 122,000 m³ (Garvin 1990).

The primary operations on this waste are shown schematically in Fig. 3B.2. The initial steps of neutralization and settling result in the formation of a sludge and a supernatant liquid in the storage tanks. Throughout the following discussion, this liquid is always referred to as "the supernate." Referring to Fig. 3B.2, the supernate is then concentrated by evaporation, and the concentrate is cooled, causing salt cake (salt crystals) to separate out of the saturated salt solution. The supernate has thus been converted to saturated salt solution and salt cake.

The sludge, which represents about 11 vol% of the total waste in the tanks, is composed largely of the precipitated hydroxides of iron, aluminum, manganese, and other metals; it contains about 60-65% of the total radioactivity, including most of the Sr-90 and small amounts of actinides, principally isotopes of uranium, plutonium, and curium, that were not recovered from the fuel during reprocessing. The largest portion of the actinide radioactivity is due to the plutonium isotopes Pu-238 and Pu-241. The sludge is kept essentially separate from the salt solution and salt cake by tankage selection and transfer operations.

The major components of the supernate, and hence of the salt solution and salt cake, are sodium nitrate, sodium aluminate, and sodium hydroxide, together with most of the Cs-137 and smaller amounts of other radionuclides. Almost all of the radioactivity in the salt solution and salt cake is due to Cs-137 and its short-lived daughter Ba-137m.

Generation of new HLW is expected to continue at the rate of about 10,000 m³/year, depending on future production requirements (Baxter 1986a). Currently there are 51 waste tanks, in sizes ranging up to 4,900 m³, with a combined capacity of about 222,000 m³ (Boore 1986).

Immobilization of essentially all of the radioactivity of the interim HLW in borosilicate glass will begin in 1993. This will take place in the Defense Waste Processing Facility (DWPF), which is now under construction at SRS.

As of March 1991, design of the DWPF was >99% complete, and construction was 99% complete. The facility includes storage buildings where the filled canisters of HLW can be stored on-site until a repository becomes available.

3B.3.1 HLW Inventories at SRS

The volumes of the interim HLW phases in storage at the end of CY 1988 were as follows: sludge 14,100 m³; salt cake 50,000 m³; and liquid 64,300 m³. Table 3B.9 shows the current and projected cumulative volumes of the liquid, sludge, salt cake, and glass, based on estimated future production, through the year 2020. Glass volumes in Table 3B.9 are based on the same canister production schedule that was shown in the immobilized HLW section of this report (Chapter 3); about 5,280 canisters are projected by the end of year 2020, with a glass volume of 0.626 m³ per

canister (Garvin 1989). Tables 3B.10 and 3B.11, from the same source, show the estimated radioactivity and the total thermal power in the interim waste forms and in the HLW glass projected from the present through the year 2020. These projections assume that the amount of radioactivity in the glass product in a given year is the same as the average radioactivity in the tank farm, and therefore should not be used for year-by-year estimation of actual radioactivity per canister. The long-term estimates of cumulative radioactivity in the glass, however, should be useful. Table 3B.12 shows approximate chemical compositions of the interim waste forms and glass (IDB 1989, based on Garvin 1989).

3B.3.2 Treatment and Conversion to Glass

Glass produced at the DWPF will be made from a blend composed of (1) washed sludge, (2) washed precipitate produced by treatment of salt solution to precipitate the cesium, and (3) glass frit. Figure 3B.3 is a simplified representation of sludge and precipitate feed preparation and glass production. The salt solution that goes to the precipitation process contains the redissolved salt cake; thus the washed precipitate will contain essentially all the radioactivity originally in the supernate.

3B.3.2.1 Sludge Feed Preparation

In the tank farm, sludge with a minimum average age of 5 years is washed with caustic to remove aluminum (this step is omitted for low-aluminum sludges) and then washed with water to remove nitrate salts. The washed sludge is then pumped to the DWPF feed collection tank as a 13% solids slurry.

3B.3.2.2 Precipitate Feed Preparation

The combined salt solution and salt cake from a supernate aged to an acceptable Ru-106 limit are dissolved with water to form an alkaline solution saturated with sodium salts. This solution is diluted with recycled wash water and then treated with sodium tetraphenylborate (NaTPB) and sodium titanate. The NaTPB reacts with potassium and cesium to form precipitates of potassium and cesium tetraphenylborates. Strontium and plutonium are adsorbed into the precipitate phase by the sodium titanate. The solids, containing essentially all of the radioactivity, are then concentrated by filtration. The decontaminated filtrate is mixed with cement and fly ash to form a stable saltstone for engineered storage as LLW. The concentrated solids fraction from filtration is washed and filtered again and collected in a storage tank for transfer to the DWPF. The process described gives decontamination factors of about 4×10^4 for cesium, 200 for strontium, and 500 for plutonium (Doherty 1986). The concentrated and washed tetraphenylborate-titanate precipitate thus contains virtually

all of the radionuclide activity originally present in the supernate. However, it has only about one-tenth the original volume.

The precipitate is further treated in the DWPF by hydrolysis with formic acid to isolate and remove aromatic species derived from tetraphenylborate. This step permits collection of the aromatics as a separate benzene-rich phase that does not go to the melter but is incinerated separately, thus simplifying and reducing the size of the melter off-gas system.

3B.3.2.3 Melter Feed Preparation

The washed sludge stream and the washed hydrolyzed precipitate are mixed in a slurry adjustment tank and concentrated by boiling with formic acid. Mercury is reduced to its elemental state and is separated and collected. The adjusted slurry is mixed with glass frit slurry and concentrated by evaporation to a 40 wt % slurry which then goes to the melter feed tank (Baxter 1986a).

3B.3.2.4 Melter Operation

The electrically-heated melter is a refractory-lined steel vessel with a diameter of 2.6 m, a height of 3.2 m, and a shell thickness of 3.8 cm. The melter is operated with a "cold cap" composed of calcine and frit about 15 cm thick on top of the melt surface; the feed slurry is introduced on top of this cap to facilitate evaporation of water. The glass melt beneath the cold cap is at about 1050°–1170°C, enabling the cap to melt from below. Residence time in the melter is about 65 h, based on a nominal pour rate of 100 kg/h and a nominal melt weight of 6,500 kg.

3B.3.2.5 Canister Filling, Sealing, and Decontamination

Pouring is accomplished by lowering the pour spout pressure to 740 mm Hg while maintaining the vapor space above the melt at 750 mm Hg. The time required to fill a canister is about 17 h. The canister is then rotated from beneath the melter pour spout, and the canister shrink fit closure plug is inserted into the canister nozzle sleeve. The canister is then cooled to a surface temperature of 100°C over a period of about 31 h. Shrinkage of the canister nozzle and sleeve during cooling seals the closure plug tightly. The outer surface of the canister is then decontaminated by frit blasting, the plug and sleeve are pushed into the nozzle, and sealing of the closure is completed by welding the plug in place (Baxter 1986a).

3B.3.3 Radioactivity and Thermal Power Per Canister

The best available data (DPSP-80-1033, Rev. 91) were used to estimate the radionuclide composition of the most highly radioactive glass that will be made; that is, the glass made from sludge aged 5 years and supernate aged for an

average of 15 years (Baxter 1986c). This composition, shown in Table 3B.13, was used to calculate the radioactivity and thermal power per canister for decay times up to 10^6 years; these are the results that were presented in Sect. 3. As stated there, the radioactivity and thermal power at the time of filling, based on ORIGEN2 calculations, were 234,400 Ci and 709 W per canister. Detailed radionuclide analyses of individual feed batches to the DWPF will not be available until about one year before each batch is processed. Recent curie balances at SRS indicate that the glass produced during the first year of vitrification (1992) will have a radioactivity of about 154,000 Ci/canister and a heat generation rate of about 418 W per canister (Garvin 1990).

3B.3.4 Volume Conversion Factors to Glass

In DOE/RL-86-10, it was estimated that the sludge, salt, and liquid in storage at SRS at the end of 1986 are equivalent to about 4,900 canisters when converted to glass (DOE 1986). The IDB for 1987 gives the volumes of these wastes in storage at SRS at the end of 1986 as 128,000 m³ (IDB 1987). Referring to Table 3B.9, the volume in storage at the end of 1988 was 128,400 m³. The volume of glass in 4,900 canisters is about 3,100 m³. This gives a volume reduction factor from interim forms to glass of about 41. The volume of saltstone (LLW) is not included in this calculation.

3B.3.5 Assessment of Data

Using the best information available at present, it is believed that the maximum values of radioactivity and thermal power per canister will not exceed 234,400 Ci and 709 W. These estimates are based on sludge cooled 5 years and supernate cooled 15 years, which is the design basis for the DWPF. For repository design and other purposes, it would be useful to have a schedule of the radionuclide content of the vitrification plant feed as a function of the year of operation; but, as indicated above, these estimates will not be available until about one year before each batch is processed.

3B.4 HANFORD SITE

Acidic high-level wastes generated at Hanford by the reprocessing of defense reactor fuel have been made alkaline with caustic soda and stored in underground concrete tanks lined with carbon steel. Separation of solid and liquid phases occurs during storage. Storage tanks of both the single-shell tank (SST) and double-shell (DST) type are in use. All of the new waste goes to DSTs: no waste is currently being added to SSTs.

Prior to 1985, the HLW at Hanford was processed to remove most of the strontium and cesium; the separated strontium and cesium were encapsulated as strontium

fluoride and cesium chloride, and the capsules were placed in interim storage. During 1985, some of the capsules were distributed to industrial users under DOE agreements. Separation of Sr and Cs from the HLW was discontinued in 1985 and is not expected to be reinstated. Thus any quantities of these nuclides produced in the future will appear in HLW tank inventories. There are no definite plans at present to convert the Sr-Cs capsules to glass, but this possibility is still under consideration. It has been definitely decided that the Sr-Cs will go to a repository, but no decision has been made as to whether the capsules will be opened and their contents vitrified along with defense HLW, or whether the capsules will be encased in overpacks and shipped to a repository in that form (HANF 1989). Additional details are given in Sect. 3B.4.5.

3B.4.1 HLW Inventories at Hanford

The alkaline HLW stored at Hanford (245,000 m³ at the end of 1989) has been accumulating since 1944 and was generated by reprocessing fuel from production reactors for the recovery of plutonium, uranium, and other elements for defense and other federal programs. Tables 3B.14 and 3B.15 summarize the current and future volumes and radioactivities of these wastes, projected to the year 2020. Data in these tables are from Hanford's 1990 submittal to the IDB (Turner 1990). Volumes and radioactivities have not been adjusted downward to reflect the production of HLW glass, because the required year-by-year projections for glass are not yet available. Representative chemical compositions of interim wastes are shown in Table 3B.16.

The waste inventories at Hanford that are reported to the IDB as HLW are reported in terms of four phases. The first three of these (liquid, sludge, and salt cake) are terms used to describe the inventories held in SSTs. The fourth (slurry) is used to describe the total averaged inventory held in DSTs. Although the entire contents of both SSTs and DSTs are reported as part of the HLW submittal to the IDB, they actually include wastes in the TRU and LLW categories as well.

Most of the old waste (the waste in SSTs) has been treated to remove radioactive strontium and cesium. The Sr-90 and Cs-137 have been converted to solid strontium fluoride and cesium chloride and, together with their daughters, have been placed in double-wall stainless-steel capsules and stored in water basins. These capsules represent an interim HLW form that is considered separately from the tanked waste inventory.

Twenty-eight DSTs are in service with a total capacity of 118,400 m³. As of December 31, 1988, 78,000 m³ of DST waste were in storage. More DST wastes are expected, the primary source being the PUREX plant operations. Neutralized current acid waste (NCAW) from the PUREX plant will be the primary high-heat source of feed to vitrification. NCAW can be stored in any of four DSTs (called aging waste tanks) that are specifically

designed for this self-boiling waste. The other 24 DSTs are called non-aging waste tanks and are designed to contain wastes of lower heat generation rates, such as neutralized cladding removal waste (NCRW), complexant concentrate (CC), and plutonium finishing plant (PFP) waste. These wastes will also provide feed to the vitrification plant and have been described in Sect. 3.

The Evaporator-Crystallizer Facility plays an important part in Hanford's waste management treatment facilities. Evaporation of liquid radioactive waste and mixed waste in this facility achieve an annual waste volume reduction of 5 to 10 million gallons. This has reduced the storage space requirements for double-shell tanks by more than 100 million gallons, thus maximizing the use of available DST tanks.

At the end of 1988 there were approximately 139,000 m³ of damp salt cake and sludge containing 134 million curies of radionuclides stored in 149 underground single-shell tanks. These tanks range in capacity from 208 to 3,785 m³ each. The salt cake and sludge contain about 26,800 m³ of interstitial liquor as of December 31, 1988. Pumpable liquid is being transferred to double-shell tanks; this operation, referred to as interim stabilization, is scheduled to be completed by September 1996 (HANF 1989).

3B.4.2 Treatment and Conversion to Glass

As pointed out in HANF 1989, most of the radioactive wastes in interim tank storage at Hanford are either HLW or TRU mixed with relative large quantities of inert elements such as sodium, aluminum, iron, or zirconium. Because of the volume of inert elements, virtually all of the waste in storage and that to be generated will require some form of "pretreatment" processing prior to vitrification. The pretreatment process will fractionate the waste into a high-volume, low-activity stream (for disposal in near-surface grout vaults) and a low-volume, high-activity stream suitable for vitrification and disposal in a geologic repository.

Figure 3B.4 shows a schematic flowsheet of the existing and future HLW treatment and vitrification facilities at Hanford. The current plan is that the Hanford Waste Vitrification Plant (HWVP) will have four feeds: NCAW, NCRW, CC, and PFP. According to the present plan, these feeds will be run separately, producing canistered glasses of various levels of radioactivity and thermal power. As previously indicated, the NCAW produces the highest thermal power per canister, and is the reference feed for HWVP design purposes. The CC, NCRW, and PFP wastes will produce glass of much lower thermal power. The Hanford "B" plant will be used to pretreat the waste before it is sent to the HWVP. The treatment includes removal of sulfate (to produce glass of acceptable quality) and partial removal of Na, Al, F, and Zr to reduce glass volume

(Wolfe 1985a,b). Vitrification is scheduled to begin in 1999.

3B.4.2.1 Melter Feed Preparation

The slurry receipt and adjustment tank will be used as the main feed tank for the feed waste slurries containing the high-level and TRU wastes to be vitrified. In this tank the slurry solids content will be adjusted by evaporation to 12–19 wt %. Treatment with formic acid will also take place at this point to adjust the rheology and redox state for vitrification. The treated feed will then be blended with glass frit and further concentrated by evaporation in the slurry mix evaporator to about 30–49 wt % solids. When it has been determined by sampling that the composition meets all specified requirements, the concentrated slurry will be transferred to the melter feed tank for vitrification (Jain and Barnes 1989).

3B.4.2.2 On-Site Storage of Glass-Filled Canisters

An on-site storage facility will be constructed at Hanford with sufficient capacity to store 2,000 canisters of HLW glass. At the present time, the building is planned to have a heat removal capacity of 1,000 W per canister. This is higher than the currently-planned maximum loading of about 860 W per canister and will allow for the possible future vitrification of higher-heat wastes such as might be encountered if the decision is made to vitrify the contents of the strontium and cesium capsules.

3B.4.3 Radioactivity and Thermal Power Per Canister

Hanford has provided estimated maximum and nominal radionuclide compositions of the NCAW melter feeds. These are shown in Tables 3B.17 and 3B.18 in terms of curies and grams per canister. The maximum case represents the current estimate of the maximum radioactivity and thermal power for HLW glass produced at Hanford (Mitchell and Nelson 1988). As stated in Sect. 3, the maximum case results in a glass with an activity of 298,000 Ci/canister and a thermal power of 869 W/canister at the time of filling, based on a glass load of 1,650 kg (0.62 m³).

3B.4.4 Volume Conversion Factors to Glass

Hanford projections of glass production for NCAW are based on a waste oxides feed to vitrification of 29 kg per metric ton of uranium processed, and on the assumption of 25 wt % waste oxides loading in the glass. Given these bases, the volumetric reduction factor for NCAW to glass is about 30:1 based on feed to the pretreatment plant.

Glass production factors for the NCRW waste type are based on limiting the zirconium oxide loading in glass to 15 wt %, whereas for the PFP and CC waste types production

factors are based on 25 wt % total waste oxides loading in glass (White 1986, Watrous 1991).

3B.4.5 Sr-90 and Cs-137 Capsules

Until 1985, the defense HLW at Hanford was processed to separate the bulk of the strontium and cesium in concentrated form from the remainder of the HLW. The Sr-90 (half-life 28.5 years) and Cs-137 (half-life 30.0 years) were converted to solid forms as strontium fluoride and cesium chloride and placed in double-walled capsules for storage in water basins. Each capsule had an external diameter of about 6.7 cm and an overall length of about 53 cm. Other dimensional data are shown in Fig. 3B.5. Separation and encapsulation were completed during 1984 and resulted in a total of 640 Sr-90 capsules and 1577 Cs-137 capsules (White 1986, Turner 1990). The capsules include the short-lived daughter isotopes Y-90 and Ba-137m that are in transient equilibrium with the parent radionuclides.

A number of capsules were distributed with the intention of utilizing the contents for useful purposes, but most of the capsules have been returned to Hanford for disposal as HLW. Some of the returned capsules were dismantled for various reasons, and at the end of 1989 there were 597 Sr-90 and 1,350 Cs-137 capsules on hand for disposal (Wojtasek 1989, Turner 1990). A firm decision has been made that the contents of these capsules will go to a repository, but it has not been decided whether the capsules will be placed in overpack canisters for repository disposal or will have their contents reprocessed and disposed of as vitrified HLW.

3B.4.5.1 Radioactivity of Sr-90 and Cs-137 Capsules

The data used here for total radioactivity are from the 1990 IDB (Integrated Data Base) submittals from Hanford (Turner 1990). At the end of 1989, the 597 strontium capsules had a total radioactivity of 5.451×10^7 Ci, and the 1,350 cesium capsules had a total radioactivity of 1.187×10^8 Ci. Thus, at the end of 1989, each Sr-90 capsule had an average radioactivity of 9.13×10^4 Ci, and each Cs-137 capsule had an average radioactivity of 8.79×10^4 Ci. Total thermal outputs at the end of 1989 were 182.5 kW and 293.7 kW for all the Sr-90 and Cs-137 capsules, respectively. These were calculated from the curie quantities. The radioactivities and thermal outputs stated here include those of the daughter isotopes. Complete radioactivity and thermal power projections for the capsules through the end of year 2020 are given in Tables 3B.19 and 3B.20.

3B.4.5.2 Overpack Dimensions

As stated previously, it is not known yet whether the strontium and cesium capsules will be placed in overpack

canisters for shipment to a repository. However, Fig. 3B.6 shows two proposed designs (thin wall and cast steel) for overpack canisters for the capsules (White 1986). The design currently proposed in the Hanford reference plan is the thin-wall canister, which uses a carbon steel container with an outside diameter of 0.3 m (about 12 in.) and an overall length of 2.7 m (about 9 ft). An internal rack supports several (3 or 4) strontium or cesium capsules along the axis of the canister so that the decay heat is distributed over the entire area of the canister, thus avoiding excessive heat fluxes to the surrounding medium. This proposed canister design has not been finalized and is open to future changes.

3B.4.5.3 Number of Overpack Canisters

If overpack canisters are used, the number of canisters that will be required depends on the heat content of the capsules at the time of filling and the thermal limit, which is based on geologic repository heat load limits. The estimated thermal limits per overpack are 1.17 Kw for strontium and 0.8 kW for cesium; these figures include the daughter isotopes. The daughter isotopes Y-90 and Ba-137m account for about 83% of the heat load for a strontium capsule and about 77% of the heat load for a cesium capsule. Assuming that capsule overpacking is used and is completed at the end of the year 2010 (Turner 1990), the total thermal power of the capsules at the end of year 2010 will be 111 kW for the 597 strontium capsules and 181 kW for the 1,350 cesium capsules, including the daughter isotopes in each case.

Assuming further that the strontium and cesium capsules will not be intermingled and that a given canister must contain an integral number of capsules, then, to stay within the heat load limits stated, each overpack canister will contain four strontium or cesium capsules. This results in a total of 94 strontium canisters and 226 cesium canisters. Overpacking at a later date could result in a reduction in the total number of canisters because of decay. Also, as mentioned earlier, some capsules now on hand could be opened before disposal begins.

3B.4.6 Assessment of Data

Although Hanford's estimates of waste activity are necessarily based on projections 10 years in the future, the upper-bound estimate for the activity of vitrified NCAW waste appears to have been conservatively calculated. More definitive data on the expected radionuclide compositions of the various HLW glasses produced in the 1999-2007 period would be useful but will probably not be available until actual tankage allocations have been determined.

Quantitative data on strontium and cesium capsules are well established. Final plans for their disposal have not been made, but it has been decided that they will go to the repository in some form.

3B.5 IDAHO NATIONAL ENGINEERING LABORATORY

The HLW generated and stored at INEL results primarily from the reprocessing of spent fuels from naval propulsion nuclear reactors and reactor testing programs; a small amount is produced from the reprocessing of fuel from nondefense research reactors. The composition of the acidic liquid initially produced depends on the type of fuel processed; the most common composition contains fluorides and nitrates of aluminum and zirconium in nitric and hydrofluoric acids. The principal radionuclides are the fission products, such as Sr-90 and Cs-137, and transuranium elements. These acidic liquid wastes are stored in large, underground, doubly contained, stainless steel tanks, and are subsequently processed to yield a calcine. This is a dry powder consisting predominantly of aluminum oxide, zirconium oxide, and calcium fluoride; it contains the radionuclides that were present in the acidic liquid. The calcine is stored retrievably in stainless steel bins housed in reinforced concrete vaults, and is segregated into different layers within the bins according to the composition of the waste from which it was derived (Knecht et al. 1985).

3B.5.1 HLW Inventories at INEL

Tables 3B.21 and 3B.22 summarize the current and projected future volumes and radioactivities of the liquid and calcine stored at INEL, projected to the year 2020. The amount of waste generated, total waste curie content, and heat generation rate are lower than prior projections because of a projected decrease in reprocessing rate through the year 2020 (Berreth 1989, 1990). Table 3B.21 also shows the canister production schedule on which the estimates are based; immobilization is assumed to start in year 2012 at a rate of 500 canisters/year and reaches a steady rate of 1,000 canisters/year in year 2014. Production of immobilized waste is expected to continue after year 2020.

The estimates shown in Table 3B.22 of radioactivity in the immobilized form were based on the assumption that the glass-ceramic produced each year would be made from calcine having the same average radionuclide composition as the average calcine inventory in storage in that year. This assumption is unlikely to hold precisely true in practice, so the estimates in Table 3B.22 should not be used for near-term estimates of the total radioactivity in the immobilized form.

Tables 3B.23 and 3B.24 show representative chemical compositions of various HLW liquids and calcines produced at INEL (IDB 1989).

3B.5.2 Treatment and Conversion to Glass-Ceramic

As already indicated, immobilization of HLW at INEL is scheduled to start in the year 2012. Various alternatives for immobilization are being studied, and no decision has been made as to what form the immobilized waste will be in for repository disposal. Figure 3B.7 shows a schematic flowsheet illustrating some of the possibilities for treatment and immobilization. As the flowsheet shows, the conversion of liquid waste to glass without passing through the calciner is one of the alternatives under study. Both glass and glass-ceramic compositions are being considered as the final waste form for the calcine and liquid HLW. Previous studies have indicated that use of the glass-ceramic form could reduce the total volume of immobilized waste by as much as 60% compared to the glass form (Knecht et al. 1985, Berreth and Knecht 1986). This potential reduction in the total number of canisters of immobilized HLW is the primary reason for favoring the glass-ceramic form.

Based on the glass-ceramic form, the future annual production of immobilized HLW is now estimated to be in the range of 450 to 650 canisters/year. The larger quantity corresponds to fuel dissolution using cadmium and boron for criticality control, while the smaller does not, relying on the use of critically safe dissolvers instead. These canister production rates are based on the annual generation of new waste. If stored calcine from old waste is also retrieved and immobilized, an additional 350 to 550 canisters of glass-ceramic would be produced annually, resulting in a total production of about 1,000 canisters/year. Other options described in WINCO-1031, such as neutralization, are potentially capable of reducing the annual volume of immobilized HLW, but are not as far developed as the ceramic forms described here, and would also produce large volumes of low-level waste. Also, these latter options at present are based on rather advanced processes and therefore are not as technically favorable as the glass-ceramic forms for near-term implementation. The remaining options shown in WINCO-1031 have been shown to be less promising and are not being developed any further (Knecht 1986a). The estimates in Chapter 3 of this report were based on 1,000 canisters/year of the glass-ceramic form, since that appears currently to be the most favorable case.

3B.5.3 Radioactivity and Thermal Power Per Canister

The estimated radioactivity and thermal power per canister shown in Sect. 3 of this report were based on an estimated radionuclide composition of 3-year old calcine taken from a 1982 INEL study (IDO 1982). This composition, shown in Table 3B.25, is the most recent available. To complete the calculation, it was also necessary to estimate the mass of calcine per canister. Based on the high-density glass-ceramic form, calculations indicated that about 1,277 kg of calcine could be incorporated in a

canister load. Assumptions used in making this estimate were as follows: (1) the volume of glass-ceramic per canister is 0.57 m³, (2) the density of the glass-ceramic is 3.2 g/cm³, and (3) the amount of calcine in the glass-ceramic is 70% by weight. These assumptions were based on data in Berreth and Knecht 1984, Knecht et al. 1985, Knecht 1986a, and Berreth 1986a. The radionuclide content per canister that resulted from the foregoing assumptions is shown in Table 3B.26; this is the composition that was used in Sect. 3 to develop radioactivity and thermal power per canister. For purposes of this report, this composition is estimated to represent the maximum radioactivity per canister as well as we can estimate it at the present time.

3B.5.4 Volume Conversion Factor to Ceramic

The bulk density of calcine is about 1.4 g/cm³ (Berreth 1986b). Thus the 1,277 kg of calcine in a canister load would have a volume of 0.91 m³ as calcine. Since its volume as glass-ceramic was estimated to be 0.57 m³, the volume reduction factor from calcine to glass-ceramic is about 1.6:1. The volume reduction factor for liquid HLW to calcine is about 6:1 (Mairson et al. 1986), and the overall volume reduction factor (liquid to glass-ceramic) is estimated to be about 9.6:1. Because of differences in flowsheets, this value should not be compared with those at WVDP, SRS, and HANF.

3B.5.5 Assessment of Data

Because of security restrictions, INEL has not released any estimates of radionuclide content per canister based on either glass or glass-ceramic. The estimates of radionuclide composition in this report, which served as the basis for ORIGEN decay calculations to determine maximum radioactivity and thermal power as a function of time, were based on 1982 data. Thus the maximum radioactivity and decay heat loads per canister shown in Sect. 3 for decay times up to 10⁶ years should be considered preliminary. Also, final decisions on processing options for INEL have not yet been made; thus the processing route, schedule, and canistered waste characteristics presented here are based on assumptions that are subject to change.

3B.6 REFERENCES FOR APPENDIX 3B

Barnes et al. 1986. S. M. Barnes, L. L. Petkus et al., *Reference Vitrification Process and Equipment Description with Results from Checkout Testing*, DOE/NE/44139-32 (DE87009317), November 1986.

Baxter 1983. Richard G. Baxter, *Description of Defense Waste Processing Facility Reference Waste Form and Canister*, DuPont Report No. DP-1606, Rev. 1, August 1983.

Baxter 1986a. R. G. Baxter, "Design and Construction of the Defense Waste Processing Facility Project at the Savannah River Plant," presented at Waste Management '86, Tucson, Arizona, March 2-6, 1986.

Baxter 1986c. Telephone conversation, R. G. Baxter and R. Salmon, May 16, 1986.

Baxter 1987. Letter from R. G. Baxter, SRS, to Royes Salmon, ORNL, February 18, 1987.

Berreth 1986a. Letter from J. R. Berreth, INEL, to J. E. Solecki, DOE, BERR-21-86, dated March 14, 1986.

Berreth 1986b. Telephone conversation, J. R. Berreth (INEL) and R. Salmon (ORNL), June 17, 1986.

Berreth 1986c. Letter from J. R. Berreth, INEL, to K. J. Notz, ORNL, December 1, 1986.

Berreth and Knecht 1984. J. R. Berreth and D. A. Knecht, "Defense High-Level Waste Management at the Idaho Chemical Processing Plant," Journal of Nuclear Materials Management, Processing Issue, Vol. XIII, pp. 7-13, July 1984.

Berreth and Knecht 1986. *Potential Process Options to Minimize Immobilized HLW Volume at the Idaho Chemical Processing Plant*, WINCO-M-10079, September 1986.

Berreth 1987. Letters from J. Berreth, INEL, to J. E. Solecki, DOE/IDO, March 19, 1987 and April 1, 1987.

Berreth 1989. J. R. Berreth, Westinghouse Idaho Nuclear Company, Inc., Idaho Falls, Idaho, letter to B. J. Mikkola, U.S. Department of Energy, Idaho Operations Office, Idaho Falls, Idaho, "FY-89 Integrated Data Base Information," Berr-11-89, dated Mar. 17, 1989.

Berreth 1990. Letter from J. R. Berreth, INEL, to M. J. Bonkowski, DOE Idaho Operations Office, "FY 90 Integrated Data Base Information," dated March 28, 1990, with revisions by D. Knecht, June 15, 1990.

Bixby 1987. Letter from W. W. Bixby, West Valley Project Office, to S. N. Storch, ORNL, February 27, 1987.

Boore 1986. W. G. Boore et al., *Radioactive Waste Spill and Cleanup on Storage Tank at the Savannah River Plant*, DP-1722, March 1986.

Crocker 1989. *West Valley Vitrification Process Mass Balance, Revision 7*, October 10, 1989.

Crocker 1990. Telecommunications from Bob Crocker, WVDP, to S. N. Storch, ORNL, May 23 and June 11, 1990.

DOE 1985. *An Evaluation of Commercial Repository Capacity for the Disposal of Defense High-Level Waste — Responses to Comments*, DOE/DP-0027, December 1985.

DOE 1986. *Perspective on Methods to Calculate a Fee for Disposal of Defense High-Level Waste in Combined (Civilian/Defense) Repositories*, DOE/RL-86-10, U.S. Department of Energy, Richland Operations Office, Richland, Washington (1986).

Doherty 1986. J. P. Doherty, "Process Innovations to Minimize Waste Volumes at Savannah River," presented at AIChE meeting in Boston, MA, August 24-27, 1986.

DPSP 80-1033. *Basic Data Report, Defense Waste Processing Facility, Savannah River Plant*, DPSP 80-1033, Rev. 91, April 1985.

DPSP 1982. *Waste Management Programs Report for December 1982*, E. I. duPont de Nemours & Co., Savannah River Plant, Report No. DPSP 82-21-12, December 1982.

DWMP 1983. U.S. DOE, *The Defense Waste Management Plan*, DOE/DP-0015 (June 1983).

Eisenstatt 1986. *Description of the West Valley Demonstration Project Reference High-Level Waste Form and Canister*, WVDP-056, July 1986.

Fehrerger 1985. D. J. Fehrerger, *An Evaluation of Radionuclide Concentrations in High-Level Radioactive Wastes*, U.S. Nuclear Regulatory Commission, NUREG-0946, October 1985.

GAO 1989. *Nuclear Waste: DOE's Program to Prepare High-Level Radioactive Waste for Final Disposal*, U.S. General Accounting Office, GAO/RCED-90-46 FS, November 1989.

Garvin 1989. R. G. Garvin, Westinghouse Savannah River Company, Savannah River Site, Aiken, South Carolina, memo to M. G. O'Rear, U.S. Department of Energy, Savannah River Operations Office, Aiken, South Carolina, "Data for Integrated Data Base," dated June 30, 1989.

Garvin 1990. R. G. Garvin letter to A. L. Watkins, DOE Savannah River Office, May 22, 1990.

HANF 1988. *Hanford Site Waste Management Plan*, U.S. Department of Energy Report DOE/RL-88-33, December 1988.

- HANF 1989. *Hanford Site Waste Management Plan*, U.S. Department of Energy Report DOE/RL-89-32, December 1989.
- Hannum 1983. W. H. Hannum, *Analysis of the Terminal Waste Form Selection for the West Valley Demonstration Project*, DOE/NE/44139-T3, January 1983.
- IDB 1987. *Integrated Data Base for 1987: Spent Fuel and Radioactive Waste Inventories, Projections, and Characteristics*, prepared for U.S. Department of Energy by Oak Ridge National Laboratory, DOE/RW-0006, Rev. 3 (September 1987).
- IDB 1988. *Integrated Data Base for 1988: Spent Fuel and Radioactive Waste Inventories, Projections and Characteristics*, prepared for U.S. Department of Energy by Oak Ridge National Laboratory, DOE/RW-0006, Rev. 4, September 1988.
- IDB 1989. *Integrated Data Base for 1989: Spent Fuel and Radioactive Waste Inventories, Projections, and Characteristics*, prepared for U.S. Department of Energy by Oak Ridge National Laboratory, DOE/RW-0006, Rev. 5, (November 1989).
- IDO 1982. *Environmental Evaluation of Alternatives for Long-term Management of Defense High-level Radioactive Wastes at the Idaho Chemical Processing Plant*, Department of Energy Idaho Operations Office Report IDO-10105, September 1982.
- Jain and Barnes 1989. V. Jain and S. M. Barnes, *Waste Description and Previtritification Treatment Activities at WVDP, HWVP, and DWPF*, DOE/NE/44139-57, February 1989.
- Knecht et al. 1985. D. A. Knecht et al., *Scoping Studies to Reduce ICPP High-Level Radioactive Waste Volumes for Final Disposal*, prepared for U.S. Department of Energy by Westinghouse Idaho Nuclear Co., WINCO-1031, August 1985.
- Knecht 1986a. Letter from D. A. Knecht, INEL, to Royes Salmon dated April 11, 1986.
- Maestas 1990. E. Maestas, DOE-West Valley Project Office, West Valley, New York, letter to S. N. Storch, Oak Ridge National Laboratory, Oak Ridge, Tennessee, "Reissue of the West Valley Demonstration Project (WVDP) Data Update for the DOE 1989 Integrated Data Base Report," dated May 3, 1990.
- Mairson et al. 1986. R. C. Mairson, B. R. Wheeler, and J. B. Whitsett, "High Level Waste Management at the ICPP," Tucson Symposium on Waste Management, March 1986.
- Martin 1986. H. D. Martin, R. G. Baxter, et al., *Costs of SRS High-Level Waste Processing in the Defense Waste Processing Facility*, DPST-86-313, December 1986.
- Mitchell 1986. D. E. Mitchell, *Hanford Waste Vitrification Plant, Preliminary Description of Waste Form and Canister*, RHO-RE-SR-55P, August 1986.
- Mitchell and Nelson 1988. D. E. Mitchell and J. L. Nelson, *Hanford Waste Vitrification Plant Preliminary Description of Waste Form and Canister — FY 1988 Update*, WHC-EP-0008, Rev. 1, June 1988.
- Mitre 1984. *An Evaluation of Commercial Repository Capacity for the Disposal of Defense High-Level Waste*, DOE/DP-0020 (Draft), prepared by the Mitre Corporation for the U.S. Department of Energy, July 1984.
- NFS 1973. *Safety Analysis Report of the Nuclear Fuel Services Reprocessing Plant at West Valley, New York*, Docket N. 50-201, NFS Incorporated, Rockville, Maryland (1973).
- O'Rear 1989. Data Transmittal from M. G. O'Rear, DOE Savannah River, to Jerry Klein, ORNL, July 10, 1989.
- Rockwell 1985. *Hanford Defense Waste Disposal Alternatives: Engineering Support Data for the Hanford Defense Waste Environmental Impact Statement*, prepared by Rockwell International for U.S. DOE, DOE/RL/01030-1, July 1985 (Draft).
- Roddy et al. 1985. J. W. Roddy et al., *Physical and Decay Characteristics of Commercial LWR Spent Fuel*, Oak Ridge National Laboratory Report No. ORNL/TM-9591/V1, September 1985.
- Rykken 1985. Personal communication, L. E. Rykken (WVDP), to Herschel W. Godbee (ORNL), April 9, 1985.
- Rykken 1986a. Letter from L. E. Rykken to R. Salmon, April 11, 1986.
- Rykken 1986d. Telephone conversation, L. E. Rykken and R. Salmon, October 2, 1986.
- Rykken 1987. Telephone conversation, L. E. Rykken and R. Salmon, March 25, 1987.

SRS 1987. Comments transmitted to DOE and ORNL by SRS representatives, March 10, 1987.

Staples, Knecht, and Berreth 1986. B. A. Staples, D. A. Knecht, and J. R. Berreth, *Technology for the Long-Term Management of Defense HLW at the ICPP*, Westinghouse Idaho Report WINCO-1038, June 1986.

Turner 1990. Letter from D. A. Turner, Westinghouse Hanford, to R. E. Gerton, DOE/RL, April 16, 1990.

USDOE 1985. *The Defense Waste Management Plan, February 1985 Update*, prepared by the Mitre Corporation for U.S. Department of Energy (Draft).

Watrous 1986. Telephone conversations, R. A. Watrous (Rockwell Hanford) and Royes Salmon (ORNL), May-August 1986.

Watrous 1991. Telephone conversation, R. A. Watrous and R. Salmon, July 1991.

White 1986. Letter from J. D. White, DOE/RL, to W. R. Bibb, ORNL, dated July 3, 1986.

Wilde 1986a. Letter from R. T. Wilde, Rockwell Hanford Operations, to M. W. Shupe, DOE/RLO, dated April 18, 1986.

Wilde 1986b. Letter from R. T. Wilde, Rockwell Hanford Operations, to Herschel W. Godbee, ORNL, dated May 5, 1986.

Wojtasek 1989. R. D. Wojtasek, Westinghouse Hanford Company, Richland, Washington, memo to R. E. Gerton, U.S. Department of Energy, Richland Operations Office, Richland, Washington, "Information for the National Integrated Data Base: High-Level Waste Submittal," dated Apr. 3, 1989.

Wolfe 1985a. Personal communication, B. A. Wolfe, Rockwell Hanford Operations, to Royes Salmon, ORNL, November 8, 1985.

Wolfe 1985b. Personal communication, B. A. Wolfe, Rockwell Hanford Operations, to Royes Salmon, November 12, 1985. (Project Status, HWVP, Nov. 6, 1985).

WVDP 1989. West Valley Demonstration Project Annual Report, WVDP/89AR, January 1989.

WVDP 1990. West Valley Demonstration Project Annual Report, WVDP/90AR, January 1990.

Fig 3B.1. HLW treatment and vitrification at WVDP. Source: IDB 1989.

3B-15

Fig. 3B.2. Simplified representation of primary HLW operations at Savannah River Site.

3B-16

Fig. 3B.3. HLW processing at Savannah River Site after startup of DWPF.

3B-17

Fig. 3B.4. Hanford site: schematic flowsheet of existing and future HLW processing facilities. Adapted from HANF 1989.

	FORM	LOADING	PERCENT OF THEORETICAL DENSITY BASED ON TOTAL VOID SPACE OF CAPSULE	TEMPERATURE			
				AIR		WATER	
				CENTER LINE	SURFACE	CENTER LINE	SURFACE
STRONTIUM FLUORIDE	COMPACTED POWDER	150 kCi (MAX)	68	860°C	430°C	680°C	71°C
CESIUM CHLORIDE	MELT-CAST	70 kCi	65	450°C	200°C	327°C	58°C

	CAPSULE									
	INNER					OUTER				
	MATERIAL	WALL THICKNESS	OUTSIDE DIAMETER	TOTAL LENGTH	TOTAL CAP THICKNESS	MATERIAL	WALL THICKNESS	OUTSIDE DIAMETER	TOTAL LENGTH	TOTAL CAP THICKNESS
STRONTIUM FLUORIDE	HASTELLOY C-276 (UT)	0.305 (UT)	5.72	48.39	1.02	STAINLESS STEEL 316-L (UT)	0.277 (UT)	6.67	51.05	1.02
CESIUM CHLORIDE	STAINLESS STEEL 316-L (UT)	0.241 (UT)	5.72	50.10	1.02	STAINLESS STEEL 316-L (UT)	0.277 (UT)	6.67	52.77	1.02

NOTE: ALL DIMENSIONS ARE IN cm

Fig. 3B.5. Dimensional data for strontium and cesium capsules. Source: White 1986.

	OVERPACK DIMENSIONS		
	OVERALL LENGTH (m)	OUTSIDE DIAMETER (m)	APPROXIMATE REQUIRED
THIN WALL CANISTER	2.7	0.3	513
MASSIVE CAST STEEL OVERPACK	1.2	0.6	1300

PS8310-21

Fig. 3B.6. Overpack concepts for strontium and cesium capsules. Source: White 1986.

Fig. 3B.7. HLW treatment and vitrification at INEL. Source: IDB 1989.

Table 3B.1. West Valley reprocessing summary, 1966-1972^a

Reactor	Type	Fuel	Cladding	Campaigns	Uranium (metric tons)	Exposure (MWd)
Yankee-Rowe	PWR	UO ₂	SS	4	88	1,412,000
CON ED/Indian Pt-1	PWR	UO ₂	SS	2	23	426,000
CVNPA	PWR	UO ₂	Zr	1	4	35,000
COMM ED/Dresden-1	BWR	UO ₂	Zr	2	72	661,000
CP/Big Rock Pt	BWR	UO ₂	SS	2	24	252,000
PG&E/Humboldt Bay	BWR	UO ₂	Zr	1	21	220,000
NSP/Pathfinder	BWR	UO ₂	SS	1	10	15,000
PR/Bonus	BWR	UO ₂	SS	1	4	11,000
NPR	Graphite	Metal	Zr	11	380	774,000
CON ED/Indian Pt-1	PWR	ThO ₂ /UO ₂ ^c	Zr	1	16 ^b	260,000
SEFOR	Test	PuO ₂ /UO ₂		1	20 ^d	
Total				27	662	4,066,000

^aSource: NFS 1973^bUranium plus thorium.^c93% U-235.^dPlutonium only processed.

Table 3B.2. Historical and projected total cumulative volume, radioactivity, and thermal power of interim forms of HLW stored at WVDP^a

End of calendar year	Cumulative quantities		
	Volume (10 ³ m ³)	Radioactivity (10 ⁶ Ci)	Thermal power (10 ³ W)
1989	1.9	27.9	83.2
1990	1.6	27.2	81.2
1991	1.4	26.6	79.3
1992	1.2	26.0	77.5
1993	1.0	23.0	69.6
1994	0.2	4.5	13.4
1995 ^b	0.0	0.0	0.0

^aThis table includes all forms of HLW at WVDP except vitrified HLW. The table is based on information in Table 2.1 of IDB 1990, with modifications due to differences in the assumed vitrification schedules. In the schedule assumed here, which is the same as that shown in Tables 3.2.2 and 3B.3, the total HLW at the site is assumed to be 9.1% vitrified at the end of 1993, 81.8% vitrified at the end of 1994, and 100% vitrified at the end of 1995. The schedule used in IDB 1990 was based on 16.7% vitrified by the end of 1993, 83.3% vitrified by the end of 1994, and 100% vitrified by the end of 1995.

^bQuantities of interim HLW after completion of vitrification are zero.

Table 3B.3. Projected volume, radioactivity, and thermal power of HLW glass produced at WVDP^a

End of calendar year	Volume (10 ³ m ³)	Radioactivity (10 ⁶ Ci)	Thermal power (10 ³ W)
1992 ^b	0.000	0.0	0.0
1993	0.016	2.3	6.9
1994	0.146	20.2	60.4
1995	0.179	24.2	72.2
1996	0.179	23.6	70.5
1997	0.179	23.0	68.7
1998	0.179	22.5	67.2
1999	0.179	22.0	65.6
2000	0.179	21.5	64.2
2001	0.179	21.0	62.7
2002	0.179	20.5	61.2
2003	0.179	20.0	59.9
2004	0.179	19.5	58.6
2005	0.179	19.1	57.2
2006	0.179	18.6	55.9
2007	0.179	18.2	54.6
2008	0.179	17.8	53.5
2009	0.179	17.4	52.2
2010	0.179	17.0	51.0
2011	0.179	16.6	49.9
2012	0.179	16.2	48.8
2013	0.179	15.8	47.6
2014	0.179	15.5	46.6
2015	0.179	15.1	45.5
2016	0.179	14.8	44.4
2017	0.179	14.4	43.6
2018	0.179	14.1	42.6
2019	0.179	13.8	41.6
2020	0.179	13.4	40.6

^aThis is based on the assumption that vitrification at WVDP begins in 1993 and is completed in 1995 (Maestas 1990), and on the vitrification schedule assumed in Tables 3.2.2 and 3B.2. The total volume of glass produced (179 m³) is based on the mass of glass (484,000 kg) shown in the Revision 7 Mass Balance (Crocker 1989), and on the estimated glass density (2.70 g/cm³) shown in Eisenstatt 1986. Radioactivity and thermal power are based on ORIGEN2 calculations using the Revision 7 Mass Balance.

^bGlass quantities prior to 1993 are zero.

Table 3B.4. Chemical composition of alkaline liquid HLW
(from reprocessing via a PUREX flowsheet) at WVDP^a

Compound	Wet basis (wt %)	Dry basis (wt %)	Total (kg)
NaNO ₃	21.10	53.38	602,659
NaNO ₂	10.90	27.57	311,326
Na ₂ SO ₄	2.67	6.75	76,261
NaHCO ₃	1.49	3.77	42,557
KNO ₃	1.27	3.21	36,274
Na ₂ CO ₃	0.884	2.24	25,249
NaOH	0.614	1.55	17,537
K ₂ CrO ₄	0.179	0.45	5,113
NaCl	0.164	0.42	4,684
Na ₃ PO ₄	0.133	0.34	3,799
Na ₂ MoO ₄	0.0242	0.06	691
Na ₃ BO ₃	0.0209	0.05	597
CsNO ₃	0.0187	0.05	534
NaF	0.0176	0.04	503
Sn(NO ₃) ₄	0.00858	0.02	245
Na ₂ U ₂ O ₇	0.00809	0.02	231
Si(NO ₃) ₄	0.00805	0.02	230
NaTcO ₄	0.00620	0.02	177
RbNO ₃	0.00417	0.01	119
Na ₂ TeO ₄	0.00287	0.007	82
AlF ₃	0.0027	0.0068	77
Fe(NO ₃) ₃	0.00151	0.004	43
Na ₂ SeO ₄	0.00053	0.0013	15
LiNO ₃	0.00049	0.0012	14
H ₂ CO ₃	0.00032	0.00080	9
Cu(NO ₃) ₃	0.00021	0.00053	6
Sr(NO ₃) ₂	0.00014	0.00035	4
Mg(NO ₃) ₂	0.00007	0.00018	2
Subtotal	39.53	100.00	1,129,038
H ₂ O (by difference)	60.47	0.00	1,727,116
Grand total	100.00	100.00	2,856,154

^aSource: IDB 1990.

Table 3B.5. Chemical composition of alkaline sludge HLW
(from reprocessing via a PUREX flowsheet) at WVDP^a

Compound	Weight, kg
Fission products	
Ge(OH) ₃	0.2
SrSO ₄	217
Y(OH) ₃	103
Zr(OH) ₄	964
Ru(OH) ₄	458
Rh(OH) ₄	79
Pd(OH) ₄	34
AgOH	0.7
Cd(OH) ₂	1.7
In(OH) ₃	0.3
Sn(OH) ₄	2.5
Sb(OH) ₃	0.7
BaSO ₄	303
La(OH) ₃	185
Ce(OH) ₃	354
Pr(OH) ₃	170
Nd(OH) ₃	621
Pm(OH) ₃	1.5
Sm(OH) ₃	143
Eu(OH) ₃	7.5
Gd(OH) ₃	1.7
Tb(OH) ₃	0.3
Dy(OH) ₃	0.2
Subtotal	3,648.3
Actinides	
UO ₂ (OH) ₂	3,087
NpO ₂	35
PuO ₂	37
AmO ₂	27
CmO ₂	0.4
Subtotal	3,186.4
Others	
Fe(OH) ₃	66,040
FePO ₄	6,351
Al(OH) ₃	5,852
AlF ₃	613
MnO ₂	4,581
CaCO ₃	3,208
SiO ₂	1,263
Ni(OH) ₂	1,088

Table 3B.5 (continued)

Compound	Weight, kg
Others (continued)	
MgCO ₃	826
Cu(OH) ₂	376
Zr(OH) ₄	964 ^b
Zn(OH) ₂	128
Cr(OH) ₃	65
Hg(OH) ₃	23
Subtotal	91,378
Grand total	98,213

^aSource: Maestas 1990.^bExcludes fission product zirconium.

Table 3B.6. Chemical composition of acid liquid HLW
(from reprocessing via a THOREX flowsheet) at WVDPA

Compound	Wt %	Total, kg
Th(NO ₃) ₄	36.42	31,054
Fe(NO ₃) ₃	9.92	8,462
Al(NO ₃) ₃	4.90	4,175
HNO ₃	3.29	2,805
Cr(NO ₃) ₃	2.25	1,918
Ni(NO ₃) ₂	0.93	79
H ₃ BO ₃	0.56	480
NaNO ₃	0.27	227
KNO ₃	0.22	191
Na ₂ SO ₄	0.21	180
Na ₂ SiO ₃	0.15	126
KMnO ₄	0.11	98
Nd(NO ₃) ₃	0.086	73
Mg(NO ₃) ₂	0.067	57
Na ₂ MoO ₄	0.063	54
NaCl	0.059	50
Ce(NO ₃) ₄	0.050	43
Ru(NO ₃) ₄	0.049	42
ZrO ₂	0.041	35
Ca(NO ₃) ₂	0.035	30
CsNO ₃	0.033	28
Ba(NO ₃) ₂	0.032	27
La(NO ₃) ₃	0.026	22
Pr(NO ₃) ₃	0.025	21
Sr(NO ₃) ₂	0.019	16
Y(NO ₃) ₃	0.016	14
Sm(NO ₃) ₃	0.016	14
Zr(NO ₃) ₄	0.014	12
Na ₃ PO ₄	0.014	12
NaTcO ₄	0.013	11
Rh(NO ₃) ₄	0.013	11
Zn(NO ₃) ₂	0.012	10
Pd(NO ₃) ₄	0.0094	8
UO ₂ (NO ₃) ₂	0.0070	6
RbNO ₃	0.0070	6
Na ₂ TeO ₄	0.0059	5
Co(NO ₃) ₂	0.0035	3
Na ₂ SeO ₄	0.0012	1
NaF	0.0012	1
Eu(NO ₃) ₃	0.0012	1
Np(NO ₃) ₄	0.0011	0.9

Table 3B.6 (continued)

Compound	Wt %	Total, kg
Cu(NO ₃) ₂	0.00094	0.8
Sn(NO ₃) ₃	0.00082	0.7
Pa(NO ₃) ₄	0.00082	0.7
Pu(NO ₃) ₄	0.00082	0.7
Gd(NO ₃) ₃	0.00047	0.4
Cd(NO ₃) ₂	0.00035	0.3
Sb(NO ₃) ₃	0.00012	0.1
AgNO ₃	0.000094	0.08
In(NO ₃) ₃	0.000047	0.04
Ge(NO ₃) ₄	0.000023	0.02
Pm(NO ₃) ₂	0.000011	0.01
Tb(NO ₃) ₃	0.0000047	0.004
Dy(NO ₃) ₃	0.0000023	0.002
Solids	59.95	51,125
H ₂ O (by difference)	40.05	34,148
Total	100.00	85,273

^aSource: IDB 1990.

Table 3B.7. Reference 1989 radionuclide composition of HLW at WVDp^{a,b}

Radionuclide	Alkaline waste (PUREX)		Acid waste (THOREX)	Total, Ci
	Liquid, ^c Ci	Sludge, Ci	Liquid, Ci	
H-3	8.23E+01	0.00E+00	1.47E+00	8.38E+01
C-14	1.37E+02	0.00E+00	1.30E-01	1.37E+02
Fe-55	0.00E+00	4.49E+02	2.53E+02	7.02E+02
Co-60	0.00E+00	3.17E+00	7.68E+02	7.71E+02
Ni-59	0.00E+00	8.56E+01	2.03E+01	1.06E+02
Ni-63	8.69E+02	5.22E+03	2.45E+03	8.54E+03
Se-79	5.68E+01	0.00E+00	3.35E+00	6.01E+01
Sr-90	2.69E+03	6.28E+06	4.23E+05	6.70E+06
Y-90	2.69E+03	6.28E+06	4.23E+05	6.70E+06
Zr-93	2.56E-01	2.56E+02	1.62E+01	2.72E+02
Nb-93m	1.73E-01	1.71E+02	1.11E+01	1.82E+02
Tc-99	1.60E+03	0.00E+00	1.04E+02	1.70E+03
Ru-106	1.40E-02	1.40E+01	7.96E-02	1.41E+01
Rh-106	1.40E-02	1.40E+01	7.97E-02	1.41E+01
Pd-107	1.09E-02	1.09E+01	1.14E-01	1.10E+01
Cd-113m	2.09E+00	2.09E+03	3.25E+01	2.12E+03
Sn-121m	1.69E-02	1.69E+01	5.75E-01	1.75E+01
Sn-126	1.01E-01	1.01E+02	3.11E+00	1.04E+02
Sb-125	2.31E+01	7.13E+03	1.36E+02	7.29E+03
Sb-126	1.41E-02	1.41E+01	4.35E-01	1.46E+01
Sb-126m	1.01E-01	1.01E+02	3.11E+00	1.04E+02
Te-125m	5.67E+00	1.75E+03	3.34E+01	1.79E+03
I-129	2.10E-01	0.00E+00	1.80E-01	3.90E-01
Cs-134	5.07E+03	0.00E+00	1.13E+02	5.18E+03
Cs-135	1.56E+02	0.00E+00	5.47E+00	1.61E+02
Cs-137	6.77E+06	0.00E+00	4.43E+05	7.22E+06
Ba-137m	6.41E+06	0.00E+00	4.19E+05	6.83E+06
Ce-144	1.45E-06	6.38E-01	9.63E-03	6.48E-01
Pr-144	1.45E-06	6.39E-01	9.64E-03	6.48E-01
Pm-146	3.27E-02	1.05E+01	3.47E-01	1.09E+01
Pm-147	2.58E+02	8.37E+04	4.12E+03	8.80E+04
Sm-151	4.92E-01	7.96E+04	4.67E+03	8.43E+04
Eu-152	3.92E-02	3.24E+02	4.14E+01	3.65E+02
Eu-154	1.13E+01	9.34E+04	1.99E+03	9.54E+04
Eu-155	1.56E+00	2.33E+04	5.55E+02	2.38E+04

Table 3B.7 (continued)

Radionuclide	Alkaline waste (PUREX)		Acid waste (THOREX)	Total, Ci
	Liquid, ^c Ci	Sludge, Ci	Liquid, Ci	
Tl-207	1.87E-08	8.55E-04	8.19E+00	8.19E+00
Tl-208	7.31E-02	1.04E-00	2.18E+00	3.29E+00
Pb-209	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Pb-211	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Pb-212	2.04E-01	2.88E+00	6.08E+00	9.17E+00
Bi-211	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Bi-212	2.04E-01	2.88E+00	6.08E+00	9.17E+00
Bi-213	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Po-212	1.30E-01	1.85E+00	3.90E+00	5.88E+00
Po-213	1.38E-04	1.93E-03	2.03E-01	2.05E-01
Po-215	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Po-216	2.04E-01	2.88E+00	6.08E+00	9.17E+00
At-217	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Rn-219	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Rn-220	2.04E-01	2.88E+00	6.08E+00	9.17E+00
Fr-221	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Fr-223	2.59E-10	1.18E-05	1.13E-01	1.13E-01
Ra-223	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Ra-224	2.04E-01	2.88E+00	6.08E+00	9.17E+00
Ra-225	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Ra-228	0.00E+00	5.09E-09	1.52E+00	1.52E+00
Ac-225	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Ac-227	1.88E-08	8.57E-04	8.22E+00	8.22E+00
Ac-228	0.00E+00	5.09E-09	1.52E+00	1.52E+00
Th-227	1.85E-08	8.45E-04	8.10E+00	8.10E+00
Th-228	2.04E-01	2.88E+00	6.08E+00	9.17E+00
Th-229	1.41E-04	1.97E-03	2.08E-01	2.10E-01
Th-230	7.57E-06	1.46E-02	4.38E-02	5.84E-02
Th-231	6.41E-03	8.94E-02	5.17E-03	1.01E-01
Th-232	0.00E+00	5.87E-09	1.64E+00	1.64E+00
Th-234	5.71E-02	7.97E-01	7.11E-05	8.54E-01
Pa-231	4.06E-07	2.98E-04	1.52E+01	1.52E+01
Pa-233	3.25E-06	2.31E+01	3.02E-01	2.34E+01
Pa-234m	5.71E-02	7.97E-01	7.11E-05	8.54E-01
U-232	3.04E-01	4.26E+00	2.66E+00	7.23E+00
U-233	4.98E-01	6.94E+00	2.09E+00	9.53E+00

Table 3B.7 (continued)

Radionuclide	Alkaline waste (PUREX)		Acid waste (THOREX)	Total, Ci
	Liquid, ^c Ci	Sludge, Ci	Liquid, Ci	
U-234	2.81E-01	3.97E+00	2.21E-01	4.47E+00
U-235	6.41E-03	8.94E-02	5.17E-03	1.01E-01
U-236	1.91E-02	2.67E-01	9.80E-03	2.96E-01
U-238	5.71E-02	7.97E-01	7.11E-05	8.54E-01
Np-236	0.00E+00	9.35E+00	1.23E-01	9.47E+00
Np-237	3.25E-06	2.31E+01	3.02E-01	2.34E+01
Np-239	0.00E+00	3.39E+02	7.83E+00	3.47E+02
Pu-236	6.56E-03	8.28E-01	1.09E-02	8.46E-01
Pu-238	1.24E+02	7.82E+03	4.69E+02	8.41E+03
Pu-239	2.54E+01	1.61E+03	1.54E+01	1.65E+03
Pu-240	1.87E+01	1.18E+03	8.09E+00	1.21E+03
Pu-241	1.26E+03	7.99E+04	7.36E+02	8.19E+04
Pu-242	2.54E-02	1.61E+00	1.19E-02	1.65E+00
Am-241	6.53E+00	5.32E+04	2.44E+02	5.34E+04
Am-242	0.00E+00	2.89E+02	6.66E+00	2.95E+02
Am-242m	0.00E+00	2.90E+02	6.70E+00	2.97E+02
Am-243	0.00E+00	3.39E+02	7.83E+00	3.47E+02
Cm-242	0.00E+00	2.39E+02	5.53E+00	2.45E+02
Cm-243	0.00E+00	1.34E+02	2.18E-01	1.34E+02
Cm-244	0.00E+00	7.63E+03	1.22E+01	7.64E+03
Cm-245	0.00E+00	8.62E-01	2.00E-02	8.82E-01
Cm-246	0.00E+00	9.87E-02	2.29E-03	1.01E-01
Total	1.32E+07	1.30E+07	1.72E+06	2.79E+07

^aSource: Maestas 1990. Compositions shown are as of the end of year 1989.

^bIncludes all radionuclides >0.1 Ci prior to year 3090.

^cThe small quantity of loaded zeolite ($0.031 \times 10^3 \text{ m}^3$ containing $1.9 \times 10^6 \text{ Ci}$) is included with the liquid.

Table 3B.8. West Valley Demonstration Project: estimated radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	0.1104E-02	0.2760E+01	0.9313E-04
Co-60	0.2679E-02	0.3030E+01	0.4666E-01
Ni-59	0.5491E+01	0.4160E+00	0.1650E-04
Ni-63	0.4895E+00	0.3020E+02	0.3039E-02
Se-79	0.1980E+00	0.1380E-01	0.3431E-05
Sr-90	0.1928E+03	0.2630E+05	0.3048E+02
Y-90	0.4833E-01	0.2630E+05	0.1456E+03
Zr-93	0.4257E+03	0.1070E+01	0.1242E-03
Nb-93m	0.2529E-02	0.7150E+00	0.1265E-03
Tc-99	0.2524E+02	0.4280E+00	0.2144E-03
Ru-106	0.1655E-04	0.5540E-01	0.3290E-05
Rh-106	0.1556E-10	0.5540E-01	0.5307E-03
Pd-107	0.8416E+02	0.4330E-01	0.2563E-05
Cd-113m	0.3845E-01	0.8340E+01	0.1402E-01
Sn-121m	0.1160E-02	0.6860E-01	0.1373E-03
Sn-126	0.1441E+02	0.4090E+00	0.5095E-03
Sb-125	0.2769E-01	0.2860E+02	0.8929E-01
Sb-126	0.6852E-06	0.5730E-01	0.1057E-02
Sb-126m	0.5206E-08	0.4090E+00	0.5201E-02
Te-125m	0.3885E-03	0.7000E+01	0.5876E-02
Cs-134	0.1569E-01	0.2030E+02	0.2063E+00
Cs-135	0.5505E+03	0.6340E+00	0.2113E-03
Cs-137	0.3252E+03	0.2830E+05	0.3126E+02
Ba-137m	0.4981E-04	0.2680E+05	0.1051E+03
Ce-144	0.8023E-06	0.2560E-02	0.1696E-05
Pr-144	0.3387E-10	0.2560E-02	0.1879E-04
Pm-146	0.9566E-04	0.4260E-01	0.2146E-03
Pm-147	0.3721E+00	0.3450E+03	0.1236E+00
Sm-151	0.1258E+02	0.3310E+03	0.3876E-01
Eu-152	0.8267E-02	0.1430E+01	0.1080E-01
Eu-154	0.1389E+01	0.3750E+03	0.3350E+01
Eu-155	0.2014E+00	0.9370E+02	0.6806E-01
Tl-207	0.1690E-09	0.3220E-01	0.9444E-04
Tl-208	0.4312E-10	0.1270E-01	0.2985E-03
Pb-209	0.1815E-09	0.8250E-03	0.9475E-06
Pb-211	0.1308E-08	0.3230E-01	0.9666E-04
Pb-212	0.2540E-07	0.3530E-01	0.6712E-04
Bi-211	0.7718E-10	0.3230E-01	0.1287E-02
Bi-212	0.2409E-08	0.3530E-01	0.5995E-03

Table 3B.8 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Bi-213	0.4265E-10	0.8250E-03	0.3464E-05
Po-212	0.1274E-18	0.2260E-01	0.1196E-02
Po-213	0.6231E-19	0.7860E-03	0.3972E-04
Po-215	0.1095E-14	0.3230E-01	0.1440E-02
Po-216	0.1013E-12	0.3530E-01	0.1443E-02
At-217	0.5124E-15	0.8250E-03	0.3516E-04
Rn-219	0.2482E-11	0.3230E-01	0.1339E-02
Rn-220	0.3826E-10	0.3530E-01	0.1339E-02
Fr-221	0.4653E-11	0.8250E-03	0.3180E-04
Fr-223	0.1117E-10	0.4320E-03	0.1120E-05
Ra-223	0.6306E-06	0.3230E-01	0.1149E-02
Ra-224	0.2216E-06	0.3530E-01	0.1210E-02
Ra-225	0.2104E-07	0.8250E-03	0.5778E-06
Ra-228	0.2550E-04	0.5970E-02	0.4595E-06
Ac-225	0.1421E-07	0.8250E-03	0.2878E-04
Ac-227	0.4464E-05	0.3230E-03	0.1562E-06
Ac-228	0.2662E-08	0.5970E-02	0.5153E-04
Th-227	0.1034E-05	0.3180E-01	0.1159E-02
Th-228	0.4306E-04	0.3530E-01	0.1153E-02
Th-229	0.3877E-02	0.8250E-03	0.2521E-04
Th-230	0.1169E-01	0.2360E-03	0.6670E-05
Th-231	0.6657E-09	0.3540E-03	0.1984E-06
Th-232	0.5880E+05	0.6450E-02	0.1559E-03
Th-234	0.1356E-06	0.3140E-02	0.1271E-05
Pa-231	0.1264E+01	0.5970E-01	0.1796E-02
Pa-233	0.4422E-05	0.9180E-01	0.2081E-03
Pa-234m	0.4571E-11	0.3140E-02	0.1550E-04
U-232	0.1270E-02	0.2720E-01	0.8721E-03
U-233	0.3666E+01	0.3550E-01	0.1031E-02
U-234	0.2640E+01	0.1650E-01	0.4746E-03
U-235	0.1637E+03	0.3540E-03	0.9259E-05
U-236	0.1700E+02	0.1100E-02	0.2976E-04
U-238	0.9337E+04	0.3140E-02	0.7954E-04
Np-236	0.2823E+01	0.3720E-01	0.7494E-04
Np-237	0.1302E+03	0.9180E-01	0.2802E-02
Np-239	0.5861E-05	0.1360E+01	0.3283E-02
Pu-236	0.6209E-05	0.3300E-02	0.1147E-03
Pu-238	0.1904E+01	0.3260E+02	0.1079E+01
Pu-239	0.1028E+03	0.6390E+01	0.1967E+00

Table 3B.8 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Pu-240	0.2053E+02	0.4680E+01	0.1455E+00
Pu-241	0.3076E+01	0.3170E+03	0.9815E-02
Pu-242	0.1668E+01	0.6370E-02	0.1879E-03
Am-241	0.6117E+02	0.2100E+03	0.6967E+01
Am-242	0.1435E-05	0.1160E+01	0.1315E-02
Am-242m	0.1204E+00	0.1170E+01	0.4616E-03
Am-243	0.6820E+01	0.1360E+01	0.4366E-01
Cm-242	0.2912E-03	0.9630E+00	0.3544E-01
Cm-243	0.1021E-01	0.5270E+00	0.1931E-01
Cm-244	0.3707E+00	0.3000E+02	0.1048E+01
Cm-245	0.2015E-01	0.3460E-02	0.1147E-03
Cm-246	0.1279E-02	0.3930E-03	0.1285E-04
Total	0.7029E+05	0.1096E+06	0.3260E+03

^aThis table represents the radionuclide content of a canister containing 1,900 kg of HLW glass having the radionuclide composition described in the WVDP Mass Balance Revision 7 (Crocker 1989). Radioactivity shown is as of the end of year 1989. The glass composition used is the WVDP Reference 4 glass (Crocker 1989).

Table 3B.9. Current and projected volumes of HLW at SRS^a

End of calendar year	Volume, 10 ³ m ³					Total
	Liquid	Sludge	Salt cake	Precipitate	Glass	
1989	53.3	13.76	54.8	0.13	0.0	122.0
1990	59.1	14.84	54.2	0.35	0.0	128.6
1991	57.0	16.46	50.9	1.21	0.0	125.6
1992	55.0	17.17	48.6	1.40	0.09	122.2
1993	53.7	16.77	44.7	1.67	0.28	117.1
1994	48.4	15.85	44.4	1.27	0.51	110.4
1995	46.9	14.91	41.3	1.55	0.77	105.4
1996	47.0	13.98	36.2	1.33	1.03	99.5
1997	46.7	13.06	32.6	1.71	1.26	95.3
1998	43.5	12.15	31.1	1.89	1.50	90.2
1999	43.4	11.22	27.4	1.36	1.73	85.1
2000	43.6	10.30	24.7	0.81	1.94	81.4
2001	39.3	9.41	23.6	0.16	2.15	74.7
2002	42.8	8.53	19.8	0.10	2.37	73.6
2003	44.9	7.91	19.2	0.19	2.58	74.8
2004	41.4	8.12	16.5	0.26	2.77	69.1
2005	41.2	8.32	14.8	0.21	2.94	67.4
2006	39.2	8.46	14.7	0.00	3.11	65.4
2007	39.6	8.60	16.0	0.00	3.28	67.5
2008	40.8	8.74	17.2	0.00	3.30	70.0
2009	42.9	8.88	18.1	0.00	3.30	73.2
2010	42.8	9.02	19.7	0.00	3.30	74.9
2011	42.2	9.16	21.4	0.00	3.30	76.1
2012	42.9	9.30	21.4	0.12	3.30	77.0
2013	42.2	9.44	21.4	0.24	3.30	76.6
2014	42.9	9.58	21.4	0.35	3.30	77.6
2015	42.2	9.72	21.4	0.47	3.30	77.1
2016	42.9	9.87	21.4	0.59	3.30	78.1
2017	42.2	10.00	21.4	0.71	3.30	77.6
2018	42.9	10.15	21.4	0.82	3.30	78.6
2019	42.2	10.29	21.4	0.94	3.30	78.2
2020	42.9	10.43	21.4	1.06	3.30	79.1

^aSource: Garvin 1990.

Table 3B.10. Radioactivity of HLW at SRS^a

End of calendar year	Radioactivity, 10 ⁶ Ci					Total
	Liquid	Sludge	Salt cake	Precipitate	Glass	
1989	94.6	351.2	152.8	0.31	0.0	598.9
1990	90.5	326.3	146.3	7.82	0.0	570.9
1991	76.9	330.1	136.2	21.92	0.0	565.2
1992	78.7	371.1	124.2	29.04	22.6	625.6
1993	77.2	367.6	115.0	29.74	52.0	641.7
1994	68.1	385.9	118.4	23.46	87.7	683.7
1995	67.5	359.1	115.1	20.83	150.0	712.7
1996	60.8	371.7	123.2	13.74	211.0	780.4
1997	59.6	359.7	114.0	19.03	263.4	815.8
1998	59.5	445.5	110.6	11.32	310.8	937.7
1999	54.8	400.3	106.9	8.65	345.6	916.3
2000	50.7	378.2	103.0	7.62	372.2	911.8
2001	50.7	443.4	99.4	7.12	392.3	993.8
2002	49.1	462.4	95.6	6.78	410.8	1,024.8
2003	46.9	465.4	91.8	6.49	425.7	1,036.4
2004	44.7	467.2	88.1	6.22	438.3	1,044.7
2005	42.5	464.2	84.4	5.96	449.3	1,046.4
2006	34.5	319.8	80.4	5.71	458.7	899.2
2007	29.8	262.6	80.2	0.00	467.4	840.1
2008	26.9	243.5	79.9	0.00	460.4	810.7
2009	24.6	234.7	79.6	0.00	449.8	788.8
2010	23.0	234.4	79.2	0.00	439.5	776.2
2011	21.4	234.6	78.6	0.00	429.3	762.0
2012	20.2	235.7	77.2	0.77	419.4	753.3
2013	18.9	235.9	75.8	1.51	409.7	741.8
2014	18.0	240.1	74.3	2.22	400.2	734.8
2015	17.0	240.7	72.8	2.90	390.9	724.3
2016	16.2	245.2	71.3	3.55	381.9	718.2
2017	15.4	246.2	69.7	4.17	373.1	708.5
2018	14.8	250.9	68.2	4.76	364.5	703.2
2019	14.1	251.9	66.6	5.32	356.0	694.0
2020	13.7	256.5	65.1	5.85	347.8	688.9

^aSource: Garvin 1990. Radioactivity includes actinides as well as fission products.

Table 3B.11. Thermal power of HLW at SRS^a

End of calendar year	Thermal power, 10 ³ W					Total
	Liquid	Sludge	Salt cake	Precipitate	Glass	
1989	217.7	1105.8	349.5	0.7	0.0	1673
1990	207.4	1036.1	334.8	17.9	0.0	1596
1991	176.7	1044.6	312.0	50.2	0.0	1583
1992	194.3	1196.5	284.9	66.5	56.9	1799
1993	191.7	1171.8	265.6	68.1	132.6	1830
1994	172.7	1237.9	274.9	53.7	223.1	1963
1995	171.2	1147.0	267.7	47.7	412.5	2046
1996	157.9	1199.5	288.1	31.5	603.0	2280
1997	155.6	1157.2	267.0	43.6	763.0	2387
1998	163.8	1524.1	259.9	26.1	908.0	2882
1999	151.8	1353.8	251.5	20.1	1018.8	2796
2000	142.0	1277.4	242.9	17.8	1103.5	2783
2001	147.8	1537.4	235.0	16.6	1170.7	3107
2002	146.8	1600.4	226.7	15.9	1227.4	3217
2003	142.3	1604.5	218.2	15.2	1276.2	3256
2004	137.6	1602.9	209.6	14.6	1318.1	3283
2005	132.3	1586.1	201.1	14.0	1355.0	3289
2006	102.4	1024.8	191.2	13.4	1387.2	2720
2007	86.5	827.6	190.5	0.0	1415.8	2520
2008	77.6	768.1	189.8	0.0	1397.3	2433
2009	71.2	745.4	188.9	0.0	1368.1	2374
2010	66.9	750.3	187.8	0.0	1339.9	2345
2011	62.8	751.3	186.6	0.0	1310.3	2312
2012	59.9	766.6	183.3	1.8	1281.5	2293
2013	56.7	771.5	179.9	3.6	1253.2	2265
2014	54.6	788.8	176.4	5.3	1225.6	2251
2015	51.9	793.9	172.8	6.9	1198.8	2224
2016	50.3	810.5	169.2	8.4	1172.5	2211
2017	48.1	815.5	165.6	9.9	1146.9	2186
2018	46.9	832.2	161.9	11.3	1121.9	2175
2019	45.0	836.5	158.3	12.6	1097.5	2150
2020	44.1	852.4	154.6	13.9	1073.7	2139

^aSource: Garvin 1990. Radioactivity includes actinides as well as fission products.

Table 3B.12. Savannah River Site: estimated chemical compositions of interim HLW^a

Liquid		Sludge	
Component	Wt %	Component	Wt %
Ag	Trace	Fe(OH) ₃	11.8
Hg	Trace	MnO ₂	2.0
Pb	Trace	UO ₂ (OH) ₂	1.3
U	Trace	Al(OH) ₃	13.7
F ⁻	.003	AlO(OH)	5.2
Fe	Trace	CaCO ₃	1.5
Cl ⁻	0.023	CaSO ₄	0.2
OH ⁻	1.63	CaC ₂ O ₄	0.2
NO ₂ ⁻	1.10	Ni(OH) ₂	0.8
NO ₃ ⁻	9.63	HgO	0.4
Al(OH) ₄ ⁻	4.54	SiO ₂	0.2
CO ₃ ²⁻	0.72	ThO ₂	1.8
CrO ₄ ²⁻	0.014	Ce(OH) ₃	0.2
SO ₄ ²⁻	0.22	ZrO(OH) ₃	0.2
PO ₄ ³⁻	0.12	Cr(OH) ₃	0.2
NH ₄ ⁺	Trace	Mg(OH) ₂	0.2
Na ⁺	11.0	NaNO ₃	1.1
H ₂ O	71.0	NaOH	1.3
	—	Zeolite	1.5
	100.0	Others	1.2
		H ₂ O	55.0
			—
			100.0
Density (25° C), g/mL	1.1		1.4

Table 3B.12 (continued)

Salt Cake		Precipitate ^b	
Component	Wt %	Component	Wt %
NaNO ₃	65.4	K(C ₆ H ₅) ₄ B	9.0
NaNO ₂	0.9	NaNO ₃	0.7
NaOH	3.4	Others	1.8
NaAl(OH) ₄	7.8	H ₂ O	88.5
Na ₂ CO ₃	2.7		—
Na ₂ SO ₄	9.4		100.0
Na ₃ PO ₄	Trace		
NaF	0.2		
Na ₂ C ₂ O ₄	0.1		
Insolubles	3.7		
H ₂ O	6.4		
	—		
	100.0		
Density (25° C), g/mL	1.9		1.05

^aSource: Garvin 1990.^bPrecipitate (non-Newtonian fluid) from the in-tank precipitation process.

Table 3B.13. Savannah River Site: radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Cr-51	0.1008E-20	0.9312E-16	0.1996E-19
Co-60	0.1502E+00	0.1699E+03	0.2619E+01
Ni-59	0.3163E+00	0.2397E-01	0.9519E-06
Ni-63	0.4824E-01	0.2975E+01	0.3000E-03
Se-79	0.2439E+01	0.1699E+00	0.4232E-04
Rb-87	0.9961E+01	0.8719E-06	0.7278E-09
Sr-89	0.1470E-08	0.4267E-04	0.1473E-06
Sr-90	0.3426E+03	0.4675E+05	0.5426E+02
Y-90	0.8795E-01	0.4786E+05	0.2653E+03
Y-91	0.3085E-07	0.7568E-03	0.2715E-05
Zr-93	0.4443E+03	0.1117E+01	0.1298E-03
Zr-95	0.4680E-06	0.1005E-01	0.5084E-04
Nb-94	0.5147E-03	0.9646E-04	0.9830E-06
Nb-95	0.5407E-06	0.2115E-01	0.1013E-03
Nb-95m	0.3272E-09	0.1247E-03	0.1730E-06
Tc-99	0.1816E+03	0.3079E+01	0.1545E-02
Ru-103	0.5217E-12	0.1684E-07	0.5827E-10
Ru-106	0.6729E+00	0.2252E+04	0.1339E-00
Rh-103m	0.5028E-15	0.1636E-07	0.3761E-11
Rh-106	0.6346E-06	0.2259E+04	0.2167E+02
Pd-107	0.2863E+02	0.1473E-01	0.8732E-06
Ag-110m	0.2647E-04	0.1258E+00	0.2098E-02
Cd-113	0.1472E+00	0.5009E-13	0.8420E-16
Cd-115m	0.4763E-13	0.1213E-08	0.4518E-11
Sn-121m	0.1336E-02	0.7902E-01	0.1581E-03
Sn-123	0.3101E-04	0.2549E+00	0.7951E-03
Sn-126	0.1556E+02	0.4415E+00	0.5508E-03
Sb-124	0.4071E-11	0.7123E-07	0.9445E-09
Sb-125	0.8226E+00	0.8496E+03	0.2656E+01
Sb-126	0.7365E-06	0.6159E-01	0.1138E-02
Sb-126m	0.5619E-08	0.4415E+00	0.5622E-02

Table 3B.13 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-126m	0.1532E-01	0.2760E+03	0.2320E+00
Te-127	0.4555E-07	0.1202E+00	0.1622E-03
Te-128m	0.1302E-04	0.1228E+00	0.6597E-04
Te-129	0.1457E-18	0.3053E-11	0.1089E-13
Te-129m	0.1576E-15	0.4749E-11	0.8316E-14
Cs-134	0.2606E+00	0.3372E+03	0.3433E+01
Cs-135	0.8633E+02	0.9943E-01	0.3319E-04
Cs-136	0.1068E-43	0.7838E-39	0.1066E-41
Cs-137	0.4989E+03	0.4341E+05	0.4802E+02
Ba-136m	0.3195E-49	0.8607E-38	0.1040E-40
Ba-137m	0.7724E-04	0.4155E+05	0.1632E+03
Ba-140	0.1404E-40	0.1024E-35	0.2853E-38
La-140	0.7734E-42	0.4304E-36	0.7205E-38
Ce-141	0.1260E-14	0.3591E-10	0.5250E-13
Ce-142	0.4005E+03	0.9609E-05	0.0000E+00
Ce-144	0.3093E+01	0.9869E+04	0.6547E+01
Pr-143	0.1780E-38	0.1198E-33	0.2291E-37
Pr-144	0.1306E-03	0.9869E+04	0.7255E+02
Pr-144m	0.6545E-06	0.1187E+03	0.4063E-01
Nd-144	0.4110E+03	0.4860E-09	0.0000E+00
Nd-147	0.1570E-48	0.1261E-43	0.3038E-46
Pm-147	0.2609E+02	0.2419E+05	0.8679E+01
Pm-148	0.4243E-15	0.6975E-10	0.5364E-12
Pm-148m	0.4722E-13	0.1009E-08	0.1277E-10
Sm-147	0.8796E+02	0.2000E-05	0.2738E-07
Sm-148	0.1916E+02	0.5788E-11	0.6901E-13
Sm-149	0.7420E+01	0.1781E-11	0.0000E+00
Sm-151	0.9418E+01	0.2478E+03	0.2906E-01
Eu-152	0.2132E-01	0.3688E+01	0.2790E-01
Eu-154	0.2295E+01	0.6196E+03	0.5543E+01
Eu-155	0.1021E+01	0.4749E+03	0.3455E+00
Eu-156	0.9489E-36	0.5231E-31	0.5392E-33

Table 3B.13 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Tb-160	0.9923E-10	0.1120E-05	0.9110E-08
Tl-208	0.3829E-11	0.1128E-02	0.2645E-04
U-232	0.6256E-03	0.1339E-01	0.4301E-03
U-233	0.1636E-03	0.1584E-05	0.4605E-07
U-234	0.5485E+01	0.3428E-01	0.9875E-03
U-235	0.7278E+02	0.1573E-03	0.4122E-05
U-236	0.1742E+02	0.1128E-02	0.3054E-04
U-238	0.3122E+05	0.1050E-01	0.2663E-03
Np-236	0.1323E-05	0.1744E-07	0.3514E-10
Np-237	0.1263E+02	0.8904E-02	0.2722E-03
Pu-236	0.2297E-03	0.1221E+00	0.4249E-02
Pu-237	0.7401E-15	0.8941E-11	0.3292E-14
Pu-238	0.8667E+02	0.1484E+04	0.4919E+02
Pu-239	0.2076E+03	0.1291E+02	0.3979E+00
Pu-240	0.3809E+02	0.8681E+01	0.2704E+00
Pu-241	0.1620E+02	0.1670E+04	0.5176E-01
Pu-242	0.3206E+01	0.1224E-01	0.3616E-03
Am-241	0.3210E+01	0.1102E+02	0.3661E+00
Am-242	0.1776E-07	0.1436E-01	0.1628E-04
Am-242m	0.1488E-02	0.1447E-01	0.5709E-05
Am-243	0.2902E-01	0.5788E-02	0.1860E-03
Cm-242	0.1057E-04	0.3495E-01	0.1288E-02
Cm-243	0.1078E-03	0.5565E-02	0.2039E-03
Cm-244	0.1329E+01	0.1076E+03	0.3763E+01
Cm-245	0.3910E-04	0.6715E-05	0.2225E-06
Cm-246	0.1739E-05	0.5342E-06	0.1747E-07
Cm-247	0.7116E-08	0.6604E-12	0.2107E-13
Cm-248	0.1614E-09	0.6864E-12	0.8533E-13
Totals	0.3427E+05	0.2344E+06	0.7093E+03

^aQuantities shown are for sludge-precipitate glass and are based on Baxter 1988, assuming sludge aged an average of 5 years and supernate aged an average of 15 years, with a canister load of 3,710 lb of glass (1,682 kg). Radionuclide contents are at time of filling canister.

Table 3B.14. Hanford Site: historical and projected volumes of high-level wastes^a

End of calendar year	Volume, 10 ³ m ³					Total
	Liquid	Sludge	Salt cake	Slurry	Sr and Cs capsules	
1988	26.8	46.0	93.0	77.7	0.004	244
1989	26.5	46.0	93.0	79.3	0.004	245
1990	25.8	46.0	93.0	94.2	0.004	259
1991	23.2	46.0	93.0	84.0	0.004	246
1992	19.5	46.0	93.0	85.2	0.004	244
1993	16.1	46.0	93.0	94.5	0.004	250
1994	14.3	46.0	93.0	91.0	0.004	244
1995	11.5	46.0	93.0	94.5	0.004	245
1996	11.3	46.0	93.0	99.0	0.004	249
1997	11.3	46.0	93.0	99.4	0.004	250
1998	11.3	46.0	93.0	100.9	0.004	251
1999	11.3	46.0	93.0	103.8	0.004	254
2000	11.3	46.0	93.0	108.3	0.004	259
2001	11.3	46.0	93.0	108.9	0.004	259
2002	11.3	46.0	93.0	113.0	0.004	263
2003	11.3	46.0	93.0	109.3	0.004	260
2004	11.3	46.0	93.0	110.0	0.004	260
2005	11.3	46.0	93.0	110.3	0.004	261
2006	11.3	46.0	93.0	110.7	0.004	261
2007	11.3	46.0	93.0	111.0	0.004	261
2008	11.3	46.0	93.0	111.4	0.004	262
2009	11.3	46.0	93.0	111.8	0.004	262
2010	11.3	46.0	93.0	112.1	0.004	262
2011	11.3	46.0	93.0	112.5	0.004	263
2012	11.3	46.0	93.0	112.9	0.004	263
2013	11.3	46.0	93.0	113.3	0.004	264
2014	11.3	46.0	93.0	113.6	0.004	264
2015	11.3	46.0	93.0	114.0	0.004	264
2016	11.3	46.0	93.0	114.4	0.004	265
2017	11.3	46.0	93.0	114.7	0.004	265
2018	11.3	46.0	93.0	115.1	0.004	265
2019	11.3	46.0	93.0	115.5	0.004	266
2020	11.3	46.0	93.0	115.8	0.004	266

^aSource: Turner 1990. Liquid, sludge, and salt cake refer to the contents of single-shell tanks. Slurry represents the entire contents of double-shell tanks. Quantities shown in this table have not been adjusted downward to reflect the vitrification that is expected to begin in 1999, that is, the volumes shown here are calculated as if no vitrification occurred.

Table 3B.15. Hanford Site: historical and projected radioactivity of high-level wastes^a

End of calendar year	Radioactivity, 10 ⁶ Ci					Total
	Liquid	Sludge	Salt cake	Slurry	Sr and Cs capsules	
1988	23.3	121.4	12.6	111.1	177.1	446
1989	22.6	118.5	12.3	89.8	173.0	416
1990	21.5	115.7	12.1	75.3	169.1	394
1991	18.9	113.0	11.8	77.1	165.1	386
1992	15.5	110.3	11.5	88.1	161.3	387
1993	12.5	107.7	11.2	100.1	157.6	389
1994	10.9	105.1	11.0	111.3	154.0	392
1995	8.5	102.7	10.7	110.7	150.5	383
1996	8.2	100.2	10.5	115.4	147.0	381
1997	8.0	97.8	10.2	111.4	143.6	371
1998	7.8	95.5	10.0	107.9	140.3	362
1999	7.7	93.3	9.8	104.7	137.0	353
2000	7.5	91.1	9.5	101.8	133.9	344
2001	7.3	88.9	9.3	99.1	130.8	335
2002	7.2	86.8	9.1	96.5	127.8	327
2003	7.0	84.7	8.9	94.1	124.8	320
2004	6.8	82.7	8.7	91.7	122.0	312
2005	6.7	80.8	8.5	89.5	119.2	305
2006	6.5	78.9	8.3	87.3	116.4	297
2007	6.4	77.0	8.1	85.3	113.7	291
2008	6.2	75.2	7.9	83.2	111.1	284
2009	6.1	73.4	7.7	81.3	108.6	277
2010	5.9	71.7	7.5	79.4	106.0	271
2011	5.8	70.0	7.4	77.5	103.6	264
2012	5.7	68.3	7.2	75.7	101.2	258
2013	5.5	66.7	7.0	74.0	98.9	252
2014	5.4	65.1	6.9	72.3	96.6	246
2015	5.3	63.6	6.7	70.6	94.4	241
2016	5.2	62.1	6.6	69.0	92.2	235
2017	5.1	60.6	6.4	67.4	90.1	230
2018	4.9	59.2	6.3	65.8	88.0	224
2019	4.8	57.8	6.1	64.3	86.0	219
2020	4.7	56.4	6.0	62.8	84.0	214

^aSource: Turner 1990. Liquid, sludge, and salt cake refer to the contents of single-shell tanks. Slurry represents the entire contents of double-shell tanks. Quantities shown in this table have not been adjusted downward to reflect the vitrification that is expected to begin in 1999, that is, the quantities shown are calculated as if no vitrification occurred.

Table 3B.16. Representative chemical composition of current and future HLW at HANF^a

Component	Composition, wt %			
	Liquid ^b	Sludge ^b	Salt cake ^b	Slurry ^c
NaNO ₃	20.8	25.3	81.5	14.8
NaNO ₂	15.8	3.8	1.7	5.6
Na ₂ CO ₃	0.6	2.2	0.5	1.9
NaOH	6.2	5.3	1.5	7.0
NaAlO ₂	12.5	1.2	1.4	6.0
NaF	—	—	—	0.4
Na ₂ SO ₄	—	1.0	1.3	0.3
Na ₃ PO ₄	2.3	15.8	1.6	0.8
KF	—	—	—	0.4
FeO(OH)	—	1.3	—	0.2
Organic carbon	0.17	—	—	1.2
NH ₄ ⁺	—	—	—	0.08
Al(OH) ₃	—	2.9	—	4.9
SrO·H ₂ O	—	0.1	—	—
Na ₂ CrO ₄	1.3	—	—	—
Cr(OH) ₃	—	0.2	—	0.02
Cd(OH) ₂	—	0.1	—	—
Ni(OH) ₂	—	—	—	<0.1
BiPO ₄	—	0.5	—	—
Cl ⁻	—	0.1	—	—
Ni ₂ Fe(CN) ₆	—	0.6	—	—
P ₂ O ₅ ·24WO ₂ ·44H ₂ O	—	<0.1	—	—
ZrO ₂ ·2H ₂ O	—	0.5	—	0.2
Fission products	—	—	—	<0.01
H ₂ O	40.2	33.6	10.5	56.2
Other	<0.1	5.5	—	<0.01
Hg ⁺	—	0.12 ppm	—	—
Total	100.0	100.0	100.0	100.0
Density, g/mL	1.6	1.7	1.4	~ 1.3

^aSource: IDB 1989, Wojtasek 1989.^bStored in single-shell tanks.^cStored in double-shell tanks.

Table 3B.17. Hanford Site: radionuclide content per HLW canister, NCAW glass, maximum case^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	5.64E-02	1.41E+02	4.765E-03
Ni-59	1.80E+00	1.36E-01	5.402E-06
Co-60	3.79E-03	4.29E+00	6.615E-02
Ni-63	2.54E-01	1.57E+01	6.236E-03
Se-79	5.60E-02	3.90E-03	9.711E-07
Sr-89	2.24E-06	6.52E-02	2.254E-04
Sr-90	3.06E+02	4.18E+04	4.852E+01
Y-90	7.68E-02	4.18E+04	2.317E+02
Y-91	2.96E-05	7.26E-01	2.608E-03
Nb-93m	2.04E-03	5.77E-01	1.023E-04
Zr-93	5.13E+02	1.29E+00	1.499E-04
Zr-95	1.28E-04	2.76E+00	1.398E-02
Nb-95	1.45E-04	5.67E+00	2.720E-02
Tc-99	5.51E+02	9.35E+00	4.689E-03
Ru-103	9.23E-09	2.98E-04	9.971E-07
Rh-103m	8.27E-12	2.69E-04	6.192E-08
Ru-106	1.49E+00	4.99E+03	2.967E-01
Rh-106	1.40E-06	4.99E+03	4.786E+01
Pd-107	7.91E+01	4.07E-02	2.413E-06
Ag-110m	3.35E-04	1.59E+00	2.655E-02
Cd-113m	6.73E-02	1.46E+01	2.458E-02
In-113m	1.51E-09	2.52E-02	5.871E-05
Sn-113	2.51E-06	2.52E-02	4.198E-06
Cd-115m	3.45E-10	8.78E-06	3.275E-08
Sn-119m	1.21E-03	5.42E+00	2.802E-03
Sn-121m	1.79E-03	1.06E-01	2.124E-04
Sn-123	3.52E-04	2.89E+00	9.027E-03
Sn-126	1.62E+01	4.60E-01	5.738E-04
Sb-124	2.00E-09	3.50E-05	4.648E-07
Sb-126	7.75E-07	6.48E-02	1.197E-03
Sb-126m	5.86E-09	4.60E-01	5.858E-03
Sb-125	1.70E+00	1.76E+03	5.503E+00
Te-125m	2.38E-02	4.29E+02	3.606E-01

Table 3B.17 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-127	1.12E-06	2.95E+00	3.984E-03
Te-127m	3.18E-04	3.00E+00	1.614E-03
Te-129	1.79E-14	3.75E-07	1.340E-09
Te-129m	1.91E-11	5.77E-07	1.012E-09
I-129	9.23E-02	1.63E-05	7.541E-09
Cs-134	9.27E-01	1.20E+03	1.221E+01
Cs-135	2.18E+02	2.51E-01	8.378E-05
Cs-137	5.86E+02	5.10E+04	5.642E+01
Ba-137m	8.96E-05	4.82E+04	1.893E+02
Ce-141	3.97E-10	1.13E-05	1.655E-08
Ce-144	9.34E+00	2.98E+04	1.977E+01
Pr-144	3.94E-04	2.98E+04	2.191E+02
Pr-144m	1.97E-06	3.58E+02	1.225E-01
Pm-147	4.28E+01	3.97E+04	1.424E+01
Pm-148m	6.18E-10	1.32E-05	1.674E-07
Sm-151	3.18E+01	8.36E+02	9.803E-02
Eu-152	1.58E-02	2.74E+00	2.073E-02
Gd-153	3.26E-06	1.15E-02	1.039E-05
Eu-154	1.24E+00	3.36E+02	3.006E+00
Eu-155	8.83E-01	4.11E+02	2.990E-01
Tb-160	9.74E-09	1.10E-04	8.961E-07
U-234	7.71E-01	4.82E-03	1.388E-04
U-235	9.11E+01	1.97E-04	5.160E-06
U-236	7.34E+00	4.75E-04	1.287E-05
U-238	1.11E+04	3.72E-03	9.437E-05
Np-237	2.82E+02	1.99E-01	6.083E-03
Pu-238	4.48E-02	7.68E-01	2.546E-02
Pu-239	2.27E+01	1.41E+00	4.346E-02
Pu-240	2.38E+00	5.42E-01	1.688E-02
Pu-241	2.50E-01	2.58E+01	7.999E-04
Pu-242	3.43E-02	1.31E-04	3.869E-06
Am-241	1.68E+02	5.77E+02	1.917E+01
Am-242	5.12E-07	4.14E-01	4.700E-04

Table 3B.17 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Am-243	3.39E-01	6.76E-02	2.173E-03
Cm-242	1.51E-04	4.99E-01	1.839E-02
Cm-244	1.54E-01	1.25E+01	4.374E-01
Total	1.40E+04	2.98E+05	8.687E+02

^aThis table identifies the maximum expected activity of HWVP canisters at the time of vitrification. The maximum is principally based on close-coupling the final accumulated tank of NCAW (21 months from fuel discharge to HWVP). Canister contains 1,650 kg of HLW glass (85% fill). Source: Mitchell and Nelson 1988.

Table 3B.18. Hanford Site: radionuclide content per HLW canister, NCAW glass, nominal case^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Fe-55	7.20E-03	1.80E+01	6.083E-04
Ni-59	1.44E+00	1.09E-01	4.330E-06
Co-60	1.33E-03	1.50E+00	2.313E-02
Ni-63	1.96E-01	1.21E+01	4.806E-03
Se-79	4.52E-02	3.15E-03	7.843E-07
Sr-89	1.84E-17	5.35E-13	1.850E-15
Sr-90	2.18E+02	2.98E+04	3.459E+01
Y-90	5.48E-02	2.98E+04	1.652E+02
Y-91	5.63E-15	1.38E-10	4.957E-13
Nb-93m	2.18E-03	6.16E-01	1.092E-04
Zr-93	4.18E+02	1.05E+00	1.220E-04
Zr-95	1.36E-13	2.92E-09	1.479E-11
Nb-95	1.72E-13	6.73E-09	3.229E-11
Tc-99	4.43E+02	7.51E+00	3.767E-03
Ru-103	1.04E-22	3.37E-18	1.126E-20
Rh-103m	9.34E-26	3.04E-18	6.988E-22
Ru-106	1.25E-02	4.18E+01	2.486E-03
Rh-106	1.17E-08	4.18E+01	4.009E-01
Pd-107	5.87E+01	3.02E-02	1.790E-06
Ag-110m	4.67E-07	2.22E-03	3.708E-05
Cd-113m	3.93E-02	8.53E+00	1.436E-02
In-113m	6.04E-15	1.01E-07	2.353E-10
Sn-113	1.01E-11	1.01E-07	1.683E-11
Cd-115m	1.26E-22	3.20E-18	1.192E-20
Sn-119m	1.52E-06	6.80E-03	3.516E-06
Sn-121m	1.31E-03	7.76E-02	1.555E-04
Sn-123	4.44E-09	3.65E-05	1.140E-07
Sn-126	1.29E+01	3.65E-01	4.553E-04
Sb-124	6.57E-19	1.15E-14	1.527E-16
Sb-126	6.10E-07	5.10E-02	9.424E-04
Sb-126m	4.65E-09	3.65E-01	4.648E-03
Sb-125	2.46E-01	2.54E+02	7.942E-01
Te-125m	3.44E-03	6.20E+01	5.212E-02
Te-127	2.48E-12	6.55E-06	8.846E-09
Te-127m	7.06E-10	6.66E-06	3.583E-09
Te-129	1.49E-30	3.14E-23	1.120E-25

Table 3B.18 (continued)

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Te-129m	1.60E-27	4.82E-23	8.440E-26
I-129	7.31E-02	1.29E-05	5.968E-09
Cs-134	7.19E-02	9.31E+01	9.476E-01
Cs-135	1.75E+02	2.02E-01	6.742E-05
Cs-137	4.15E+02	3.61E+04	3.994E+01
Ba-137m	6.32E-05	3.40E+04	1.335E+02
Ce-141	1.03E-26	2.93E-22	4.284E-25
Ce-144	2.51E-02	8.00E+01	5.307E-02
Pr-144	1.06E-06	8.00E+01	5.881E-01
Pr-144m	5.29E-09	9.60E-01	3.285E-04
Pm-147	5.62E+00	5.21E+03	1.869E+00
Pm-148m	2.92E-23	6.23E-19	7.889E-21
Sm-151	2.65E+01	6.98E+02	8.185E-02
Eu-152	8.09E-03	1.40E-00	1.059E-02
Gd-153	3.83E-09	1.35E-05	1.220E-08
Eu-154	5.37E-01	1.45E+02	1.297E+00
Eu-155	2.94E-01	1.37E+02	9.965E-02
Tb-160	8.41E-17	9.49E-13	7.730E-15
U-234	7.31E-01	4.57E-03	1.316E-04
U-235	8.83E+01	1.91E-04	5.003E-06
U-236	6.51E+00	4.21E-04	1.141E-05
U-238	1.04E+04	3.51E-03	8.904E-05
Np-237	2.21E+02	1.56E-01	4.769E-03
Pu-238	2.59E-02	4.43E-01	1.468E-02
Pu-239	1.88E+01	1.17E+00	3.606E-02
Pu-240	1.72E+00	3.93E-01	1.224E-02
Pu-241	1.22E-01	1.26E+01	3.907E-04
Pu-242	1.99E-02	7.61E-05	2.248E-06
Am-241	8.27E+01	2.84E+02	9.436E+00
Am-242	2.73E-07	2.21E-01	2.509E-04
Am-243	1.90E-01	3.79E-02	1.218E-03
Cm-242	5.50E-05	1.82E-01	6.707E-03
Cm-244	6.22E-02	5.03E+00	1.760E-01
Total	1.26E+04	1.37E+05	3.892E+02

^aThis table identifies the nominal expected activity of HWVP canisters at the time of vitrification. Canister contains 1,650 kg of HLW glass (85% fill). Source: Mitchell and Nelson 1988.

Table 3B.19. Hanford Site: total radioactivity and thermal power of 597 strontium capsules projected through year 2020^a

End of calendar year	Radioactivity, Ci			Thermal power, 10 ³ W		
	Sr-90	Y-90	Total	Sr-90	Y-90	Total
1991	2.599E+07	2.599E+07	5.198E+07	30.1	143.9	174.0
1992	2.538E+07	2.538E+07	5.075E+07	29.4	140.5	169.9
1993	2.478E+07	2.478E+07	4.956E+07	28.7	137.1	165.8
1994	2.420E+07	2.420E+07	4.839E+07	28.1	133.9	162.0
1995	2.363E+07	2.363E+07	4.726E+07	27.4	130.8	158.2
1996	2.307E+07	2.307E+07	4.614E+07	26.7	127.7	154.4
1997	2.253E+07	2.253E+07	4.506E+07	26.1	124.7	150.8
1998	2.200E+07	2.200E+07	4.400E+07	25.5	121.8	147.3
1999	2.148E+07	2.148E+07	4.296E+07	24.9	118.9	143.8
2000	2.098E+07	2.098E+07	4.195E+07	24.3	116.1	140.4
2001	2.048E+07	2.048E+07	4.097E+07	23.7	113.4	137.1
2002	2.000E+07	2.000E+07	4.000E+07	23.2	110.7	133.9
2003	1.953E+07	1.953E+07	3.906E+07	22.6	108.1	130.7
2004	1.907E+07	1.907E+07	3.814E+07	22.1	105.6	127.7
2005	1.862E+07	1.862E+07	3.725E+07	21.6	103.1	124.7
2006	1.819E+07	1.819E+07	3.637E+07	21.1	100.7	121.8
2007	1.776E+07	1.776E+07	3.552E+07	20.6	98.3	118.9
2008	1.734E+07	1.734E+07	3.468E+07	20.1	96.0	116.1
2009	1.693E+07	1.693E+07	3.387E+07	19.6	93.7	113.3
2010	1.653E+07	1.653E+07	3.307E+07	19.2	91.5	110.7
2011	1.615E+07	1.615E+07	3.229E+07	18.7	89.4	108.1
2012	1.577E+07	1.577E+07	3.153E+07	18.3	87.3	105.6
2013	1.539E+07	1.539E+07	3.079E+07	17.8	85.2	103.0
2014	1.503E+07	1.503E+07	3.007E+07	17.4	83.2	100.6
2015	1.468E+07	1.468E+07	2.936E+07	17.0	81.3	98.3
2016	1.433E+07	1.433E+07	2.867E+07	16.6	79.3	95.9
2017	1.400E+07	1.400E+07	2.799E+07	16.2	77.5	93.7
2018	1.367E+07	1.367E+07	2.734E+07	15.8	75.7	91.5
2019	1.335E+07	1.335E+07	2.669E+07	15.5	73.9	89.4
2020	1.303E+07	1.303E+07	2.606E+07	15.1	72.1	87.2

^aSource: Turner 1990 and ORNL calculations. Curies and watts shown are totals for 597 capsules.

Table 3B.20. Hanford Site: total radioactivity and thermal power of 1350 cesium capsules projected through year 2020^a

End of calendar year	Radioactivity, Ci			Thermal power, 10 ³ W		
	Cs-137	Ba-137m	Total	Cs-137	Ba-137m	Total
1991	5.826E+07	5.512E+07	1.134E+08	64.4	216.1	280.5
1992	5.693E+07	5.386E+07	1.108E+08	62.9	211.2	274.1
1993	5.563E+07	5.263E+07	1.083E+08	61.5	206.3	267.8
1994	5.436E+07	5.143E+07	1.058E+08	60.1	201.6	261.7
1995	5.312E+07	5.025E+07	1.034E+08	58.7	197.0	255.7
1996	5.191E+07	5.910E+07	1.010E+08	57.3	192.6	249.9
1997	5.072E+07	4.798E+07	9.871E+07	56.0	188.1	244.1
1998	4.957E+07	4.689E+07	9.645E+07	54.8	183.9	238.7
1999	4.843E+07	4.582E+07	9.425E+07	53.5	179.6	233.1
2000	4.733E+07	4.477E+07	9.210E+07	52.3	175.6	227.9
2001	4.625E+07	4.375E+07	8.999E+07	51.1	171.6	222.7
2002	4.519E+07	4.275E+07	8.794E+07	49.9	167.6	217.5
2003	4.416E+07	4.177E+07	8.593E+07	48.8	163.8	212.6
2004	4.315E+07	4.082E+07	8.397E+07	47.7	160.1	207.8
2005	4.217E+07	3.989E+07	8.205E+07	46.6	156.4	203.0
2006	4.120E+07	3.898E+07	8.018E+07	45.5	152.8	198.3
2007	4.026E+07	3.809E+07	7.835E+07	44.4	149.3	193.7
2008	3.934E+07	3.722E+07	7.656E+07	43.5	145.9	189.4
2009	3.844E+07	3.637E+07	7.481E+07	42.5	142.6	185.1
2010	3.757E+07	3.554E+07	7.310E+07	41.5	139.4	180.9
2011	3.671E+07	3.473E+07	7.143E+07	40.6	136.2	176.8
2012	3.587E+07	3.393E+07	6.980E+07	39.6	133.1	172.7
2013	3.505E+07	3.316E+07	6.821E+07	38.7	130.0	168.7
2014	3.425E+07	3.240E+07	6.665E+07	37.8	127.0	164.8
2015	3.347E+07	3.166E+07	6.513E+07	37.0	124.2	161.2
2016	3.270E+07	3.094E+07	6.364E+07	36.1	121.3	157.4
2017	3.196E+07	3.023E+07	6.219E+07	35.3	118.6	153.9
2018	3.123E+07	2.954E+07	6.077E+07	34.5	115.8	150.3
2019	3.052E+07	2.887E+07	5.938E+07	33.7	113.2	146.9
2020	2.982E+07	2.821E+07	5.803E+07	32.9	110.6	143.5

^aSource: Turner 1990 and ORNL calculations. Curies and watts shown are totals for 1350 capsules.

Table 3B.21. Idaho National Engineering Laboratory: historical and projected volumes of interim and immobilized high-level wastes^a

End of calendar year	Volume, 10 ³ m ³			
	Liquid	Calcine	Glass- ceramic	Total
1989	8.5	3.5	0.0	12.0
1990	7.6	3.7	0.0	11.3
1991	7.0	4.1	0.0	11.1
1992	8.0	4.1	0.0	12.1
1993	8.5	4.4	0.0	12.9
1994	6.4	4.9	0.0	11.3
1995	7.5	5.0	0.0	12.5
1996	5.9	5.4	0.0	11.3
1997	6.9	5.4	0.0	12.3
1998	5.8	5.9	0.0	11.7
1999	6.4	5.9	0.0	12.3
2000	6.1	6.0	0.0	12.1
2001	7.1	6.4	0.0	13.5
2002	7.4	6.9	0.0	14.3
2003	7.7	7.3	0.0	15.0
2004	8.2	7.8	0.0	16.0
2005	8.5	8.3	0.0	16.8
2006	8.7	8.8	0.0	17.5
2007	9.0	8.8	0.0	17.8
2008	9.2	8.8	0.0	18.0
2009	5.8	9.3	0.0	15.1
2010	4.7	10.0	0.0	14.7
2011	3.7	10.70	0.0	14.4
2012	2.7	10.96	0.28	13.94
2013	2.0	11.12	0.63	13.75
2014	2.3	10.95	1.03	14.28
2015	4.5	10.28	1.60	16.38
2016	3.4	10.11	2.17	15.68
2017	2.3	9.91	2.74	14.95
2018	3.3	9.43	3.31	16.04
2019	0.2	8.98	3.88	13.06
2020	0.3	8.62	4.45	13.37

^aSource: Berreth 1990. Quantities shown are based on the assumptions that immobilization starts in year 2012 and that the glass-ceramic form is used. Each canister is assumed to contain 0.57 m³ of ceramic (1,825 kg), with a calcine loading of 70 wt % or 1,277 kg; this is the equivalent of 0.91 m³ of calcine prior to immobilization. The reader is cautioned that these projections are based on estimates and assumptions that are subject to change.

Table 3B.22. Idaho National Engineering Laboratory: historical and projected radioactivity and thermal power of interim and immobilized high-level wastes^a

End of calendar year	Liquid		Calcine		Glass-Ceramic		Total	
	10 ⁶ Ci	10 ³ W	10 ⁶ Ci	10 ³ W	10 ⁶ Ci	10 ³ W	10 ⁶ Ci	10 ³ W
1989	11.5	34.3	56.9	164.9	0.00	0.0	68.4	199.2
1990	6.6	19.3	59.5	172.9	0.00	0.0	66.1	192.2
1991	11.4	33.5	62.8	182.3	0.00	0.0	74.2	215.8
1992	10.7	31.0	61.0	177.3	0.00	0.0	71.7	208.3
1993	13.2	39.5	67.8	196.9	0.00	0.0	81.0	236.4
1994	8.4	24.7	72.7	211.1	0.00	0.0	81.1	235.8
1995	7.3	21.2	75.9	220.4	0.00	0.0	83.2	241.6
1996	17.2	55.6	79.2	230.0	0.00	0.0	96.4	285.6
1997	15.7	47.8	79.2	230.0	0.00	0.0	94.9	277.8
1998	11.4	33.9	84.9	246.7	0.00	0.0	96.3	280.6
1999	5.7	16.7	87.0	252.8	0.00	0.0	92.7	269.5
2000	3.6	10.6	92.0	267.2	0.00	0.0	95.6	277.8
2001	14.5	45.9	97.8	284.0	0.00	0.0	112.3	329.9
2002	19.2	57.4	107.0	310.6	0.00	0.0	126.2	368.0
2003	21.8	65.3	119.1	345.9	0.00	0.0	140.9	411.2
2004	22.0	67.9	113.3	387.9	0.00	0.0	135.3	455.8
2005	23.6	72.1	141.1	410.6	0.00	0.0	164.7	482.7
2006	26.8	80.3	145.0	421.9	0.00	0.0	171.8	502.2
2007	24.0	70.7	140.3	408.3	0.00	0.0	164.3	479.0
2008	19.8	57.7	138.8	404.1	0.00	0.0	158.6	461.8
2009	13.9	42.5	147.4	429.0	0.00	0.0	161.3	471.5
2010	15.9	49.6	161.5	471.7	0.00	0.0	177.4	521.3
2011	15.1	45.9	178.1	521.3	0.00	0.0	193.2	567.2
2012	15.9	49.3	187.7	549.5	7.80	22.9	211.4	621.7
2013	12.8	39.0	196.7	576.8	17.74	52.0	227.2	667.8
2014	18.0	55.5	193.3	565.2	28.89	84.5	240.2	705.2
2015	25.1	77.0	182.0	533.4	45.23	132.0	252.3	742.4
2016	18.3	55.7	183.0	534.2	62.40	182.1	263.7	772.0
2017	10.0	51.4	178.8	521.5	78.62	229.4	267.4	802.3
2018	13.2	40.3	170.7	497.4	95.65	278.7	279.6	816.4
2019	2.0	5.9	160.3	466.7	110.04	320.4	272.3	793.0
2020	3.7	11.3	153.7	448.8	126.86	370.3	284.3	830.4

^aSource: Berreth 1990 and ORNL calculations based on the assumption that the glass-ceramic produced each year has the same radionuclide composition as the average calcine in storage in that year. Each canister is assumed to contain 0.57 m³ of ceramic (1,825 kg), with a calcine loading of 70 wt % or 1,277 kg; this is the equivalent of 0.91 m³ of calcine prior to immobilization. The reader is cautioned that the assumptions and estimates used here are subject to change.

Table 3B.23. Representative chemical composition of current and future HLW liquid at INEL^a

Component	Composition, wt %			
	Zirconium fluoride	Sodium bearing	Nonfluoride	Fluorinel
Al	1.3	0.8-1.6	1.51	0.742
B	0.15	0.005-0.01	0.003	0.241
Ca	-	0.03-0.2	0.27	-
Cl ⁻	-	0.06-0.1	0.023	-
Cd	-	-	1.42	-
Cr	-	-	0.036	0.0087
F ⁻	3.4	0.005-0.06	0.032	5.99
Fe	0.04	0.05-0.09	0.19	0.023
H ⁺	1.12	0.03-0.15	0.12	0.18
K	1.12	0.03-0.15	0.33	-
Mg	-	-	0.062	-
Mn	-	-	0.048	0.0004
Na	0.12	2.1-4.0	1.31	-
Ni	-	-	0.016	0.0049
NO ₃ ⁻	13.7	19.4-23.3	23.1	11.47
SO ₄ ²⁻	-	0.33-0.5	0.65	1.52
Zr	2.47	-	-	3.80
H ₂ O	76.6	76.6-69.2	70.9	76.0
	100.0	100.0	100.0	100.0
Density, g/mL	1.2	1.2-1.3	1.2	1.2

^aSource: IDB 1989.

Table 3B.24. Composition of typical HLW calcines produced at INEL^a

Component	Type of calcine and composition, wt %			
	Alumina	Zirconia	Zirconia-Fluorinel	sodium
Al ₂ O ₃	82-95	13-17	6	12-14
Na ₂ O	1-3	-	--	0-5
ZrO ₂	-	21-27	23	20-26
CaF ₂	-	50-56	56	48-53
CaO	-	2-4	4	2-4
Nitrate	5-9	0.5-2	0.5-2	0.5-4
B ₂ O ₃	0.5-2	3-4	4	3-4
CdO	-	-	6	-
Fission products and actinides	≤1	≤1	≤1	≤1

^aSources: Staples, Knecht, and Berreth, 1986; Knecht 1991.

Table 3B.25. Idaho National Engineering Laboratory:
radionuclide concentrations in 3-year-cooled calcined wastes^a

Radionuclide	Radioactivity (Ci/kg)
Se-79	6.4E-05
Rb-87	3.6E-09
Sr-90	1.3E+01
Y-90	1.3E+01
Zr-93	3.1E-04
Nb-93m	7.5E-05
Tc-99	2.1E-03
Ru-106	9.7E-01
Rh-106	9.7E-01
Pd-107	2.0E-06
Sn-126	3.2E-05
Sb-126m	3.2E-05
Sb-126	3.2E-05
Cs-134	3.3E+00
Cs-135	7.5E-05
Cs-137	1.3E+01
Ba-137m	1.2E+01
Ce-144	8.2E+00
Pr-144	8.2E+00
Pm-147	1.2E+01
Sm-151	1.7E-01
Eu-154	1.8E-01
U-233	1.2E-12
U-234	4.3E-10
U-235	1.8E-09
U-236	1.0E-08
U-237	4.8E-12
U-238	1.0E-14
Np-237	4.8E-08
Pu-238	7.0E-02
Pu-239	7.0E-04
Pu-240	6.5E-04
Pu-241	1.6E-01
Pu-242	1.8E-06
Am-241	9.1E-04
Am-243	8.3E-06
Cm-242	6.5E-04
Cm-244	5.2E-04
Total	8.5E+01

^aSource: IDO 1982. Radioactivities are stated as curies per kg of calcine.

Table 3B.26. Idaho National Engineering Laboratory: radionuclide content per HLW canister^a

Radionuclide	Mass (g/canister)	Radioactivity (Ci/canister)	Thermal power (W/canister)
Se-79	0.1173E+01	0.8173E-01	0.2035E-04
Rb-87	0.5252E+02	0.4597E-05	0.3843E-08
Sr-90	0.1217E+03	0.1660E+05	0.1927E+02
Y-90	0.3051E-01	0.1660E+05	0.9204E+02
Zr-93	0.1575E+03	0.3959E+00	0.4600E-04
Nb-93m	0.3387E-03	0.9577E-01	0.1697E-04
Tc-99	0.1582E+03	0.2682E+01	0.1346E-02
Ru-106	0.3701E+00	0.1239E+04	0.7365E-01
Rh-106	0.3479E-06	0.1239E+04	0.1188E+02
Pd-107	0.4965E+01	0.2554E-02	0.1514E-06
Sn-126	0.1440E+01	0.4086E-01	0.5097E-04
Sb-126m	0.5201E-09	0.4086E-01	0.5203E-03
Sb-126	0.4887E-06	0.4086E-01	0.7552E-03
Cs-134	0.3256E+01	0.4214E+04	0.4290E+02
Cs-135	0.8316E+02	0.9577E-01	0.3197E-04
Cs-137	0.1908E+03	0.1660E+05	0.1837E+02
Ba-137m	0.2848E-04	0.1532E+05	0.6017E+02
Ce-144	0.3282E+01	0.1047E+05	0.6947E+01
Pr-144	0.1386E-03	0.1047E+05	0.7700E+02
Pm-147	0.1653E+02	0.1532E+05	0.5499E+01
Sm-151	0.8250E+01	0.2171E+03	0.2546E-01
Eu-154	0.8513E+00	0.2299E+03	0.2056E+01
U-233	0.1583E-06	0.1532E-08	0.4456E-10
U-234	0.8785E-04	0.5491E-06	0.1582E-07
U-235	0.1063E+01	0.2299E-05	0.6020E-07
U-236	0.1973E+00	0.1277E-04	0.3459E-06
U-237	0.7507E-13	0.6130E-08	0.1158E-10
U-238	0.3797E-04	0.1277E-10	0.3235E-12
Np-237	0.8693E-01	0.6130E-04	0.1874E-05
Pu-238	0.5221E+01	0.8939E+02	0.2963E+01
Pu-239	0.1437E+02	0.8939E+00	0.2754E-01
Pu-240	0.3642E+01	0.8300E+00	0.2585E-01
Pu-241	0.1983E+01	0.2043E+03	0.6336E-02
Pu-242	0.6018E+00	0.2299E-02	0.6788E-04
Am-241	0.3385E+00	0.1162E+01	0.3861E-01
Am-243	0.5315E-01	0.1060E-01	0.3407E-03
Cm-242	0.2510E-03	0.8300E+00	0.3059E-01
Cm-244	0.8201E-02	0.6640E+00	0.2322E-01
Total	0.8315E+03	0.1088E+06	0.3393E+03

^aQuantities are at time of filling canister and are based on 3-year-old calcine immobilized in glass-ceramic with a load of 1,277 kg of calcine per canister (1,825 kg of glass-ceramic per canister). Based on IDO 1982 and Berreth 1986c.

DISTRIBUTION

DOE — Office of Civilian Radioactive Waste Management, Washington, DC 20585

J. W. Bartlett
J. C. Bresee
S. Brocoun
*A. B. Brownstein
*J. H. Carlson
*M. Conroy
W. J. Danker
J. B. Easterling
H. J. Hale
V. King
*P. Kumar
W. Lake

*W. Lemeschewsky
R. Milner
*N. Montgomery
*T. D. Nguyen
M. L. Payton
J. Roberts
S. Rousso
J. D. Saltzman
W. Sprecher
*V. Trebules
*J. R. Williams

DOE — Office of Nuclear Energy, Washington, DC 20545

F. J. Goldner
E. F. Mastal

J. L. Smiley
J. A. Turi

DOE — Office of Defense Programs, Washington, DC 20545

E. A. Jordon
A. L. Taboas

R. Cooperstein
G. H. Daly

DOE — Office of Environmental Restoration, 1000 Independence Ave., Washington, DC 20585

W. W. Bixby

R. P. Whitfield

DOE — Office of Environmental Restoration and Waste Management, Washington, DC 20585-0002

J. A. Coleman
L. Duffy
C. Frank
R. I. Greenberg
E. A. Jordan

A. F. Kluk
M. J. Lilly
W. E. Murphie
G. P. Turi
W. E. Wisenbaker

DOE — Energy Information Administration, 1707 H. Street, NW, Washington, DC 20006

*J. A. Disbrow
*K. Gibbard

*B. K. O'Brien

DOE — Environmental Protection Agency, Washington, DC 20460

J. W. Gunther
M. Halper

M. Oge

DOE — Albuquerque Operations Office, P.O. Box 5400, Albuquerque, NM 87115

*K. Golliher

D. M. Lund

DOE — Chicago Operations Office, Bldg. 201, 9800 S. Cass Ave., Argonne, IL 60439

M. Klimas

S. A. Mann

DOE — Idaho Operations Office, 785 DOE Place, Idaho Falls, ID 83402

M. Fisher
R. Sandwina

K. Svinicki
T. J. Zielinski

DOE — Idaho Operations, West Valley Project Office, P. O. Box 191, West Valley, NY 14171-0191

T. J. Rowland

W. S. Ketola

DOE — Yucca Mountain Site Characterization Project Office, 101 Convention Center Drive, Suite 200, Las Vegas, NV 89109

*M. Blanchard
*C. Gertz
*R. Dyer
*D. Harrison-Giesler

*E. Petrie
*D. Royer
*W. Simecka

DOE — Oak Ridge Operations Office, P.O. Box 2001, Oak Ridge, TN 37831

R. Collier

DOE — Richland Operations Office, P.O. Box 550, Richland, WA 99352

D. C. Langstaff

DOE — Richland Operations, Shippingport Project Office, P.O. Box 335, Shippingport, PA 15077

J. J. Schreiber

DOE — Savannah River Operations Office, P.O. Box A, Aiken, SC 29801

J. M. Gallagher

W. D. Pearson

DOE — WIPP Project Office, P.O. Box 3090, Carlsbad, NM 88220

W. Walker

U.S. Nuclear Regulatory Commission, 1717 H Street NW, Washington, DC 20555

Director, Div. of High-level Waste Mgmt., NMSS, OWFN, MS 4-H-3
Director, Div. of Low-Level Waste Mgmt. and Decommissioning, NMSS, OWFN, MS 5-E-4
Director, Div. of Safeguards and Transportation, NMSS, OWFN, MS 4-E-4
Director, Div. of Industrial and Medical Nuclear Safety, OWFN, MS 6-H-3
Irradiated Fuel Section, Source Containment & Devices Branch

U. S. Environmental Protection Agency, 401 M Street SW., Washington, DC 20460

Analysis and Support Division, Martin Halper (ANR-461)
Criteria and Standards Division, James W. Gunther, (ANR-460W)
Office of Radiation Programs, M. Oge, (ANR-450W)

Office of Technology Assessment, 600 Pennsylvania Avenue., SW., Washington, DC 20510

Defense Nuclear Facilities Safety Board, 625 Indiana Ave., NW, Suite 700, Washington, DC 20004

L. A. Ettlinger

Argonne National Laboratory, 9700 S. Cass Ave., Argonne, IL 60439

S. Y. Chen
J. Peterson

M. J. Steindler

Brookhaven National Laboratory, Upton, Long Island, NY 11973

B. Barletta
R. Fullwood

C. Sastre
R. Youngblood

Lawrence Livermore National Laboratory, P.O. Box 808, Livermore, CA 94550

*J. Blink
*T. Buschek
L. J. Jardine
H. Leider

W. R. Lloyd
R. B. Stout
R. Van Konynenberg

Los Alamos National Laboratory, P.O. Box 1663, Los Alamos, NM 87545

D. T. Oakley
K. K. S. Pillay
L. Walker

J. L. Warren
W. J. Whitty

Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, TN 37831

J. M. Begovich
J. T. Bell
A. G. Croff
C. W. Forsberg
W. Fulkerson
R. K. Genung
H. W. Godbee
D. S. Joy
*J. A. Klein
S. B. Ludwig
A. P. Malinauskas

K. J. Notz (50 copies)
C. V. Parks
R. L. Pearson
*R. B. Pope
R. R. Rawl
W. J. Reich
R. Salmon
S. N. Storch
T. D. Welch
R. M. Westfall

Pacific Northwest Laboratory, P.O. Box 999, Richland, WA 99352

S. R. Bierman
M. E. Cunningham
R. Einziger
U. Jenquin
*A. T. Luksic
J. McElroy

R. W. McKee
G. Mellinger
R. I. Smith
V. W. Thomas
S. Vance
R. C. Walling

Sandia National Laboratories, P.O. Box 5800, Albuquerque, NM 87185

*G. Allen
R. S. Case
A. W. Dennis
T. Hunter
P. C. Reardon

L. E. Romesberg
*E. Ryder
*T. Sanders
*R. Sandoval

Westinghouse/Hanford Company, P.O. Box 1970, Richland, WA 99352

G. M. Bain
C. S. Clark
F. M. Coony
J. A. Nelson

O. A. Turner
R. A. Watrous
R. T. Wilde
B. A. Wolfe

Westinghouse/Idaho Nuclear Company, P.O. Box 4000, Idaho Falls, ID 83404

D. A. Knecht

G. G. Simpson

Westinghouse/WIPP Project, P.O. Box 2078, Carlsbad, NM 88220

K. S. Donovan

J. Lippis

Westinghouse/Savannah River Company, P.O. Box 616, Aiken, SC 29802

P. D. D'Entremont
W. T. Goldston

S. L. Marra
M. J. Plodinec

Advanced Nuclear Fuels Corporation, P.O. Box 90777, Bellevue, WA 98009

J. D. Floyd

Advanced Nuclear Fuels Corporation, 2101 Horn Rapids Road, Richland, WA 99352

D. L. Adams
K. R. Merckx

M. Valentine

Analysas Corporation, 151 Lafayette Drive, Oak Ridge, TN 37830

Debbie Gifford

Anatech Research Corp., P. O. Box 9165, La Jolla, CA 92038

A. J. Zangari

Arizona State University, Nuclear Engineering Science Program, Tempe, AZ 85287-5706

J. W. McKIveen

Automated Sciences Group, 800 Oak Ridge Turnpike, Oak Ridge, TN 37830

*C. G. Cowart
E. A. Dorsey

R. S. Moore (10 copies)
D. Murphy

Babcock and Wilcox, P.O. Box 10935, Lynchburg, VA 24506

P. Childress
J. Matheson
L. Walton

D. A. Wesley
G. Vames

Bechtel National, Inc., P.O. Box 3965, San Francisco, CA 94119

Chin W. Ma

R. J. Tosetti

Bechtel National, 800 Oak Ridge Turnpike, Oak Ridge, TN 37831-0350

P. Bradbury
R. A. Langley

R. Philippon
A. Yazdi

E. J. Bentz & Associates, Inc., 7915 Richfield Road, Springfield, VA 22153

R. Peterson

Brown Boveri Nuclear Services, Inc., 9190 Red Branch Road, Columbia, MD 21045

B. Snyder

Carolina Power and Light Co., Rt. 1, P.O. Box 327, New Hill, NC 27562

R. Wild

Chem-Nuclear Systems, Inc., 220 Stoneridge Drive, Columbia, SC 29210

P. A. Craig

Clark County — Nuclear Waste Division, 301 E. Clark, Suite 570, Las Vegas, NV 89101

J. Duke

Columbia University, Department of Applied Physics, New York, NY 10027

L. Lidofsky

Combustion Engineering, 1000 Prospect Hill Road, Windsor, CT 06095

W. Burns

D. Hayduk

Dataphile, 9121 Garrison Rd., Knoxville, TN 37931

K. E. Jones

David Address Associates, Inc., 11008 Harriett Lane, Kensington, MD 20895

*D. Address

Duke Engineering & Services, P. O. Box 1004, Charlotte, NC 28201

D. Anthony
*R. Eble (3 copies)
*V. H. Pierce

*A. Segrest
*J. Stringer

Ebasco Environmental, 143 Union Boulevard, Suite 1010, Lakewood, CO 80228-1824

W. T. Gregory

Edison Electric Institute, 1111 19th Street NW, Washington, DC 20036

S. Kraft

EG&G Idaho, Inc., P.O. Box 1625, Idaho Falls, ID 83415

M. T. Magleby
R. R. Piscitella

R. L. Scott
H. Worle

Energy Resources International, Inc., 1015 18th St., N.W., Suite 500, Washington, DC 20036

M. H. Schwartz

EPRI — Electric Power Research Institute, P.O. Box 10412, Palo Alto, CA 94303

C. Braun
O. Ozer
R. L. Yang

R. W. Lambert
R. Williams

E. R. Johnson Associates, Inc., 10461 White Granite Drive, Suite 204, Oakton, VA 22124

*B. M. Cole
E. R. Johnson
D. C. Jones
*D. A. Williamson

*N. B. McLeod
M. White
*R. R. MacDonald
*W. Teer

Fluor Daniel, Inc., 3333 Michelson Drive, Irvine, CA 92730

G. E. Benedict
S. Little

T. O. Mallonee
B. R. McElmurry

Foster Wheeler Development Corporation, 12 Peach Tree Hill Road, Livingston, NJ 07039

S. E. Himsl

Georgia Institute of Technology, Nuclear Engineering Program, Atlanta, GA 30332-0405

S. I. Abdel-Khalik

D. B. Bullen

General Atomics, P. O. Box 85608, MS 15-219A, San Diego, CA 92186-9784

J. Boshoven

R. M. Grenier

General Electric Nuclear Energy, 175 Curtner Ave., San Jose, CA 95125

S. O. Akerlund

R. E. Spicka

H & R Technical Associates, 575 Oak Ridge Turnpike, Oak Ridge, TN 37830

W. R. Rhyne

International Atomic Energy Agency, P. O. Box 250, A-1400 Vienna, Austria

M. J. Bell
A. Grigoriev

B. F. Judson

Institute of Nuclear Energy Research, P. O. Box 3-7, Lung-Tan, 32500 TAIWAN

L. C. Liu

Kaiser Engineering, P.O. Box 23210, Oakland, CA 94623

H. Isakari

Korea Atomic Energy Research Institute, P. O. Box 7, Dalduk-Danji, Taejeon, Korea 305-606

Jong Won Choi

Los Alamos Technical Associates, 6501 Americus Parkway, Suite 900, Albuquerque, NM 87110

D. Guerin

Louisiana State University, Nuclear Science Center, Baton Rouge, LA 70803-5820

E. N. Lambremont

M. L. Williams

Massachusetts Institute of Technology, Dept. of Nuclear Engineering, 77 Massachusetts Avenue., Bldg. 24-102, Cambridge, MA 02139

M. S. Kazimi
N. C. Rasmussen

N. E. Todreas

Monsanto Research Corporation, Mound Facility, Miamisburg, OH 45342

J. W. Doty

National TSING HUA University, Dept. of Nuclear Engineering, 101 Section 2, Kuang-Fu Rd., Hsinchu 30043, TAIWAN

S - P. Teng

North Carolina State University, Dept. of Nuclear Engineering, 1110 Burlington Laboratories, Raleigh, NC 27695-7909

D. J. Dudziak
T. S. Elleman

P. J. Turinsky

Northeast Technology Corp., 26 Pearl Street, UPO Box 4178, Kingston, NY 12401

K. O. Lindquist

Northeast Utilities, 107 Selden St., P. O. Box 270, Berlin, CT 06037

G. Betancourt

B. Isakson

Nuclear Assurance Corp., 6251 Crooked Creek Road, Norcross, GA 30092

M. Aucoin
I. F. Stuart

Information Center

NUS Corporation, 910 Clopper Road, Gaithersburg, MD 20878

C. Heath

Y. S. Kim

Nutech Engineers, 340 Six Branches Court, Roswell, GA 30076

J. D. Rollins

Nutech Engineers, 14 S. Martinvale Lane, San Jose, CA 95119

N. W. Edwards

Oklahoma Dept. of Libraries, U.S. Government Infor. 200 N.E. 18th St., Oklahoma City, OK 73105-3298

S. Beleu

Oregon State University, Dept. of Nuclear Engineering, Radiation Center C116, Corvallis, OR 97331-5902

A. H. Robinson

J. C. Ringle

PAI Corporation, P.O. Box 1135, Oak Ridge, TN 37830

D. L. Phung

F. E. Harrington

Pacific Nuclear Fuel Services, Inc., 145 Martinvale Lane, San Jose, CA 95119

L. M. Stamm

Pacific Nuclear Systems, 1010 South 336th Street, Federal Way, WA 98003

C. J. Temus

Public Utility Commission of Texas, 7800 Shoal Creek Boulevard, Suite 400N, Austin, TX 78757

J. McArthur

Purdue University, School of Nuclear Engineering, West Lafayette, IN 47907

V. H. Ransom

Radioactive Exchange, P.O. Box 9528, Washington, DC 20016

E. L. Helminski

Raytheon, Yucca Mountain Project, 101 Convention Center Dr., Suite P-250, Las Vegas, NV 89109

R. C. DeKlever

REECO, P.O. Box 14400, Las Vegas, NV 89114

G. W. Kendall

Rensselaer Polytechnic Institute, Dept. of Nuclear Engineering & Engineering Physics, Troy, NY 12180-3590

R. C. Block

D. R. Harris

RE/SPEC Inc., 4775 Indian School Road NE, Suite 300, Albuquerque, NM 87110

S. Petney

Rochester Gas and Electric Corp., 89 East Avenue, Rochester, NY 14649

A. Fuierer

Rockwell International Corporation, P. O. Box 800, Richland, WA 99352

K. A. Gasper

Rockwell International Corp., 6633 Canoga Avenue, Canoga Park, CA 91303

J. Guon

Rogers & Associates Engineering Corporation, P. O. Box 330, Salt Lake City, UT 84110-0330

B. Bailey

San Francisco State University, 1600 Holloway Ave., San Francisco, CA 94132

Y. T. Mandra

Science Applications International Corp., 4900 Hopyard Road, Suite 310, Pleasanton, CA 94588

J. Cohen

C. Smith

Science Applications International Corp., 800 Oak Ridge Turnpike, Oak Ridge, TN 37830

E. Bergman

R. Best

Science Applications International Corp., 20030 Century Boulevard, Suite 201, Germantown, MD 20874

R. E. Yoder

Science Applications International Corp., 101 Convention Center Drive, Las Vegas, NV 89109

W. MacNabb

R. D. Zimmerman

*R. Morissette (3 copies)

*M. Voegele (3 copies)

J. Nelson

Science Applications International Corp., Atlanta, GA

L. Danese

Southwest Research Institute, 6220 Culebra Rd., San Antonio, TX 78228

H. Manaktala

State of Washington, Department of Social and Health Services, Olympia, WA 98504

E. L. Gronemeyer

State of Nevada, Division of Health, 505 East King St., Carson City, NV 89710

S. Marshall

Texas A&M University, Dept. of Nuclear Engineering, 129 Zachry Engineering Center, College Station, TX 77843

K. L. Peddicord

J. W. Poston

The Ohio State University, Nuclear Engineering Program, 206 West 18th Avenue, Columbus, OH 43210

A. W. Fentiman

D. W. Miller

The Pennsylvania State University, Nuclear Engineering Dept., 231 Sackett Building, University Park, PA 16802

E. H. Klevans

TLG Engineering, 31 N. Pond Rd., Chesshire, CT 06410

A. Levin

TRW Environmental Safety Systems Inc., 2650 Park Tower Drive, Suite 800, Vienna, VA 22180-7306

*W. Bailey (3 copies)

*P. Lovett

*E. Bogart

*J. Mayer

*G. Carruth (5 copies)

J. Miller

*J. R. Clark

*L. Rickertson

*W. R. Griffin (3 copies)

*W. Stanley

*C. Kerrigan (17 copies; 1 controlled)

G. Vawter (2 copies)

TRW Environmental Safety Systems Inc., 600 Maryland Avenue SW, Suite 695, Washington, DC 20024

P. M. Krishna

TRW Environmental Safety Systems Inc., 101 Convention Center Drive, Suite P-110, Las Vegas, NV 89109

*H. Benton (2 copies)

J. Gauthier

J. Berg

H. Hartman

J. K. Clark

R. Jiu

*T. Doering (2 copies)

R. Low

M. Fortsch

S. Pahwa (2 copies)

*L. D. Foust

*W. D. Schutt

University of Arizona, Department of Nuclear and Energy Engineering, Tucson, AZ 85721

S. C. Crow

M. E. Wacks

University of California, Dept. of Nuclear Engineering, Berkeley, CA 94720

D. R. Olander

S. G. Prussin

T. H. Pigford

University of California, School of Engineering and Applied Science, 405 Hilgard Ave., Los Angeles, CA 90024-1597

P. Friedmann

D. Okrent

University of Cincinnati, Dept. of Mechanical, Industrial and Nuclear Engineering, Mail Location 72, Cincinnati, OH 45221

R. Eckart

University of Colorado, Campus Box 184, Boulder, CO 80309-0184

Government Publications

University of Florida, Dept. of Nuclear Engineering Sci., 202 Nuclear Sciences Ctr, Gainesville, FL 32611

J. S. Tulenko

University of Illinois, Dept. of Nuclear Engineering, Urbana, IL 61801

R. A. Axford

B. G. Jones

University of Lowell, Dept. of Chemical and Nuclear Engineering, One University Avenue, Lowell, MA 01854

J. P. Phelps

University of Michigan, 108 Cooley Lab, North Campus, Dept. of Nuclear Engineering, Ann Arbor, MI 48109-2104

W. R. Martin

University of Missouri-Columbia, Nuclear Engineering Program, 333 Electrical Engineering Bldg., Columbia, MO 65211

J. F. Kunze

University of Missouri-Rolla, Department of Nuclear Engineering, 102 Fulton Hall, Rolla, MO 65401-0249

A. E. Bolon

N. Tsoulfanidis

University of Nevada — Las Vegas, J. R. Dickinson Library, 4505 Maryland Pkwy, Las Vegas, NV 89154

K. Schott

University of New Mexico, Dept. of Chemical and Nuclear Engineering, Farris Engineering Center, Room 209, Albuquerque, NM 87131

F. L. Williams

University of Tennessee, Knoxville, Nuclear Engineering Dept., 315 Pasqua Building, Knoxville, TN 37996-2300

H. L. Dodds

T. W. Kerlin

E. M. Katz

L. F. Miller

University of Wisconsin-Extension, Wisconsin Geological and Natural History Survey, 3817 Mineral Point Road, Madison, WI 53705

T. J. Evans

University of Wisconsin-Madison, Dept. of Nuclear Engineering and Engineering Physics, 151 Engineering Research Building, Madison, WI 53706

M. W. Carbon

M. L. Corradini

Westinghouse Electric Corp., Nuclear Waste Technology, 1501 Ardmore Blvd., Pittsburgh, PA 15221

R. Bahorich
P. J. Kersting

B. R. Nair
W. J. Severson

West Valley Demonstration Project, P.O. Box 191, West Valley, NY 14171-0191

R. L. Crocker
E. Maestas

J. Pope

West Valley Nuclear Services Company Inc., P. O. Box 191, West Valley, NY 14171-0191

G. G. Baker
S. M. Barnes
R. L. Crocker
R. G. Gessner
D. H. Kurasch

R. E. Lawrence
R. A. Palmer
D. K. Ploetz
S. J. Szalinski
WVDP Technical Library

Weston, 955 L'Enfant Plaza, SW, Eighth Floor, Washington, DC 20024

*P. A. Bolton
*E. Ching
A. Mobasheran

*N. A. Norman
*M. Rahimi
W. Wowak

ORNL Central Research Library
ORNL Laboratory Records Department
ORNL Laboratory Records, RC
ORNL Y-12 Technical Library, Document Reference Section
ORNL Patent Section

Given distribution as shown in document DOE/OSTI-4500-R75 under categories UC-810, 812, and 820 (299 copies including those for NTIS):

UC-810 High Level Radioactive Waste Management
UC-812 Spent Fuel Storage
UC-820 Transport of High Level Radioactive Waste

Extra copies for OSTI to fill orders (15).

NOTE: Asterisk by name means controlled distribution.