NOV 28 1992

DOE/PC/89785--12

DE-FG22-89PC89785-12

DOE/PC/89785--12

DE93 006014

THE SINGLE ELECTRON CHEMISTRY OF COALS

Report for the period
July 1 to September 30, 1992

John W. Larsen
and Thomas P. Eskay

Department of Chemistry
Lehigh University
Bethlehem, PA 18015

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PREPARED FOR THE UNITED STATES DEPARTMENT OF ENERGY

Received by OSTI

JAN 20 1993

MASTER

doc

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

OBJECTIVES AND SCOPE OF WORK

Objectives

To investigate and characterize the single electron reactions of alkyl and alkoxy aromatic compounds in order to determine the role these reactions play in the chemistry of coal.

Scope of Work

1. Attempts will be made to demonstrate that the radicals from inertinite maceral group will initiate the polymerization of 4-vinylpyridine.
2. The molecule, N,N-diphenyl-phenylenediamine, will be deposited in coals to characterize their native free radicals.
3. Tetracyanoquinodimethane (TCNQ) and tetracyanoethylene (TCNE) will be used to characterize the numbers and strengths of single electron donors in coals.

Introduction

Depolymerization of coals at low temperatures may offer advantages over thermal bond cleavage. Because bond cleavage energies of radical cations are lower than the corresponding homolytic bond cleavage energies of the same bond^{1,2}, generation of radical cations in coal may make possible depolymerization at lower temperatures.

We seek to investigate the above possibility using single molecules containing functional groups common in coals. Since the generation of a radical cation requires the removal of an electron from a neutral molecule, a primary focus of the study will be finding oxidants that will remove an electron from compounds with structural similarity to those typically found in coals. The study will also be concerned with the decomposition of radical cations and the products formed as a result of the decomposition.

Results and Discussions

In our last report³ we described that treatment of bibenzyl and neo-pentylbenzene with $\text{Fe(III)(1,10-phenanthroline)}_3(\text{ClO}_4)_3$ (**FE(III)(PHEN)**) in refluxing CH_3CN (82°C) failed to produce substantial bond cleavage β to the aromatic ring. Because bond cleavage was not observed, we have continued our study by moving to compounds which have lower ionization potentials as well as studying other oxidants.

Oxidation of 4,4'-dimethoxybibenzyl (DMBB) with excess **FE(III)PHEN** was carried out in refluxing CH_3CN (see experimental section for details) Gas chromatography analysis of the reaction mixture revealed 77 % of the starting material, **DMBB**, remained with 5 %(weight percent) of the cleavage product p-anisaldehyde

Equation 1

(**PA**) (eq 1.) (see Table I in the data and results section). A mass balance revealed that 23 % of the starting material is not detected by GC analysis. This loss of material is most likely due to the dimerization of **DMBB** radical cations formed. This is a known process and we have demonstrated that dimerization occurs in the single electron oxidation of benzyl phenyl ether.⁴

To investigate if trace amounts of H_2O or O_2 were responsible for the formation of **PA**, reactions were performed in the presence of air or added H_2O . The results of these experiments are given in Tables II-IV in the data and results section. The same result was observed for the reaction whether or not a stream of air was bubbled through the solution. In the presence of added H_2O , small amounts of two additional unidentified products were observed by GC. The water reactions also indicate a slight increase in the

amount of unreacted **DMBB** recovered, although the amount of **PA** produced was unchanged when compared to reactions run under anhydrous conditions.

The oxidation of **DMBB** was attempted using other known single electron oxidants. When **DMBB** was treated with tris(p-bromophenyl)aminium hexachloroantimonate in refluxing CH_3CN , 64 ± 1 % of the **DMBB** was recovered after 18 hours and no cleavage products were detected by GC analysis. When **DMBB** was refluxed in CH_3CN in the presence of excess ammonium cerium(IV) nitrate 98 % conversion of starting material was observed and 24 % **PA** was formed (Table V in the Data and Results Section). The low mass balances observed in these reactions by GC analysis may indicate the formation of dimers or polymers from **DMBB** radical cations generated. Further analysis of the reaction products and studies of other oxidants are currently being carried out.

Experimental Section

All compounds were commercially available, except for **FE(III)(PHEN)** and **DMBB** (synthesis below), and were used as purchased or repurified as follows. Anhydrous acetonitrile, p-anisaldehyde, and tris(4-bromophenyl)aminium hexachloroantimonate (Aldrich). Phenanthrene (Baker) the GC internal standard was recrystallized from absolute ethanol prior to use. **DMBB** and **FE(III)(PHEN)** were dried to constant weight in vacuum (0.4 torr.) across from P_2O_5 .

Ultraviolet spectra were obtained on a Perkin-Elmer Lambda 5 UV/VIS spectrophotometer between the wavelengths of 200 and 900 nm. with a slit width of 2 nm. The solvent for both reference and sample was anhydrous acetonitrile using 1 cm. quartz cells.

NMR spectra were obtained on a JEOL FX90Q FT NMR spectrometer.

Iron(III)(1,10-phenanthroline)₃(perchlorate)₃

Iron(III)(1,10-phenanthroline)₃(perchlorate)₃ was prepared by the method described in the previous paper.⁴ Iron(II) sulfate heptahydrate 1.34 g. (.0048 moles) and 2.59 g. (.014 moles) of 1,10-phenanthroline were warmed with stirring in 29 ml. of water. When all the solid had dissolved, the resulting deep red solution was filtered into a flask which was immersed in an ice bath. 7.2 ml. of 1M H_2SO_4 was then added to the solution. After cooling in an ice bath, 4.58 g. (.019 moles) of lead(IV) oxide was added and the

solution was stirred for approximately 10 minutes until all the iron(II) (red) was converted to iron(III) (blue). Excess lead(IV) oxide was filtered off by using a sintered funnel and 1.76 g. (.014 moles) of sodium perchlorate was added to the filtrate. The solution was stirred and a blue precipitate appeared. The precipitate was filtered and recrystallized from dilute perchloric acid (1:1 v:v). After drying (see above) it was characterized by UV spectroscopy λ_{max} 590 nm. in acetonitrile (lit. 592⁵). The synthesis was completed as quickly as possible since the complex decomposes in aqueous solution.

4,4'-dimethoxybibenzyl

DMBB was synthesized using a procedure by Trahanovsky⁶. 0.84 g. (0.035 moles) of magnesium and 10 ml. of anhydrous diethyl ether were placed into a flame dried two neck 50 ml. flask under N₂ containing a magnetic stirrbar and equipped with a flame dried addition funnel. A 10 ml. diethyl ether solution containing 10 g. (0.069 moles) of p-methoxybenzyl chloride (Aldrich) was added to the addition funnel and a dropwise addition of the solution was carried out over the period of one hour. The resulting solution was allowed to stir overnight. The reaction was quenched with the dropwise addition of H₂O after which 3 M. HCl was added until the aqueous layer was acidic to litmus paper. The aqueous layer was extracted with ether (3 X 15 ml.) and a white solid present remained in this layer. The solid was filtered from the ethereal layer taken up in CH₂Cl₂ and filtered to remove residual magnesium salts. The CH₂Cl₂ was removed by rotary evaporation and the solid

was recrystallized from absolute ethanol to give 2.1 g.(25%)of product. M.p. 125-127°C (lit.⁶ 125-126°C). NMR (CDCl₃ , TMS internal standard) 7.2-6.95 (m, 4H), 6.9-6.7 (m, 4H), 3.75 (s, 6H), 2.75 (s, 4H).

Reactions DMBB with FE(III) (PHEN)

The following is a representative procedure. An oven dried 25 ml. two neck flask containing a magnetic stirrbar was flushed with N₂ while hot, capped, and cooled to room temperature. the flask was weighed and moved into a glovebox (N₂ atmosphere) where DMBB was placed into the flask. The flask was removed from the glovebox and weighed to give 0.05031 g. (0.208 mmoles) DMBB. The flask was then equipped with an oven dried condenser under N₂ and was wrapped in Al foil. The flask was then charged with 0.5 ml. of anhydrous CH₃CN. 3.789 g. of an FE(III) (PHEN) solution prepared by dissolving 0.546 g. of FE(III)PHEN in 7.904 g. of CH₃CN was then injected into the reaction flask. By weight percent 0.244 (0.274 mmoles) FE(III) (PHEN) was transferred. The flask was immediately lowered into an oil bath at 100°C and stirring was started. After 3 hrs. the reaction solution was removed from the oil bath. After cooling, 10 drops of saturated aqueous NaHCO₃ was added and the solution was allowed to stir for 0.5 hrs. 0.0516 g of phenanthrene (the internal standard) was added to the solution and the solution was diluted with 10 ml. CH₃CN. A portion of this solution was removed, diluted with 10 ml. CH₃CN. dried over MgSO₄, and used in the GC analysis.

Reactions containing water were carried out in the same manner with the addition of 0.5 ml. of H₂O prior to the addition of **FE(III)(PHEN)** solution. Reactions of **DMBB** with other oxidants were carried out in an analogous manner as described for **FE(III)(PHEN)** reactions.

Chromatography Analysis and Methods

Gas chromatography was carried out using a (10 m. x 0.53 mm. ID) Supelco Nukol column in a Hewlett-Packard 5890 Series II gas chromatograph equipped with an FID detector and on column injector port. The carrier gas was He at a flow rate of 10 cm³/min. determined at the detector at an oven temperature of 75°C. The analysis was carried out using the following program and an injection size of 0.3 ul.: oven temperature 75°C for 0.5 min., 10°C/min increase to 200°C, 200°C for 35 min. Injector: 75°C for 0.5 min., 40°C/min. to 300°C, 300°C for 15 min. The detector temperature was 250°C.

All reported GC data were response factor corrected using phenanthrene as an internal standard. At least 3 injections of a response factor standard were made for each analysis except where otherwise noted.

Data and Results

Table I. Reaction **DMBB** with **FE(III) (PHEN)**. 73 % recovery of **DMBB** after 3 hrs. reflux in CH_3CN . Mass balance $77 \pm 3 \%$ ^a.

Product	Weight %
PA	5 %

Table II. Reaction **DMBB** with **FE(III) (PHEN)** in the presence of air. 69 % recovery of **DMBB** after 16 hrs. reflux in CH_3CN . Mass balance $72 \pm 2 \%$ ^a.

Product	Weight %
PA	3 %

Table III. Reaction **DMBB** with **FE(III) (PHEN)** in the presence of water. 79 % recovery of **DMBB** after 3 hrs. reflux in CH_3CN . Mass balance $85 \pm 4 \%$ ^a. Small quantities of 2 additional products were detected by GC analysis.

Product	Weight %
PA	6 %

Table IV. Reaction **DMBB** with a 250 % molar excess **FE(III) (PHEN)** in the presence of water with. 81 % recovery of **DMBB** after 60 hrs. at room temperature in CH_3CN . Mass balance $87 \pm 3 \%$ ^a. Small quantities of 2 additional products were detected by GC analysis.

Product	Weight %
PA	6 %

Table V. Reaction **DMBB** with $(\text{NH}_4)_2\text{Ce}(\text{IV})(\text{NO}_3)_6$. 98 % **DMBB** reacted after 18 hrs. refluxing in CH_3CN . Mass balance 23 ± 1 %^a.

Product	Weight %
PA	24 %

a. Error given as the standard deviation of scatter in GC runs.

References

1. D. M. Camaioni, J. A. Franz, *J. Org. Chem.*, **1984**, 1601.
2. A. Okamoto, D. R. Arnold, *Can. J. Chem.*, **1985**, 2340.
3. Larsen, J. W., Eskay T. P., *The Single Electron Chemistry of Coals*, A paper prepared for the Department of Energy, # DE-FG22-89PC89785-11.
4. Larsen, J. W., Eskay T. P., *The Single Electron Chemistry of Coals*, A paper prepared for the Department of Energy, # DE-FG22-89PC89785-10.
5. Wong, C. L., Kocki, J. K., *J. Amer. Chem. Soc.*, **1979**, 5593.
6. Trahanovsky, W. S., Brixius, D. W., *J. Amer. Chem. Soc.*, **1973**, 6778.

**DATE
FILMED**

2 / 17 / 93