LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Studies of High-K Isomers at TRIUMF-ISAC

M. B. Smith, P. M. Walker, G. C. Ball, J. J. Carroll, P. E.
Garrett, G. Hackman, R. Propri, F. Sarazin, H. C.
Scraggs

March 12, 2004

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.

STUDIES OF HIGH- K ISOMERS AT TRIUMF-ISAC

M.B. Smith,¹ P.M. Walker^{1,2} G.C. Ball,¹ J.J. Carroll,³ P.E. Garrett,⁴ G. Hackman,¹
R. Propri,³ F. Sarazin,¹ H.C. Scraggs¹

¹*TRIUMF, 4004 Wesbrook Mall, Vancouver, BC, Canada V6T 2A3*

²*Department of Physics, University of Surrey, Guildford, Surrey GU2 7XH, UK*

³*Department of Physics and Astronomy, Youngstown State University, OH 44555, USA*

⁴*Lawrence Livermore National Laboratory, Livermore, CA 94551, USA*

The 8π spectrometer, an array of 20 Compton-suppressed HPGe detectors situated at TRIUMF, has been reconfigured as a high-efficiency device for γ -ray spectroscopy with radioactive beams [1], and recently augmented by the addition of the 20-plastic-scintillator SCEPTAR array and a moving tape transport. This powerful combination will be used for $\beta - \gamma$ coincidence measurements following the implantation of radioactive beams from the ISAC facility, produced by bombarding 500 MeV protons with refractory production targets. By detecting β and γ radiation, while removing long-lived isobars with the tape transport, isomers and β decays are characterized by their half-lives and decay schemes.

One of the first experimental programs to use the new detection system is a study of high- K isomers in neutron-rich nuclei with $A = 170 - 190$. Many such isomers, which arise because of the K -selection rule, are known [2] in this region of the nuclear chart (albeit closer to stability) and their properties elucidate basic nuclear-structure information, particularly on the interplay of single-particle and collective excitations. The initial focus at TRIUMF is on the doubly-mid-shell region around ^{170}Dy [3].

As a precursor to the radioactive-beam experiments, the decay of the remarkable $^{178m2}\text{Hf}$ isomer ($T_{1/2} = 31$ years) has been investigated with a 15 kBq source placed at the center of the 8π spectrometer [4]. High-multipolarity $M4$ and $E5$ transitions, which represent the first definitive evidence for γ -ray emission directly from the isomer, have been identified, together with other low-intensity transitions. Branching ratios for these latter transitions elucidate the spin-dependence of the mixing between the two $K^\pi = 8^-$ bands. The $M4$ and $E5$ transitions are the first strongly K -forbidden examples with such high multipolarities, and demonstrate a consistent extension of K -hindrance systematics. The ^{178}Hf results, and the status of the 8π -SCEPTAR experiments, will be discussed.

References

- [1] C.E. Svensson *et al.*, Nucl. Instrum. Methods Phys. Res. B **204**, 660 (2003).
- [2] P.M. Walker and G.D. Dracoulis, Nature **399**, 35 (1999).
- [3] P.H. Regan *et al.*, Phys. Rev. C **65**, 037302 (2002).
- [4] M.B. Smith *et al.*, submitted to Phys. Rev. C.