SPECIFICITY OF MUTATION INDUCED BY IONIZING
RADIATION AND OXIDIZING FREE RADICALS

DOE/ER/60678--5

DE93 006077

Progress Report

for Period April 15, 1992 - April 14, 1993

Bernard S. Strauss

University of Chicago
Chicago, Illinois 60637

January 1992

Prepared for

DOE/ER/60678-5

JAN 20 1993

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

U.S. DEPARTMENT OF ENERGY
AGREEMENT NO. DE-FG02-88ER60678

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DE-FG02-88ER60678. A004

B. Strauss P.I.

Factors Affecting Mutational Specificity Induced by Ionizing Radiation and Oxidizing Radicals

Technical Progress Report.

Period Covered: Feb 1, 1992- October 15, 1992

1. Reminder of Objectives (Copied from the report of the A. 003 year)

" The ABSTRACT of our 1988 Proposal continues to give the objectives of our work.

We propose to analyze the factors affecting the specificity of mutational change as induced by ionizing radiation and oxidizing radicals. We want to understand not only the rules that affect base substitution, but also the mechanism(s) by which additions and deletions are produced, since deletions are a common consequence of radiation. We wish to carry out this analysis in an *in vitro* mutation system that permits us to analyze the role of base sequence, of polymerase and of mutagenic agent. Our system is designed to screen out most direct breaks as a cause of mutation and should indicate the changes resulting from base damage to the DNA. We want to answer the following questions:

- a) What types of base substitution are produced by ionizing radiation and oxidizing radicals?
- b) Are deletions and/or additions produced?
- c) Is there a difference in the type of mutation produced dependent on the polymerase used? Do mammalian polymerases plus their accessory factors result in different patterns of mutation?
- d) What is the mechanism by which base damage is converted to mutation?"

2. Specific Progress since the last report (2/1/92)

A. We published a general view of the mutation process in tumors (reprint accompanying report).

Strauss, B. 1992. The origin of point mutations in human tumor cells. *Cancer Research* 52: 249-253.

A brief report of our recent work on the production of frameshift mutation *in vitro* was included in a symposium on DNA repair (in press). The material in this paper is covered in the present report and a copy of the manuscript is appended.

Strauss, B., Sagher, D. and Turkington, E. The biological role of the different DNA repair systems: The association of replication and repair. Alfred Benzon Symposium 35. DNA Repair Mechanisms. (Eds. V. Bohr, K. Wasserman, K. Kraemer and J.H. Thaysen) Munksgaard Int. Publishers, Copenhagen *in press*

B. The following information about ionizing radiation-induced damage to DNA is based on an electrophoretic method for distinguishing single stranded circles and linear molecules of the same size.

1. Transfection activity and survival of intact single stranded DNA as a function of dose of ^{60}Co γ rays.

We interpret this data to mean that a major portion of the inactivating effect of gamma irradiation on single stranded DNA comes from damage to the DNA molecule that does NOT involve breakage of the phosphodiester chain.

2. M13mp2 RF and M13mp2 single stranded DNA irradiated with ^{60}Co γ rays at the same concentration were analyzed for biological activity (transfection activity) and for strand breakage by gel electrophoresis. Strand breaks convert double stranded supercoiled molecules into relaxed forms and single strand circles into linear molecules. A dose of 100 Grays resulted in the conversion of a double stranded preparation from a visually estimated initial value of 90 percent supercoils and 10 percent relaxed molecules to approximately 25 % supercoiled and 75 percent relaxed molecules. In contrast the same dose to single stranded DNA resulted in the conversion of a preparation of 97 % circular and 3 % linear to 69% circular and only 31% linear. We conclude that the supercoiled double-

stranded DNA is more sensitive to radiation-induced cleavage than is single stranded DNA.

C. In order to study frameshift mutation we have obtained a mutant M13mp2 molecule containing an extra T (+1 frameshift) to give the sequence:

5' GCC GTC GTT(T)TTA CAA CGT 3'

Virus carrying this sequence give colorless plaques when plated with appropriate host bacteria on appropriate medium. Frame-shift revertants give blue plaques. The experimental advantage in use of this molecule for mutation studies is that it is possible to go from colorless (original) to blue plaques (frameshift mutant), a change which is easy to detect. We used ultraviolet irradiation to test the method since the sequence of 5 T's should present a good target for the formation of pyrimidine dimers. The following data shows the production of mutants *in vitro* after irradiation of single stranded (+T) DNA, formation of purified hybrid by the method described in our previous Progress Report (A003) and synthesis using two related enzymes.

This experiment shows that Sequenase, a genetically engineered T7 DNA polymerase devoid of 3'→5' exonuclease activity, induces frameshift mutations whereas the parent T7 DNA polymerase does not. Sequence analysis of the mutants obtained indicates that most occur in the region of T's as expected.

Some of the mutants result from both frame shift and base substitution. For example, the sequence 5' GCC GTC GTT(T)TTA CAA CGT 3' has been altered to produce 5' GCC GTC GTT ATA CAA CGT 3' and 5' GCC GTC GTT CTA CAA CGT 3'. Such changes can be used to deduce the probable position of the dimer and the mechanism of the change. "Transient misalignment" as suggested by Kunkel is a likely explanation. Additional substrates, with different changes that can be corrected by frameshift are being studied.

We want to know what properties of DNA polymerase are important in determining the ability to make frameshift mutations. Sequenase differs from the wild-type T7 DNA polymerase in at least two properties: the activity of the 3'-->5' exonuclease and the rate of polymerization. Loss of exonuclease activity is associated with increased rate of polymerization. We have obtained from S. Tabor and C. Richardson a set of genetically altered polymerases. We have made the following observations

ENZYME	Polymerase Activity*	Exonuclease Activity ^	Control template	UV template
Δ28	880	<0.001	0.21%	1.15%
Δ6	580	<0.001	0.20%	0.79%
his ¹²³ -->Glu	350	0.3	0.29%	0.47
no synthesis			0.11%	<0.1%

*^ from Tabor and Richardson, J. Biol. Chem. 264: 6447 (1989). Activity is given relative to native T7 DNA polymerase set at 100.

Δ6 = deletion of (Lysine 118- His 123)

Δ28 = deletion of (Lys 118 - Arg 145) = Sequenase

These data, along with other observations showing, for example, that the Taq DNA polymerase (devoid of 3'-->5" exonuclease activity) does not make frame shifts suggests that low exonuclease activity alone is not responsible for the frameshifts.

IWe are inquiring as to the relation between the changes observed *in vitro* and *in vivo* events. For this purpose we are constructing *Escherichia coli* mutator strains which do not repair frameshift damage and which carry repair deficiency genes in order to produce and examine frameshift mutants produced in virus carrying the same frameshifts used for the *in vitro* experiments.

D. We are extending these studies to understand the nature of damage induced by ionizing radiation and oxidizing agents. These experiments build on the observations we recently reported (Mutation Res. 251: 187 (1991)).

1) We produce gapped hybrid is made by hybridizing double stranded fragment with single stranded (irradiated) DNA, removing single stranded DNA with hydroxylapatite and then removing linear fragments with the *M luteus* ATP dependent nuclease as described in our last Progress Report. We found the following:

Yield of hybrid (% of starting material)

Control	300 Gray	600 Gray
57%	29 %	18%

These data imply that as suggested by us (Mutation Res. 251: 187 (1991)), the dose-effect relationship observed in *in vitro* mutagenesis by γ -rays can be obscured by changes in the yield of hybrid.

2) DNA synthesis catalyzed by Sequenase on irradiated hybrid is greatly increased over the amount on control DNA indicating the presence of additional primer sites. These data indicate that circular gapped hybrid can be formed from irradiated DNA containing some internal nicks terminated at 3'OH sites which serve as sites for the initiation of DNA synthesis in addition to initiation at the 3'OH end of the gap.

3) Use of equal amounts of hybrid formed from a substrate designed to test for frameshifts showed no evidence of γ -ray induced frameshifts after DNA synthesis as compared to those induced by UV. We are surprised at this result since our previous studies indicated the region of the substrate containing an extra T to be a hotspot for frameshift mutations.

Frameshift Mutants observed on a substrate with an extra T at position 70.

<u>DOSE</u>	<u>No synthesis</u>	<u>Sequenase</u>	<u>T7 polymerase</u>
0	0.017 [1/5728]	0.13 [7/5421]	0.03 [1/3430]
300 Gray	0	0.21 [5/2372]	0.09 [1/1145]
600 Gray	0.06 [1/1736]	0.15 [3/2026] 0.1 [1/1035]	< 0.05 0.09 [3/3323]
1 KJoule (UV)		2.23 [18/807]	

Values in Percent mutants. Values in [] indicate number of mutants per total number of plaques

**DATE
FILMED**

2 / 12 / 93