LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Multinomial Combinatorial Group Representations of the Octahedral and Cubic Symmetries

K. Balasubramanian

December 29, 2003

Journal of Mathematical Chemistry

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

Multinomial Combinatorial Group Representations of the Octahedral and Cubic Symmetries

K. Balasubramanian^{*}

*Center for Image Processing and Integrated computing, University of
California Davis, Livermore, CA 94550; University of California*

Chemistry and Material Science Directorate

Lawrence Livermore National Laboratory

Livermore, California 94550;

*and Glenn T Seaborg Center, Lawrence Berkeley Laboratory, University
of California, Berkeley, CA 94720*

We consider the full multinomial combinatorics of all irreducible representations of the octahedral (cubic) symmetry as a function of partitions for vertex, face and edge colorings. Full combinatorial tables for all irreducible representations and all multinomial partitions are constructed. These enumerations constitute multinomial expansions of character-based cycle index polynomials, and grow in combinatorial complexity as a function of edge or vertex coloring partitions.

^{*} Address correspondence at kbala@ucdavis.edu or fax:925-422-6810

1. Introduction.

Applications of combinatorics and graph theory to chemical problems and spectroscopy have been the topic of numerous studies in the last several decades [1-33]. These applications have ranged from simple enumeration of chemical isomers, isomerization reactions, enumeration of polyhedral ligand substitutions, chirality, use of mark groups, double cosets, combinatorics of configuration interaction computations, many-electron correlations, NMR, ESR, and so on. Both mathematical and chemical areas have enjoyed significant advances and growth as a result of this cross-fertilization. Pólya's classical paper [6,7] on a combinatorial enumeration technique, now well known as, Pólya's theorem derived its motivation from chemistry. Enumerations of chemical isomers [1-7,8-18] including those of fullerene cages [31] have in turn been benefited by Pólya's technique and its generalization.

Advances have been made in the mathematical context by Williamson [22,23] and Merris [24], and in the chemical context by the current author [20] to generalize Pólya's theorem to all irreducible representations of the group. Generalization of De Bruijn's theorem to all characters has also been accomplished by the author [9]. The current author [19,20] for the first time provided a geometric, chemical or spectroscopic interpretation to the generalization of this technique to all irreducible representations of the groups. While such techniques have been presented they were often restricted to a few kinds of substituents or colors or ligands. The algebra of applying these irreducible representations based combinatorics becomes complex as a function of different types of partitions. Thus these techniques have been often applied to cases of 2 colors for which the expressions become binomial expansions, or three or even four types of such colors. Yet a full combinational classification of these irreducible patterns requires exhaustive types of all possible distinct colors. For example, the edges of an octahedron or a cube can be colored at most with 12 different kinds of colors, vertices of a cube with eight different types of colors and so on. The computational complexity also grows exponentially as the color type partition, as represented by Young's diagram, becomes complex with many rows. Hence exhaustive generation of such combinatorial numbers can be challenging and may provide additional insight.

Many practical applications [20,32] often require such multinomially driven combinatorial expansions. Consider for example, the naturally occurring isotope of Bismuth, ^{209}Bi , has a nuclear spin of $9/2$. This means that there are 10 distinct orientations of the magnetic spin vector, which we represent by $m_f = -9/2, -7/2, -5/2, -3/2, -1/2, 1/2, 3/2, 5/2, 7/2, \text{ and } 9/2$. One can visualize each of these orientations as a distinct type of color and thus the nuclear spin statistics of bismuth clusters would require decanomial combinatorics. The challenge is simply from combinatorial explosions that occur quite rapidly with cluster size. Fullerene cages also present considerable challenge when extended for multinomial combinatorics.

In this work we consider exhaustive multinomial combinatorics of the octahedron and its close relative, the cube. We have constructed exhaustive combinatorial tables for all irreducible representations of the cube vertex (octahedral face) colorings with 8 different types of colors represented by the Young's tableau of 8 for all irreducible representations, edge colorings with young tableau of 12, and vertex colorings of octahedron (face colorings of cube), all of which were accomplished using multinomial combinatorics. This also yields some well-known results on chirality and ligand partitions [25,26,29] as special cases and isomer counts for all types of partitions exhaustively.

2. Multinomial Combinatorics

Let $[n]$ be an ordered partition of n into p parts such that

$$n_1 \geq 0, n_2 \geq 0, \dots, n_p \geq 0, \quad \sum_{i=1}^p n_i = n$$

Then a multinomial expansion in λ 's is defined as

$$(\lambda_1 + \lambda_2 + \dots + \lambda_p)^n = \sum_{[n]} \binom{n}{n_1 \ n_2 \ \dots \ n_p} \lambda_1^{n_1} \lambda_2^{n_2} \dots \lambda_p^{n_p}$$

where $\binom{n}{n_1 \ n_2 \ \dots \ n_p}$ are defined as multinomial coefficients, and the sum is over all such ordered partitions, also known as compositions of the integer n into p parts. The multinomial coefficients can be proven to satisfy

$$\binom{n}{n_1 \ n_2 \ \dots \ n_p} = \frac{n!}{n_1! n_2! \dots n_p!}$$

The sum of all coefficients in the multinomial expansion is p^n . It can also be proven that

$$\binom{n+q}{n_1 \ n_2 \ \dots \ n_p} = \sum_{[k]=n} \binom{n}{k_1 \ k_2 \ \dots \ k_p} \binom{q}{n_1-k_1 \ n_2-k_2 \ \dots \ n_p-k_p}$$

,

where $[k]$ stands for all ordered partitions of m such that $k_1+k_2+\dots+k_p=n$, with k_i non-negative integers.

The well-known Pólya's theorem [7] provides a generating function for multinomial expansions in terms of the ordinary cycle index of a group, which is simply a sum of all orbit structures of permutations of a group when it acts on a set D divided by $|G|$, the order of the group. Williamson [22,23], Merris [24] and the present author [19,20] have independently generalized the cycle index, and in particular, the current author provided a physical and geometrical interpretation of the numbers enumerated.

Define the symmetry operator T_G^χ as

$$T_G^\chi = \frac{1}{|G|} \sum_{g \in G} \chi(g) P(g)$$

where $\chi(g)$ be the character value of $g \in G$ for an irreducible representation Γ in the group G and $P(g)$ is a permutation operator for g . One can define a weighted permutation operator by introducing weights, which also appear in the multinomial expansion. Let D be the set of vertices, edges or faces and R be a set of colors and let us assign a weight w for each color r in R . Then the weight of a function f from D to R is defined as

$$W(f) = \prod_{i=1}^n w(f(i))$$

A permutational operator for each weight W is denoted by $P_w(g)$. In a matrix representation of $P_w(g)$ we can define a tensor version of Pólya's theorem for all irreducible representations in the group. In this representation the trace of $P_w(g)$ is

$$Tr(P_w(g)) = \sum_f \binom{g}{f} W(f),$$

where the sum is over all f for which $gf = f$, we thus obtain

$$T_G^{W,\chi} = \frac{1}{|G|} \sum_{g \in G} \chi(g) P_W(g)$$

$$\therefore \text{Tr} T_G^{W,\chi} = \frac{1}{|G|} \sum_{g \in G} \chi(g) \text{Tr}[P_W(g)] = \frac{1}{|G|} \sum_{g \in G} \chi(g) \sum_f^{(g)} W(f)$$

The above generalized version Pólya's theorem can be conveniently expressed in terms of generalized character cycle index (GCCCI) P_G^χ for every irreducible representation can be defined as

$$P_G^\chi = \frac{1}{|G|} \sum_{g \in G} \chi(g) s_1^{b_1} s_2^{b_2} \dots s_n^{b_n},$$

where the sum is over all elements of the group and $s_1^{b_1} s_2^{b_2} \dots s_n^{b_n}$ is the cyclic polynomial representation if g in G generates b_1 cycles of length 1, b_2 cycles of length 2, ..., b_n cycles of length n when g acts on the D . The difference between Pólya's cycle index and the one above is that for each irreducible representation, we have a cycle index due to the character values multiplying the polynomial.

A multinomial combinatorial function can be obtained for each irreducible representation by substituting every s_k in the above expression. Let n be the number of elements in the set D . Then let R be a set of n distinct types of colors such as white, green, yellow, purple, magenta, green, red, and so on. Let $\lambda_1, \lambda_2, \dots, \lambda_n$ be distinct weights for these colors. The multinomial combinatorial function is obtained by substituting every s_k in the GCCCI by $\lambda_1^k + \lambda_2^k + \dots + \lambda_n^k$. Symbolically,

$$GF^\chi = P_G^\chi (s_k \rightarrow \sum_i \lambda_i^k)$$

Thus the above substitution generates a powerful multinomial expansion where coefficients have rich combinatorial significance for each irreducible representation of

the group. For example, for the identity representation the GF enumerates achiral isomers or equivalence classes in Pólya's term. For the anti-symmetric irreducible representation (the one which has -1 character values for all improper rotations), the index enumerates all chiral isomers. In general, for any irreducible representation Γ whose character is χ , the GF^χ enumerates equivalence classes of functions from the set D to R such that they transform according to the irreducible representation Γ . We believe that this is the most powerful and general interpretation under which all enumerative combinatorics is encompassed.

The terms in the multinomial expansion would have all ordered partitions of n into various parts. First partitions of n are enumerated as provided by Young's diagram. Then for each partition all ordered tuples arising from that partition by allowing permutations of the entries correspond to the terms in the multinomial. This generates all compositions of n into p parts, and p is varied from 0 to n , to represent all terms in the multinomial expansion. The whole generating function then contains terms some of which are combinatorially equivalent. For example, the terms

$\lambda_1^4 \lambda_2^2 \lambda_3 \lambda_4^1$, $\lambda_1^4 \lambda_3^2 \lambda_4 \lambda_6^1$, $\lambda_2^4 \lambda_4^2 \lambda_6 \lambda_7^1$, etc., are all equivalent to each other in combinatorial terms and their coefficients in the multinomial expansion would be the same. Thus the unique terms of the multinomial GF^χ 's are simply represented by the Young diagrams of partition n .

3. Application to the Octahedron (Cube)

The octahedron and its relative cube present one of the most interesting cases of multinomial combinatorics since the three-dimensional octahedral group O_h of 48 operations presents interesting combinatorics, especially for the edge coloring character combinatorics. Thus as a first case, we shall consider the edge colorings of the cube and octahedron for all irreducible representations of the group. Let D be the set of 12 edges and R be a set of 12 distinct colors such as white, yellow, blue, green, red, purple, magenta, cyan, and so on. There are exactly 12^{12} such possible colorings of the edges of the cube. These 12^{12} colorings would transform as various irreducible representations of the O_h group when appropriately symmetry-adapted. In particular the number of totally

symmetric representations, that is, number of maps among 12^{12} maps that transform as the A_{1g} representation of the group is the number of equivalence classes or positional (achiral) isomers for the various distribution of colors. Likewise, the number of functions that transform as the A_{1u} representation, which has the character values of -1 for all improper rotations correspond to the patterns of chiral edge colors among 12^{12} maps, such that one function is not transformable into another chiral representation by the operation of the O_h group. Note that unlike ordinary isomer enumeration, where one considers only the rotational subgroup, we consider the entire group so that chiral, achiral, and in general all irreducible representations are covered.

We shall first construct the various cycle indices for the set D of edges. Note that the polynomials in the cycle index would depend on the set D. Hence for the 12 edges, the polynomial for the A_{1u} irreducible representation of the O_h group is

$$P_G^{A_{1u}} = \frac{1}{48} [s_1^{12} - 3s_1^4 s_2^4 + (3-1)s_2^6 + (-6+6)s_1^2 s_2^5 + (-6+6)s_4^3 + 8s_3^4 - 8s_6^2]$$

where we have shown the terms that cancel out for clarity. By replacing every s_k by

$$\sum_i \lambda_i^k$$

in the above expression, we obtain,

$$P_{O_h}^{A_{1u}} = \frac{1}{48} [(\lambda_1 + \lambda_2 + \dots + \lambda_{12})^{12} - 3(\lambda_1 + \lambda_2 + \dots + \lambda_{12})^4 (\lambda_1^2 + \lambda_2^2 + \dots + \lambda_{12}^2)^4 + 2(\lambda_1^2 + \lambda_2^2 + \dots + \lambda_{12}^2)^6 + 8(\lambda_1^3 + \lambda_2^3 + \dots + \lambda_{12}^3)^4 - 8(\lambda_1^6 + \lambda_2^6 + \dots + \lambda_{12}^6)^2]$$

The coefficient of a term $\lambda_1^{b_1} \lambda_2^{b_2} \dots \lambda_{12}^{b_{12}}$ enumerates number of times the irreducible representation A_{1u} occurs in the equivalence classes of functions that contain b_1 colors of type 1, b_2 colors of type 2, ..., b_{12} colors of the type 12. Since A_{1u} representation has all -1 character values for the improper rotations this also corresponds to the number of chiral edge colorings of the cube with the given distribution of colors.

Table 1 shows all the cycle index polynomials of all irreducible representations for the O_h group for edge colorings. As seen from Table 1 another example of the GCCI would be that of the T_{1g} irreducible representation. The multinomial expansion is given as follows:

$$P_{O_h}^{T_{1g}} = \frac{1}{48} \left[3(\lambda_1 + \lambda_2 + \dots + \lambda_{12})^{12} - 6(\lambda_1 + \lambda_2 + \dots + \lambda_{12})^4 (\lambda_1^2 + \lambda_2^2 + \dots + \lambda_{12}^2)^4 \right. \\ \left. - 12(\lambda_1 + \lambda_2 + \dots + \lambda_{12})^2 (\lambda_1^2 + \lambda_2^2 + \dots + \lambda_{12}^2)^5 + 12(\lambda_1^4 + \lambda_2^4 + \dots + \lambda_{12}^4)^3 \right]$$

Again the coefficient of a typical term $\lambda_1^{b_1} \lambda_2^{b_2} \dots \lambda_{12}^{b_{12}}$ enumerates number of times the irreducible representation T_{1g} occurs in the equivalence classes of functions that contain b_1 colors of type 1, b_2 colors of type 2, ..., b_{12} colors of the type 12.

Table 2 shows our complete combinatorial results for the edge colorings of the octahedron or cube for all irreducible representations. As noted above there are many more terms in the above expansions than the ones displayed in Table 2, but we show only the unique terms as determined by the Young diagrams of the integer 12. For example,

the very first diagram represents 12 terms, viz., $\lambda_1^{12} \text{ or } \lambda_2^{12} \dots \lambda_{12}^{12}$. Likewise each

Young diagram represents multiple ordered partitions, and we have shown only one of them since coefficients are same for all of these. As seen from Table 2 some fascinating results are revealed about the cube for all irreducible representations of the O_h group.

First non-zero A_{1u} representation occurs for the partition $[10+2]$, which has one chiral

coloring. This would correspond to the term $\lambda_1^{10} \lambda_2^2$. Many other results are

combinatorially complex. An important result that enables us to verify the correctness of all numbers is that the sum of all numbers for each irreducible representation multiplied by the number of times that partition occurs for the irreducible representation in a given column in Table 2 is given by

$$GF^\chi = P_G^\chi (s_k \rightarrow 12)$$

For example, the above result for the A_{1u} representation gives

$$P_{O_h}^{A_{1u}} = \frac{1}{48} \left[(12^{12} - 3 \times 12^4 \times 12^4 + 2 \times 12^6 + 8 \times 12^4 - 8 \times 12^2) \right] = 185752092669$$

We have shown in Table 3 all such frequencies thus obtained for all the irreducible representations. The last few rows for the edge colorings according to the irreducible representation Γ can be obtained. For example, the last row is always given by

$$\frac{1}{48} [12! \times \dim(\Gamma)]$$

The one previous to the last row is given by

$$\frac{1}{48} [12! \times \dim(\Gamma) / 2]$$

The last but one row is given by

$$\frac{1}{48} [12! \times \dim(\Gamma) / 4]$$

The last minus third one row is given by

$$\frac{1}{48} [12! \times \dim(\Gamma) / 6]$$

These results follow from the fact that the multinomial expansions that contain the terms corresponding to last 4 partitions for edge colorings appear only in the lead term and thus we have the result.

Table 4 shows the vertex colorings of the cube and also the face colorings of octahedron. As seen from Table 4, first A_{1u} non-zero representation appears for the partition 4+4, and then 5+2+1 suggesting that to produce a chiral structure by vertex coloring of a cube, one needs at least 4 colors of one kind and 4 colors of another kind, a result that is well known. The last 2 rows of numbers and the first 2 rows of numbers are directly predictable. The first row is non-zero for only A_{1g} or totally symmetric representation. The last row is always given by

$$\frac{1}{48} [8! \times \dim(\Gamma)]$$

The one previous to the last row is given by

$$\frac{1}{48} [8! \times \dim(\Gamma) / 2]$$

While some of these results are well known for the A_{1g} representation in the context of isomer enumeration a full table for all irreducible representations for all partitions has not

appeared before in the literature, and thus our tables provide complete multinomial combinatorics of ligand partitions or edge partitions.

In order to complete the combinatorics of the octahedron and cube we have also considered the face colorings of the cube, which are equivalent to vertex colorings of the octahedron. Table 5 shows our complete results for this case for all irreducible representations. All of the results that we have obtained for the A_{1g} representation are well-known results for the vertex coloring isomers of the octahedron. The results for the A_{1u} representation are for the chiral isomers, while other results for all colorings that transform according to the given representation in the column.

The character cycle indices contain significantly richer and important information that could lead to powerful results in number theory and other branches of mathematics. The substitution of s_k by $\lambda_1^k + \lambda_2^k + \dots + \lambda_n^k$ is one of the possibilities suggested by Pólya. If we replaces s_k in GCCI by other symmetric functions [34] for example,

$$\lambda_1^k \lambda_2^k + \lambda_1^k \lambda_3^k + \dots + \lambda_1^k \lambda_n^k + \lambda_2^k \lambda_3^k + \dots + \lambda_2^k \lambda_n^k + \lambda_{n-1}^k \lambda_n^k = \sum \lambda_1^k \lambda_2^k$$

the generalized cycle indices produce new combinatorics for each irreducible representation. We may also consider the replacement by symmetric functions of the kind

$$\sum \lambda_1^k \lambda_2^k \lambda_3^k \text{ where the sum involves all possible symmetric combinations involving}$$

three terms at a time, and so on. In general it can be envisaged that the method could be generalized to a multinomial symmetric function with a multinomial expansion, which we call multinomial S-functions. Such topics can be studied separately.

Another special case of multinomial combinatorics with character theory is to choose weights $1, w, w^2, \dots, w^n$. Then we obtain powerful multinomial generators in w 's. If we stop with weights $1, w$, and w^2 this corresponds to filling the orbitals with zero, 1 or 2 electrons as by Pauli exclusion principle only at most 2 electrons are allowed [21]. In this case it is a special application to enumeration of electronic configurations that comply with the Pauli exclusion principle. In the general case, for example, the GF for the A_{2g} representation of the edge group of the cube for this case is given by

$$P_{O_h}^{A_{2g}} = \frac{1}{48} \left[(1 + w + w^2 + w^3 + \dots w^{12})^{12} - 3(1 + w + w^2 + w^3 + \dots w^{12})^4 (1 + w^2 + w^4 + w^6 + \dots w^{24})^4 + \right. \\ \left. + 2(1 + w^2 + w^4 + w^6 + \dots w^{24})^6 + 8(1 + w^3 + w^6 + w^9 + \dots w^{36})^4 - 8(1 + w^{12} + w^{24} + w^{36} + \dots w^{72})^2 \right]$$

Other variations that lead to infinite power series in multinomial weights are also feasible. There is much to be desired in such powerful multinomial combinatorial theory.

4. Conclusion

In this work we obtained the full combinatorial tables for all irreducible representations of the octahedral group for the vertex, edge and face colorings of the cube and octahedron using multinomial combinatorics. The tables provided complete enumeration numbers for all possible partitions in the young diagram for the colorings that transform according to the various irreducible representations of the group. As special cases, the chiral edge, vertex, and face colorings for all young diagrams were generated. The results for other irreducible representations were interpreted. We believe that there is rich combinatorics in GCCIs when replacements are made for each term by a multinomial of symmetric functions as opposed to an ordinary Pólya replacement. Such topics will be studied in the future, as they seem to result in novel mathematical identities in number theory.

5. Acknowledgment

This research was performed under the auspices of the US department of Energy by the University of California, LLNL under contract number W-7405-Eng-48 while the work at UC Davis was supported in part by US National Science Foundation grant CHE-0236434.

6. References

- [1] K. Balasubramanian, *Chemical Rev.* 85 (1985) 599.
- [2] K. Balasubramanian, *Theor. Chim. Acta.* 51(1979)37.
- [3] A. T. Balaban, “Chemical Applications of Graph Theory”, Academic Press (New York, NY 1976).

- [4] K. Balasubramanian, *Indian J. Chem.* 16B (1978) 1094.
- [5] K. Balasubramanian, *Theor. Chim. Acta* 53, (1979)129.
- [6] G. Pólya. - R.C. Read, *Combinatorial Enumeration of Groups, Graphs, and Chemical Compounds*, Springer (New York - Berlin, 1987).
- [7] G. Pólya. *Acta. Math.* 68 (1937)145.
- [8] J.H. Redfield, *Amer. J. Math.* 49 (1927)433.
- [9] K. Balasubramanian, *J. Math Chem.* 14 (1993)113.
- [10] R. W. Robinson, F. Harary and A. T. Balaban, *Tetrahedron* 32 (1976)355.
- [11] D. H. Rouvray, *Chem. Soc. Rev.* 3 (1974)355.
- [12] D. H. Rouvray, *Endeavour* 34 (1975) 28.
- [13] R. W. Robinson, *J. Comb. Theory* 9 (1970) 327.
- [14] F. Harary and E. M. Palmer, “Graphical Enumeration”, Academic press (New York, NY 1973)
- [15] A. T. Balaban, E. M. Palmer and F. Harary, *Rev. Roum. Chim.* 22 (1977) 517.
- [16] D. J. Klein and A. H. Cowley, *J. Am. Chem. Soc.* **100** (1978) 2593.
- [17] W. G. Klemperer, *J. Chem. Phys.* 56 (1972) 5478.
- [18] K. Balasubramanian, *Annals of New York Academy Sciences* 319(1979)33.
- [19] K. Balasubramanian, *Theor. Chim. Acta*, 59 (1981) 237.
- [20] K. Balasubramanian, *Journal of Chemical Phys.* 74 (1981) 6824.
- [21] K. Balasubramanian, *Int. J. Quant. Chem.* 20(1981)1255.
- [22] S. G. Williamson, *J. Comb. Theory* 8 (1970) 163.
- [23] S. G. Williamson, *J. Comb. Theory* 11 (1971) 122.
- [24] R. Merris, *Linear Algebra and Application*, 29 (1980)255.
- [25] R. B. King, *J. Math. Chem.* 1 (1987)15.
- [26] R. B. King, *J. Math. Chem.* 1 (1987) 55.
- [27] R. B. King and D. H. Rouvray, *Theor. Chim. Acta.* 69 (1986) 1.

- [29] R. B. King, in *Chemical Explanation: Characteristics, Development, Autonomy*, ed. Earley, J. E., *Ann. N. Y. Acad. Sci.*, 988 (2003)158.
- [30] S. Fujita, "Symmetry and Combinatorial Enumeration in Chemistry ",Springer-Verlag (New York 1991).
- [31] K. Balasubramanian, *Chemical Physics Letters* 183 (1991)292.
- [32] K. Balasubramanian, *Journal of Chemical Physics* 95 (1991)8273.
- [33] V. Krishnamurthy, "Combinatorics: Theory and Applications", Affiliated East west press (New Delhi, India 1985).
- [34] D. E. Littlewood, "Theory of Group Representation and Matrix Product of Groups", (Clarendon, Oxford 1940).

Table 1 The GCCIs table for the edge colorings of the cube

	1^{12}	$1^4 2^4$	2^6	2^6	$1^2 2^5$	$1^2 2^5$	4^3	4^3	3^4	6^2
	1	3	3	1	6	6	6	6	8	8
A_{1g}	1	1	1	1	1	1	1	1	1	1
A_{2g}	1	1	1	1	-1	-1	-1	-1	1	1
A_{2u}	1	-1	1	-1	1	-1	-1	1	1	-1
A_{1u}	1	-1	1	-1	-1	1	1	-1	1	-1
E_g	2	2	2	2	0	0	0	0	-1	-1
E_u	2	-2	2	-2	0	0	0	0	-1	1
T_{1g}	3	-1	-1	3	-1	-1	1	1	0	0
T_{2g}	3	-1	-1	3	1	1	-1	-1	0	0
T_{1u}	3	1	-1	-3	1	-1	1	-1	0	0
T_{2u}	3	1	-1	-3	-1	1	-1	1	0	0

Table 2 Octahedral Edge (Cube Edge)Combinatorics for All Irreducible Representations *

Partition	A_{1g}	A_{2g}	$A_{2u}(A_{1u})$	E_g	E_u	T_{1g}	T_{2g}	$T_{1u}=T_{2u}$
1	0	0	0	0	0	0	0
1	0	0	1	0	0	1	1
4	1	1	5	2	2	5	4
9	4	4	11	6	10	15	15
18	9	9	27	18	26	32	31
24	14	14	38	28	42	52	52
30	20	18	46	34	50	60	58
4	3	2	7	4	7	8	9
18	13	12	31	24	37	42	43
47.	42	38	89	76	118	123	127

92	82	78	174	156	238	248	252
126	116	110	242	220	336	346	352
76.	61	59	137	118	174	189	186
182.	162	158	344	316	478	498	502
318	288	282	606	564	842	872	868
372	342	336	714	672	1014	1044	1050
408.	388	376	790	746	1134	1154	1166
606	576	564	1182	1128	1704	1734	1746
768	720	711	1488	1422	2130	2172	2169

29	29	26	58	52	81	81	84
130	125	120	255	240	365	370	375
336	336	324	672	648	984	984	996
590	580	570	1170	1140	1720	1730	1740
702	702	684	1404	1368	2070	2070	2088
514	494	486	1008	972	1466	1486	1494
1178	1158	1142	2336	2284	3442	3462	3478
1764	1734	1716	3498	3432	5166	5196	5214
2328	2328	2292	4656	4584	6912	6912	6948
2922	2892	2868	5814	5736	8628	8658	8682

1792	1732	1718	3524	3436	5148	5208	5202
3522	3462	3438	6984	6876	10338	10398	10422
4422	4332	4308	8754	8616	12918	13008	13002
5838	5778	5742	11616	11484	17262	17322	17358
*	7740	7740	7668	15468	15324	23064	23064	23136
248	249	246	498	492	741	741	744
996	996	984	1992	1968	2964	2964	2976
2316	2316	2304	4632	4608	6924	6924	6936
3474	3474	3456	6948	6912	10386	10386	10404

3490	3470	3450	6960	6900	10370	10390	10410
6948	6948	6912	13896	13824	20772	20772	20808
8700	8670	8640	17370	17280	25950	25980	26010
11568	11568	11532	23136	23064	34632	34632	34668
10548	10398	10362	20856	20724	31122	31182	31218
17394	17334	17286	34728	34572	51906	51966	52014
23136	23136	23064	46272	46128	69264	69264	69336
26172	25992	25938	52164	51876	77808	77988	77982
34776	34656	34584	69432	69168	103824	103944	104016

1980	1980	1980	3960	3960	5940	5940	5940
6936	6936	6924	13872	13848	20784	20784	20796
13860	13860	13860	27720	27720	41580	41580	41580
17334	17334	17316	34668	34632	51966	51966	51984
20808	20808	20772	41616	41544	62352	62352	62388
34668	34668	34632	69336	69264	103932	103932	103968
46200	46200	46200	92400	92400	138600	138600	138600
52050	51990	51930	104040	103860	155850	155910	155970
69336	69336	69264	138672	138528	207864	207864	207936

104088	103968	103872	208056	207744	311712	311832	311928
156300	155940	155820	312240	311640	467460	467820	467820
13860	13860	13860	27720	27720	41580	41580	41580
41580	41580	41580	83160	83160	124740	124740	124740
69300	69300	69300	138600	138600	207900	207900	207900
103968	103968	103932	207936	207864	311832	311832	311868
138600	138600	138600	277200	277200	415800	415800	415800
207936	207936	207864	415872	415728	623664	623664	623736
312000	311880	311760	623880	623520	935400	935520	935640

83160	83160	83160	166320	166320	249480	249480	249480
207900	20790	207900	415800	415800	623700	623700	623700
277200	277200	277200	554400	554400	831600	831600	831600
415800	415800	415800	831600	831600	1247400	1247400	1247400
623736	623736	623664	1247472	1247328	1871064	1871064	1871136
415800	415800	415800	831600	831600	1247400	1247400	1247400
831600	831600	831600	1663200	1663200	2494800	2494800	2494800
1247400	1247400	1247400	2494800	2494800	3742200	3742200	3742200
1663200	1663200	1663200	3326400	3326400	4989600	4989600	4989600

Table 3 Frequencies of all irreducible representations in the edge combinatorics of octahedron(cube) shown in Table 2 for all multinomial terms^a

A_{1g}	185788177224
A_{2g}	185752217877
$A_{1u}=A_{2u}$	185752092669
E_g	371504434605
E_u	371504184051
T_{1g}	557256402066
T_{2g}	557256401934
$T_{1u}=T_{2u}$	557256029616

^a Sum of frequencies x dimension of the reps for all irreducible reps is verified to be 12^{12} .

Table 4 Cube Vertex(Octahedral face)Combinatorics for All Irreducible Representations^a

Partition	A _{1g}	A _{2g}	A _{2u}	A _{1u}	E _g	E _u	T _{1g}	T _{2g}	T _{1u}	T _{2u}
1	0	0	0	0	0	0	0	0	0
1	0	1	0	0	0	0	1	1	0
3	0	1	0	2	1	0	3	2	1
3	0	3	0	2	2	2	5	5	2
6	0	3	1	4	2	2	6	5	3
3	0	3	0	2	2	2	5	5	2
6	1	6	1	7	7	8	13	13	8
10	3	10	3	11	11	14	21	21	14
16	5	11	6	21	17	20	31	28	23
17	7	17	7	23	23	30	40	40	30
10	4	10	4	14	14	18	24	24	18
22	13	22	13	35	35	48	57	57	48
*	30	18	30	18	46	46	64	76	76	64
*	42	28	42	28	70	70	98	112	112	98

68	44	58	46	112	104	144	168	162	150
38	32	38	32	70	70	102	108	108	102
76	64	76	64	140	140	204	216	216	204
114	96	114	96	210	210	306	324	324	306
140	140	140	140	280	280	420	420	420	420
216	204	216	204	420	420	624	636	636	624
420	420	420	420	840	840	1260	1260	1260	1260
840	840	840	840	1680	1680	2520	2520	2520	2520

^aYoung diagrams with * must be conjugated to obtain the partition. These are shown in this way to save space.

Table 5 Octahedral Vertex (cube face) Combinatorics for All Irreducible Representations^a

Partition	A_{1g}	A_{2g}	A_{2u}	A_{1u}	E_g	E_u	T_{1g}	T_{2g}	T_{1u}	T_{2u}
	1	0	0	0	0	0	0	0	0	0
	1	0	0	0	1	0	0	0	1	0
	2	0	0	0	2	0	0	1	1	1
	2	1	1	0	2	0	0	1	2	1
	2	1	1	0	3	0	1	1	3	2
	3	2	2	0	5	1	2	3	5	4
	5	2	2	1	7	2	3	6	6	6
	4	4	4	1	8	2	6	6	9	9
	6	5	5	2	11	5	9	10	13	12
	9	9	9	6	18	12	21	21	24	24
	15	15	15	15	30	30	45	45	45	45

^aYoung diagrams with * must be conjugated to obtain the partition. These are shown in this way to save space.