Annual Progress Report 1991-1992

THEORETICAL STANDARDS IN X-RAY SPECTROSCOPIES

Univ. of Washington, Seattle

TABLE OF CONTENTS

ABSTRACT	1
I. PROJECT DESCRIPTION	2
II. RESULTS of CURRENT CONTRACT YEAR	3
A. Summary of Proposed Work	3
B. Multiple-scattering - Scattering Matrix Approach	3
C. Multiple-scattering - Separable Approximation	3
D. Other X-ray Spectroscopies	7
III. PROPOSED RESEARCH 2nd YEAR - <i>Removed</i>	9
IV. BIBLIOGRAPHY	14

ABSTRACT

We propose to extend our state-of-the-art, *ab initio* XAFS (X-ray absorption fine structure) codes, FEFF. Our current work has been highly successful in achieving accurate, user-friendly XAFS standards, exceeding the performance of both tabulated standards and other codes by a considerable margin. We now propose to add the capability to treat more complex materials. This includes multiple-scattering, polarization dependence, an approximate treatment of XANES (x-ray absorption near edge structure), and other improvements. We also plan to adapt FEFF to other spectroscopies, e.g., photoelectron diffraction (PD) and diffraction anomalous fine structure (DAFS).

MASTER

SEP 01 1992

PROJECT DESCRIPTION: THEORETICAL STANDARDS IN XAFS

The major accomplishment of our previous one-year grant was the successful development of accurate, reliable theoretical XAFS (X-ray Absorption Fine Structure) standards. These standards consist of *ab initio* codes as well as tables of XAFS phases and amplitudes for the entire periodic table. They are described in the following publications:^{1,2}

¹ Theoretical X-ray Absorption Fine Structure Standards, J.J. Rehr, J. Mustre de Leon, S.I. Zabinsky, and R.C. Albers, J. Am. Chem. Soc **113**, 5135 (1991).

² *Ab initio* curved-wave x-ray-absorption fine structure, J. Mustre de Leon, J.J. Rehr, S.I. Zabinsky and R.C. Albers, Physical Review B **44**, 4146 (1991).

These "FEFF" codes are in use world-wide and have had an immediate and substantial impact on XAFS analysis techniques. The main reason for the superiority of our codes to others, is that our theoretical model is more complete. Another reason is that the code is both fast and user-friendly, yielding reliable results for all users. Tests in a wide variety of materials^{1,2} and by independent researchers³ indicate that our theoretical XAFS standards are superior to existing tabulated standards and are comparable to experimental standards. They yield interatomic distances to an accuracy of order 0.02 Å or better and amplitudes and coordination numbers of order 20% or better. The name FEFF refers to the effective back-scattering amplitude $f_{eff}(\pi, k)$ that replaces the plane wave scattering amplitude $f(\pi, k)$ in the standard XAFS equation,

$$\chi_l(E) = - \sum N S(E) \frac{|f_{eff}(\pi, k)|}{k R^2} \sin(2kR + 2\delta_c + \Phi) e^{-2R/\lambda} e^{-2\sigma^2 k^2}. \quad (1)$$

Here $f_{eff}(\pi, k) = |f_{eff}(\pi, k)| e^{i\Phi}$ is the effective curved-wave scattering amplitude for a given m-leg path of total length 2R and degeneracy N, δ_c the l-wave central-atom phaseshift, λ the photoelectron mean-free path, σ the rms fluctuation in the half-path length R, and S(E) an amplitude reduction factor.

We have endeavored to make the FEFF codes and tables readily available to all interested scientists. Copies of the codes are available for DOE researchers from the Office of Scientific and Technical Information, or from users at the National Synchrotron Light Source at Brookhaven. Our codes are also licensed and distributed for general academic and corporate use by the University of Washington Office of Technology Transfer.

II. RESULTS of CURRENT BUDGET PERIOD

A. Summary of Proposed Work

The work of our current grant is aimed at extensions of our codes, in particular multiple-scattering and XANES, that are necessary to treat more complex materials. Though adequate for the first few coordination shells, a serious limitation of our original code was the restriction to single-scattering contributions in near neighbor environments. Thus many systems, indeed, perhaps half of the systems of current interest could not be addressed with our current standards, good as they have proved to be. The primary goal of the first year of our proposal has been to add multiple-scattering capabilities to the code, and we have succeeded in this effort. Our primary goal for the continuation of this grant in the second year is to add polarization dependence and an approximate treatment of XANES (x-ray absorption near edge structure). A second goal is to adapt FEFF to other spectroscopies, e.g., photoelectron diffraction (PD) and diffraction anomalous fine structure (DAFS). Although we have made some progress on 2nd year topics, the proposed work is essentially unchanged from our original proposal and it is repeated for completeness in Sec. III of this report.

B. Multiple Scattering - Scattering Matrix Approach

We have successfully implemented the scattering matrix approach of Rehr and Albers⁴ for fast accurate multiple scattering calculations. In this method

each multiple scattering contribution can be expressed in a form analogous to that with the plane-wave approximation, but with the usual scattering amplitude $f(\theta)$ replaced by low order "scattering-matrices," $F_{\lambda,\lambda'}$. The new method was an important breakthrough for the following reason: although the general multiple-scattering formalism for calculating XAFS is straightforward to write down, calculations heretofore have been time-consuming, with compute times growing as a large power of the maximum angular momentum $l_{max} \sim 20$. Attempts to speed up "exact" codes have been made by others;⁵ however, we found that such attempts are complicated to code, converge slowly at high energies, and are limited to paths of low order - triple scattering or less.

Our new multiple scattering code, FEFF (Version 5) executes several steps sequentially: i) the construction of an overlapped atom coulomb potential, scattering potentials, and partial wave phase shifts; ii) the calculation of all multiple scattering paths up to a maximum total path length; iii) the calculation of the effective scattering matrices $F_{\lambda,\lambda'}$, and iv) the calculation of the contribution to the XAFS spectrum $\chi(k)$ for each path. Notable in our code is a new approach for estimating multiple scattering Debye-Waller factors based on a new, multiple scattering correlated Debye-Model. Although most appropriate for homogeneous materials, the model takes local atomic masses into account and has been found useful as a first approximation in more complex materials. Also notable is an improved treatment of inelastic losses at low energies, based on an approximation of electron-hole excitations due to Quinn.⁶ An example of our results for Cu is given in Fig. 1 on the next page.

The example above has only a limited amount of multiple scattering. One of our goals, however, was to develop a computational method for multiple scattering which could compete with band-structure approaches. What was not clear was how many multiple-scattering paths were needed to obtain convergence. Metallic fcc Cu provides a severe test case: because of the close packed structure there

are are many thousands of important multiple scattering corrections. Indeed, we found that on the order of a million paths would be needed to approximate the band-structure results for Cu. To simplify the calculations we introduced a simple cutoff procedure based on the separable approximation, which eliminates all paths with amplitudes below a given value. In this way we were able to reduce the total number of paths needed for convergence to about 10,000. A manuscript describing our work is in preparation. A brief description will be given at the March 1992 Meeting of the American Physical Society. The abstract published

Cu 190K: Comparison of Data and Feff 5 (32 Paths)

Fig. 1. Contribution to the K-shell XAFS of Cu from 32 multiple scattering paths with a total path length less than $4 R_{nn}$, as calculated by the scattering matrix approach⁴ with (6×6) matrices (dotted lines) and all exact single-scattering paths up to the 6th shell, and for comparison, the results of experiment.

in Bull. Am. Phys. Soc. **37**, 475 (1992) is given below:

Ab initio XAFS Multiple-scattering Calculations

J.J. Rehr, S.I. Zabinsky and R.C. Albers

A fast, accurate multiple-scattering approach is implemented for calculations of x-ray absorption fine structure (XAFS) in molecules and solids. Our method is based on our *ab initio* single-scattering XAFS code FEFF, together with an automated path finder and the efficient scattering-matrix approach of Rehr and Albers. Converged results are obtained for each scattering path, usually with (6×6) matrices. Multiple-scattering Debye Waller factors are included based on the correlated Debye model. Results for fcc Cu are presented which are in excellent agreement with experiment for the first six shells with all 169 unique paths. However, more than 100,000 paths are needed for convergence to bandstructure. Criteria are discussed which significantly reduce the number of paths needed to achieve full multiple scattering accuracy.

C. Multiple Scattering - Separable Approximation

The lowest order approximation in our approach is the "separable approximation" for the electron propagator, and is equivalent to the point scattering approximation. In this approximation, the scalar (1×1) scattering matrices are simply effective curved wave scattering amplitudes $f(\theta, k, R)$ for general angles θ , and are as easy to calculate as curved wave back-scattering amplitudes. Thus as a first step, we proposed to implement the separable approximation with our single-scattering cluster code FEFF (Version 4). This approximation formed the basis for two Ph.D. theses completed during the past year: one was work of Dr. Brian Houser,⁷ (now at Central Washington Univ.) a former grad student of Prof. R. Ingalls at the University of Washington. A second thesis was completed by Dr. Gerd Pfeiffer⁸ (now at MPI Stuttgart), a former grad student of Prof.

D. Sayers, of N. Carolina St. University. We had speculated that the separable approximation might suffice for many applications. This would considerably simplify multiple-scattering calculations, and it was relatively easy to code. However, the results not quite as good as one might hope, with errors on the order of 10-30%. Moreover, the next level of approximation, namely low order scattering matrices yielded nearly exact results. Still the separable approximation can be advantageous: it is very fast and yields a reliable indicator of the importance of a given path, thereby permitting one to ignore a large number of unimportant paths. This work is described in a forthcoming paper in Physical Review B, in press (1992) - see abstract below.

The Separable Spherical Wave Approximation: Application to XAFS
Multiple Scattering in ReO_3

B. Houser, R. Ingalls and J.J. Rehr

Rehr and Albers have shown that the exact XAFS propagator may be expanded in a separable matrix form, and that the lowest order term in the expansion yields XAFS formulas containing spherical wave corrections, yet retaining the simplicity of the plane wave approximation. This *separable spherical wave approximation* was used to model the multiple scattering contributions to the XAFS spectrum of rhenium trioxide. We report a modest improvement over the plane wave approximation.

D. Photoelectron Diffraction, DAFS, and Other Spectroscopies

We had also proposed to extend our codes to other spectroscopies. Although not the primary focus of our work during the first year, we have made steady progress toward this goal. In Photoelectron Diffraction we have collaborated with Prof. F. Brown of the University of Washington and with Prof. C. Fadley⁹ of U. Hawaii in implementing the Rehr-Albers approach. Our work with Brown¹⁰ will shortly be published in Physical Review B, in press (1992), (see abstract below).

Photoelectron Diffraction Study of Ultrathin Films of AgBr on Ag(111)

A.-B. Yang, F.C. Brown, J.J. Rehr, M.G. Mason, and Y.T. Tan

Angle-resolved photoemission from the core levels of Br has been used to study the Stranski-Krastanov (SK) layer which precedes the growth of islands for molecular-beam epitaxy of AgBr onto Ag(111). Experimental photoelectron diffraction results compare favorably with a spherical wave theory of the differential cross section for photoemission. This theory uses a local cluster of atoms whose structure is consistent with LEED observations. A simple model is suggested which leads to the formation of (strained) ionic crystal islands. By making use of a large observed core-level shift, the island component of thick films can be separated from emission due to the thin SK layer. It is suggested that photoelectron diffraction for the island component would be especially interesting in connection with termination of the (111) ionic surface.

Our work on DAFS is being carried out in collaboration with Prof. L. Sorensen of the University of Washington and Dr. C. Bouldin of NIST. A manuscript is in preparation and will shortly be submitted to Physical Review Letters. The work will also be described at the March 1992 Meeting of the American Physical Society, in a paper, Simple Theory of Diffraction Anomalous Fine Structure (DAFS), by L.B. Sorensen, H. Stragier, J.O. Cross, J.J. Rehr, and C.E. Bouldin, Bull. Am. Phys. Soc, **37** 474 (1992).¹¹

Finally we have written for NATURE a News and Views article on the fine structure predicted by S. Koonin¹² to accompany β -emission. This new spectroscopy could be used to map Hydrogen in solids and may have implications for searches for heavy neutrinos with solid-state detectors. The reference¹³ is NATURE, **354**, 436 (1991).

IV. BIBLIOGRAPHY

1. J.J. Rehr, J. Mustre de Leon, S.I. Zabinsky, and R.C. Albers, J. Am. Chem. Soc. **113**, 5135 (1991).
2. J. Mustre de Leon, J.J. Rehr, S.I. Zabinsky, and R.C. Albers, Phys. Rev. B **44**, 4146 (1991).
3. M. Vaarkamp, E.A. Stern, N. Binsted, R.J. Oldman and D.C. Konigsberger, Eindhoven preprint February 1992.
4. J.J. Rehr and R.C. Albers, Phys. Rev. B **41**, 8139 (1990).
5. N. Binsted, S. J. Gurman and J.W. Cambell, EXCURV86 "SERC Daresbury Laboratory Program;" S.J. Gurman, N. Binsted and I. Ross, J. Phys. C **17**, 143 (1984).
6. J.J. Quinn, Phys. Rev. **126**, 1453 (1962).
7. Study of Bond Angle Determination by XAFS using the Separable Spherical Wave Approximation, B.D. Houser, Ph.D. Thesis, Univ. of Washington (1991); B. Houser, R. Ingalls and J.J. Rehr, Phys. Rev. B, in press (1992).
8. G. Pfeiffer, Ph.D. Thesis, N. Carolina State Univ. (1991).
9. Application of a Novel Multiple Scattering Approach to Photoelectron Diffraction and Auger Electron Diffraction, A.P. Kaduwela, Ph. D. Thesis, U. Hawaii (1991).
10. A.-B. Yang, F.C. Brown, J.J. Rehr, M.G. Mason, and Y.T. Tan, Phys. Rev. B, in press (1992).
11. L. Sorensen, C. Bouldin and J.J. Rehr, (manuscript in preparation. 1991).
12. S.E. Koonin, NATURE, **354**, 468 (1991).
13. J.J. Rehr, NATURE, **354**, 436 (1991).
14. P.J. Durham, J.B. Pendry, and C.H. Hodges, Comp. Phys. Comm. **25**, 193 (1982).
15. Ph. Saintavit, J. Petaiu, M. Benfatto, and C.R. Natoli Phys. Rev. B (to be published).
16. R.C. Albers, A.K. McMahan, and J.E. Müller, Phys. Rev. **31**, 3435 (1985); we are collaborating with R.C. Albers, and T. Redondo of Los Alamos National Laboratories to obtain these potentials.
17. J. Mustre de Leon, Ph.D. Thesis, Univ. of Washington 1989.
18. P.A. Lee and G. Beni, Phys. Rev. B **15**, 2862 (1977).

19. P. Horsch, W. von der Linden and W.-D. Lukas, Solid State Commun. **62**, 359 (1987).
20. See for example, D.B. Tran Thoai and W. Ekardt, Solid State Commun. **40**, 269 (1981); D. Lu and J.J. Rehr, Phys. Rev. B **37**, 6126 (1988); J.J. Rehr, E.A. Stern, R.L. Martin and E.R. Davidson, Phys. Rev. B **17** 560 (1978); S.-H. Chou, J.J. Rehr, E.A. Stern and E.R. Davidson, Phys. Rev. B **35**, 2604 (1987); L. Hedin, Physica B **158**, 344 (1989).
21. J.J. Rehr, R.C. Albers, C.R. Natoli, and E.A. Stern, Phys. Rev. B **34** 4350 (1986); J. Osterwalder, E.A. Stewart, D. Cyr, C.S. Fadley and J.J. Rehr, Phys. Rev. B **35**, 9859 (1987).
22. J. Mustre de Leon, J.J. Rehr, C.R. Natoli, C.S. Fadley, and J. Osterwalder, Phys. Rev. B **39**, 5632 (1989).
23. J.J. Barton and D.A. Shirley, Phys. Rev. B **32**, 1906 (1985).
24. See I. Arcon, A. Kodre, D. Glavic and M. Hribar, J. de Physique, Colloque C9, 1005 (1987), and references therein.
25. L. Sorensen and C. Bouldin (unpublished).
26. H. Stragier, J.O. Cross, J.J. Rehr, L.B. Sorensen, C.E. Bouldin, and J. C. Woicik, submitted to Physical Review Letters (1992).

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

END

DATE
FILMED

10 / 8 / 92