Copy 10 of ten

UGRL 18712

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

FINAL REPORT

Lawrence Livermore Laboratory

Subcontract 8273105

by

C. F. Hooper, Jr.

Professor of Physics

Principal Investigator

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

RECEIVED
JAN 10 1964

I. Introduction

This is the final report of work done under Lawrence Livermore Laboratory subcontract No. 8273105. As such it covers work already reported in the form of manuscript entitled Stark Broadening in Hot Dense Laser-Produced Plasmas, as well as that accomplished subsequently. A slightly revised version of that manuscript is attached as Appendix A; the paper itself is scheduled to appear in the October issue of Physical Review A.

There are 3 topics of primary importance in this report: (1) The calculation of electric-microfield-distribution functions for plasmas having ions and electrons at different kinetic temperatures, (2) The calculation of electric microfield distributions for one-temperature plasmas having two species of ions present in varying ratios, (3) The extension of the line-shape (profile) calculation to higher-series (Lyman) members.

In addition, brief mention will be made of 2 other problems: (4) The calculation of electric microfield distributions for very dense plasmas, and (5) The derivation of a new line-broadening theory appropriate for plasmas in which the component particles are strongly coupled and in which the electrons are at least partially degenerate.

II. Technical Results

(1) The first two-temperature microfield calculations have just been completed. Figures 1 through 4 show the results for two representative situations. Each of the graphs is labeled with the quantities $\underline{a}$, χ , Z_1 , R , and T_R . The quantity $\underline{a}$ is related to the plasma parameter, Γ :

$$\Gamma = \frac{\underline{a}^2}{3} ,$$

where,

$$\Gamma \equiv \frac{e^2/r_D}{kT}$$

$$r_D = (3/4\pi n_e)^{1/3} ;$$

n_e is the electron density, χ gives the net radiator charge and Z_1 is the charge of the perturbing ion. R is equal to zero in all these figures which implies that there is only one type of ion perturber (of charge Z_1). T_R is the ratio of electron to ion kinetic temperatures. Figures 1 and 2 refer to the field distribution at an argon XVIII ion in a fully stripped deuterium or hydrogen plasma. As T_R increases from 0.25 to 4.00, the distribution becomes more sharply peaked and shifts to smaller field strengths. As the plasma becomes more strongly coupled (as a increases) the effect of T_R variations is amplified. Figures 3 and 4 are similar to 1 and 2 except that they represent field distributions at an aluminum XIII radiator that is immersed in a plasma containing only Al XIII ions. The comments made about figures 1 and 2 also apply here. However, by overlaying figures 2 and 4, it can be seen that the Al XIII results are more sensitive to T_R variation. This is due to the fact that the correlation between radiator and perturbing ions ($\sim Z_1^2$) is greater in this case.

(2) Figures 5 and 6 give the field distributions at an Argon XVIII radiator immersed in a plasma containing both Argon XVIII ions and fully stripped hydrogen. Three values of R ($= N^{XVIII}/N^I$) are chosen. T_R is set equal to unity. The results in figure 5 are for $a = 0.2$ and in figure 6 for $a = 0.4$. Looking at figure 5, we see that the variation in R produces large changes in the field distribution. Furthermore, this change between $R = 0.0$ and 0.1 is surprisingly large whereas between $R = 0.1$ and ∞ it is relatively small. Continuing on to figure 6, we see that there is again a variation in $P(\epsilon)$ with R . Although the sensitivity changes somewhat, what is most striking is that the qualitative order of the curves is reversed. This behavior is due to the fact that the correlations between the ions depends on both R and a , but not with the same functional dependence. It should also be pointed out that for plasmas whose ion species have similar charges, the R sensitivity is virtually non-existent. Appendix B shows the functional forms involved in our calculations; these are the modifications necessary to convert the formalism in Appendix A to that appropriate for both two

temperatures and two ion-perturber species.

(3) Although no numerical calculations have been made for higher-series numbers, our programs are being modified so that we may do so. We expect to produce higher-series-number line shapes within the next month or so.

(4) We are currently calculating microfield distributions for strongly coupled plasmas, $\Gamma \gtrsim 1$. To assess the validity of these calculations we are comparing with Monte Carlo results being generated by H. E. DeWitt of LLL. Initial results are very promising.

(5) Our efforts to generate a line-broadening theory suitable for strongly coupled degenerate plasmas centers on extending a recently developed theory. See the two papers attached - appendices C and D. It is too early to gauge the final promise of this technique.

Copy to Bureau

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 6