THE RESPONSE OF ESTUARINE FISH EMBYROS
TO ENVIRONMENTAL TEMPERATURE SHOCK
Annual Progress Report - 1/14/75 through 1/13/76

NOTICE

PORTIONS OF THIS REPORT ARE ILLEGIBLE. It
has been reproduced from the best available
copy to permit the broadest possible avail-
ability.

John Mark Dean

Belle W. Baruch Institute for
Marine Biology and Coastal Research and Department of Biology
University of South Carolina
Columbia, South Carolina 29208

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Date Published--April 1, 1976

Prepared for the U. S. Energy Research and Development Administration Under
Contract AT(38-1)-869

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

SUMMARY

The results have demonstrated that the most sensitive stages to acute thermal shock were the 1-2 and 2-4 cell stages of periods of cell division. Late embryos and larvae were more tolerant of temperature and chemical stress than were juveniles or adults.

The level and duration of exposure was not critical at less than 40°C for 24°C acclimated fish.

Increased incubation temperatures shortened the time to hatching and fish reared at high but non-lethal temperatures had higher feeding rates than those reared below an apparently sensitive environmental level of 24°C.

The results of the exposure of different developmental stages to chlorine and temperature at different levels and for different durations are still undergoing analysis. Preliminary interpretation is possible of the 0 and 7 day larvae stage where the effects were obvious but more subtle interaction effects could occur at earlier developmental stages and not be apparent without the further statistical analysis that is underway.

INTRODUCTION

The objectives of this study were to determine the effect of acute temperature shock on; defined developmental stages of the mummichog, Fundulus heteroclitus; the period of starvation death; and feeding and growth of larvae; In addition, we measured the interaction of temperature shock/developmental stage and chlorine exposure and whether

the effect on hatching was a function of the acute exposure or the duration of exposure.

MATERIALS AND METHODS

Fish were collected in the creeks of North Inlet near Georgetown, South Carolina and We Creek near North Edisto Inlet, South Carolina. Males and females were injected with gonadotropin to induce spawning. The males and females received 50 units of the gonadotropin on two successive days and on the fourth day the eggs and sperm were stripped from the fish into an 8-inch fingerbowl containing filtered sea water and the eggs and sperm were mixed and stirred well immediately. During this process some of the eggs stuck to the bottom but gentle movement with the water or a finger removed them from the glass. Fifteen minutes after complete mixing, fertilization had taken place as evidenced by the formation of the chorionic membrane and perivitelline space. The fertilized eggs were placed in a hatching jar with a continuously circulating supply of filtered sea water at 24°C and 25 o/oo S. About 1,000 fertilized eggs were placed in the hatching jar and a similar procedure was followed to establish the beginning stock for each of the experiments described in the report.

In order to monitor the developmental stages, several of the eggs in the incubation jar were removed with a pipette and placed in a depression slide covered with a slide cover and their development was observed under a microscope. When the desired stage of development had been reached, eggs were removed from the incubation jar, placed in a fingerbowl, and the appropriate number of experimental embryos removed and

placed in exposure chambers.

The exposure chambers were placed in a temperature-controlled bath and the appropriate handling controls were placed in exposure chambers in a control temperature bath. After exposure, the exposure chambers and handling-controls were removed and placed in the incubation aquarium for the remainder of the period of the experiment.

Our experience has shown that after 13 days of development at 22-24°C Fundulus were ready to hatch and it was possible to induce hatching in the majority of the eggs by changing water conditions at that time. When we wanted to determine the percentage of hatching or mortality, the exposure chambers were removed at 13 days and placed in an aquarium that contained unfiltered sea water. Hatching usually occurred within a period of several hours. Twenty-four hours after hatching induction, the exposure chambers were removed and the number of viable larvae, the number of eggs that did not hatch, and the obviously dead eggs were counted.

The post hatch thermal sensitivity was determined by exposing groups of 15 fish to the experimental temperature for set periods of time (a dose of time/temperature) and replicates were done at each temperature. The TD_{50} was calculated by plotting the % mortality versus time on probit paper.

If we wanted to induce hatching to get a known age of Fundulus, it was only necessary to add several liters of unfiltered sea water to the circulating incubation jar system and, once again, within several hours virtually all of the Fundulus that were going to hatch had hatched.

Newly-hatched larvae (N=20) were placed in individual finger bowls, kept at different temperatures, and presented with known numbers of newly-hatched brine shrimp Artemia. Each morning the number of Artemia remaining in the finger bowl were counted and in this fashion the number of brine shrimp eaten the the individual fish larvae could be determined. On day 14, the oxygen consumption of the larval fish was measured and after the respiration study was completed the individuals were weighed and frozen for caloric determinations to be done in the future. These experiments were done at 2° increments between 18 and 30°C.

The temperature/developmental stage/chlorine interaction exposure was accomplished in the same basic fashion as the previous experiments with the exception that the chlorine exposure was done in a constantly flowing exposure chamber with chlorine introduced into the chamber. The end point for the developmental stage exposure was hatching and the end point for the larval stages was mortality measured 24 hours after exposure.

RESULTS

Developmental Exposures

Different developmental stages of Fundulus heteroclitus (mummichog) embryos were exposed to 40°C for 5 minutes (Figure 1, Table 1) and there were two distinct results. First, exposure of fertilized eggs 60 minutes after fertilization, just prior to the first cell division, resulted in an extremely high mortality and a high incidence of abnormal

development. Interestingly enough, exposure 90 minutes after fertilization, between the first and second cell division, resulted in about the same incidence of mortality as the 60 minute exposure but a much higher percentage of abnormal development. When the exposure occurred 2 hours after fertilization, the mortality effect was virtually eliminated, but there was still some abnormal development. Exposure at the onset of circulation showed no significant difference between controls and experimentals for mortality or development. Similar results were obtained when embryos were exposed 10 days after fertilization.

Embryonic Response to Acute Thermal Shock

Fish were stripped and eggs and sperm mixed to follow the early developmental effects of acute exposure for different periods of time at 40°C. Ten of the developing embryos were exposed to the acute temperature shock 90 minutes after fertilization, at the time when the majority of the embryos in the process of division from the 1 to 2 cell stage, for 4, 5, 6, 7, 8, 9, and 10 minutes with a handling control (Table 2).

One hour and forty-five minutes after fertilization, the 4-minute exposure group was obviously in the 2-cell stage but did not show good cell resolution. The 5-minute group was in the 2-cell stage, well defined, and resembled the controls. The 6-minute group was in the 2-cell stage, irregular in shape, and dark. The 7-minute group was in the 2-cell stage, but the cells were large, ballooned and appeared abnormal. One could not determine the individual cells in the 8-minute group, and a similar phenomenon was apparent in the 9-minute exposure group; that is, a loss of cell staging capability. The 10-minute exposure group appeared very abnormal. It looked like the cells were breaking down and

there did not appear to be good cell membrane morphology. The controls showed a very clean two cell stage with good membrane definition.

After 2-1/2 hours the control cells were all clean and in the 4-cell stage. The 4-minute exposure group appeared normal. The 5-minute exposure group showed most of the embryos in the 4-cell stage with one showing an abnormal development. The 6-minute group all had the 4-cell stage development but the development appeared disoriented. The 7-minute group showed a high level of variability, some multiple cells, some arrested cells, and some cells with abnormal shapes. The 8 and 9-minute groups had a similar appearance with abnormal types of cells, and some clumping. The 9 and 10-minute group also showed some beginning of reorganization after the loss of staging observed in the 1 hour and the 45 minute period. However, they still had very poor geometry.

After 3-1/2 hours the controls were at the 16-cell stage and had excellent geometry. The cells were uniform in size and shape. The 4-minute exposure group appeared similar. The 5-minute group had half of the embryos showing some abnormal types of development, and a similar observation could be made of the 6-minute exposure group. The 7-minute exposure had some embryos that appeared as the controls but there was a great deal of variability. The 8-minute group showed a few 16-cells, but there were a large number of the cells with strange shapes and geometry. The 9-minute group had a great deal of clumping and aberrant cell shapes and spatial relationships. The 10-minute group showed abnormal clumping, geometry, development, and cell sizes.

Four hours and 45 minutes after fertilization, the controls were in the 32-64-cell stage of development and showed excellent geometry.

The 4-minute exposure group could not be differentiated from the control.

The 5-minute exposure group appeared predominantly as did the controls and a few abnormal cells continued development. The 6 minute exposure

group appeared similar to the controls with a few embryos showing

a double layering rather than the single layer. The 7-minute group

was highly variable with different sized cells and arrangements

of cells. The 8-minute group had large numbers of abnormally developing

cells. The 9-minute group showed double groups of cells with clumping

and a large number of different sizes of cells. In the 10-minute group,

virtually all the embryos appeared abnormal, but they were still

dividing and progressing.

The final results as measured by hatching, of that experiment were:

the control had 90% hatch success; the 4-minute exposure was 82%; the

5-minute exposure was 83%; the 6-minute exposure was 71%; the 7-minute

exposure was 62%; the 8-minute exposure was 78%; and 10-minute was

63% (Table 3). The 9-minute exposure group was lost during the course

of the experiment. Thus, it would appear that the abnormal develop-

mental stages observed during the early periods of embryonic development

were carried through and expressed as mortality in the hatchling.

Observations would indicate that if the exposures were done between

the 1-cell to 2-cell and 2-cell to 4-cell divisions that the incidence

of abnormality increased and almost equaled the mortality expressed in

the total group.

Duration of Exposure

The duration of exposure experiment confirmed the findings of the acute exposure experiment; that is, the early cell division embryonic stages were most sensitive to acute temperature shock and that sensitivity was expressed as soon as 4 minutes after the beginning of the acute exposure period (Table 2). If exposure did not occur until much later in development, there was no significant difference between the experimental groups and the controls.

During the course of this experiment it was apparent that the later stages of embryonic development had a lower thermal sensitivity.

Post-Hatch Thermal Sensitivity

We had observed the variation in sensitivity to thermal shock in Fundulus larvae before, and wanted to test it more definitively. It is not an easy experiment to do as the larval fish are difficult to handle and, as has become apparent through this work, the age of the larvae can be a significant factor. The results were quite dramatic (Table 4). The least sensitive fish to thermal shock were the youngest larvae and the most sensitive were the adults of the species. All the fish demonstrated a graded response to temperature. At temperatures of less than 40°C the response total was measured over a period of days in the larvae and juveniles.

Starvation Death

Starvation death (SD) has been utilized in other studies to measure the effects of embryonic stress. In this study SD was found to be temperature dependent, that is the higher rearing temperature the faster the time to

starvation death. However, SD times within experimental groups was so variable that it was not feasible to attempt using it as a definitive measure of the effect of the embryonic shock, and the results obtained were ecologically meaningful. All the fish that hatched when presented with food and the opportunity to feed did so no matter what their embryonic exposure.

Feeding Rate

We did, however, observe rather dramatic changes associated with transformation from the yolk-sac larval stage to the resorbed yolk-sac post-larval stage. This time period is also temperature dependent. At normal developmental temperatures, that is about 24°C, resorption occurs at about 6-7 days.

The results showed that there was a real change in feeding activity associated with the transformation from utilization of the yolk sac to absolute dependence on external food sources (Figures 2-8). This response is more obvious and exaggerated at higher incubation temperatures as demonstrated by a comparison to 22° with 26°. The total feeding activity is also temperature dependent. There is a significant difference between the activity of groups above and below 24°C (Figure 9). The change in feeding activity at 6-7 days was identified as a critical development stage and utilized in other sensitivity experiments.

Temperature/Developmental Stage/Chlorine Interaction

During the course of the experiments it was apparent that the later stages of development had a lower thermal sensitivity than the early stages. The opportunity existed to test the interaction of temperature,

developmental stage and chemical stress. This was done in cooperation with Doug Middaugh of the EPA Bears Bluff Field Station, South Carolina. Previous experiments had demonstrated sensitive stages of early cell division and closure of the blastopore to the gastrula stage and, as less sensitive periods, the onset of circulation and exposure immediately post-hatch. It was decided to test the interaction of temperature and chlorine during these specific developmental stages.

The experiment involved 6 developmental stages of Fundulus (1-2 cell, gastrulation, onset of circulation, 10-day embryo, 0-day post-hatch larvae, and 7-day post-hatch larvae). The levels of chlorine exposure, measured as total residual chlorine, were 0.05 ppm, 0.1 ppm, 0.5 ppm and 1.0 ppm. The exposure temperatures for each of the developmental stages at each of the chlorine levels were 24°C (ambient), 29°C (+5° ΔT), and 34°C (+10°ΔT). Thirty defined stage embryos or larvae were placed in 2 sets of exposure cups to provide a replicate and exposed for a duration of 7.5, 15, 30 and 60 minutes at all treatments.

The results showed that exposures to chlorine at high levels and for longer periods of time were most toxic to later embryos and larvae rather than earlier stages and there was significant variation in the sensitivity of the larvae. That is, the 7-day old larvae showed greater sensitivity at all levels and periods than did the 0-day larvae. This is consistent with the previous developmental data.

When embryos were exposed at 24°C, the response appeared to be exclusively due to the chlorine effect (Figure 10). A lethal effect was observed at 1 ppm exposure for 15 minutes with 7-day larvae (1/15/7) and that was

the first combination of factors that demonstrated an effect. During the 30-minute exposure, the 1/30/7 was the only group that showed a response but the 1/60/0 (0-day larvae) and the 1/60/0 and the 1/60/7 were affected.

The groups exposed at 29°C gave similar but graded results. The 1/7.5/7 was the first group to respond followed by 1/15/7 and 1/30/0 and 1/30/7. The 60-minute exposure at 29°C was the first to show a response at the 0.5 o/oo chlorine level (0.5/60/0 and 0.5/60/7) as well as 1/60/0 and 1/60/7.

The exposure at 34°C, 10° above the ambient temperature for the fish gave dramatic results. The 1/7.5/7 and 1/7.5/0 showed a distinct response as did the 1/15/7 and 1/15/0. The 1/30/7 and 1/30/0 had almost total mortality and once again 0.5/30/7 and 0.5/30/0 showed a definite response. The 0.5/60/7 and 0.5/60/0 were markedly affected and consistent with the results obtained at 1/60/7 and 1/60/0. It appeared that there might be an effect on the 1/60/10-day old embryo, but the results were not as striking as the others.

The results are summarized in Figure 10 and showed an increase in percentage mortality with increased periods of exposure at the higher temperatures and higher chlorine levels during increased durations. In addition, the newly hatched larvae were most sensitive, not the embryos, and, as far as chlorine toxicity was concerned, it was not the younger developmental stages but rather the older ones that showed the most obvious response.

A sensitive statistical analysis of this data is under way and the preliminary results indicate a temperature effect can be measured on the developmental stages. However, it is in the 0 and 7 day larvae that all the interactions became significant and the synergistic results are definitive.

DISCUSSION

The results are being prepared for submission to the open literature. There is a very real effect of exposure of early developmental stages to thermal shock. I have compared these results with Schubel's published results, and discussed them with Schubel. I think the fundamental difference between our results is the fact that most of his exposures were not done at the sensitive developmental stages of the fish he has tested. This is understandable because of the difficulty in handling the early stages of the experimental fish he has used. If you accept that, both our studies are consistent.

Of course, what we now can see is that it is critical to know where the embryos are distributed in time and space relative to the possibility of exposure to thermal stress. Thus the ecology of the beast becomes critical. Will there be the possibility of the embryos being entrained in a power station? And, if so, these data suggest there is a relatively short period when that would have a large negative effect. On the other hand, the post-larvae and early juveniles seem to be more vulnerable to the possibility of entrainment and impingement and there is a longer period when exposure could be detrimental.

Once again, a consideration of the temporal aspects of fish larvae distribution becomes critical. Many of the important coastal species of fish are spawned in near coastal waters, and migrate into estuaries in the winter. There is an increased demand placed on nearshore and estuarine waters for utilization by systems that could provide time and space opportunities for stress exposures, such as floating nuclear plants, OCS development and deep water terminals.

It would be desirable to know stress response relationships on a species specific basis. Since this is unlikely, it will at least be necessary to identify key group characteristics that will permit some generalizations to be developed. I think this might be possible. For example, in the southeastern estuaries, croakers, mullet, spot, menhaden and some flounder are important cohabitant species that occupy the estuaries as larvae in the winter. In contrast, larvae of silverside, mummichogs, anchovies and some of the sea bass are dominant in the summer during equivalent developmental stages. By determining the stress response of various developmental stages of representatives of each group, some guidelines could be developed. Some of these groups are very difficult to use experimentally and it might be necessary to make assumptions about their specific responses. However, even if they can't be used experimentally, it will be necessary to know their temporal distribution in the estuary.

I would suggest that there are several existing problems with broad implications that are suggested by this work.

It is important to know where the developmental stages of the fish are in time and space.

It is important to know whether the cohabitant fish respond in a similar fashion to thermal stress, and, whether generalizations be made for winter and summer cohabitants.

It is important to know whether the interactions for temperature and chlorine and stage of development are generalized responses for fish that could be subjected to that stress.

Table 1. Results of Exposure of Eggs for 5 Min. @ 40°C.

Time of Exposure After Fertilization	60 Min.		90 Min.		120 Min.		Onset Of Circulation		10 Days	
	Control	Exp.	Control	Exp.	Control	Exp.	Control	Exp.	Control	Exp.
Number of Eggs	689	318	421	191	453	209	409	209	420	161
Abnormal Development Number (%)	12(.1)	31(10)	8(2)	40(21)	3(.6)	26(12)	8(2)	4(2)	0	3(2)
Hatch Success (%)	404(59)	72(23)	324(81)	44(23)	277(61)	155(74)	238(59)	138(66)	354(60)	107(66)

Hatching success and abnormal development of Fundulus heteroclitus after exposure to 40°C
for 5 minutes at defined developmental stages

Table 2. Response of 1st cell embryos to acute thermal shock.

	Time After Fertilization				% Hatch
	1 hr 45 min	2 1/2 hrs	3 1/2 hrs	4 hrs 45 min	
4	+++	+++	+++	++++	82
5	++++	+++	+++	+++	83
6	++	++	++	++	71
7	++	+	+	+	62
8	-	-	+	+	78
9	-	-	-	-	(last group)
10	--	--	-	--	63
Controls	++++	++++	++++	++++	90

Subjective evaluation scale

- ++++ all appear normal
- +++ a few abnormal
- ++ about 1/2 normal
- + about 1/3 normal
- no normal
- great variation in appearance of cells

Table 3. Results of Exposure for Extended Periods of Time @ 40°C.

Results are percent of shocked embryos hatching.

Period of Exposure In Minutes	Control	4	5	6	7	8	10	15	16	30	60
1 Cell/2 Cells - % hatch (N)	90 33	82 29	83 30	71 14	62 34	78 29	63 46				
Onset of Circulation (N)	59 30		66 30				58 30	57 30		57 30	65 30
10 Days Past Fertilization (N)	60 30		66 30				59 30	74 30		59 30	66 30

Table 4. Thermal Shock Post Hatch (LD₅₀) Results are based on dose exposures.

	<u>39°C</u>	<u>40°C</u>	<u>40.5°C</u>	<u>41°C</u>	<u>41.5°C</u>
Day of Hatch			240 Min.	790 Min.	
2 Day Old Fish			150 Min.	45 Min.	7.5 Min.
4 Day Old Fish		5 Hours	60 Min.	< 20 Min.	
6 Day Old Fish		>5 Hours	< 30 Min.	20 Min.	7.0 Min.
Adult	9 Min.	7.25 Min.		4.5 Min.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

*:0

Figure 8

Figure 9

(1.0 ppm TRC/15 min exposure/7 day larvae)

concentration of chlorine (TRC) in
ppm/duration of exposure/developmental stages