DOE/ER/53275--4

DE92 040891

Department of Physics
University of California, San Diego
La Jolla, CA 92093

Annual Summary of Progress
November 1991 - November 1992

"Investigations in Anomalous Transport and Ignition
Physics

DOE DE-FG03-88ER53275

Principal Investigators:

Patrick H. Diamond
Marshall N. Rosenbluth

September 1992

MASTER

THIS DOCUMENT IS UNCLASSIFIED

PROGRESS REPORT FOR DOE CONTRACT DE-FG03-88ER53275

In the 1991-1992 funding period, the major topics of emphasis in the U.C.S.D. research program were Drift Wave Turbulence Theory, the $L \rightarrow H$ Transition, Active Control, miscellaneous topics related to magnetic fluctuations and Dynamo Theory, and energetic particle driven Alfvén waves.

In Drift Wave Turbulence Theory, significant progress was made in understanding disparate scale interactions. In particular, a two-paper series on a two nonlinearity model was produced. These uncovered a blocking of the oft-invoked "inverse cascade" and demonstrated the importance of nonlocal transfer in k space due to $E \times B$ nonlinearity. Work on coherent structures helped elucidate their relevance to anomalous transport. In particular, studies of a simple 2-D Hasegawa-Wakatani model revealed that in the adiabatic regime structures form, but have little or no impact on transport predictions, ala quasilinear theory. However in the hydrodynamic regime, transport is significantly lower than either the prediction of quasilinear theory or standard resonance broadened quasilinear theory. This reduction in transport may be due to trapping in structures. We hope to elucidate this matter further by comparing the results of a precise direct interaction approximation code with the exact numerical simulations to isolate whether the failure is in resonance broadening theory in particular or in weak coupling random phase approximation, in general. Recent work has focused on the nonlinear dynamics of convective cells

(trapped ion modes). In particular it has been demonstrated that decay instability, driven by short wavelength drift wave turbulence, may energize trapped ion modes more rapidly than linear growth processes. Also, a type of two-dimensional structure which is formed naturally by the Reynold's stresses of the drift waves acting on the long wavelength cells has been identified. This tends to trap the long wavelength cells between bipolar structures of the drift wave envelope.

Major progress occurred in the past year on the theory of flow generation and the $L \rightarrow H$ transition. A general theory which encompasses both dynamo-like instabilities and direct generation of shear and curvature was developed. This theory was demonstrated to include previous results by us and the Maryland group. It was also well established that flow curvature (not shear!) is the dominant stabilizing mechanism, in agreement with recent experimental observations from DIII-D and JET. A self-consistent model of the flow turbulence evolution in a simplified context was developed and shown to demonstrate bifurcation-like phenomena. Furthermore, the ongoing research on flow generation has uncovered an important relation between vorticity transport and particle transport.

Another topic of research was in the area of active control. In particular, the theory of active control of plasma rotation by Ion Bernstein wave injection was developed. The competing roles of Reynold's stress flow

redistribution and direct momentum input due to acceleration of ions by high frequency waves was elucidated. Ongoing work is focused on extending this quasilinear calculation to the highly nonlinear edge. Also, an extensive study of helicity injection by Alfvén waves in a high temperature plasma was carried out. In particular, it was demonstrated that the efficiency of kinetic Alfvén wave helicity injection is significantly lower than that predicted by MHD calculations.

Another topic of research concerning magnetic fluctuation dynamics and dynamo theory. In the realm of dynamo theory, a simple analytical physical interpretation of the interesting result of Vainshtein and collaborators that magnetic diffusion is suppressed in 2-D MHD was developed. Extensions to 3-D reduced MHD and 3-D MHD are in progress. Also, a more complete and inclusive kinetic theory of the effects of high frequency microturbulence on low frequency dissipative instabilities was developed. A paper on this topic is in preparation at this time.

In the past year significant progress on the theory of toroidal Alfvén eigen modes was occurred. In particular, a precise calculation of the linear damping of toroidal Alfvén eigen-modes by coupling to the continuum occurred and was published in Physical Review Letters. Extensions which include FLR effects are in progress.

In the coming year, this group expects to focus on the development and dynamics of the $L \rightarrow H$ bifurcation, both with

and without ion orbit loss, the theory of Flow Generation and Flow Profile Modification in Toroidal Geometry, the nonlinear theory of Flow Generation by Ion Bernstein Waves, the development of a more accurate resonance broadened quasilinear theory (which reflects potential vorticity conservation constraints in transport predictions), comparisons of structure dynamics with closure code predictions, and further work on toroidal Alfvén wave eigenmodes, including kinetic corrections.

END

**DATE
FILMED**

11 / 23 / 92