DOE/ER/60673-T5

Neutron Scatter Studies of Chromatin Structures
Related to Functions

DOE/ER/60673--T5

DE93 002520

DE-FG03-88ER60673

E. Morton Bradbury, Principal Investigator

Technical Progress Report
for the period from November 1, 1991 to May 15, 1992

Summary

In our original application we proposed to study using neutron scattering the structure of chromatin, DNA/protein complex such as transcription factor TFIIIA and its DNA binding site, the effect of VLR (Very Lysine Rich) histones H1/H5 on chromatin and the effect of histone modification on the structure of chromatin. Our primary source of neutron for the scattering has been the Manuel Lujan Neutron Scattering Center at Los Alamos National Laboratory. This facility has been shut down due to safety problems at LAMPF since August 1991. It is scheduled to be on line June 1992. We have been granted 3 days of neutron beam time using the LQD during this cycle to study the structure of TFIIIA/DNA complex under different ionic strengths. Our alternate neutron source at Institut Laue-Langevin, Grenoble, France has been down since April 1991. It is not expected to be operational until at least 1994. A new 30m SANS equipment at the Cold Neutron Research Facility (CNRF) of National Institute of Standard and Technology (NIST) in Gaithersburg, Maryland has been on line since March 1992. We have done preliminary measurements of chromatin core particles and reconstituted mononucleosomes at NIST. The configuration of the current equipment is suitable for R_g (radius of gyration) measurements. We plan to use the allocated 3 days of neutron beam time at NIST to study the effect of VLR histone on the path of DNA extended beyond the nucleosome. Another new 30m SANS station with a wider range of momentum transfer Q will be available in September 1992. This new equipment will be more suitable in calculating the shapes and sizes of the DNA and protein components in chromatin containing fully acetylated histones as well as other chromatin variables such as histone phosphorylation and histone subtypes.

Despite of setbacks in the lack of neutrons for the proposed experiments, we have made considerable progress in chromatin reconstitution with the VLR histone H1/H5 (Appendix 1) and in understanding the dynamics of nucleosomes (Appendix 2 and 4). A ferromagnetic fluid was developed to align biological molecules for structural studies using small-angle-neutron-scattering (Appendix 3). We have also identified and characterized an intrinsically bent DNA region flanking the RNA polymerase I binding site of the ribosomal RNA gene in *Physarum Polycephalum*. Finally projects in progress are in the areas of studying the interactions of histone H4 amino-terminus peptide 1-23 and acetylated 1-23 peptide with DNA using thermal denaturation; study of GGAAT repeats found in human centromeres using high resolution Nuclear Magnetic Resonance and nuclease sensitivity assay; and the role of histones and other sperm specific proteins with sperm chromatin.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

Progress report

Chromatosome positioning on assembled long chromatin. (Appendix 1):

Using tandemly repeated 172 bp (172 x 45) and 207 bp (207 x 18) DNA from the 5S rDNA nucleosome positioning sequence, linker histones H1 or H5 was assembled on the reconstituted chromatin and subjected to detailed analysis by a combination of restriction endonucleases. In the absence of linker histone, nucleosomes were found to be positioned predominantly in the major site as previously reported by Simpson. We have noticed however there are a group of minor sites 10 to 15 bp away from the major site. Addition of linker histones did not alter the underlying DNA signals that induce the very regular spacing of the sequence.

Mobility of positioned nucleosomes on 5S rDNA. (Appendix 2):

Characterization of nucleosomes reconstituted onto the 5S rDNA sequence using 2-D nucleoprotein gel electrophoresis clearly indicated the nucleosome positioning to be a dynamic process. The histone octamers mobility was temperature dependent and was reversibly inhibited by Mg^{2+} .

Intrinsically bent DNA flanks both sides of a RNA polymerase I transcription start site.

We have identified two intrinsically bent DNA regions which flank the transcription start site of the rDNA gene from *Physarum polycephalum*. These sequences were isolated and analysed by a circular permutation polyacrylamide gel electrophoresis assay. The two sequences differ from each other in regard to their bending behaviors to temperature, degree of bend as well as sensitivities to varying levels of ethidium bromide. The location of these strongly bent regions on either side of a RNA polymerase I transcription start site suggests important roles for such structures in chromatin structure and transcription initiation.

Orienting rigid and flexible biological assemblies in ferrofluids for small-angle neutron scattering studies. (Appendix 3)

Small-angle scattering from macromolecules in solution is widely used to study their structures, but the information content is often limited because the molecules are generally randomly oriented and hence the data are spherically averaged. The use of oriented rodlike structures for scattering, as in fiber diffraction, greatly increases the amount of structural detail that can be obtained. A new technique using a ferromagnetic fluid has been developed to align elongated structures independent of their intrinsic magnetic properties. This technique is ideal for small-angle neutron scattering because the scattering from the ferrofluid particles can be reduced significantly by matching the neutron scattering length density of the particle to a D_2O solvent ("contrast matching"). The net result is scattering primarily from the ordered biological assembly in a solution environment that can be adjusted to physiological pH and ionic strength. Scattering results from ordered tobacco mosaic virus, tobacco rattle virus, and chromatin fibers were used for this study.

Mobile Nucleosomes - A general behavior. (Appendix 4)

We have previously reported a mobility of positioned nucleosomes on sea urchin 5S rDNA. In this study we demonstrate the temperature dependence and the range of this mobility on 5S rDNA constructs. We find that this dynamic behavior also applies to bulk mononucleosomes and nucleosomes reconstituted onto sequences of the Alu family of ubiquitous repeats. We conclude that short range sliding is potentially a general phenomenon that is dependent on the underlying sequence and its position on the histone octamer. The nucleoprotein gel analysis also reveals the dramatic effect on gel electrophoretic migration caused by the location of the histone octamer on DNA fragments. The usefulness of this technique for studying nucleosome positioning and its dynamics is demonstrated.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Experiments in progress:

Structure of TFIIIA/DNA complex at different ionic strengths using small angle neutron scattering.

3 days of neutron beam time has been granted for this study at the Manuel Lujan Neutron Scattering Center at Los Alamos National Laboratory on August 1992.

Structure of reconstituted nucleosome containing H1/H5 using neutron scattering.

3 days of neutron beam time has been allocated for this study at NIST using the 30m SANS-NG7 equipment.

Structure of fully acetylated nucleosome using neutron scattering.

A proposal for beam allocation for the new 30m SANS-NG3 has been submitted for a detail study of the fully acetylated core particle.

Structure of a human centromeric repeat GGAAT using high resolution NMR and nuclease sensitivity assay.

We have identified a loop structure containing non Watson-Crick basepairing in a human centromeric repeating motif GGAAT using NMR and nuclease sensitivity studies.

Identification of a 19K protein associated with sperm heads.

An abundant 19K protein associated with sperm heads has been identified to belong to a family of seminal plasma inhibins. Inhibins are specifically processed from a 92K precursor protein known as semonogelin. This specific 19K peptide is uniquely associated with the sperm heads and not associated with the seminal fluids. The localization and potential function of this peptide is being investigated.

Publications from 11/01/91 to 5/14/ 1992 funded directly from the grant:

1. Meersseman, G., Pennings, S. and Bradbury, E.M. (1991) Chromatosome Positioning on Assembled Long Chromatin, Linker Histones Affect Nucleosome Placement on 5S rDNA. *J. Mol. Biol.* **220**, 89-100. (Appendix 1)
2. Pennings, S., Meersseman, G. and Bradbury, E.M. (1991) Mobility of Positioned Nucleosomes on 5S rDNA. *J. Mol. Biol.* **220**, 101-110. (Appendix 2)
3. Schroth, G.P., Siino, J.S., Cooney, C.A., Th'ng, J.P.H., Ho, P.S. and Bradbury, E.M. (1992) Intrinsically Bent DNA Flanks Both Sides of a RNA Polymerase I Transcription Start Site. *J. Biol. Chem.* **267** (in press).
4. Meersseman, G., Pennings, S. and Bradbury, E.M. (1992) Mobile Nucleosomes - A General Behavior. *EMBO* (in press) (Appendix 4).
5. Sosnick, T., Charles, S., Stubbs, G., Yau, P., Bradbury, E.M., Timmins, P. and Trehwella, J. (1991) Orienting Rigid and Flexible Biological Assemblies in Ferrofluid for Small-Angle Neutron Scattering Studies. *Biophysical J.* **60**, 1178-1189. (Appendix 3)

Justifications for Equipment Purchase:

Equipment requested: Pharmacia Oligonucleotide Synthesizer (Gene Assembler Special).
List Price: 19,500

Justifications: This oligonucleotide synthesizer will be used mainly to produce deuterated oligonucleotides for use with NMR and neutron scattering studies. It will also be used to generate nondeuterated primers and DNA sequences in the study of chromatin reconstitution, nucleosome positioning and the structure of oligonucleosomes.

END

**DATE
FILMED**

12 / 11 / 92