NOTICE

**CERTAIN DATA
CONTAINED IN THIS
DOCUMENT MAY BE
DIFFICULT TO READ
IN MICROFICHE
PRODUCTS.**

PROGRESS REPORT

UCLA

DOE GRANT No. DE-FG03-90ER20011.

"Carbon dioxide and the stomatal control of water balance and photosynthesis in higher plants"

Funding period: July 1st, 1990 to June 30, 1992

A. Publications.

I. BOOKS.

Taiz, L and Zeiger, E. 1991. Plant Physiology. Benjamin Cummings, Menlo Park, CA, 559 pp.

II. FULL PAPERS.

1. Mawson, B.T., Zeiger, E. Blue-light modulation of chlorophyll a fluorescence transients in guard cell chloroplasts. Plant Physiol., in press.
2. Cornish, K., Radin, J.W., Turcotte, E.L., Lu, Z., Zeiger, E. Enhanced photosynthesis and stomatal conductance of Pima cotton (Gossypium barbadense) bred for increased yield. Plant Physiol., in press.
3. Hercovitz, S., Tallman, G., Zeiger, E. Long-term survival of Vicia guard cell protoplasts in culture. Submitted to Plant Science
4. Karlsson, P-E., Bogomolni, R.A., Zeiger, E. High performance liquid chromatography analysis of pigments present in guard cell protoplasts and mesophyll tissue from Vicia faba. Submitted to Photochem. Photobiol.

III. ABSTRACTS

5. Talbott, L.D. and Zeiger, E. 1991. HPLC analysis of Vicia guard cells indicates that products from photosynthetic carbon fixation and starch hydrolysis have an osmotic role during stomatal opening under blue and red light. Plant Physiol. 96S, 119.
6. Zeiger, E. and Srivastava, A. 1991. Fast fluorescence quenching from isolated guard cells chloroplasts from Vicia faba is induced by blue and not by red light. Plant Physiol. 96S, 42.
7. Srivastava, A. and Zeiger, E. 1991. Guard cells from Pima cotton (Gossypium barbadense) lines bred for heat tolerance have enhanced rate of respiration and proton pumping. Plant Physiol. 96S, 30.

B. SCIENTIFIC PROGRESS.

I. Instrumentation and methodology Efforts on this granting period (July, 1990 to date) have primarily focused on the development of an experimental approach capable of providing high resolution information on the modulation of osmoregulation in guard cells in response to CO₂. Awarded start-up funds at UCLA made it possible to acquire new instrumentation including a patch clamp set up, a digital image analysis system and a HPLC system.

Availability of these systems has made it possible to refine available methodology for the study of mechanisms underlying the stomatal responses to CO_2 .

Mechanical isolation of epidermal peels. We have developed a mechanical method to isolate large numbers of epidermal peels by blending leaf segments in a medium containing salts and ficoll (43). Mechanical isolation yields intact, physiologically competent guard cells and allows to process about 100 leaves/hr/person. This method provides sufficient epidermal peels from Vicia faba to isolate sufficient numbers of guard cell chloroplasts for biochemical or fluorescence studies (ref. 6, above). In addition, mechanical blending allows to isolate epidermal peels from leaves which cannot be peeled manually, a feature that has made it possible to work with epidermal peels of cotton for the first time (ref. 7, above).

Isolation of guard cell chloroplasts. We have developed a method that yields intact, purified preparations of guard cell chloroplasts. Guard cell protoplasts are isolated by mechanical blending and ruptured by passing by a 10 μm nylon mesh. After washing, the intact guard cell chloroplasts are purified on a Percoll gradient. The preparation is free from contaminant mesophyll chloroplasts but contains some small guard cell vacuoles and a few guard cell nuclei. Guard cell chloroplasts suspended in 330 mM sorbitol, 50 mM Hepes/KOH pH 7.8, 1 mM MgCl_2 , 1 mM MnCl_2 and 2 mM EDTA were used for HPLC and fluorescence analysis.

Measurement of Stomatal Apertures with a Digital Image Analysis System. We have also installed a digital imaging analysis system which permits fast and accurate measurements of physical dimensions of stomata, protoplasts and chloroplasts. The system consists of an Olympus microscope connected with a Javelin CCD digital imaging camera. Image processing is handled with an IBM PC based image analysis board from Metrabyte Corporation and Jandel Scientific's image analysis software. The system allows measurement of stomatal apertures, pore area and stomatal density. It can also be used to measure swelling of isolated protoplasts. The method is faster and more versatile than micrometer methods and permits easy storage and retrieval of image data. Aperture measurements are accurate to within 0.01 μm . Figure 1 shows a digitized image of a stomatal complex of Vicia faba.

Development of an HPLC Method for Quantitation of metabolites in Guard Cells.

High performance liquid chromatography is an analytical technique which permits direct measurement of metabolites in guard cells. Separation of the compound of interest is accomplished in small volume columns which combine traditional chromatographic chemistries with large surface area, high resolution separations. Quantification of the separated metabolites is accomplished with flow-through detectors tailored to the compound of interest. We have developed HPLC methods for the direct quantitation of the three major classes of metabolites of interest in guard cell physiology; organic acids, carbohydrates, and phosphorylated intermediates of glycolysis.

a) Sample preparation. Thus far, we have conducted experiments with sonicated (mesophyll free) epidermal peels and isolated guard cell chloroplasts. Work with epidermal peels allows concurrent measurement of stomatal aperture and metabolite levels. Peels are prepared and pre-incubated in the dark. A timecourse of stomatal opening under different light qualities is shown in fig. 2. After the experimental treatment, the peels are rapidly frozen at -80°C to stop further reaction and to rupture the guard cells. Cell sap is expressed from the peels at 5°C and used directly for HPLC analysis. This procedure allows both rapid preparation and a minimum of sample handling. The cell sap from 4-6 mg dry weight of peels (3-4 leaves) diluted to $30\ \mu\text{l}$ provides enough sample for two repeats each of organic acid and carbohydrate determinations.

b) Organic acid measurement. Anion exchange chromatography can be used for low pressure and HPLC determinations of organic acids. Our analysis employs a $2.6 \times 220\text{mm}$ polypore H chromatographic column from Brownlee Associates. A $10\ \mu\text{l}$ sample is loaded onto the column and separated by isocratic elution with 0.018M sulfuric acid at 75°C . A UV absorbance detector operating at 210nm is used to quantitate organic acids according to calibration curves prepared with known standards. Sensitivity limits range between $0.5\text{--}1.0\ \text{nmoles}/\mu\text{l}$, depending on the specific acid, and total separation time is approximately 16 minutes.

Figure 3 shows a separation of

acid standards.

c) Carbohydrate measurements. We use a SugarPak 6.5x300mm cation exchange column from Waters Chromatography, which provides a good separation of major monosaccharides present as well as important di- and trisaccharides. A 5 μ l sample is loaded onto the column and eluted with 50mM CaEDTA buffer at 90° C. A major technical difficulty in previous carbohydrate separations has been the difficulty in detection. We employ a Dionex PADII electrochemical detector which combines high specificity for carbohydrate compounds with high sensitivity (0.2-0.4 nmoles/ μ l). Total separation time is approximately 22 minutes. Figure 4 shows a separation of a standard carbohydrate mixture.

d) Phosphorylated intermediates. We are currently developing HPLC analysis of various key phosphorylated carbohydrates that are intermediates in important metabolic pathways in guard cells. This separation uses a Dionex Carbo-Pak 4x250mm anion exchange column. Samples are loaded onto the column and eluted with a 27.5 to 57.5% gradient of 750 mM Na-acetate in 150mM NaOH. For quantification we use the PADII electrochemical detector, taking advantage

of the carbohydrate character of many of these compounds. Detection limits are from 0.4 to 5 nmoles/ μ l, depending upon the size of the molecule and number of phosphate groups. Total separation time is 40 minutes. Figure 5 shows a separation of a standard phospho-sugar mix. Results from HPLC analysis of guard cells from epidermal peels and isolated, intact guard cell chloroplasts are described below.

II. Conceptual Developments. Radiolabel experiments with guard cell protoplasts irradiated with red light in the presence of $^{14}\text{CO}_2$ have resolved labeled 3-PGA within 5 s of exposure to the radiolabel (39). Patch clamp experiments with guard cell protoplasts

have shown red light-stimulated, DCMU-sensitive outward currents at the guard cell plasma membrane, resulting from the activity of a proton pumping ATPase modulated by photosynthetic processes in the guard cell chloroplast (4). These data, obtained in our laboratory with DOE support, indicate that different osmoregulatory pathways might modulate stomatal movements in response to different stimuli. Current DOE-supported research in our laboratory is primarily devoted to the characterization of the role of carbon metabolism in guard cells on the supply of osmotica for stomatal movements, and on regulatory mechanisms which transduce signal perception into osmotic changes in guard cells.

Light quality modulates the activity of guard cell chloroplasts

Studies of chlorophyll fluorescence transients with single guard cells have shown that transients associated with photosynthetic carbon fixation are expressed in guard cells under green or red light but are inhibited when blue light is used as the actinic light (ref. 1, above). These results indicate that the controversy involving the issue of photosynthetic carbon fixation in guard cells might be partly due to intrinsic regulatory processes of guard cell photosynthesis which are tightly correlated with environmental conditions. In order to increase our understanding of the response, we isolated guard cell chloroplasts from Vicia faba and studied their fluorescence transients in the absence of interactions with the cytosol. Obtained results show that the transients have different characteristics when stimulated with blue and red light (ref. 6, above), indicating that the guard cell chloroplasts have an intrinsic response to blue light, presumably mediated by carotenoids. These results suggest that carbon metabolism in the guard cell chloroplast might be regulated by light quality, and these responses might have a bearing on sensory transduction of signals modulating stomatal movements.

Long-term survival of guard cell protoplasts kept in cell culture

Guard cell protoplasts kept in cell culture in a minimal medium under red light irradiation survive for several weeks (ref. 3, above), indicating that photosynthetic carbon metabolism can render the guard cells autotrophic. Characterization of a cell culture system with guard cell protoplasts should be valuable for metabolic studies of guard cell properties.

Time Course of Accumulation of Organic Acids and Carbohydrates in Guard Cells

We used HPLC analysis to probe the accumulation of sugars and organic acids in guard cells opening under red and blue light. The analysis allows us to directly determine the type of osmoticum accumulated in each light regime and provides means to identify the metabolic pathway acting as the source of osmoticum (ref. 5, above).

Sonicated epidermal peels of *Vicia faba* were incubated in 1mM KCl and 0.1mM CaCl_2 . After a 2 hr dark adjustment period, the peels were placed in fresh medium, divided into three portions and placed into a temperature controlled bath at 25°C. One set of peels was kept in darkness, one set was exposed to $10 \mu \text{mol m}^{-2} \text{s}^{-1}$ blue light, and the third set was exposed to $125 \mu \text{mol m}^{-2} \text{s}^{-1}$ red light. At appropriate times, samples were

removed for assay. Figure 6 shows an example of carbohydrate and organic acid accumulation during opening stimulated by blue light. The major feature is a transient accumulation of malate accompanied by a steady accumulation of sucrose. Under red light (Fig. 7), there is no malate accumulation and opening is accompanied mainly by an increase in sucrose.

Certain compounds can be used as indicators of the operation of particular metabolic pathways. Table I tracks the changes of some of these diagnostic compounds under red and blue light-stimulated opening. Malate is used as a counter ion for potassium. In our experimental conditions, malate accumulation occurs only during the first 30 minutes of blue light opening and not at all during red light opening. Sucrose accumulation could result from the mobilization of starch reserves or the fixation of carbon through the Calvin cycle. Maltose and maltotriose are two and three glucose oligosaccharides which are products of starch hydrolysis and occur in no other biochemical pathway. Table I shows that both maltose and maltotriose levels increase during blue light opening but not during red light opening. These results indicate that opening under low fluence rates of blue light depended on osmotica (malate and sucrose) from starch hydrolysis. In contrast, opening under red light depends on sugar production by the PCR.

Results with Isolated Chloroplasts

The HPLC technique can also be used to assay the contents of isolated *Vicia faba* guard cell chloroplasts. Initial

Figure 6 Changes in Osmotica During Blue Light-Induced Opening

Figure 7 Changes in Osmotica During Red Light-Induced Opening

TABLE I Levels of Certain Diagnostic Compounds in Guard Cells During Opening in Differing Light Qualities (% of initial level)

Compound	30 min			60 min		
	Dark	Red	Blue	Dark	Red	Blue
Malate	97	92	189	75	89	151
Maltose	22	16	159	14	34	569
Maltotriose	10	10	636	169	42	233

results of an experiment in which isolated guard cell chloroplasts were exposed to 30 min of red light or 30 minutes of red together with 5 minutes of blue light given during the last 5 minutes of the red light treatment are shown in table II. Both the internal contents of the chloroplasts and the compounds exported into the medium were assayed. In red light, both internal and external glucose levels rise, while maltose levels drop. Added blue light further increases glucose levels and causes the appearance of maltose, diagnostic of starch breakdown. Interestingly, blue light also promotes the appearance of malate in the guard cell chloroplast. These results suggest that the chloroplast has an intrinsic blue light response and also suggests that the chloroplast might be a source of organic acid during stomatal opening.

TABLE II Levels of Certain Diagnostic Compounds in Guard Cell Chloroplasts after Differing Light Treatments (% of level in dark control)

Compound	30 min Red		30 min Red + Blue	
	Internal	Medium	Internal	Medium
Malate	50	117	443	102
Maltose	54	118	110	156
Glucose	132	452	384	686

Genetic control of stomatal movements in Pima cotton

Studies with Pima cotton (*Gossypium barbadense*) have characterized a series of Pima lines bred for higher yield potential and heat tolerance, which show increasingly higher stomatal conductances as a function of the extent of selection (ref. 2, above). Crosses between old, low conductance lines with advanced, high conductance ones yield lines with intermediate conductance values, indicating that these differences are genetically determined.

Cotton epidermal peels cannot be obtained manually but use of the mechanical blending method yields large amounts of epidermal peels with functional guard cells. Availability of epidermal peels made it possible to compare guard cell properties in old and advanced lines, and results from these studies have shown that respiratory rates and rates of proton pumping are higher in guard cells from the advanced lines (ref. 7, above). Thus, prevailing genetic differences between lines are also expressed at the guard cell level.

Analysis of the different stomatal responses in the Pima system has shown that the response to temperature has been primarily affected by the genetic modifications ensuing from the breeding process. Reported studies of ABA metabolism in relationship to stomatal conductances indicate that the higher conductances seen in cotton leaves at high temperatures could ensue from higher rates of ABA catabolism (44). In addition, the stomatal response to CO₂ in cotton is sensitized by ABA (45). Availability of the Pima system could thus provide a genetic tool to analyze prevailing relations between the stomatal responses to ABA, CO₂ and temperature. Proposed experiments to investigate these properties are discussed below.

**DATE
FILMED**

8/25/92